

Crystal structure of (1*S*,2*R*,6*R*,7*R*,8*S*,12*S*)-4,10,17-triphenyl-15-thia-4,10-diazapentacyclo-[5.5.5.0^{1,16}.0^{2,6}.0^{8,12}]heptadeca-13,16-diene-3,5,9,11-tetrone *p*-xylene hemisolvate

Wayland E. Nolan,* Neil J. Kroll, Matthew P. Huisenga, Ruixian A. Yue, Simon B. Lang, Nathan D. Klein and Kenneth J. Tritch

Received 12 November 2014
Accepted 16 November 2014

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

Department of Chemistry, University of Minnesota, Minneapolis, MN 55455-0431, USA. *Correspondence e-mail: nolan001@umn.edu

Keywords: crystal structure; cycloaddition; autoxidation; double addition; thiophene


CCDC reference: 1034481

Supporting information: this article has supporting information at journals.iucr.org/e

The title tetrone compound, C₃₂H₂₂N₂O₄S·0.5C₈H₁₀, is the major product (50% yield) of an attempted Diels–Alder reaction of 2-(α -styryl)thiophene with *N*-phenylmaleimide (2 equivalents) in toluene. Recrystallization of the resulting powder from *p*-xylene gave the title hemisolvate; the *p*-xylene molecule is located about an inversion center. In the crystal, the primary tetrone contacts are between a carbonyl O atom and the four flagpole H atoms of the bicyclo[2.2.2]octene core, forming chains along [001].


1. Chemical context


The title compound, (3), is the first reported double-Diels–Alder adduct obtained from a one-pot reaction of a 2-vinylthiophene (Fig. 1). This methodology may have use in the synthesis of novel ligands, zeolites, or polyamides.



Diels–Alder methodology: Reactions between vinyl-heterocycles and dienophiles have been useful in natural product synthesis and in the development of potential medicinal compounds (Booth *et al.*, 2005; Kanai *et al.*, 2005). Reported heterocycles include indole, pyrrole (Le Strat *et al.*, 2005; Nolan *et al.*, 2013), furan (Brewer *et al.*, 1971; Brewer & Elix, 1975*b*; Davidson & Elix, 1970), benzofuran, and benzothiophene (Marrocchi *et al.*, 2001; Pihera *et al.*, 1999). A Diels–Alder reaction was attempted between 2-(α -styryl)thiophene (1) (Tasch *et al.*, 2013) and *N*-phenylmaleimide (2) in an effort to expand this methodology (Fig. 2). Based on work by Watson (2012), the expected products were adduct (4), aromatized adduct (5), or (6) *via* ene addition of (2) to (4). Given the scope of simpler products from these reactions, it was surprising to obtain tetrone (3) in such a high yield.

Mechanism: Mechanisms proposed for double adducts (7) (Lovely *et al.*, 2007) and (8) (Nolan *et al.*, 1993) suggest a





Figure 1


The molecular structure of compound (3), with atom labelling (non-labelled atoms in the *p*-xylene solvent molecule are related to the labelled atoms by inversion symmetry). Displacement ellipsoids are drawn at the 50% probability level.

Diels–Alder reaction (Fig. 3), with loss of H₂ by an unknown pathway, and then a second cycloaddition. Noland *et al.* (1993) observed that formation of (8) was accelerated by exposure to oxygen, and aromatization to (9) was favored over (8) in acid. Brewer & Elix (1975*a*) reported a double adduct (10) and a hydroperoxy intermediate thereof; they proposed loss of H₂ in an autoxidation followed by elimination of H₂O₂, a pathway that fits both observations made by the Noland group. The crystal structures of (3) and the hydroperoxide (11) (Noland *et al.*, 2014), and preliminary HRMS and ¹H NMR evidence that (12) is an intermediate to (3), all support the mechanism proposed by Brewer & Elix (1975*a*).

Applications: Compounds related to (3) are used as bridging ligands in organometallic complexes (see: §4. *Database survey*), synthesis of zeolites (Cantín *et al.*, 2006; Inagaki *et al.*, 2013), and polyamides (Faghihi & Shabaniyan, 2010). Most examples are derived from dianhydride (13) (Hu, 2008) or a similar substrate, reacting with ammonia or primary amines, limiting variability to imido substitution. Domino methodology has been developed that could give more diverse functionality (Strübing *et al.*, 2005).


Figure 2


Synthesis of the title compound (3). Structures (4)–(6) were the expected products.


Figure 3

Contextual compounds. Double adducts (7) and (8) were previously reported. In acid, aromatized adduct (9) was favored over double addition. Double adduct (10) is the closest reported kin of (3). Recently reported (11) supports the proposed mechanism. Hydroperoxide (12) is a likely intermediate to (3). Dianhydride (13) is commonly used for ligand synthesis.

2. Structural commentary

In compound (3) (Fig. 1), the *N*-phenyl rings (C24–C29) and (C30–C35) are twisted out of the plane of their respective succinimido rings, (N4/C3/C2/C6/C5) and (N10/C9/C8/C12/C11), by 54.83 (8) and 54.97 (8)°, respectively, with the same chirality, giving helical character along the major axis (C27 to C33). Figs. 4 and 5 show a left-handed molecule. The bicyclo[2.2.2]octene rings have a typical boat shape. The other rings are nearly planar; the r.m.s. deviations from their respective mean planes are 0.026 and 0.030 Å for the succinimido rings (N4/C3/C2/C6/C5) and (N10/C9/C8/C12/C11),


Figure 4

A molecule of (3) viewed near [114], normal to the pyrrolo[3,4-*g*]-isoindole ring system. The styrylthiophene unit (C21, C18, C17, C16, S15, C14) is forward. The *N*-phenyl rings are twisted so C26 and C32 are forward, C29 and C35 are behind.

Table 2
Experimental details.

Crystal data	
Chemical formula	C ₃₂ H ₂₂ N ₂ O ₄ S·0.5C ₈ H ₁₀
<i>M_r</i>	583.65
Crystal system, space group	Monoclinic, <i>P</i> 2 ₁ / <i>c</i>
Temperature (K)	123
<i>a</i> , <i>b</i> , <i>c</i> (Å)	10.5944 (14), 26.529 (4), 10.4286 (14)
β (°)	99.675 (2)
<i>V</i> (Å ³)	2889.4 (7)
<i>Z</i>	4
Radiation type	Mo <i>K</i> α
μ (mm ⁻¹)	0.16
Crystal size (mm)	0.45 × 0.22 × 0.22
Data collection	
Diffraction	Bruker APEXII CCD
Absorption correction	Multi-scan (<i>SADABS</i> ; Sheldrick, 1996)
<i>T_{min}</i> , <i>T_{max}</i>	0.685, 0.746
No. of measured, independent and observed [<i>I</i> > 2 σ (<i>I</i>)] reflections	33190, 6576, 5803
<i>R_{int}</i>	0.025
(<i>sin</i> θ / λ) _{max} (Å ⁻¹)	0.648
Refinement	
<i>R</i> [<i>F</i> ² > 2 σ (<i>F</i> ²)], <i>wR</i> (<i>F</i> ²), <i>S</i>	0.040, 0.109, 1.00
No. of reflections	6576
No. of parameters	388
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{max}$, $\Delta\rho_{min}$ (e Å ⁻³)	0.38, -0.36

Computer programs: *APEX2* and *SAINT* (Bruker, 2007), *SHELXS97*, *SHELXL2014* and *SHELXTL2008* (Sheldrick, 2008), *Mercury* (Macrae *et al.*, 2008), *enCIFer* (Allen *et al.*, 2004) and *pubCIF* (Westrip, 2010).

interpenetrating nets (Song *et al.*, 2012); and a coumarin-derived double-Diels–Alder adduct (16) (Nicolaidis *et al.*, 1997).

5. Synthesis and crystallization

2-(α -Styryl)thiophene (200 mg, Tasch *et al.*, 2013) and *N*-phenylmaleimide (372 mg, 2 equiv.) were partially dissolved in toluene (5 mL). The resulting mixture was refluxed open to air for 100 h. Upon cooling to room temperature, the resulting suspension was separated by column chromatography (SiO₂, hexane:ethyl acetate, gradient from 1:0 to 1:1). The desired fraction (*R_f* = 0.09 in 1:1) was concentrated at reduced pressure giving compound (3) as a white powder (287 mg, 50%, m.p. 554–555 K). ¹H NMR (500 MHz, CD₂Cl₂) δ 7.498 (*dd*, *J* = 8.0, 1.5 Hz, 2H, H19, H23), 7.388 (*tt*, *J* = 7.0, 2.5 Hz, 4H, H26, H28, H32, H34), 7.374 (*td*, *J* = 5.0, 1.5 Hz, 2H, H20, H22), 7.351 (*tt*, *J* = 7.0, 1.5 Hz, 2H, H27, H33), 7.263 (*tt*, *J* = 4.5, 1.5 Hz, 1H, H21), 6.987 (*dd*, *J* = 7.0, 1.5 Hz, 4H, H25, H29, H31, H35), 6.600 (*d*, *J* = 6.0 Hz, 1H, H14), 6.446 (*d*, *J* = 6.5 Hz, 1H, H13), 4.607 (*t*, *J* = 3.3 Hz, 1H, H7), 3.435 (*d*, *J* = 8.5 Hz, 2H, H2, H12), 3.379 (*dd*, *J* = 8.3, 3.3 Hz, 2H, H6, H8); ¹³C NMR (126 MHz, CD₂Cl₂) δ 175.04 (C5, C9), 172.73 (C3, C11), 136.98 (C18), 135.21 (C16), 132.06 (C24, C30), 129.66 (C26, C28, C32, C34), 129.42 (C27, C33), 129.34 (C20, C22), 128.42 (C21), 127.71 (C17), 126.92 (C25, C29, C31, C35), 126.71 (C14), 126.63 (C19, C23), 126.16 (C13), 62.32 (C1), 47.35 (C2, C12), 41.72 (C6, C8), 40.47 (C7); IR (KBr, cm⁻¹) 3065 (C–H), 2926 (C–H), 2853

(C–H), 1717 (C=O), 1497 (C=C), 1379 (C=C), 1188 (C–N), 743, 727; MS (ESI, PEG, *m/z*) [*M*+H]⁺ calculated for C₃₂H₂₂N₂O₄S 531.1373, found 531.1383.

Recrystallization from many solvent combinations was attempted. The first good crystals were obtained from toluene:1,2-dichloroethane (DCE) [ratio 19:1]. These were empirically (3)·0.5C₇H₈·0.5DCE, with toluene on inversion centers and DCE on twofold axes; both solvents were disordered. Recrystallization from *p*-xylene gave orderly crystals of (3) by suction filtration after 5 days of slow evaporation at room temperature. No conditions were found that gave neat crystals of (3).

6. Refinement

Crystal data, data collection, and structure refinement details are summarized in Table 2. C-bound H atoms were placed in calculated positions and refined as riding atoms, with C–H = 0.095–0.098 Å and with *U*_{iso}(H) = 1.5*U*_{eq}(C) for methyl H atoms and = 1.2*U*_{eq}(C) for other H atoms.

Acknowledgements

The authors thank Victor G. Young Jr (X-Ray Crystallographic Laboratory, University of Minnesota) for assistance with the crystal structure and helpful consultation, and the Wayland E. Noland Research Fellowship Fund at the University of Minnesota Foundation for generous financial support of this project.

References

- Allen, F. H., Johnson, O., Shields, G. P., Smith, B. R. & Towler, M. (2004). *J. Appl. Cryst.* **37**, 335–338.
- Booth, R. J., Lee, H. H., Kraker, A., Ortwine, D. F., Palmer, B. D., Sheehan, D. J. & Toogood, P. L. (2005). US Patent 20050250836.
- Brewer, J. D., Davidson, W. J., Elix, J. A. & Leppik, R. A. (1971). *Aust. J. Chem.* **24**, 1883–1898.
- Brewer, J. D. & Elix, J. A. (1975a). *Aust. J. Chem.* **28**, 1059–1081.
- Brewer, J. D. & Elix, J. A. (1975b). *Aust. J. Chem.* **28**, 1083–1096.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cantín, A., Corma, A., Diaz-Cabanas, M. J., Jordá, J. & Moliner, M. (2006). *J. Am. Chem. Soc.* **128**, 4216–4217.
- Davidson, W. J. & Elix, J. A. (1970). *Aust. J. Chem.* **23**, 2119–2131.
- Faghihi, K. & Shabaniyan, M. (2010). *J. Chil. Chem. Soc.* **55**, 491–496.
- Groom, C. R. & Allen, F. H. (2014). *Angew. Chem. Int. Ed.* **53**, 662–671.
- Hu, T. (2008). *Acta Cryst.* **E64**, o1021.
- Inagaki, S., Tsuboi, Y., Nishita, Y., Syahyiah, T., Wakihara, T. & Kubota, Y. (2013). *Chem. Eur. J.* **19**, 7780–7786.
- Kanai, F., Murakata, C., Tsujita, T., Yamashita, Y., Mizukami, T. & Akinaga, S. (2005). US Patent 20050070591 A1.
- Le Strat, F., Vallette, H., Toupet, L. & Maddaluno, J. (2005). *Eur. J. Org. Chem.* pp. 5296–5305.
- Liu, Z.-M., Liu, Y., Zheng, S.-R., Yu, Z.-Q., Pan, M. & Su, C.-Y. (2007). *Inorg. Chem.* **46**, 5814–5816.
- Lovely, C. J., Du, H., Sivappa, R., Bhandari, M. R., He, Y. & Dias, H. V. R. (2007). *J. Org. Chem.* **72**, 3741–3749.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.

- Marrocchi, A., Minuti, L., Taticchi, A. & Scheeren, H. W. (2001). *Tetrahedron*, **57**, 4959–4965.
- Nicolaides, D. N., Bezergiannidou-Balouctsi, C., Awad, R. W., Litinas, K. E., Malamidou-Xenikaki, E., Terzis, A. & Raptopoulou, C. P. (1997). *J. Org. Chem.* **62**, 499–502.
- Noland, W. E., Gullickson, G. C., Dickson, R. R., Groess, L. L. & Tritch, K. J. (2014). *Acta Cryst.* **E70**, 192–195.
- Noland, W. E., Konkel, M. J., Tempesta, M. S., Cink, R. D., Powers, D. M., Schlemper, E. O. & Barnes, C. L. (1993). *J. Heterocycl. Chem.* **30**, 183–192.
- Noland, W. E., Lanzatella, N. P., Dickson, R. R., Messner, M. E. & Nguyen, H. H. (2013). *J. Heterocycl. Chem.* **50**, 795–808.
- Pihera, P., Dvořáková, H. & Svoboda, J. (1999). *Collect. Czech. Chem. Commun.* **64**, 389–407.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Song, X.-Z., Qin, C., Guan, W., Song, S.-Y. & Zhang, H.-J. (2012). *New J. Chem.* **36**, 877–882.
- Strübing, D., von Wangelin, A. J., Neumann, H., Gördes, D., Hübner, S., Klaus, S., Spannenberg, A. & Beller, M. (2005). *Eur. J. Org. Chem.* pp. 107–113.
- Tasch, B. O. A., Bensch, L., Antovic, D. & Müller, T. J. J. (2013). *Org. Biomol. Chem.* **11**, 6113–6118.
- Watson, L. J. (2012). *Pericyclic Reactions of Vinyl-Heteroaromatics: Multi-Component Domino and Sequential Processes*. Dissertation, Newcastle University, England.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.
- Yu, Z.-Q., Pan, M., Jiang, J.-J., Liu, Z.-M. & Su, C.-Y. (2012). *Cryst. Growth Des.* **12**, 2389–2396.
- Zhang, Y. (2012). *Acta Cryst.* **E68**, m1275.
- Zhang, Z.-J., Shi, W., Niu, Z., Li, H.-H., Zhao, B., Cheng, P., Liao, D.-Z. & Yan, S.-P. (2011). *Chem. Commun.* **47**, 6425–6427.

supporting information

Acta Cryst. (2014). E70, 550-554 [doi:10.1107/S1600536814025094]

Crystal structure of (1*S*,2*R*,6*R*,7*R*,8*S*,12*S*)-4,10,17-triphenyl-15-thia-4,10-diazapentacyclo[5.5.5.0^{1,16}.0^{2,6}.0^{8,12}]heptadeca-13,16-diene-3,5,9,11-tetrone *p*-xylene hemisolvate

Wayland E. Noland, Neil J. Kroll, Matthew P. Huisenga, Ruixian A. Yue, Simon B. Lang, Nathan D. Klein and Kenneth J. Tritch

Computing details

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT* (Bruker, 2007); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL2014* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL2008* (Sheldrick, 2008), *enCIFer* (Allen *et al.*, 2004) and *publCIF* (Westrip, 2010).

(1*S*,2*R*,6*R*,7*R*,8*S*,12*S*)-4,10,17-Triphenyl-15-thia-4,10-diazapentacyclo[5.5.5.0^{1,16}.0^{2,6}.0^{8,12}]heptadeca-13,16-diene-3,5,9,11-tetrone *p*-xylene hemisolvate

Crystal data

C₃₂H₂₂N₂O₄S·0.5C₈H₁₀

M_r = 583.65

Monoclinic, *P*2₁/*c*

a = 10.5944 (14) Å

b = 26.529 (4) Å

c = 10.4286 (14) Å

β = 99.675 (2)°

V = 2889.4 (7) Å³

Z = 4

F(000) = 1220

D_x = 1.342 Mg m⁻³

Melting point: 554 K

Mo *K*α radiation, λ = 0.71073 Å

Cell parameters from 2771 reflections

θ = 3.1–27.4°

μ = 0.16 mm⁻¹

T = 123 K

Plate, colourless

0.45 × 0.22 × 0.22 mm

Data collection

Bruker APEXII CCD
diffractometer

Radiation source: sealed tube

φ and ω scans

Absorption correction: multi-scan
(*SADABS*; Sheldrick, 1996)

T_{min} = 0.685, *T_{max}* = 0.746

33190 measured reflections

6576 independent reflections

5803 reflections with *I* > 2σ(*I*)

R_{int} = 0.025

θ_{\max} = 27.4°, θ_{\min} = 2.0°

h = -13→13

k = -34→34

l = -13→13

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.040

wR(*F*²) = 0.109

S = 1.00

6576 reflections

388 parameters

0 restraints

Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0547P)^2 + 1.9057P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.38 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.36 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.47103 (13)	0.62662 (5)	0.07444 (13)	0.0168 (3)
C2	0.56246 (13)	0.67315 (5)	0.10206 (13)	0.0171 (3)
H2	0.6037	0.6805	0.0246	0.021*
O3	0.74841 (10)	0.63409 (4)	0.23636 (11)	0.0275 (2)
C3	0.66409 (13)	0.66474 (5)	0.22147 (13)	0.0188 (3)
N4	0.64349 (11)	0.70008 (4)	0.31639 (11)	0.0183 (2)
O5	0.51276 (10)	0.76804 (4)	0.33509 (10)	0.0237 (2)
C5	0.54585 (13)	0.73378 (5)	0.27202 (13)	0.0174 (3)
C6	0.48591 (13)	0.71923 (5)	0.13506 (13)	0.0168 (3)
H6	0.4931	0.7475	0.0733	0.020*
C7	0.34372 (13)	0.70480 (5)	0.13224 (13)	0.0171 (3)
H7	0.2938	0.7341	0.1573	0.020*
C8	0.29297 (13)	0.68806 (5)	-0.00915 (13)	0.0186 (3)
H8	0.2980	0.7166	-0.0709	0.022*
O9	0.06508 (10)	0.69309 (4)	0.00144 (11)	0.0270 (2)
C9	0.15643 (13)	0.66936 (5)	-0.02085 (13)	0.0202 (3)
N10	0.15215 (11)	0.61919 (4)	-0.06089 (11)	0.0195 (2)
O11	0.28909 (10)	0.56023 (4)	-0.12507 (11)	0.0262 (2)
C11	0.27107 (13)	0.60134 (5)	-0.08300 (13)	0.0196 (3)
C12	0.36946 (13)	0.64258 (5)	-0.04520 (13)	0.0181 (3)
H12	0.4127	0.6511	-0.1206	0.022*
C13	0.53628 (13)	0.57771 (5)	0.05823 (14)	0.0215 (3)
H13	0.5845	0.5721	-0.0095	0.026*
C14	0.52206 (14)	0.54274 (6)	0.14507 (15)	0.0246 (3)
H14	0.5584	0.5101	0.1437	0.030*
S15	0.43063 (4)	0.56125 (2)	0.26360 (4)	0.02309 (10)
C16	0.40457 (13)	0.62075 (5)	0.19169 (13)	0.0166 (3)
C17	0.33937 (12)	0.66069 (5)	0.22472 (13)	0.0168 (3)
C18	0.27549 (13)	0.66453 (5)	0.34017 (13)	0.0186 (3)
C19	0.33888 (16)	0.64756 (7)	0.46108 (15)	0.0296 (3)
H19	0.4224	0.6336	0.4683	0.035*
C20	0.28055 (19)	0.65094 (8)	0.57056 (17)	0.0400 (4)
H20	0.3243	0.6393	0.6522	0.048*
C21	0.15919 (18)	0.67111 (7)	0.56115 (18)	0.0378 (4)

H21	0.1191	0.6731	0.6359	0.045*
C22	0.09641 (16)	0.68835 (7)	0.44274 (18)	0.0332 (4)
H22	0.0127	0.7021	0.4361	0.040*
C23	0.15467 (15)	0.68579 (6)	0.33309 (16)	0.0262 (3)
H23	0.1115	0.6987	0.2526	0.031*
C24	0.71510 (13)	0.70018 (6)	0.44597 (13)	0.0202 (3)
C25	0.71973 (16)	0.65636 (7)	0.51885 (16)	0.0303 (3)
H25	0.6760	0.6269	0.4835	0.036*
C26	0.78922 (18)	0.65597 (8)	0.64436 (17)	0.0413 (4)
H26	0.7930	0.6262	0.6953	0.050*
C27	0.85297 (16)	0.69899 (8)	0.69534 (16)	0.0401 (5)
H27	0.9007	0.6985	0.7810	0.048*
C28	0.84726 (15)	0.74241 (8)	0.62212 (16)	0.0344 (4)
H28	0.8905	0.7719	0.6579	0.041*
C29	0.77860 (13)	0.74341 (6)	0.49609 (15)	0.0244 (3)
H29	0.7753	0.7732	0.4452	0.029*
C30	0.03938 (13)	0.58848 (5)	-0.06870 (14)	0.0201 (3)
C31	-0.02016 (15)	0.58564 (6)	0.03952 (15)	0.0261 (3)
H31	0.0120	0.6044	0.1156	0.031*
C32	-0.12704 (15)	0.55524 (6)	0.03589 (17)	0.0299 (3)
H32	-0.1686	0.5531	0.1096	0.036*
C33	-0.17302 (15)	0.52803 (6)	-0.07528 (17)	0.0284 (3)
H33	-0.2459	0.5070	-0.0776	0.034*
C34	-0.11321 (15)	0.53137 (6)	-0.18289 (16)	0.0288 (3)
H34	-0.1456	0.5127	-0.2591	0.035*
C35	-0.00598 (15)	0.56179 (6)	-0.18087 (15)	0.0247 (3)
H35	0.0352	0.5642	-0.2549	0.030*
C36	-0.0645 (2)	0.53975 (8)	0.43607 (17)	0.0449 (5)
H36	-0.1088	0.5675	0.3918	0.054*
C37	0.0665 (2)	0.54081 (8)	0.46333 (18)	0.0451 (5)
H37	0.1114	0.5690	0.4375	0.054*
C38	0.1347 (2)	0.50084 (8)	0.52868 (17)	0.0449 (5)
C39	0.2786 (2)	0.50191 (11)	0.5586 (2)	0.0633 (7)
H39A	0.3098	0.5334	0.5262	0.095*
H39B	0.3067	0.4997	0.6529	0.095*
H39C	0.3131	0.4733	0.5163	0.095*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S15	0.0287 (2)	0.01758 (17)	0.02321 (18)	0.00367 (13)	0.00515 (14)	0.00323 (13)
O3	0.0221 (5)	0.0302 (6)	0.0284 (6)	0.0087 (4)	-0.0012 (4)	-0.0075 (4)
O5	0.0243 (5)	0.0242 (5)	0.0216 (5)	0.0044 (4)	0.0006 (4)	-0.0065 (4)
O9	0.0203 (5)	0.0227 (5)	0.0357 (6)	0.0046 (4)	-0.0021 (4)	-0.0033 (4)
O11	0.0239 (5)	0.0247 (5)	0.0301 (6)	-0.0002 (4)	0.0047 (4)	-0.0097 (4)
N4	0.0175 (5)	0.0207 (6)	0.0162 (5)	0.0013 (4)	0.0013 (4)	-0.0022 (4)
N10	0.0186 (6)	0.0191 (6)	0.0198 (6)	0.0000 (4)	0.0000 (4)	-0.0019 (4)
C1	0.0176 (6)	0.0177 (6)	0.0150 (6)	0.0006 (5)	0.0022 (5)	-0.0018 (5)

C2	0.0174 (6)	0.0183 (6)	0.0159 (6)	0.0003 (5)	0.0034 (5)	-0.0014 (5)
C3	0.0174 (6)	0.0210 (7)	0.0181 (6)	0.0002 (5)	0.0034 (5)	-0.0027 (5)
C5	0.0164 (6)	0.0186 (6)	0.0173 (6)	-0.0013 (5)	0.0026 (5)	-0.0002 (5)
C6	0.0181 (6)	0.0162 (6)	0.0159 (6)	0.0000 (5)	0.0024 (5)	-0.0003 (5)
C7	0.0169 (6)	0.0164 (6)	0.0170 (6)	0.0015 (5)	0.0003 (5)	-0.0004 (5)
C8	0.0197 (7)	0.0177 (6)	0.0169 (6)	0.0007 (5)	-0.0014 (5)	0.0014 (5)
C9	0.0208 (7)	0.0199 (7)	0.0177 (6)	0.0017 (5)	-0.0031 (5)	0.0010 (5)
C11	0.0202 (7)	0.0236 (7)	0.0146 (6)	-0.0002 (5)	0.0014 (5)	-0.0008 (5)
C12	0.0205 (6)	0.0193 (6)	0.0141 (6)	-0.0002 (5)	0.0016 (5)	-0.0008 (5)
C13	0.0193 (7)	0.0233 (7)	0.0214 (7)	0.0032 (5)	0.0025 (5)	-0.0059 (5)
C14	0.0254 (7)	0.0206 (7)	0.0268 (7)	0.0061 (6)	0.0013 (6)	-0.0054 (6)
C16	0.0168 (6)	0.0174 (6)	0.0147 (6)	-0.0006 (5)	0.0004 (5)	0.0012 (5)
C17	0.0151 (6)	0.0187 (6)	0.0161 (6)	-0.0007 (5)	0.0011 (5)	-0.0001 (5)
C18	0.0207 (7)	0.0172 (6)	0.0196 (6)	0.0010 (5)	0.0085 (5)	-0.0019 (5)
C19	0.0259 (8)	0.0418 (9)	0.0214 (7)	0.0042 (7)	0.0052 (6)	-0.0001 (6)
C20	0.0424 (10)	0.0570 (12)	0.0221 (8)	0.0027 (9)	0.0095 (7)	-0.0010 (8)
C21	0.0425 (10)	0.0432 (10)	0.0336 (9)	-0.0048 (8)	0.0231 (8)	-0.0092 (7)
C22	0.0270 (8)	0.0328 (9)	0.0434 (10)	0.0008 (7)	0.0167 (7)	-0.0083 (7)
C23	0.0234 (7)	0.0256 (7)	0.0306 (8)	0.0031 (6)	0.0076 (6)	-0.0033 (6)
C24	0.0147 (6)	0.0295 (7)	0.0160 (6)	0.0036 (5)	0.0016 (5)	-0.0030 (5)
C25	0.0306 (8)	0.0330 (8)	0.0261 (8)	0.0064 (7)	0.0014 (6)	0.0023 (6)
C26	0.0407 (10)	0.0557 (12)	0.0258 (8)	0.0200 (9)	0.0012 (7)	0.0096 (8)
C27	0.0257 (8)	0.0732 (14)	0.0190 (7)	0.0187 (8)	-0.0037 (6)	-0.0078 (8)
C28	0.0160 (7)	0.0587 (11)	0.0277 (8)	0.0028 (7)	0.0009 (6)	-0.0198 (8)
C29	0.0161 (6)	0.0345 (8)	0.0232 (7)	0.0002 (6)	0.0044 (5)	-0.0076 (6)
C30	0.0173 (6)	0.0183 (6)	0.0234 (7)	0.0013 (5)	-0.0006 (5)	-0.0011 (5)
C31	0.0245 (7)	0.0279 (8)	0.0256 (7)	-0.0005 (6)	0.0036 (6)	-0.0059 (6)
C32	0.0253 (8)	0.0324 (8)	0.0339 (8)	-0.0010 (6)	0.0103 (6)	-0.0038 (7)
C33	0.0202 (7)	0.0230 (7)	0.0416 (9)	-0.0018 (6)	0.0037 (6)	-0.0036 (6)
C34	0.0266 (8)	0.0259 (8)	0.0316 (8)	-0.0029 (6)	-0.0017 (6)	-0.0077 (6)
C35	0.0259 (7)	0.0253 (7)	0.0219 (7)	-0.0006 (6)	0.0012 (6)	-0.0033 (6)
C36	0.0680 (14)	0.0399 (10)	0.0260 (8)	0.0267 (10)	0.0057 (8)	0.0051 (7)
C37	0.0680 (14)	0.0381 (10)	0.0294 (9)	0.0164 (9)	0.0082 (9)	0.0007 (8)
C38	0.0593 (12)	0.0504 (11)	0.0240 (8)	0.0221 (10)	0.0037 (8)	-0.0016 (8)
C39	0.0584 (14)	0.0869 (19)	0.0425 (12)	0.0185 (13)	0.0027 (10)	0.0007 (12)

Geometric parameters (Å, °)

S15—C14	1.7635 (16)	C20—C21	1.381 (3)
S15—C16	1.7495 (14)	C20—H20	0.9500
O3—C3	1.1986 (17)	C21—C22	1.379 (3)
O5—C5	1.2081 (17)	C21—H21	0.9500
O9—C9	1.2094 (18)	C22—C23	1.389 (2)
O11—C11	1.2028 (17)	C22—H22	0.9500
N4—C3	1.4069 (17)	C23—H23	0.9500
N4—C5	1.3865 (17)	C24—C25	1.385 (2)
N4—C24	1.4342 (17)	C24—C29	1.387 (2)
N10—C9	1.3934 (18)	C25—C26	1.390 (2)

N10—C11	1.4008 (18)	C25—H25	0.9500
N10—C30	1.4370 (18)	C26—C27	1.386 (3)
C1—C2	1.5655 (18)	C26—H26	0.9500
C1—C12	1.5629 (18)	C27—C28	1.378 (3)
C1—C13	1.4930 (19)	C27—H27	0.9500
C1—C16	1.5181 (18)	C28—C29	1.391 (2)
C2—C3	1.5196 (18)	C28—H28	0.9500
C2—C6	1.5377 (18)	C29—H29	0.9500
C2—H2	1.0000	C30—C31	1.384 (2)
C5—C6	1.5124 (18)	C30—C35	1.383 (2)
C6—C7	1.5498 (18)	C31—C32	1.386 (2)
C6—H6	1.0000	C31—H31	0.9500
C7—C8	1.5472 (18)	C32—C33	1.383 (2)
C7—C17	1.5220 (18)	C32—H32	0.9500
C7—H7	1.0000	C33—C34	1.381 (2)
C8—C9	1.5147 (19)	C33—H33	0.9500
C8—C12	1.5351 (19)	C34—C35	1.391 (2)
C8—H8	1.0000	C34—H34	0.9500
C11—C12	1.5167 (19)	C35—H35	0.9500
C12—H12	1.0000	C36—C37	1.370 (3)
C13—C14	1.323 (2)	C36—C38 ⁱ	1.393 (3)
C13—H13	0.9500	C36—H36	0.9500
C14—H14	0.9500	C37—C38	1.395 (3)
C16—C17	1.3410 (19)	C37—H37	0.9500
C17—C18	1.4805 (18)	C38—C39	1.504 (3)
C18—C19	1.400 (2)	C39—H39A	0.9800
C18—C23	1.389 (2)	C39—H39B	0.9800
C19—C20	1.390 (2)	C39—H39C	0.9800
C19—H19	0.9500		
C14—S15—C16	90.95 (7)	C17—C18—C23	121.85 (13)
C3—N4—C5	112.96 (11)	C19—C18—C23	118.48 (13)
C3—N4—C24	122.96 (11)	C18—C19—C20	120.47 (15)
C5—N4—C24	124.07 (11)	C18—C19—H19	119.8
C9—N10—C11	112.78 (12)	C20—C19—H19	119.8
C9—N10—C30	122.91 (12)	C19—C20—C21	120.27 (17)
C11—N10—C30	124.15 (12)	C19—C20—H20	119.9
C2—C1—C12	104.78 (10)	C21—C20—H20	119.9
C2—C1—C13	114.97 (11)	C20—C21—C22	119.70 (15)
C2—C1—C16	106.80 (10)	C20—C21—H21	120.1
C12—C1—C13	114.50 (11)	C22—C21—H21	120.1
C12—C1—C16	108.69 (11)	C21—C22—C23	120.46 (16)
C13—C1—C16	106.76 (11)	C21—C22—H22	119.8
C1—C2—C3	111.49 (11)	C23—C22—H22	119.8
C1—C2—C6	109.56 (11)	C18—C23—C22	120.58 (15)
C3—C2—C6	105.13 (10)	C18—C23—H23	119.7
C1—C2—H2	110.2	C22—C23—H23	119.7
C3—C2—H2	110.2	N4—C24—C25	118.68 (13)

C6—C2—H2	110.2	N4—C24—C29	120.19 (13)
O3—C3—N4	124.15 (13)	C25—C24—C29	121.13 (14)
O3—C3—C2	128.01 (13)	C24—C25—C26	119.17 (17)
N4—C3—C2	107.84 (11)	C24—C25—H25	120.4
O5—C5—N4	124.88 (12)	C26—C25—H25	120.4
O5—C5—C6	126.30 (12)	C25—C26—C27	120.14 (18)
N4—C5—C6	108.79 (11)	C25—C26—H26	119.9
C2—C6—C5	105.11 (11)	C27—C26—H26	119.9
C2—C6—C7	110.30 (11)	C26—C27—C28	120.18 (15)
C5—C6—C7	109.48 (11)	C26—C27—H27	119.9
C2—C6—H6	110.6	C28—C27—H27	119.9
C5—C6—H6	110.6	C27—C28—C29	120.45 (17)
C7—C6—H6	110.6	C27—C28—H28	119.8
C6—C7—C8	105.66 (11)	C29—C28—H28	119.8
C6—C7—C17	108.10 (10)	C24—C29—C28	118.93 (16)
C8—C7—C17	109.92 (11)	C24—C29—H29	120.5
C6—C7—H7	111.0	C28—C29—H29	120.5
C8—C7—H7	111.0	N10—C30—C31	118.24 (12)
C17—C7—H7	111.0	N10—C30—C35	120.38 (13)
C7—C8—C9	110.18 (11)	C35—C30—C31	121.37 (14)
C7—C8—C12	110.23 (11)	C30—C31—C32	119.45 (14)
C9—C8—C12	105.01 (11)	C30—C31—H31	120.3
C7—C8—H8	110.4	C32—C31—H31	120.3
C9—C8—H8	110.4	C31—C32—C32	119.85 (15)
C12—C8—H8	110.4	C31—C32—H32	120.1
O9—C9—N10	124.61 (13)	C33—C32—H32	120.1
O9—C9—C8	126.75 (13)	C32—C33—C34	120.17 (14)
N10—C9—C8	108.64 (12)	C32—C33—H33	119.9
O11—C11—N10	124.66 (13)	C34—C33—H33	119.9
O11—C11—C12	127.20 (13)	C33—C34—C35	120.67 (14)
N10—C11—C12	108.14 (11)	C33—C34—H34	119.7
C1—C12—C8	109.78 (10)	C35—C34—H34	119.7
C1—C12—C11	111.29 (11)	C30—C35—C34	118.48 (14)
C8—C12—C11	105.22 (11)	C30—C35—H35	120.8
C1—C12—H12	110.1	C34—C35—H35	120.8
C8—C12—H12	110.1	C37—C36—C38 ⁱ	121.92 (18)
C11—C12—H12	110.1	C37—C36—H36	119.0
C1—C13—C14	115.08 (13)	C38 ⁱ —C36—H36	119.0
C1—C13—H13	122.5	C36—C37—C38	120.6 (2)
C14—C13—H13	122.5	C36—C37—H37	119.7
S15—C14—C13	115.07 (11)	C38—C37—H37	119.7
S15—C14—H14	122.5	C36 ⁱ —C38—C37	117.4 (2)
C13—C14—H14	122.5	C36 ⁱ —C38—C39	121.83 (19)
S15—C16—C1	112.12 (9)	C37—C38—C39	120.7 (2)
S15—C16—C17	130.82 (11)	C38—C39—H39A	109.5
C1—C16—C17	117.04 (12)	C38—C39—H39B	109.5
C7—C17—C16	111.89 (12)	C38—C39—H39C	109.5
C7—C17—C18	121.95 (12)	H39A—C39—H39B	109.5

C16—C17—C18	126.07 (12)	H39A—C39—H39C	109.5
C17—C18—C19	119.66 (13)	H39B—C39—H39C	109.5
S15—C14—C13—C1	-0.77 (17)	C5—C6—C7—C17	-59.61 (13)
S15—C16—C1—C2	-122.42 (10)	C6—C2—C1—C12	61.10 (13)
S15—C16—C1—C12	125.04 (10)	C6—C2—C1—C13	-172.32 (11)
S15—C16—C1—C13	1.04 (13)	C6—C2—C1—C16	-54.10 (13)
S15—C16—C17—C7	179.78 (10)	C6—C5—N4—C24	-175.22 (12)
S15—C16—C17—C18	3.1 (2)	C6—C7—C8—C9	175.86 (11)
O3—C3—N4—C5	175.46 (14)	C6—C7—C8—C12	60.43 (14)
O3—C3—N4—C24	-5.7 (2)	C6—C7—C17—C16	-58.81 (14)
O3—C3—C2—C1	64.82 (19)	C6—C7—C17—C18	118.02 (13)
O3—C3—C2—C6	-176.55 (15)	C7—C8—C12—C11	121.82 (12)
O5—C5—N4—C3	-178.39 (13)	C7—C17—C18—C19	-132.20 (14)
O5—C5—N4—C24	2.8 (2)	C7—C17—C18—C23	46.14 (19)
O5—C5—C6—C2	-179.22 (13)	C8—C7—C17—C16	56.06 (15)
O5—C5—C6—C7	-60.75 (18)	C8—C7—C17—C18	-127.11 (13)
O9—C9—N10—C11	178.47 (14)	C8—C9—N10—C11	-2.16 (15)
O9—C9—N10—C30	-5.9 (2)	C8—C9—N10—C30	173.45 (12)
O9—C9—C8—C7	59.85 (18)	C8—C12—C1—C13	170.30 (12)
O9—C9—C8—C12	178.53 (14)	C8—C12—C1—C16	51.04 (14)
O11—C11—N10—C9	-175.58 (14)	C9—N10—C11—C12	4.28 (15)
O11—C11—N10—C30	8.9 (2)	C9—N10—C30—C31	-52.23 (19)
O11—C11—C12—C1	-65.82 (18)	C9—N10—C30—C35	129.14 (15)
O11—C11—C12—C8	175.34 (14)	C9—C8—C12—C11	3.17 (13)
N4—C3—C2—C1	-115.29 (12)	C9—C8—C7—C17	59.45 (14)
N4—C3—C2—C6	3.34 (14)	C11—N10—C30—C31	122.88 (15)
N4—C5—C6—C2	-1.26 (14)	C11—N10—C30—C35	-55.75 (19)
N4—C5—C6—C7	117.21 (12)	C11—C12—C1—C13	54.22 (15)
N4—C24—C25—C26	179.77 (14)	C11—C12—C1—C16	-65.04 (14)
N4—C24—C29—C28	179.97 (13)	C12—C1—C13—C14	-120.51 (14)
N10—C9—C8—C7	-119.51 (12)	C12—C1—C16—C17	-56.16 (15)
N10—C9—C8—C12	-0.83 (14)	C12—C11—N10—C30	-171.27 (12)
N10—C11—C12—C1	114.33 (12)	C12—C8—C7—C17	-55.98 (14)
N10—C11—C12—C8	-4.51 (14)	C13—C1—C16—C17	179.84 (12)
N10—C30—C31—C32	-178.26 (14)	C13—C14—S15—C16	1.19 (13)
N10—C30—C35—C34	178.12 (13)	C14—S15—C16—C17	-179.83 (14)
C1—C2—C6—C5	118.66 (11)	C14—C13—C1—C16	-0.18 (17)
C1—C2—C6—C7	0.75 (14)	C16—C17—C18—C19	44.2 (2)
C1—C12—C8—C7	1.97 (15)	C16—C17—C18—C23	-137.49 (15)
C1—C12—C8—C9	-116.67 (12)	C17—C18—C19—C20	179.87 (16)
C1—C16—S15—C14	-1.25 (10)	C17—C18—C23—C22	179.23 (14)
C1—C16—C17—C7	1.26 (17)	C18—C19—C20—C21	0.0 (3)
C1—C16—C17—C18	-175.42 (12)	C18—C23—C22—C21	1.9 (2)
C2—C1—C13—C14	118.06 (14)	C19—C18—C23—C22	-2.4 (2)
C2—C1—C16—C17	56.38 (15)	C19—C20—C21—C22	-0.6 (3)
C2—C1—C12—C8	-62.83 (13)	C20—C19—C18—C23	1.5 (2)
C2—C1—C12—C11	-178.92 (11)	C20—C21—C22—C23	-0.3 (3)

C2—C3—N4—C5	-4.44 (15)	C24—C25—C26—C27	-0.1 (3)
C2—C3—N4—C24	174.41 (12)	C24—C29—C28—C27	0.6 (2)
C2—C6—C7—C8	-62.06 (13)	C25—C24—C29—C28	-0.4 (2)
C2—C6—C7—C17	55.57 (14)	C25—C26—C27—C28	0.3 (3)
C3—N4—C5—C6	3.62 (15)	C26—C25—C24—C29	0.1 (2)
C3—N4—C24—C25	-54.29 (19)	C26—C27—C28—C29	-0.6 (2)
C3—N4—C24—C29	125.40 (15)	C30—C31—C32—C33	0.1 (2)
C3—C2—C1—C12	177.05 (11)	C30—C35—C34—C33	0.1 (2)
C3—C2—C1—C13	-56.37 (15)	C31—C30—C35—C34	-0.5 (2)
C3—C2—C1—C16	61.85 (13)	C31—C32—C33—C34	-0.4 (2)
C3—C2—C6—C5	-1.26 (13)	C32—C31—C30—C35	0.4 (2)
C3—C2—C6—C7	-119.17 (11)	C32—C33—C34—C35	0.3 (2)
C5—N4—C24—C25	124.43 (15)	C36—C37—C38—C36 ⁱ	0.3 (3)
C5—N4—C24—C29	-55.88 (19)	C36—C37—C38—C39	179.96 (19)
C5—C6—C7—C8	-177.24 (11)	C38 ⁱ —C36—C37—C38	-0.3 (3)

Symmetry code: (i) $-x, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D—H...A</i>	<i>D—H</i>	<i>H...A</i>	<i>D...A</i>	<i>D—H...A</i>
C23—H23...O9	0.95	2.59	3.435 (2)	149
C2—H2...O5 ⁱⁱ	1.00	2.46	3.158 (2)	126
C6—H6...O5 ⁱⁱ	1.00	2.56	3.206 (2)	122
C8—H8...O5 ⁱⁱ	1.00	2.66	3.269 (2)	131
C12—H12...O5 ⁱⁱ	1.00	2.47	3.182 (2)	128
C14—H14...O11 ⁱⁱⁱ	0.95	2.50	3.413 (2)	162

Symmetry codes: (ii) $x, -y+3/2, z-1/2$; (iii) $-x+1, -y+1, -z$.