

aimed at finding unusual ways to develop emotional intelligence.

Methods: The study involved 55 participants aged 18 to 25 years (average age-20.38 ± 0.28), 23 of whom were men and 22 were women. During the study we diagnosed the level of development of emotional intelligence (Sergienko, Vetrova, 2009) and spatial abilities (Rimfeld et al., 2017), type of attachment to the loved one (Sabelnikova, Kashirsky, 2015), and the negotiating style of personality (Soldatova, Gasimov, 2019). In the VR CAVE system, a situation was simulated in which the subject had to detect the avatar and determine the emotional-facial expression displayed by it.

Results: It was shown that the level of respondent's emotional intelligence development does not determine the success of identifying the avatar's emotion in VR. The success of identifying emotions depends on the level spatial abilities development. Therefore, it is assumed that in the simulated situation, the avatar is perceived as a special spatial image, and not as a full-fledged partner for interpersonal communication.

Conclusions: Thus, the use of VR systems for training and development of emotional intelligence is not proven.

Disclosure: No significant relationships.

Keywords: development of emotional intelligence; virtual reality; emotional expression

EPP0355

Effects of Mindful Self-Compassion on Psychological Well-Being in Psychiatric Rehabilitation: A Randomized-Controlled Trial

A. Andorfer^{1,2,3*}, M. Hiebler-Ragger^{2,4}, P. Kaufmann³, E. Pollheimer³, L. Gaiswinkler¹, H.-F. Unterrainer^{1,2,5}, H.-P. Kapfhammer¹ and A. Kresse⁶

¹Medical University Graz, Department Of Psychiatry And Psychotherapeutic Medicine, Graz, Austria; ²Grüner Kreis Society, Center For Integrative Addiction Research (ciar), Vienna, Austria; ³pro mente Reha, Sonnenpark Neusiedlersee, Rust, Austria; ⁴pro mente Reha, Apr Graz, Graz, Austria; ⁵University of Vienna, Institute For Religious Studies, Vienna, Austria and ⁶Medical University Graz, Institute For Pathophysiology Und Immunology, Graz, Austria
*Corresponding author.
doi: 10.1192/j.eurpsy.2022.632

Introduction: The evidence for the positive effects of mindfulness-based interventions on psychological well-being and physical health has been convincing in recent years. As a specific form of such an intervention, the Mindful Self-Compassion (MSC) training program was developed to promote self-compassion and mindfulness. An initial study on an adapted version of the MSC training program considered it to be beneficial in psychiatric inpatient rehabilitation. **Objectives:** The present study aims to further evaluate the link between MSC and psychological symptoms as well as quality of life. **Methods:** A randomized controlled trial was conducted from September 2020 to August 2021. A total of 228 patients (64% female, 36% male) participated in a six-week psychiatric rehabilitation program to assess the impact of an adapted MSC training program compared to the control intervention of Progressive Muscle Relaxation training (PMR) on psychological well-being. Both training programs took place once a week for 75 minutes as part of a standardized inpatient rehabilitation program. The participants completed the Self-Compassion Scale (SCS), the Brief

Symptom Inventory (BSI-18), and the Short-Form-Health-Survey-12 (SF-12) pre and post intervention.

Results: At the moment, statistical analyses are being carried out. Detailed results will be presented on the poster.

Conclusions: The results of this study will contribute to rehabilitation research as they provide further insight into the role of MSC in the treatment of mental disorders. In addition, the clinical implications, and possible effects of changes in the rehabilitation program during the COVID-19 pandemic on the protocol and the results of this study will be discussed.

Disclosure: No significant relationships.

Keywords: Mindfulness; self-compassion; Randomized Controlled Trial; psychological well-being

EPP0357

Religious Coping Strategies for Religious Patients with Schizophrenia and Schizoaffective Disorder

E. Gedeveni, G. Kopeiko, O. Borisova, T. Vladimirova* and V. Kaleda

FSBSI Mental Health Research Center, Researching Group Of Specific Forms Of Mental Disorders, Moscow, Russian Federation

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.633

Introduction: Over last decades convincing evident data was accumulated about the positive correlation of religious involvement and better mental health in depression, substance abuse, suicide, stress-related disorders and dementia. The studies of the impact of religion on patients with schizophrenia and schizoaffective disorders are still insufficient and controversial.

Objectives: To investigate the impact of the religious coping strategies in patients with schizophrenia and schizoaffective disorder with focus on resolution and quality of remission in schizophrenia and schizoaffective disorder.

Methods: The pilot 1 year study covers 68 orthodox (group 1) and 55 unbeliever (group 2) outpatients with schizophrenia and schizoaffective disorder in remission on the maintenance therapy in FSBSI MHRC. The groups matched in age (18-60 y.o.), gender, treatment. The orthodox group received religious coping therapy. Number of relapses, remission quality (PANSS), quality of life (QLS), compliance (MARS) were measured 3 times (baseline visit, 6, 12 months). Statistical analysis (regression and correlation) was applied.

Results: Religious coping strategies proposed by Paragment K. (2013) were applied considering the peculiarities of value-semantic structures and selected religious values of the patients as important rehabilitation resource (Kopeyko et al, 2016). Group 1 demonstrated statistically significant better remission at 12-months point – better subjective well-being, social/functional outcomes, higher adherence to medication, less relapses, less psychotic symptoms.

Conclusions: Religion contributes to acquire adaptive functions as the meaning of life, sense of hope, spiritual comfort, supports the overcome the disease burden. Religious coping is an important tool for rehabilitation and preventing the relapses in patients with schizophrenic disorders.

Disclosure: No significant relationships.

Keywords: Religious Coping; remission in schizophrenia; schizophrenia; Schizoaffective disorder