

Poster presentation

Open Access

Protein tyrosine phosphatase nonreceptor 13 (PTPN13) is targeted by human papillomavirus 16 E6 in cervical epithelium

P Nowicki*², A Hoover¹, K De Geest², M Goodheart², M Lace³, T Haugen³, L Turek³, G Strand⁴, A Bossler⁴, A Klingelutz⁵ and J Lee⁶

Address: ¹Department of Otolaryngology, University of Iowa, Iowa City, Iowa, USA, ²Department of Obstetrics and Gynecology, Division of Gynecologic Oncology, University of Iowa, Iowa City, Iowa, USA, ³Veterans Affairs Medical Center, Iowa City, Iowa, USA, ⁴Department of Anatomic Pathology, University of Iowa, Iowa City, Iowa, USA, ⁵Department of Microbiology, University of Iowa, Iowa City, Iowa, USA and ⁶Sanford HPV Research Institute, Sanford Health, Sioux Falls, South Dakota, USA

* Corresponding author

from 11th International Conference on Malignancies in AIDS and Other Acquired Immunodeficiencies (ICMAOI): Basic, Epidemiologic, and Clinical Research
Bethesda, MD, USA. 6–7 October 2008

Published: 17 June 2009

Infectious Agents and Cancer 2009, **4**(Suppl 2):P31 doi:10.1186/1750-9378-4-S2-P31

This abstract is available from: <http://www.infectagentscancer.com/content/4/S2/P31>

© 2009 Nowicki et al; licensee BioMed Central Ltd.

Objectives

The Post Synaptic density protein-95/Drosophila disc large/Zonula occludens (PDZ) domain binding motif of human papillomavirus (HPV) E6 correlates with malignant transformation for high risk HPV. Recent evidence in our lab has shown that HPV16 E6 binds and degrades PTPN13 in a PDZ binding motif-dependent manner in HPV-positive head and neck squamous cell cancer. The loss of PTPN13 results in invasive growth when combined with Ras expression. The purpose of this study is to investigate PTPN13 expression in normal and malignant cervical epithelial cells, and to evaluate its role in cervical cancer.

Methods

PTPN13 expression was measured by Western blotting in primary cultures of normal and high-risk HPV-infected human cervix and foreskin keratinocytes, in non-tumorigenic HPV 31b positive cells, in HPV positive (CaSki, SiHa) and HPV-negative (C-33A) cervical carcinoma cell lines. Immunohistochemistry with PTPN13 (C-20): sc-1138 antibody was performed on cervical tissue specimens. Primary human cervical cultures and C33A cells were transduced with recombinant adenovirus containing HPV16 E6, E7, and mutant lacking the PDZ domain binding motif (E6Δ146–151). SiHa and Caski cells were transfected with plasmid containing PTPN13. PTPN13 loss was

induced in C33A cells using shRNA strategy. Morphologic features, in vitro growth rate, and anchorage independent growth were evaluated. RT-PCR was used to measure E6 and E6* expression.

Results

Normal cervical epithelial cells, foreskin keratinocytes and the HPV-negative C-33A cancer cell line express PTPN13. HPV-positive cancer cell lines, as well as cervical and foreskin epithelial cells containing HPV 16, 18, 31 show significant loss of PTPN13 expression. Cells transduced with HPV16 E6Δ146–151/E7 do not survive in vitro compared to cells transduced with HPV E6/E7. PTPN13 loss is E6 PDZ dependent and appears to be associated with integration of E6/E7 into the host genome. Re-establishment of PTPN13 in SiHa and Caski cells decreased colony formation efficiency in soft agar by 21 (p = 0.0031) and 68 percent (p = 0.0001) respectively. Loss of PTPN13 via shRNA in C33A cell line increases its colony formation efficiency 133 times (p = 0.0001) in soft agar.

Conclusion

PTPN13 may play an important role as a tumor suppressor in HPV-mediated cervical carcinogenesis. Understanding of this PDZ dependent mechanism may lead to development of targeted molecular therapy against HPV-positive cervical cancer.