



Published in final edited form as:

*Water Sci Technol.* 2024 May ; 89(9): 2440–2456. doi:10.2166/wst.2024.139.

## Isolation and characterization of pure cultures for metabolizing 1,4-dioxane in oligotrophic environments

Ermias Gebrekrstos Tesfamariam<sup>a</sup>, Dennis Ssekimpi<sup>a</sup>, Sarajeen Saima Hoque<sup>a</sup>, Huan Chen<sup>b</sup>, Joshua D. Howe<sup>c</sup>, Chao Zhou<sup>d</sup>, Yue-xiao Shen<sup>e</sup>, Youneng Tang<sup>a,\*</sup>

<sup>a</sup>Department of Civil and Environmental Engineering, FAMU-FSU College of Engineering, Florida State University, Tallahassee, Florida 32310, USA

<sup>b</sup>National High Magnetic Field Laboratory, Tallahassee, Florida 32310, USA

<sup>c</sup>Department of Chemical Engineering, Texas Tech University, Lubbock, Texas 79409, USA

<sup>d</sup>Geosyntec Consultants Inc., Costa Mesa, California 92626, USA

<sup>e</sup>Department of Civil, Environmental, and Construction Engineering, Texas Tech University, Lubbock, Texas 79409, USA

### Abstract

1,4-Dioxane concentration in most contaminated water is much less than 1 mg/L, which cannot sustain the growth of most reported 1,4-dioxane-metabolizing pure cultures. These pure cultures were isolated following enrichment of mixed cultures at high concentrations (20 to 1,000 mg/L). This study is based on a different strategy: 1,4-dioxane-metabolizing mixed cultures were enriched by periodically spiking 1,4-dioxane at low concentrations (1 mg/L). Five 1,4-dioxane-metabolizing pure strains LCD6B, LCD6D, WC10G, WCD6H, and WD4H were isolated and characterized. The partial 16S rRNA gene sequencing showed that the five bacterial strains were related to *Dokdonella* sp. (98.3%), *Acinetobacter* sp. (99.0%), *Afipia* sp. (99.2%), *Nitrobacter* sp. (97.9%), and *Pseudonocardia* sp. (99.4%), respectively. *Nitrobacter* sp. WCD6H is the first reported 1,4-dioxane-metabolizing bacterium in the genus of *Nitrobacter*. The net specific growth rates of these five cultures are consistently higher than those reported in the literature at 1,4-dioxane concentrations <0.5 mg/L. Compared to the literature, our newly discovered strains have lower half-maximum-rate concentrations (1.8 to 8.2 mg-dioxane/L), lower maximum specific 1,4-dioxane utilization rates (0.24 to 0.47 mg-dioxane/(mg-protein · d)), higher biomass yields

This is an Open Access article distributed under the terms of the Creative Commons Attribution Licence (CC BY-NC-ND 4.0), which permits copying and redistribution for non-commercial purposes with no derivatives, provided the original work is properly cited (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\*Corresponding author. ytang@eng.famu.fsu.edu.

#### AUTHOR CONTRIBUTIONS


Ermias Gebrekrstos Tesfamariam: methodology, investigation, formal analysis, software, visualization, writing – original draft, review and editing. Dennis Ssekimpi: investigation, formal analysis, writing – review and editing. Sarajeen Saima Hoque: investigation, formal analysis, writing – review and editing. Huan Chen: methodology, resources, writing – review and editing. Joshua D. Howe: methodology, writing – review and editing. Chao Zhou: methodology, writing – review and editing. Yue-xiao Shen: methodology, writing – review and editing. Youneng Tang: conceptualization, methodology, validation, supervision, funding acquisition, project administration, writing – review and editing.

#### CONFLICT OF INTEREST

The authors declare there is no conflict.

(0.29 to 0.38 mg-protein/mg-dioxane), and lower decay coefficients (0.01 to 0.02 d<sup>-1</sup>). These are characteristics of microorganisms living in oligotrophic environments.

## GRAPHICAL ABSTRACT



## Keywords

1,4-Dioxane; enrichment strategy; kinetics; *Nitrobacter*; oligotrophic environment; pure culture

## 1. INTRODUCTION

1,4-Dioxane (C<sub>4</sub>H<sub>8</sub>O<sub>2</sub>), a colorless cyclic ether compound (EPA 2017), was commonly used as a stabilizer of chlorinated solvents such as 1,1,1-trichloroethane (Mohr et al. 2010). It is also used as a solvent in some commercial and industrial processes and consumer products, leading to widespread groundwater contamination (Sei et al. 2013a; Pugazhendi et al. 2015; Zhang et al. 2017). According to the third Unregulated Contaminant Monitoring Rule (UCMR3), 1,4-dioxane was detected in 21% of the U.S. public water systems and exceeded the health-based reference concentration of 0.35 µg/L (EPA 2017) at 6.9% of the systems. The detection of 1,4-dioxane was ranked second among the 28 UCMR3 contaminants, only exceeded by chlorate (Adamson et al. 2017). The high mobility of 1,4-dioxane in groundwater has impacted relatively large regions with trace levels of contamination, which underscores the need to explore effective treatment methods applicable to environmentally relevant, low 1,4-dioxane concentrations of 365 µg/L (Adamson et al. 2014; Chiang et al. 2016; Wang et al. 2021).

Removal of 1,4-dioxane from contaminated water is challenging due to its unique physical and chemical properties such as high water solubility, low Henry's law constant (4.80 × 10<sup>-6</sup> atm m<sup>3</sup>/mol), low octanol–water partition coefficient (log *K<sub>ow</sub>* = -0.27), and low organic carbon partition coefficient (log *K<sub>oc</sub>* = 1.23) (Zenker et al. 2003; EPA 2017; Zhang et al. 2017). Bioremediation is a promising method for treating 1,4-dioxane-contaminated water as it is potentially cost-effective and ecofriendly. Plenty of studies have demonstrated aerobic biodegradation of 1,4-dioxane under metabolic and co-metabolic conditions. 1,4-Dioxane was co-metabolized in the presence of primary substrates such as ethane, propane,

isobutane, tetrahydrofuran, and toluene (Zenker et al. 2002; Mahendra & Alvarez-cohen 2006; Sei et al. 2013b; Hatzinger et al. 2017; Zhang et al. 2017; Barajas-Rodriguez & Freedman 2018; Rolston et al. 2019; Xiong et al. 2019a, 2020; Luo et al. 2021; Wang et al. 2021). In the presence of propane and isobutane as primary substrates, the minimum 1,4-dioxane concentrations required to sustain steady-state biomass ( $S_d^{min}$ ) for *Rhodococcus ruber* ENV425, Propanotrophic mixed culture ENV487, *Rhodococcus rhodochrous* strain 21198 were 1.0, 0.6, and 1.3 mg/L, respectively (Barajas-Rodriguez & Freedman 2018; Rolston et al. 2019). Co-metabolic bioremediation has the advantage of biomass growth supported by the primary substrate while treating 1,4-dioxane at low concentrations. However, the primary substrates can make the treatment process expensive and may lead to secondary contamination.

About 20 bacterial strains (Supporting Information Table S1) have been reported in the literature to use 1,4-dioxane as the sole carbon and energy source (Parales et al. 1994; Mahendra & Alvarez-Cohen 2005, 2006; Kim et al. 2009; Sei et al. 2013a; Huang et al. 2014; Pugazhendi et al. 2015; Chen et al. 2016; Inoue et al. 2016; Matsui et al. 2016; He et al. 2017b; Barajas-Rodriguez & Freedman 2018; Inoue et al. 2018; Yamamoto et al. 2018; Tusher et al. 2020; Ma et al. 2021; Simmer et al. 2021; Tusher et al. 2021; Dai et al. 2022; Ramos-García et al. 2022). The majority of them are gram-positive bacteria that belong to the genera *Pseudonocardia*, *Mycobacterium*, and *Rhodococcus*. The remaining are gram-negative bacteria that belong to the genera *Acinetobacter*, *Afipia*, *Ancylobacter*, *Dokdonella*, *Rhodanobacter*, *Variovorax*, and *Xanthobacter*. These strains were enriched and isolated under high concentrations of 1,4-dioxane (i.e., 20 to 1,000 mg/L). Only a few strains' 1,4-dioxane metabolism kinetics have been fully characterized (Barajas-Rodriguez & Freedman 2018; Ramos-García et al. 2022). There is a need to characterize the 1,4-dioxane metabolism kinetics of more cultures. Isolation and identification of 1,4-dioxane-metabolizing bacteria that can thrive in dilute 1,4-dioxane environments are of particular interest.

In this study, a different enrichment strategy was evaluated: 1,4-dioxane-metabolizing mixed cultures were sub-cultured by periodically spiking 1,4-dioxane at low concentrations (i.e., 1 mg/L). Three methods – centrifugation, dilution, and combination of centrifugation and dilution – were used to enrich 1,4-dioxane-metabolizing mixed cultures from two sources. After enrichment, five 1,4-dioxane-metabolizing pure cultures were isolated and characterized in terms of phylogenetics and 1,4-dioxane metabolism kinetics and intermediates.

## 2. MATERIALS AND METHODS

### 2.1. Chemicals and medium composition

All mixed cultures enrichment, growth of pure cultures, and kinetic experiments were performed in nitrate mineral salts medium. The medium was composed of 1.18 mM NaNO<sub>3</sub>, 6.1 mM K<sub>2</sub>HPO<sub>4</sub>, 3.9 mM KH<sub>2</sub>PO<sub>4</sub>, 1.28 mM Na<sub>2</sub>SO<sub>4</sub>, 0.08 mM FeSO<sub>4</sub>·7H<sub>2</sub>O, 0.15 mM MgCl<sub>2</sub>·6H<sub>2</sub>O, 0.07 mM CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.002 mM ZnCl<sub>2</sub>, 0.002 mM MnCl<sub>2</sub>·4H<sub>2</sub>O, 0.004 mM CoCl<sub>2</sub>·6H<sub>2</sub>O, 0.001 mM CuCl<sub>2</sub>·2H<sub>2</sub>O, 0.001 mM NiCl<sub>2</sub>·6H<sub>2</sub>O, 0.002 mM H<sub>3</sub>BO<sub>3</sub>, 0.004 mM Na<sub>2</sub>MoO<sub>4</sub>·2H<sub>2</sub>O, 0.001 mM Na<sub>2</sub>WO<sub>4</sub>·2H<sub>2</sub>O, 0.001 mM Na<sub>2</sub>SeO<sub>4</sub>·5H<sub>2</sub>O, and 0.001 mM KI in deionized water (Xiong et al. 2019a). The pH of the medium was adjusted to ~7.0

using 0.1 mM H<sub>2</sub>SO<sub>4</sub>. The deionized water was autoclaved at 121 °C for 20 min and cooled to room temperature before the stock chemicals were added. 1,4-Dioxane was purchased from Acros Organics as American Chemical Society (ACS) grade with >99% purity. Agar powder was obtained from Thermo Fisher Scientific. All other chemicals and solvents used in this study were purchased as ACS grade.

## 2.2. Mixed culture enrichment

Two microbial cultures originally from a landfill (L) and activated sludge of a wastewater treatment plant (W) were separately cultured by regularly adding 1,4-dioxane for more than 300 days. Each culture was maintained in three 500 mL serum bottles containing 200 mL of nitrate mineral salts medium. Before each culture was added into the serum bottle, 20 mg of the original culture was centrifuged at 5,000 × *g* for 10 minutes and washed with the nitrate mineral salts medium twice. The objective was to remove residual carbon carried over from the seed material. 1,4-Dioxane was added as a sole electron donor at 1 mg/L whenever its concentration dropped below 0.2 mg/L. The cultures were incubated aerobically at room temperature (25 °C) on a shaker rotating at 120 revolutions per minute (Sei et al. 2013a, 2013b; Nam et al. 2016; Yamamoto et al. 2018). To minimize contamination, the bottles were closed with rubber stoppers and were opened inside a biosafety cabinet every two days to allow oxygen replenishment.

Each culture was enriched by three methods, corresponding to the three serum bottles mentioned above. In the first method (i.e., centrifugation), centrifugation was applied to the cultures after every five cycles of 1,4-dioxane degradation. One bottle from each culture (LC and WC) was centrifuged at 5,000 × *g* for 10 minutes. The pellet was washed with the nitrate mineral salts medium and resuspended into 200 mL fresh medium containing 1 mg/L 1,4-dioxane (Sei et al. 2013a; Nam et al. 2016; Yamamoto et al. 2018). In the second method (i.e., dilution), one bottle from each culture (LD and WD) was sub-cultured through 2–10 times dilution with fresh nitrate mineral salts medium containing 1,4-dioxane at 1 mg/L (Kim et al. 2009; Sei et al. 2013b; Tusher et al. 2020). In the third method (i.e., centrifugation and dilution), one bottle from each culture (LCD and WCD), was sub-cultured through centrifugation followed by dilution. After every 10 cycles of 1,4-dioxane degradation, half of the culture medium was centrifuged at 5,000 × *g* for 10 minutes. The pellet was washed with and then resuspended into 200 mL fresh medium containing 1 mg/L 1,4-dioxane. The three enrichment methods were selected because they create environmental conditions similar to that of 1,4-dioxane contaminated groundwater and allow the growth of the microorganisms of interest.

## 2.3. Isolation and phylogenetic characterization of 1,4-dioxane-metabolizing pure cultures

1,4-Dioxane-metabolizing bacteria from the six culture bottles described in Section 2.2 (i.e., LC, WC, LD, WD, LCD, and WCD) were isolated by streaking 100 µL of the enriched cultures on solid agar plates supplemented with nitrate mineral salts medium containing 100 mg/L 1,4-dioxane. A 100 mg/L 1,4-dioxane concentration was used during the isolation stage to achieve good growth of colonies. The plates were sealed with a parafilm and incubated upside down at 25 °C. A total of 54 colonies were isolated from the

six microbial consortia: nine colonies for each consortium. Single colonies were inoculated into 50 mL cell culture flasks (with vented caps) containing 10 mL of nitrate mineral salts medium with 100 mg/L 1,4-dioxane. The flasks were incubated aerobically at room temperature (25 °C) on a shaker rotating at 120 revolutions per minute. The concentrations of 1,4-dioxane in the flasks were monitored to evaluate their 1,4-dioxane degradation ability. Forty-seven of the 54 colonies showed 1,4-dioxane degradation. Among them, 30 colonies were further selected, based on their faster 1,4-dioxane degradation, for phylogenetic identification. Genomic DNA were extracted using EZNA<sup>®</sup> Bacterial DNA Kit following the manufacturer's protocol. The extracted DNA were analyzed at Florida State University, Department of Biological Science, Core Facilities. Partial sequences of 16S rRNA gene were amplified from the extracted DNA by polymerase chain reaction using the 27F and 1492R primer set. The amplicons were purified and sequenced using bi-directional Sanger sequencing. The forward and backward sequences were initially analyzed using Molecular Evolutionary Genetics Analysis version 11 (MEGA11) (Tamura et al. 2021). Chimera analysis was then performed using vsearch 2.21.1. Five strains that passed the chimera check were further studied for 1,4-dioxane degradation kinetics and intermediates. The aligned sequences were compared with sequences in the National Center of Biotechnology Information database using the Basic Local Alignment Search Tool. A phylogenetic tree was constructed using MEGA11 for the five 1,4-dioxane-metabolizing strains identified in this study and the bacteria reported in the literature that could metabolically degrade 1,4-dioxane. The partial 16S rRNA sequences were deposited to the GenBank database of the National Center of Biotechnology Information.

#### 2.4. Kinetic experiments and modeling for the pure cultures

The kinetic experiments were conducted in sealed 250 mL serum bottles. The biomass from the pure cultures was centrifuged at  $5,000 \times g$  for 10 minutes. After removing the supernatant, the pelleted biomass was washed with the nitrate mineral salts medium, centrifuged, and resuspended into 100 mL fresh medium with 100 mg/L 1,4-dioxane. The initial biomass concentrations ranged from 10 to 20 mg/L. This range was chosen due to the 5 mg/L detection limit of the Pierce BCA Protein Assay Kit used. A 100 mg/L 1,4-dioxane concentration was used so that the 1,4-dioxane concentration range (0–100 mg/L) covered the median half-maximum-rate constant of 1,4-dioxane reported in the literature (42.5 mg/L) (Tang 2023). This helps to increase the accuracy of the parameter estimation. The concentrations of 1,4-dioxane and protein (representing biomass) were monitored daily.

A Monod-based kinetic model was used to describe the rates of change in the concentrations of 1,4-dioxane (Equation (1)) and biomass (Equation (2)). Descriptions of the model input and model output parameters are presented in Table 1. Equations (1) and (2) were solved numerically by a numerical code written in FORTRAN. Initial conditions included the concentrations of 1,4-dioxane and biomass that were experimentally determined at day 0 (i.e.,  $S_{d(0)}^e$  and  $X_{(0)}^e$ ). The unknown kinetic parameters  $q_d$ ,  $K_d$ ,  $Y_d$ , and  $b$  were determined by fitting the experimentally measured concentrations (i.e.,  $S_{d(t)}^e$  and  $X_{(t)}^e$ ) to the model simulated concentrations (i.e.,  $S_{d(t)}^m$  and  $X_{(t)}^m$ ) using a model-independent parameter estimator (Doherty 2019). The goodness of fit between the experiment and the model simulation was determined using the mean sum of squared relative residuals (Equation (3)). This approach

differs from previous work reported by Barajas-Rodriguez & Freedman (2018) in that the four kinetic parameters were estimated simultaneously. The estimated parameters were then evaluated by comparing a new set of experimental data with their corresponding simulated results. The conditions of the experiments used for the model evaluation were the same as the kinetics estimation, except that the initial 1,4-dioxane concentration was 10 mg/L in the evaluation experiments.

$$\frac{dS_d}{dt} = -X q_d \left[ \frac{S_d}{K_d + S_d} \right] \quad (1)$$

$$\frac{dX}{dt} = Y_d X q_d \left[ \frac{S_d}{K_d + S_d} \right] - b X \quad (2)$$

$$MSRR = \frac{1}{N} \left[ \sum_{t=0}^T \left( \frac{S_{d(t)}^m - S_{d(t)}^e}{S_{d(t)}^e} \right)^2 + \sum_{t=0}^T \left( \frac{X_{(t)}^m - X_{(t)}^e}{X_{(t)}^e} \right)^2 \right] \quad (3)$$

To evaluate the growth potential of the isolated cultures in oligotrophic environments (i.e., < 1 mg/L in this study) and compare it to that of the literature-reported cultures, the net specific growth rates of the cultures and the minimum 1,4-dioxane concentration required to sustain steady-state biomass were calculated by Equations (4) and (5) (Rittmann & McCarty 2020).

$$\mu = Y_d q_d \left[ \frac{S_d}{K_d + S_d} \right] - b \quad (4)$$

$$S_d^{min} = K_d \frac{b}{Y_d q_d - b} \quad (5)$$

## 2.5. Identification and quantification of 1,4-dioxane intermediates for the pure cultures

The experiments for identification and quantification of 1,4-dioxane intermediates were performed in triplicate for each of the five pure cultures. The biomass was centrifuged at  $5,000 \times g$  for 10 minutes. After removing the supernatant, the pelleted biomass was washed with the nitrate mineral salts medium, centrifuged, and resuspended into 200 mL fresh medium with 100 mg/L 1,4-dioxane. The bottles were incubated aerobically at room temperature (25 °C) on a shaker rotating at 120 revolutions per minute. Liquid samples

were filtered using 0.20  $\mu\text{m}$  Acrodisc<sup>®</sup> syringe filters and stored in a refrigerator at 4 °C before they were analyzed for 1,4-dioxane and its degradation intermediates following the analytical methods in Section 2.6.

## 2.6. Analytical methods

The concentrations of 1,4-dioxane in all experiments were measured using solid phase micro-extraction followed by gas chromatography/mass spectrometry, which is described in our previous publication (Xiong et al. 2019a). The quantification limit was 0.5  $\mu\text{g}$ -1,4-dioxane/L. 1,4-Dioxane intermediates were analyzed by adapting the methods in our previous publication (Xiong et al. 2019b). First, chemical oxygen demand (COD) concentrations were measured by Hach DR 3900 spectrophotometer using low range (3 to 150 mg-COD/L) and ultra-low range (1 to 60 mg-COD/L) COD kits. The COD concentration was compared to the 1,4-dioxane concentration (after being converted to mg-COD/L) for each sample to estimate whether intermediates were contributing to COD. A paired t-test at a  $p$ -value of 0.05 was used to evaluate the statistical significance of the difference between the concentrations of 1,4-dioxane and total COD. Second, the presence of organic acids, including formic, glycolic, glyoxylic, methoxyacetic, and oxalic acids, as potential 1,4-dioxane degradation intermediates were checked using a DIONEX ion chromatography system. The detection limits of the organic acids were approximately 0.1 mg/L. All aqueous samples used for COD and ion chromatography measurements were filtered by 0.20  $\mu\text{m}$  Acrodisc<sup>®</sup> syringe filters. Third, the presence of potential 1,4-dioxane degradation intermediates including the organic acids, ethylene glycol, and 1,4-dioxan-2-ol, were checked using gas chromatography/mass spectrometry operated in the scan mode.

To evaluate the biomass concentration change in all the experiments, protein concentrations were measured by a Nanodrop spectrophotometer using Pierce BCA Protein Assay Kit with Albumin Standard Ampules. The detection limit for protein measurement was 5 mg-protein/L. The concentrations of the extracted DNA were measured using the same spectrophotometer.

## 3. RESULTS AND DISCUSSION

### 3.1. The five isolated 1,4-dioxane-metabolizing strains

Six microbial consortia capable of utilizing 1,4-dioxane as a sole carbon and energy source were enriched. They corresponded to two sources (L and W) and three enrichment methods (C, D, and CD) for each source. Figure S1 shows the pictures of the six mixed cultures during the enrichment: the three cultures based on the L source were green, while the three cultures based on the W source were brown. The 1,4-dioxane degradation profiles of different generations of the consortia are presented in Figure S2. The six mixed cultures were able to degrade 1,4-dioxane from 1 mg/L to below 0.01 mg/L. Figure S3 compares the 1,4-dioxane utilization rates of the cultures. At the beginning of the enrichment, the 1,4-dioxane utilization rates of the six cultures slightly varied between  $2.1 \times 10^{-3}$  and  $2.9 \times 10^{-3}$  mg-dioxane/(mg-protein  $\cdot$  d). During the 300 days of enrichment, the two CD-based cultures (LCD and WCD) had the highest utilization rates, approximately five times higher than their initial rates. On the other hand, the utilization rates of the two C-based (LC and

WC) and D-based (LD and WD) cultures increased by two and four times, respectively, compared to their initial rates. The lower utilization rates of the cultures enriched by centrifugation (C-based) could be due to the retainment of the inert biomass from one generation to the next. Therefore, dilution or combination of centrifugation and dilution should be considered during culture enrichments to enhance the growth of active biomass and minimize retainment of inert biomass.

The five isolated bacterial strains were *Dokdonella* sp. LCD6B, *Acinetobacter* sp. LCD6D, *Afipia* sp. WC10G, *Nitrobacter* sp. WCD6H, and *Pseudonocardia* sp. WD4H, respectively. They represent successful isolation from both mixed cultures and the three enrichment methods. The five identified strains were related to genera *Dokdonella* sp. (98.3%), *Acinetobacter* sp. (99.0%), *Afipia* sp. (99.2%), *Nitrobacter* sp. (97.9%), and *Pseudonocardia* sp. (99.4%) respectively. The partial 16S rRNA gene sequences of the five pure strains are deposited at the NCBI GenBank database under accession numbers OP362562, OP362563, OP362564, OP362565, and OP362566, respectively. The phylogenetic tree shown in Figure 1 presents the evolutionary relationships among the five 1,4-dioxane-metabolizing bacteria isolated in this study and bacteria reported in the literature that can metabolically degrade 1,4-dioxane. The 24 literature-reported 1,4-dioxane-metabolizing pure cultures are summarized in Table S1. The upper 14 in the tree are gram-negative bacterial strains that belong to the phylum Proteobacteria while the lower groups are gram-positive bacteria from the Actinobacteria phylum. This study isolated *Nitrobacter* sp. WCD6H as the first 1,4-dioxane-metabolizing bacterium that belongs to the *Nitrobacter* genus.

1,4-Dioxane biodegradation is initiated by enzymes encoding soluble di-iron monooxygenase (SDIMO) genes. SDIMO genes also have the ability to catalyze the initial oxidation of a wide variety of hydrocarbons (Li et al. 2014; Inoue et al. 2016; Tusher et al. 2021). To date, three SDIMO gene clusters (Group-2, Group-5, and Group-6) have been confirmed to be involved in the metabolic biodegradation of 1,4-dioxane (Table S1). Li et al. (2014) found a positive correlation between the biodegradation of 1,4-dioxane and the abundance of Group-5 SDIMO genes, tetrahydrofuran/dioxane monooxygenase (*thmA/dxmA*). Genes *thmA/dxmA* also were identified in the 1,4-dioxane-metabolizing bacteria that belong to the genera *Dokdonella*, *Afipia*, and *Pseudonocardia* (Inoue et al. 2016; Tusher et al. 2021) (Table 2). Similarly, tetrahydrofuran monooxygenase gene cluster (*thmADBC*) was expressed by *Pseudonocardia dioxanivorans* CB1190 during growth on 1,4-dioxane (Grostern et al. 2012; Sales et al. 2013). However, the molecular mechanisms of 1,4-dioxane biodegradation are only partially understood. The genes associated with the metabolism of 1,4-dioxane by *Acinetobacter*, *Nitrobacter*, and several 1,4-dioxane degrading pure cultures in the literature are unknown (Table S1). Future studies that incorporate genomic and proteomic analyses will help to further elucidate the metabolic pathways and enzymatic mechanisms underlying 1,4-dioxane degradation.

### 3.2. Kinetic parameters estimation and evaluation

The 1,4-dioxane biodegradation kinetics of the five strains were described well by the Monod-based equations with low MSRR values ranging from 0.01 to 0.11. The estimated kinetic parameters for the five strains are presented in Table 3. Figure 2 further compares



the experimental and modeling results for 1,4-dioxane and biomass concentrations in the kinetic experiments of the five strains. The parameterized model captured all major trends for the concentrations of 1,4-dioxane and biomass. In general, the variability of the kinetic parameters is quite small, especially  $Y_d$  and  $b$ . The variability for  $q_d$  and  $K_d$  are slightly greater, with approximate 2- and 4.5-fold difference between the largest and smallest values, respectively. The reason for the narrow range of kinetic parameters may be associated with the enrichment strategy. As the mixed cultures were enriched with 1 mg/L 1,4-dioxane for more than 300 days, it may have created an environment to enrich strains with similar kinetic parameters. The kinetic parameters presented in Table 3 will be discussed in this section by comparing them with corresponding values from the literature.

The net specific growth rates of the five cultures isolated in this study and two pure cultures from the literature are shown in Figure 3. *Pseudonocardia dioxanivorans* CB1190 (Barajas-Rodriguez & Freedman 2018) and *Pseudonocardia dioxanivorans* BERK-1 (Ramos-García et al. 2022) are the only pure cultures with a full set of kinetics in the literature and are used for comparison. *P. dioxanivorans* CB1190 has about 2- to 8-fold higher net specific growth rates at 1,4-dioxane concentrations  $> 0.5$  mg/L, while all the five strains in this study have about four times higher net specific growth rates at concentrations  $< 0.5$  mg/L. This supports our hypothesis that enrichment with 1,4-dioxane at lower concentration helps to isolate oligotrophs. The minimum 1,4-dioxane concentration required to sustain steady-state biomass in a continuous-flow, suspended-growth reactor ( $S_d^{min}$ ) ranged from 0.4 to 1.5 mg/L (Table 3). This is at the lower end of the range (0.4–20 mg/L) reported in the literature (Tang 2023). Based on the  $S_d^{min}$ , two of the five cultures in Table 3 are able to grow at 1,4-dioxane  $< 1$  mg/L. In an attached-growth reactor (i.e., biofilm reactor) that enhances biomass attachment and growth, the  $S_d^{min}$  values should be much lower than 0.4 mg/L (Tang 2023). Simmer et al. (2021) reported low  $S_d^{min}$  value of 0.49  $\mu$ g/L for *Rhodococcus ruber* strain 219. However, the decay coefficient was assumed to be the same as that of *P. dioxanivorans* CB1190 (Simmer et al. 2021).

Figure 4 compares the 1,4-dioxane biodegradation kinetic parameters determined in this study to the same parameters reported in the literature for all 1,4-dioxane metabolizing mixed cultures (Zenker et al. 2002; Mahendra & Alvarez-cohen 2006; Nam et al. 2016; Barajas-Rodriguez & Freedman 2018; Lee et al. 2020; Xiong et al. 2022) and pure cultures (Mahendra & Alvarez-cohen 2006; Sei et al. 2013a; Chen et al. 2016; He et al. 2017a; Barajas-Rodriguez & Freedman 2018; Inoue et al. 2018; Yamamoto et al. 2018; Ma et al. 2021; Simmer et al. 2021). Figure 5 further compares the kinetic parameters of *Afiplia* and *Pseudonocardia* isolated in this study and the literature. Only two genera are compared in Figure 5 because no kinetics data is available in the literature for biodegradation of 1,4-dioxane by *Dokdonella*, *Acinetobacter*, and *Nitrobacter*. Figures 4 and 5 show the same comparison results: compared to the 1,4-dioxane-metabolizing cultures in the literature, the five pure cultures have lower half-maximum rate constants, higher biomass yields, lower maximum specific 1,4-dioxane utilization rates, and lower decay coefficients. These are characteristics of microorganisms living in oligotrophic environments (Koch 2001; Xiong et al. 2022; Tang 2023).

As an example of these four parameters, the half-maximum-rate concentration is discussed in detail here. The half-maximum-rate concentrations obtained in this study ranged from 1.8 to 8.2 mg-dioxane/L. The 1,4-dioxane-metabolizing pure cultures reported in the literature have high half-maximum-rate concentrations ranging from 6.3 to 411 mg-dioxane/L (Mahendra & Alvarez-cohen 2006; Sei et al. 2013a; Chen et al. 2016; He et al. 2017a; Barajas-Rodriguez & Freedman 2018; Inoue et al. 2018; Yamamoto et al. 2018; Ma et al. 2021). *Nitrobacter* sp. WCD6H showed the lowest half-maximum-rate concentration of 1.8 mg-dioxane/L among the five strains isolated in this study. The half-maximum-rate concentration of *Afiplia* sp. WC10G was 8.2 mg-dioxane/L, about 3-fold lower than that of *Afiplia* sp. D1, 26 mg-dioxane/L (Sei et al. 2013a). Similarly, *Pseudonocardia* sp. WD4H had a half-maximum-rate concentration of 5.3 mg-dioxane/L. This was about 1.2 to 30-fold lower than those of *P. dioxanivorans* BERK-1 (14.7 mg-dioxane/L) (Ramos-García et al. 2022), *P. dioxanivorans* CB1190 (6.3 to 160 mg-dioxane/L) (Mahendra & Alvarez-cohen 2006; He et al. 2017a; Barajas-Rodriguez & Freedman 2018), *Pseudonocardia* sp. D17 (60 mg-dioxane/L) (Sei et al. 2013a), and *Pseudonocardia* sp. N23 (80 mg-dioxane/L) (Yamamoto et al. 2018).

The kinetic parameters presented in Table 3 were evaluated by comparing results from a second set of independent experiments with their corresponding model-simulated results. Figure 6 compares the experimentally measured data and the model simulated results for the five isolated strains. The parameterized model captured the major trends for the concentrations of 1,4-dioxane and biomass. The MSRR values obtained during model evaluation were 0.22, 0.27, 0.26, 0.22, and 0.13, respectively. These results were 2- to 13-fold larger than those obtained during parameter estimation. The slower degradation rates in the evaluation experiments than the model predictions could be due to the effect of lag phase in the biomass growth.

### 3.3. 1,4-dioxane intermediates

Figure 7 compares the 1,4-dioxane concentrations (converted from mg-1,4-dioxane/L to mg-COD/L by a conversion factor of 1.82) and the total COD of the corresponding water samples for the five isolated strains. When 1,4-dioxane was metabolized from 180 to 20 mg-COD/L, the difference between the concentrations of 1,4-dioxane and total COD was not statistically significant at a confidence level of 95%, suggesting negligible production of 1,4-dioxane intermediates. When the 1,4-dioxane was < 20 mg-COD/L (i.e., 11 mg-1,4-dioxane/L), the total COD concentration was higher than the 1,4-dioxane concentrations, suggesting that some chemicals other than 1,4-dioxane were present. The base COD value of the nitrate mineral salts medium without 1,4-dioxane was  $1.33 \pm 0.58$  mg-COD/L. The other chemicals could be 1,4-dioxane metabolism intermediates or soluble microbial products due to endogenous respiration of the bacteria (Rittmann & McCarty 2020). Further analyses of the liquid samples by the gas chromatography/mass spectrometry and ion chromatograph showed no presence of any 1,4-dioxane intermediates discussed in Section 2.6 above the detection limits. Therefore, we inferred that soluble microbial products generated from endogenous respiration of the active biomass might have contributed to the total COD during the decay phase of the experiment.

Several previous studies identified biodegradation intermediates of 1,4-dioxane. The major intermediates identified during 1,4-dioxane biodegradation by monooxygenase-expressing bacteria were ethylene glycol, glycolate, 2-Hydroxyethoxyacetic acid, and oxalate (Mahendra et al. 2007). 1,4-Dioxane degradation by *Mycobacterium* sp. PH-06 also showed 1,4-dioxane-2-ol and ethylene glycol as the major degradation intermediates (Kim et al. 2009). On the other hand, 1,4-dioxene was the only 1,4-dioxane intermediate by *Acinetobacter baumannii* DD1 (Huang et al. 2014). Chen et al. (2016) found three 1,4-dioxane intermediates: 1,4-dioxene, ethylene glycol, and oxalic acid by *Xanthobacter flavus* DT8. The differences in the identification of 1,4-dioxane intermediates indicate that the degradation pathway of 1,4-dioxane may vary depending on microbial composition.

### 3.4. Practical implications of the study

The feasibility and practical applications of 1,4-dioxane bioremediation require profound research on the kinetics, operating parameters, and sustainability of the treatment system. This research will contribute to the current needs to identify microorganisms for treating 1,4-dioxane. The kinetic constants obtained in this study could be used in the design and assessment of large scale 1,4-dioxane remediation technologies. The minimum 1,4-dioxane concentrations required to sustain steady-state biomass ( $S_d^{min}$ ) for the isolated strains are at the lower end of the range reported in the literature. However, the  $S_d^{min}$  values are still higher than most 1,4-dioxane contaminated sites. The  $S_d^{min}$  of these strains are derived based on continuous-flow stirred-tank reactor containing suspended biomass. Biofilm reactors inoculated with these microorganisms are expected to have a smaller  $S_d^{min}$  than those of suspended-growth reactors since the biofilm provides protection to the microorganisms, leading to a lower effective biomass decay rate. Incorporating adsorption and biodegradation using materials that can adsorb 1,4-dioxane, provide biomass attachment, and concentrate substrates and nutrients could also enhance the bioremediation of 1,4-dioxane.

## 4. CONCLUSIONS

1,4-Dioxane-metabolizing pure cultures reported in the literature were isolated following enrichment with 1,4-dioxane at high concentrations (20–1,000 mg/L). This study enriched 1,4-dioxane-metabolizing mixed cultures through periodically spiking 1,4-dioxane at low concentrations (1 mg/L) and then isolated five bacterial strains that could metabolize 1,4-dioxane. This strategy worked well for two mixed cultures: the first one from a landfill leachate and the second one from a municipal wastewater treatment plant. It also worked well for three enrichment methods: dilution, centrifugation, and their combination, but dilution and combination of centrifugation and dilution resulted in the highest 1,4-dioxane utilization rate for the mixed cultures. The five isolated bacterial strains were *Dokdonella* sp. LCD6B, *Acinetobacter* sp. LCD6D, *Afipia* sp. WC10G, *Nitrobacter* sp. WCD6H, and *Pseudonocardia* sp. WD4H. To the best of our knowledge, WCD6H is the first known strain in the genus *Nitrobacter* that can metabolize 1,4-dioxane. Only two 1,4-dioxane-metabolizing pure cultures in the literature have a full set of kinetic parameters for calculating the net specific growth rate. Compared to these pure cultures, the five strains isolated in our study have higher net specific growth rates when the 1,4-dioxane concentrations are <0.5 mg/L. Further comparison of the kinetic parameters between the

five strains in this study and the pure and mixed cultures reported in the literature shows that the five strains in this study have lower half-maximum-rate concentrations (1.8 to 8.2 mg-dioxane/L), higher biomass yields (0.29 to 0.38 mg-protein/mg-dioxane), lower decay coefficients (0.01 to 0.02 d<sup>-1</sup>), and lower maximum specific 1,4-dioxane utilization rates (0.24 to 0.47 mg-dioxane/(mg-protein · d)). The minimum 1,4-dioxane concentration required to sustain steady-state biomass ranged from 0.4 to 1.5 mg/L. The five strains were also similar in that they did not produce detectable 1,4-dioxane intermediates.

In order to further elucidate the metabolic pathways and enzymatic mechanisms underlying 1,4-dioxane degradation by the isolated strains, future studies incorporating genomic and proteomic analyses are warranted. These analyses will provide a more comprehensive understanding of the metabolic capabilities of the organisms and contribute to the broader knowledge of biodegradation processes. Additionally, exploring the genomic and proteomic profiles of these strains will facilitate the identification of specific genes or enzymes critical to 1,4-dioxane degradation, ultimately aiding in the development of targeted bioremediation strategies.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## ACKNOWLEDGEMENTS

Research reported in this publication was funded by the National Institute of Environmental Health Sciences of the National Institutes of Health, United States under Award Number R01ES032692. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Work performed at the National High Magnetic Field Laboratory ICR Facility was supported by the National Science Foundation Division of Chemistry and Division of Materials Research through NSF DMR-2128556, and the State of Florida.

## DATA AVAILABILITY STATEMENT

All relevant data are included in the paper or its Supplementary Information.

## REFERENCES

- Adamson DT, Mahendra S, Walker KL, Rauch SR, Sengupta S & Newell CJ 2014 A multisite survey to identify the scale of the 1,4-dioxane problem at contaminated groundwater sites. *Environmental Science & Technology Letters* 1 (5), 254–258. 10.1021/ez500092u.
- Adamson DT, Piña EA, Cartwright AE, Rauch SR, Anderson RH, Mohr T & Connor JA 2017 1,4-Dioxane drinking water occurrence data from the third unregulated contaminant monitoring rule. *Science of the Total Environment* 596–597, 236–245. 10.1016/j.scitotenv.2017.04.085.
- Barajas-Rodriguez FJ & Freedman DL 2018 Aerobic biodegradation kinetics for 1,4-dioxane under metabolic and cometabolic conditions. *Journal of Hazardous Materials* 350, 180–188. 10.1016/j.jhazmat.2018.02.030. [PubMed: 29477886]
- Chen D, Jin X, Chen J, Ye J, Jiang N & Chen J 2016 Intermediates and substrate interaction of 1,4-dioxane degradation by the effective metabolizer *Xanthobacter flavus* DT8. *International Biodeterioration & Biodegradation* 106, 133–140. 10.1016/j.ibiod.2015.09.018.
- Chiang SD, Anderson RH, Wilken M & Walecka-Hutchison C 2016 Practical perspectives of 1,4-Dioxane investigation and remediation. *Remediation* 27 (1), 7–27. 10.1002/rem.21494.

- Dai C, Wu H, Wang X, Zhao K & Lu Z 2022 Network and meta-omics reveal the cooperation patterns and mechanisms in an efficient 1,4-dioxane-degrading microbial consortium. *Chemosphere* 301, 134723. 10.1016/j.chemosphere.2022.134723. [PubMed: 35489450]
- Doherty J. 2019 PEST Model-Independent Parameter Estimation. User Manual Part I: PEST, SENSAN and Global Optimisers, 7th edn. Watermark Numerical Computing, Brisbane, Australia.
- EPA. 2017 Technical Fact Sheet – 1,4-Dioxane. United States Environmental Protection Agency Office, Office of Land and Emergency Management, Washington, D.C., USA.
- Groster A, Sales CM, Zhuang WQ, Erbilgin O & Alvarez-Cohen L 2012 Glyoxylate metabolism is a key feature of the metabolic degradation of 1,4-dioxane by *Pseudonocardia dioxanivorans* strain CB1190. *Applied and Environmental Microbiology* 78 (9), 3298–3308. 10.1128/AEM.00067-12. [PubMed: 22327578]
- Hatzinger PB, Banerjee R, Rezes R, Streger SH, Mcclay K & Schaefer CE 2017 Potential for cometabolic biodegradation of 1,4-dioxane in aquifers with methane or ethane as primary substrates. *Biodegradation* 28 (5–6), 453–468. 10.1007/s10532-017-9808-7. [PubMed: 29022194]
- He Y, Mathieu J, Yang Y, Yu P, da Silva MLB & Alvarez PJJ 2017a 1,4-Dioxane biodegradation by *Mycobacterium Dioxanotrophicus* PH-06 is associated with a group-6 soluble di-iron monooxygenase. *Environmental Science & Technology Letters* 4 (11), 494–499. 10.1021/acs.estlett.7b00456.
- He Y, Wei K, Si K, Mathieu J, Li M & Alvarez PJJ 2017b Whole-genome sequence of the 1,4-dioxane-degrading bacterium *Mycobacterium dioxanotrophicus* PH-06. *Genome Announcement* 5 (35), 1–3. 10.1128/genomeA.00625-17.
- Huang H, Shen D, Li N, Shan D, Shentu J & Zhou Y 2014 Biodegradation of 1,4-Dioxane by a novel strain and its biodegradation pathway. *Water, Air, & Soil Pollution* 225, 2135. 10.1007/s11270-014-2135-2.
- Inoue D, Tsunoda T, Sawada K, Yamamoto N, Saito Y, Sei K & Ike M 2016 1,4-Dioxane degradation potential of members of the genera *Pseudonocardia* and *Rhodococcus*. *Biodegradation* 27 (4–6), 277–286. 10.1007/s10532-016-9772-7. [PubMed: 27623820]
- Inoue D, Tsunoda T, Yamamoto N, Ike M & Sei K 2018 1,4-Dioxane degradation characteristics of *Rhodococcus aetherivorans* JCM 14343. *Biodegradation* 29 (3), 301–310. 10.1007/s10532-018-9832-2. [PubMed: 29696449]
- Kim YM, Jeon JR, Murugesan K, Kim EJ & Chang YS 2009 Biodegradation of 1,4-dioxane and transformation of related cyclic compounds by a newly isolated *Mycobacterium* sp. PH-06. *Biodegradation* 20 (4), 511–519. 10.1007/s10532-008-9240-0. [PubMed: 19085063]
- Koch AL 2001 Oligotrophs versus copiotrophs. *BioEssays* 23 (7), 657–661. [PubMed: 11462219]
- Lee KH, Wie YM & Lee YS 2020 Characterization of 1,4-dioxane biodegradation by a microbial community. *Water* 12 (3372), 1–12. 10.3390/w12123372.
- Li M, Mathieu J, Liu Y, Orden ETV, Yang Y, Fiorenza S & Alvarez PJJ 2014 The abundance of tetrahydrofuran/dioxane monooxygenase genes (*thmA/dxmA*) and 1,4-dioxane degradation activity are significantly correlated at various impacted aquifers. *Environmental Science & Technology Letters* 1, 122–127. 10.1021/ez400176h.
- Luo Y, Long X, Wang B, Zhou C, Tang Y, Krajmalnik-Brown R & Rittmann BE 2021 A synergistic platform for continuous co-removal of 1,1,1-Trichloroethane, trichloroethene, and 1,4-dioxane via catalytic dechlorination followed by biodegradation. *Environmental Science & Technology* 55, 6363–6372. 10.1021/acs.est.1c00542. [PubMed: 33881824]
- Ma F, Wang Y, Yang J, Guo H, Su D & Yu L 2021 Degradation of 1,4-dioxane by *Xanthobacter* sp. YN2. *Current Microbiology* 78 (3), 992–1005. 10.1007/s00284-021-02347-6. [PubMed: 33547937]
- Mahendra S & Alvarez-cohen L 2006 Kinetics of 1,4-dioxane biodegradation by monooxygenase-expressing bacteria. *Environmental Science & Technology* 40 (17), 5435–5442. 10.1021/es060714v. [PubMed: 16999122]
- Mahendra S & Alvarez-Cohen L 2005 *Pseudonocardia dioxanivorans* sp. nov., a novel actinomycete that grows on 1,4-dioxane. *International Journal of Systematic and Evolutionary Microbiology* 55 (2), 593–598. 10.1099/ijs.0.63085-0. [PubMed: 15774630]


- Mahendra S, Petzold CJ, Baidoo EE, Keasling JD & Alvarez-Cohen L 2007 Identification of the intermediates of in vivo oxidation of 1,4-dioxane by monooxygenase-containing bacteria. *Environmental Science & Technology* 41 (21), 7330–7336. 10.1021/es0705745. [PubMed: 18044507]
- Matsui R, Takagi K, Sakakibara F, Abe T & Shiiba K 2016 Identification and characterization of 1,4-dioxane-degrading microbe separated from surface seawater by the seawater-charcoal perfusion apparatus. *Biodegradation* 27, 155–163. 10.1007/s10532-016-9763-8. [PubMed: 27094948]
- Mohr T, Stickney J & DiGiuseppi B 2010 *Environmental Investigation and Remediation: 1,4-Dioxane and Other Solvent Stabilizers*. Taylor and Francis Group, Boca Raton, FL USA.
- Nam JH, Ventura JRS, Yeom IT, Lee Y & Jahng D 2016 Structural and kinetic characteristics of 1,4-dioxane-degrading bacterial consortia containing the phylum TM7. *Journal of Microbiology and Biotechnology* 26 (11), 1951–1964. 10.4014/jmb.1601.01095. [PubMed: 27470275]
- Parales RE, Adamus JE, White N & May HD 1994 Degradation of 1,4-dioxane by an actinomycete in pure culture. *Applied and Environmental Microbiology* 60 (12), 4527–4530. 10.1128/aem.60.12.4527-4530.1994. [PubMed: 7811088]
- Pugazhendi A, Banu JR, Dhavamani J & Yeom IT 2015 Biodegradation of 1,4-dioxane by *Rhodanobacter* AYS5 and the role of additional substrates. *Annals of Microbiology* 65, 2201–2208. 10.1007/s13213-015-1060-y.
- Ramos-García ÁA, Walecka-Hutchison C & Freedman DL 2022 Effect of biostimulation and bioaugmentation on biodegradation of high concentrations of 1,4-dioxane. *Biodegradation* 33 (2), 157–168. 10.1007/s10532-022-09971-4. [PubMed: 35102492]
- Rittmann BE & McCarty PL 2020 *Environmental Biotechnology: Principles and Applications*, 2nd edn. McGraw-Hill, New York, USA.
- Rolston HM, Hyman MR & Semprini L 2019 Aerobic cometabolism of 1,4-dioxane by isobutane-utilizing microorganisms including *Rhodococcus rhodochromus* strain 21198 in aquifer microcosms: Experimental and modeling study. *Science of the Total Environment* 694, 133688. 10.1016/j.scitotenv.2019.133688. [PubMed: 31756820]
- Sales CM, Grostern A, Parales JV, Parales RE & Alvarez-Cohen L 2013 Oxidation of the cyclic ethers 1,4-dioxane and tetrahydrofuran by a monooxygenase in two *Pseudonocardia* species. *Applied and Environmental Microbiology* 79 (24), 7702–7708. 10.1128/AEM.02418-13. [PubMed: 24096414]
- Sei K, Miyagaki K, Kakinoki T, Fukugasako K, Inoue D & Ike M 2013a Isolation and characterization of bacterial strains that have high ability to degrade 1,4-dioxane as a sole carbon and energy source. *Biodegradation* 24, 665–674. 10.1007/s10532-012-9614-1. [PubMed: 23239086]
- Sei K, Oyama M, Kakinoki T, Inoue D & Ike M 2013b Isolation and characterization of tetrahydrofuran-degrading bacteria for 1,4-dioxane-containing wastewater treatment by cometabolic degradation. *Journal of Water and Environment Technology* 11 (1), 11–19.
- Simmer RA, Richards PM, Ewald JM, Schwarz C, Silva MLB, Mathieu J, Alvarez PJJ & Schnoor JL 2021 Rapid metabolism of 1,4-dioxane to below health advisory levels by thiamine-amended *rhodococcus ruber* strain 219. *Environmental Science & Technology Letters* 8 (11), 975–980. 10.1021/acs.estlett.1c00714.
- Tamura K, Stecher G & Kumar S 2021 MEGA11: molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution* 38 (7), 3022–3027. 10.1093/molbev/msab120. [PubMed: 33892491]
- Tang Y 2023 A review of challenges and opportunities for microbially removing 1,4-dioxane to meet drinking-water and groundwater guidelines. *Current Opinion in Environmental Science & Health* 31, 100419. 10.1016/j.coesh.2022.100419. [PubMed: 36582465]
- Tusher TR, Inoue C & Chien MF 2022 Efficient biodegradation of 1,4-dioxane commingled with additional organic compound: role of interspecies interactions within consortia. *Chemosphere* 308, 136440. 10.1016/j.chemosphere.2022.136440. [PubMed: 36116621]
- Tusher TR, Shimizu T, Inoue C & Chien MF 2020 Enrichment and analysis of stable 1,4-dioxane-degrading microbial consortia consisting of novel dioxane-degraders. *Microorganisms* 8 (50), 1–16. 10.3390/microorganisms8010050.

- Tusher TR, Shimizu T, Inoue C & Chien MF 2021 Isolation and characterization of novel bacteria capable of degrading 1,4-dioxane in the presence of diverse co-occurring compounds. *Microorganisms* 9 (887), 1–14. 10.3390/microorganisms9050887.
- Wang P, Li F, Wang W, Wang R, Yang Y, Cui T, Liu N & Li M 2021 Cometabolic degradation of 1,4-dioxane by a tetrahydrofuran-growing *Arthrobacter* sp. WN18. *Ecotoxicology and Environmental Safety* 217, 112206. 10.1016/j.ecoenv.2021.112206. [PubMed: 33866286]
- Xiong Y, Mason OU, Lowe A, Zhang Z, Zhou C, Chen G, Villalonga MJ & Tang Y 2020 Investigating promising substrates for promoting 1,4-dioxane biodegradation: Effects of ethane and tetrahydrofuran on microbial consortia. *Biodegradation* 31 (3), 171–182. 10.1007/s10532-020-09901-2. [PubMed: 32361902]
- Xiong Y, Mason OU, Lowe A, Zhou C, Chen G & Tang Y 2019a Microbial community analysis provides insights into the effects of tetrahydrofuran on 1,4-dioxane biodegradation. *Applied and Environmental Microbiology* 85 (11), 1–11. 10.1128/AEM.00244-19.
- Xiong Y, Wang B, Zhou C, Chen H, Chen G & Tang Y 2022 Determination of growth kinetics of microorganisms linked with 1,4-dioxane degradation in a consortium based on two improved methods. *Frontiers of Environmental Science & Engineering* 16 (5), 62. 10.1007/s11783-022-1567-y.
- Xiong Y, Zhang Q, Wandell R, Bresch S, Wang H, Locke BR & Tang Y 2019b Synergistic 1,4-dioxane removal by non-thermal plasma followed by biodegradation. *Chemical Engineering Journal* 361, 519–527. 10.1016/j.cej.2018.12.094.
- Yamamoto N, Saito Y, Inoue D, Sei K & Ike M 2018 Characterization of newly isolated *Pseudonocardia* sp. N23 with high 1,4-dioxane-degrading ability. *Journal of Bioscience and Bioengineering* 125 (5), 552–558. 10.1016/j.jbiosc.2017.12.005. [PubMed: 29301721]
- Zenker MJ, Borden RC & Barlaz MA 2002 Modeling cometabolism of cyclic ethers. *Environmental Engineering Science* 19 (4), 215–228. 10.1089/109287502760271535.
- Zenker MJ, Borden RC & Barlaz MA 2003 Occurrence and treatment of 1,4-dioxane in aqueous environments. *Environmental Engineering Science* 20 (5), 423–432. 10.1089/109287503768335913.
- Zhang S, Gedalanga PB & Mahendra S 2017 Advances in bioremediation of 1,4-dioxane-contaminated waters. *Journal of Environmental Management* 204 (2), 765–774. 10.1016/j.jenvman.2017.05.033. [PubMed: 28625566]

**HIGHLIGHTS**


- A low 1,4-dioxane concentration-based enrichment strategy was evaluated.
- A combination of centrifugation and dilution is the best enrichment strategy.
- Five isolated strains have higher net specific growth rates compared to the literature.
- The kinetics of the five isolated strains are suitable for oligotrophic environments.
- *Nitrobacter* sp. WCD6H is the first 1,4-dioxane-metabolizing strain in *Nitrobacter*.






**Figure 1.**

Phylogenetic tree based on partial 16S rRNA gene sequences showing the evolutionary relationship among five 1,4-dioxane-metabolizing bacteria isolated in this study (LCD6B, LCD6D, WC10G, WCD6H, and WD4H) and bacteria reported in the literature that can metabolically degrade 1,4-dioxane. Note: *Nitrobacter* sp. TS12, which cannot degrade 1,4-dioxane (Tusher et al. 2022), is added to the tree to show the evolutionary relationship of WCD6H.




**Figure 2 I.** Comparison of experimental data and model-simulated results for 1,4-dioxane and biomass concentrations in the kinetic experiments used for parameter estimation. The inset graphs show fittings for 1,4-dioxane concentrations below 1 mg/L.




**Figure 3 1.**

Net specific growth rates for 1,4-dioxane-metabolizing pure cultures isolated in this study and reported in the literature. *Note: P. dioxanivorans* CB1190 (Barajas-Rodriguez & Freedman 2018) and *P. dioxanivorans* BERK-1 (Ramos-García et al. 2022) are the only pure cultures with a full set of kinetic parameters reported in the literature.




**Figure 4 l.**

Comparison of kinetic parameters of the 1,4-dioxane-metabolizing pure cultures isolated in this study and 1,4-dioxane-metabolizing pure and mixed cultures in the literature.




**Figure 5 l.**

Comparison of kinetic parameters for *Afipia* and *Pseudonocardia* isolated in this study and reported in the literature. *Note: Afipia* in the literature has no reported biomass decay value.



**Figure 6 I.** Comparison of experimental data and modeling results for 1,4-dioxane and biomass concentrations during model evaluation.



**Figure 7 I.** Comparison of 1,4-dioxane and total COD concentrations for the 1,4-dioxane-metabolizing pure cultures isolated in this study. The 1,4-dioxane concentrations were converted from mg-1,4-dioxane/L to mg-COD/L by a conversion factor of 1.82. Aqueous samples used for COD measurements were filtered by 0.20  $\mu\text{m}$  Acrodisc<sup>®</sup> syringe filters.

**Table 1 |**

Description of model parameters

Parameter	Description	Unit
(a) Model input parameters		
$N$	Total number of experimental measurements	–
$S_{d(t)}^e$	1,4-Dioxane concentration in the experiment at time $t$	mg-dioxane/L
$t$	Time	d
$T$	Time at the end of each experiment	d
$X_{(t)}^e$	Biomass concentration in the experiment at time $t$	mg-protein/L
(b) Model output parameters		
$b$	Biomass decay coefficient	1/d
$K_d$	Half-maximum-rate concentration of 1,4-dioxane	mg-dioxane/L
$MSRR$	Mean sum of squared relative residuals	–
$q_d$	Maximum specific 1,4-dioxane utilization rate	mg-dioxane/(mg-protein-d)
$S_d$ or $S_{d(t)}^m$	1,4-Dioxane concentration at time $t$ of the model	mg-dioxane/L
$S_d^{min}$	Minimum 1,4-dioxane concentration	mg-dioxane/L
$X$ or $X_{(t)}^m$	Biomass concentration at time $t$ of the model	mg-protein/L
$Y_d$	Biomass yield of 1,4-dioxane	mg-protein/(mg-dioxane)
$\mu$	Net specific growth rate of active biomass	1/d



**Table 2 |**

Summary of 1,4-dioxane-metabolizing pure cultures and their corresponding 1,4-dioxane degrading genes

Pure cultures in this study	Pure cultures in the literature	1,4-Dioxane degrading genes	Notes	References
<i>Dokdonella</i> sp. LCD6B	<i>Dokdonella</i> sp. TS32	<i>thmA/dxmA</i>	Tetrahydrofuran/dioxane monoxygenase	Tusher et al. (2021)
<i>Acinetobacter</i> sp. LCD6D	<i>Acinetobacter baumannii</i> DD1	–	No study on 1,4-dioxane degrading genes	Huang et al. (2014)
<i>Afiplia</i> sp. WC10G	<i>Afiplia</i> sp. TS43	<i>thmA/dxmA</i>	Tetrahydrofuran/dioxane monoxygenase	Tusher et al. (2021)
<i>Nitrobacter</i> sp. WCD6H	–	–	No study on 1,4-dioxane degrading genes	–
<i>Pseudonocardia</i> sp. WD4H	<i>Pseudonocardia dioxanivorans</i> CB1190	<i>thmADBC</i>	Tetrahydrofuran monoxygenase	Grosterm et al. (2012)
	<i>Pseudonocardia dioxanivorans</i> JCM 13855	<i>thmA/dxmA</i>	Tetrahydrofuran/dioxane monoxygenase	Inoue et al. (2016)
	<i>Pseudonocardia</i> sp. D17	<i>thmA/dxmA</i>	Tetrahydrofuran/dioxane monoxygenase	Inoue et al. (2016)
	<i>Pseudonocardia</i> sp. TS28	<i>thmA/dxmA</i>	Tetrahydrofuran/dioxane monoxygenase	Tusher et al. (2021)

**Table 3 |**

Summary of kinetic parameters and minimum 1,4-dioxane concentrations required for the 1,4-dioxane-metabolizing strains isolated in this study

Pure culture	$q_d \left( \frac{\text{mg dioxane}}{\text{mg protein} \cdot \text{d}} \right)$	$K_d \left( \frac{\text{mg dioxane}}{\text{L}} \right)$	$\gamma_d \left( \frac{\text{mg protein}}{\text{mg dioxane}} \right)$	$b \left( \frac{1}{\text{d}} \right)$	$S_d^{min} \left( \frac{\text{mg dioxane}}{\text{L}} \right)$	MSRR
<i>Dokdonella</i> sp. LCD6B	0.24	7.7	0.34	0.01	1.5	0.04
<i>Acinetobacter</i> sp. LCD6D	0.27	5.6	0.38	0.02	1.3	0.02
<i>Afiplia</i> sp. WC10G	0.47	8.2	0.36	0.02	1.0	0.11
<i>Nitrobacter</i> sp. WCD6H	0.24	1.8	0.30	0.01	0.4	0.08
<i>Pseudonocardia</i> sp. WD4H	0.26	5.3	0.29	0.01	0.8	0.01

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript