

Plasma Serotonin in Laying Hens (*Gallus gallus domesticus*) With and Without Foot pad Dermatitis

Daniela Alberghina , Vito Biondi, Annamaria Passantino, Fabiola Giunta and Michele Panzera

Department of Veterinary Science, University of Messina, Messina, Italy.

International Journal of Tryptophan Research
Volume 13: 1–3
© The Author(s) 2020
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1178646920927380



ABSTRACT: The aim of this study was to investigate whether plasma serotonin (5-hydroxytryptamine [5-HT]) was associated with the presence of foot pad dermatitis (FPD) in laying hens. FPD birds ($n=20$) and healthy individuals ($n=22$) were included. Plasma 5-HT was investigated. FPD laying hens showed significantly higher 5-HT levels (median = $6\mu\text{mol/L}$) compared with healthy individuals (median = $4.28\mu\text{mol/L}$, $P < .001$). When present, FPD were scored as either 1 ($n=12$) indicating mildly to moderately abnormal or 2 indicating severely abnormal ($n=8$). The subjects whose lesions scored 2 had higher plasma 5-HT levels than those whose lesions scored 1. Inflammatory mechanisms seem to be related to plasma 5-HT levels in laying hens. Assessing plasma 5-HT could be useful to evaluate chicken welfare.

KEYWORDS: Plasma serotonin, laying hens, animal welfare, inflammation

RECEIVED: January 20, 2020. **ACCEPTED:** April 20, 2020.

TYPE: Short Report

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by Funds for Financing Basic Research Activities (FFABR 2017).

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Daniela Alberghina, Department of Veterinary Science, University of Messina, Polo Universitario dell'Annunziata, 98168, Messina, Italy. Email: dalberghina@unime.it

Serotonin (5-hydroxytryptamine [5-HT]) is a neuromodulator involved in many physiological functions. 5-HT plays a modulatory role in temperature regulation, feeding, sexual behavior, painful stimuli response, flight, and stress.¹ In birds as in mammals, 5-HT is produced not only in the neuronal system but also independently in peripheral tissues.² Surprisingly only about 1% to 2% of the total amount of serotonin is produced by serotonergic neurons in the brain, whereas close to 95% of the 5-HT in the body is synthesized, stored, and released by cells in the intestinal mucosa known as enterochromaffin cells.³ 5-HT can be detected in several non-neuronal tissues of the cardiovascular and renal systems, as well as in blood.⁴ Mammalian enterochromaffin cells release 5-HT into the blood, where platelets have evolved a highly efficient 5-HT uptake and transport system.⁵ Approximately 95% of 5-HT in blood is carried in platelets.⁶ Similar to mammals' platelets, avian thrombocytes contain serotonin.⁷ In chickens, whole-blood 5-HT is mainly distributed among the following cell types: thrombocytes (50%), lymphocytes (25%–29%), and polymorphs and monocytes (19%–25%).⁸ Lymphocytes appear to possess the ability to synthesize, transport, store, and/or respond to 5-HT.⁹ Circulating 5-HT is considered an accessible model for the investigation of the central serotonin system.¹⁰ In particular, the similarity of platelets and central serotonergic synaptosomes indicates that the first ones can represent a non-invasive model to study modulation of 5-HT (release and/or uptake) in neurons.¹¹ In homeotherms, such as mammals and birds, circulating 5-HT plays a major role in reducing heat loss when the environmental temperature is lower than the core temperature.¹² The amino acid tryptophan (TRP) is used for the synthesis of central and peripheral 5-HT.¹³ Both central and peripheral serotonin levels can be influenced by TRP availability and enzymatic activity of tryptophan hydroxylase

(TPH). Stress factors, such as social stress, can influence peripheral 5-HT in chickens. In young chickens, exposure to elevated corticosterone during incubation resulted in lower whole-blood 5-HT and higher platelet 5-HT uptake as compared with controls.¹⁴


A deficient serotonergic system, potentially due to low TRP availability and/or reduced TPH activity, was found to be associated with psychopathological behaviors in various species, among which feather pecking has been described in chickens.¹⁵ Feather pecking is a repetitive oral behavior in which individuals peck repetitively at another bird's feather cover. Severe feather pecking influenced hens' plasma TRP.¹⁶ In the peripheral system, the biological roles of 5-HT in behavioral adaptation and motivational regulation are unclear.¹⁷ In the genetic selection for low productivity group hens, higher blood 5-HT levels were associated with lower survivability resulting from higher cannibalism.¹⁸ Among various physiological functions of peripheral 5-HT, it is becoming increasingly more evident that it plays an important role in many different contexts, including inflammation and immunity.¹⁹ Although footpad dermatitis (FPD) has gained considerable attention in broilers, little attention has been paid so far to FPD in laying hens.²⁰ As FPD are painful for the hens, they can have negative effects on their welfare. If the lesions become ulcerated, they can affect standing and walking ability. Therefore, the aim of this study was to conduct a pilot study to investigate the concentration of plasma 5-HT in laying hens and to evaluate the possible relation between 5-HT levels and FPD.

Materials and Methods

Subjects

Siciliana breed laying hens ($n=42$), kept in non-cage system at a commercial farm, were kept under standardized conditions.





At week 45, a general health check was carried out. Individuals without FPD ($n=22$) and with FPD ($n=20$) were included in the study. Lesion scoring was performed using the photographic system described in Welfare Quality Assessment (WQ) protocol for poultry.²¹ Individuals without FPD were assigned to group A. When present, FPD were scored as either 1, indicating mildly to moderately abnormal ($n=12$), or 2, indicating severely abnormal ($n=8$), and the subjects were assigned to groups B and C respectively. The birds were taken out of their home cage individually and, within 2 minutes, a blood sample (1.5 mL) was collected from the wing vein. The samples were stored into EDTA tubes and centrifuged (1.785g for 10 minutes). Protocols of animal husbandry and experimentation were reviewed and approved in accordance with the standards recommended by the Guide for the Care and Use of Laboratory Animals and Directive 2010/63/EU.

Plasma serotonin assessment

The plasma aliquots obtained from each sample were transported to the laboratory in a cooler with an ice block within 1 hour of being drawn. Plasma for 5-HT analysis was stored frozen at -20°C . Samples were analyzed within 1 month of collection. As 5-HT occurs at extremely high levels in birds,²² plasma was diluted in a 1:10 ratio with distilled water before 5-HT measurement. A commercially available ELISA kit (analytic range 10.2–2500 ng/mL, 5-HT ELISA^{Fast Track}, LDN GmbH & Co. KG) was used to measure plasma 5-HT levels, following the manufacturer's instructions as previously described.²³ Intra- and inter-coefficient of variation (CV) were 5.6% and 9%, respectively. The amount of 5-HT in the sample was estimated with the calibration curve, multiplied by 10 and expressed in $\mu\text{mol/L}$.




Figure 1. Plasma 5-HT levels ($\mu\text{mol/L}$) in 3 experimental groups of laying hens based on foot-pad dermatitis. 5-HT indicates 5-hydroxytryptamine.

Statistical analysis

The Mann-Whitney test was used to compare 5-HT differences between individuals with FPD and healthy individuals. The significance of the differences between the 3 groups was tested with the 1-way analysis of variance Kruskal-Wallis test on ranks with Dunn's multiple comparison post hoc test. A probability level of $P < .05$ was considered significant. Data were analyzed using the software STATISTICA 7.5 (StatSoft Inc., Tulsa, OK, USA).

Results

Plasma 5-HT differences between healthy individuals and birds with FPD were significant ($P < .01$; 4.24 ± 0.41 and $6.88 \pm 0.59 \mu\text{mol/L}$ respectively). The comparison among the groups showed significant differences (Kruskal-Wallis, $P < .01$). Group C had higher 5-HT plasma levels compared with Group A ($P < .01$; Figure 1) when all groups were compared.

Discussion

The aim of this study was to evaluate whether plasma 5-HT in laying hens could be influenced by a chronic inflammatory condition such as FPD. Serotonin in chicken has been studied in plasma²⁷⁻²⁹ and in whole blood.²⁴⁻²⁶ The use of the peripheral 5-HT system as a less invasive model for central 5-HT system activity has been proposed after discovering the correlation between whole-blood 5-HT and brain 5-HT in laying hens.¹⁸ Unfortunately 5-HT cannot be analyzed in whole blood by ELISA. As indicated in the manufacturer's description, this kit is used for the quantitative ultra-sensitive determination of 5-HT in any species and on various biological samples (serum, plasma, urine). The present study shows that high plasma 5-HT concentrations are associated with FPD in laying hens. Similar findings were encountered in horses, where subjects with bowel disorders showed higher plasma 5-HT than healthy horses.³⁰

We hypothesized that it would be more reliable to assess an inflammatory response by measuring 5-HT in plasma than in whole blood because the massive 5-HT release from platelets/thrombocytes and lymphocytes into systemic circulation concurs to generate inflammation.

Assessing plasma 5-HT could be useful to evaluate chicken welfare. Further studies and the standardization of the ELISA method are likely to guide future investigations.

Acknowledgments

We would like to thank Dr. Bruno Scalia for his advice and help in English correction and Dr. Concetta Saoca and sig. Giuseppe Bella for their help in the preparation of the manuscript.

Author Contributions

Conceived and designed the experiments: DA, AP, MP.

Performed the experiments: VB, FG.

Analyzed the data: DA.

Wrote the paper: DA.

ORCID iD

Daniela Alberghina  <https://orcid.org/0000-0003-2826-0325>

REFERENCES

- Lucki I. The spectrum of behaviors influenced by serotonin. *Biol Psychiatry*. 1998;44:151-162.
- Walther DJ, Peter JU, Winter S, et al. Serotonylation of small GTPases is a signal transduction pathway that triggers platelet α -granule release. *Cell*. 2003;115:851-862.
- Gershon MD, Tack J. The serotonin signaling system: from basic understanding to drug development for functional GI disorders. *Gastroenterology*. 2007;132:397-414.
- Hoyer D, Hannon JP, Martin GR. Molecular, pharmacological and functional diversity of 5-HT receptors. *Pharmacol Biochem Behav*. 2002;71:533-554.
- Stahl SM, Meltzer HY. A kinetic and pharmacologic analysis of 5-hydroxytryptamine transport by human platelets and platelet storage granules: comparison with central serotonergic neurons. *J Pharmacol Exp Ther*. 1978;205:118-132.
- Anderson GM, Feibel FC, Cohen DJ. Determination of serotonin in whole blood, platelet-rich plasma, platelet-poor plasma and plasma ultrafiltrate. *Life Sci*. 1987;40:1063-1070.
- Stiller RA, Belamarich FA, Shepro D. Aggregation and release in thrombocytes of the duck. *Am J Physiol*. 1975;229:206-210.
- Meyer MJ, Sturkie PD. Diurnal rhythm of histamine in blood and oviduct of the domestic fowl. *Comp Gen Pharmacol*. 1974;5:225-228.
- Chen Y, Leon-Ponte M, Pingle SC, O'Connell PJ, Ahern GP. T lymphocytes possess the machinery for 5-HT synthesis, storage, degradation and release. *Acta Physiol (Oxford)*. 2015;213:860-867.
- Karakulova UV, Selyanina NV, Sumlivaya ON, Vorobyeva NN, Okishev MA. Involvement of peripheral unit of serotonin energetic system in brain lesions of different types. *World Appl Sci*. 2013;26:374-376.
- Da Prada M, Cesura AM, Launay JM, Richards JG. Platelets as a model for neurons? *Experientia*. 1988;44:115-126.
- Maurer-Spurej E. Circulating serotonin in vertebrates. *Cell Mol Life Sci*. 2005;62:1881-1889.
- Coates MD, Mahoney CR, Linden DR, et al. Molecular defects in mucosal serotonin content and decreased serotonin reuptake transporter in ulcerative colitis and IBS. *Gastroenterology*. 2004;126:1657-1664.
- Ahmed AA, Ma W, Ni Y, Zhou Q, Zhao R. Embryonic exposure to corticosterone modifies aggressive behavior through alterations of the hypothalamic pituitary adrenal axis and the serotonergic system in the chicken. *Horm Behav*. 2014;65:97-105.
- de Haas EN, van der Eijk JAJ. Where in the serotonergic system does it go wrong? Unravelling the route by which the serotonergic system affects feather pecking in chickens. *Neurosci Biobehav Rev*. 2018;95:170-188.
- Birkel P, Kjaer JB, Szkotnicki W, Forsythe P, Harlander-Matauschek A. Acute tryptophan depletion: the first method validation in an avian species (*Gallus gallus domesticus*). *Poult Sci*. 2017;96:3021-3025.
- Cheng HW, Dillworth G, Singleton P, Chen Y, Muir WM. Effects of group selection for productivity and longevity on blood concentrations of serotonin, catecholamines, and corticosterone of laying hens. *Poult Sci*. 2001;80:1278-1285.
- Buitenhuis AJ, Rodenburg TB, Wissink PH, et al. Genetic and phenotypic correlations between feather pecking behavior, stress response, immune response, and egg quality traits in laying hens. *Poult Sci*. 2004;83:1077-1082.
- Herr N, Bode C, Duerschmied D. The effects of serotonin in immune cells. *Front Cardiovasc Med*. 2017;4:48.
- Niebuhr K, Arhant C, Lugmair A, Gruber B, Zaludik K. *Foot Pad Dermatitis in Laying Hens Kept in Non-Cage Systems in Austria*. Proceedings of 8th Poultry Welfare Symposium, Cervia, Italy, 18-22 May 2009, p8.
- Welfare Quality®. *Welfare Quality® Assessment Protocol for Poultry (Broilers, Laying Hens)*. Lelystad, The Netherlands: Welfare Quality® Consortium; 2009.
- Meyer D. *Studies on factors regulating blood serotonin levels in the domestic fowl* [PhD thesis]. New Brunswick, NJ: Rutgers University; 1973.
- Alberghina D, Tropa E, Piccione G, Giannetto C, Panzera M. Serum serotonin (5-HT) in dogs (*Canis familiaris*): preanalytical factors and analytical procedure for use of reference values in behavioral medicine. *J Vet Behav* 2019;32:72-75.
- Bolhuis JE, Ellen ED, Van Reenen CG, et al. Effects of genetic group selection against mortality on behavior and peripheral serotonin in domestic laying hens with trimmed and intact beaks. *Physiol Behav*. 2009;97:470-475.
- Rodenburg TB, Bolhuis JE, Koopmanschap RE, Ellen ED, Decuyper E. Maternal care and selection for low mortality affect post-stress corticosterone and peripheral serotonin in laying hens. *Physiol Behav*. 2009;98:519-523.
- Uitdehaag KA, Rodenburg TB, Van Reenen CG, et al. Effects of genetic origin and social environment on behavioral response to manual restraint and monoamine functioning in laying hens. *Poult Sci*. 2011;90:1629-1636.
- Buitenhuis AJ, Kjaer JB. Long term selection for reduced or increased pecking behaviour in laying hens. *Worlds Poult Sci J*. 2008;64:477-487.
- Agnvall B, Katajamaa R, Altimiras J, Jensen P. Is domestication driven by reduced fear of humans? Boldness, metabolism and serotonin levels in divergently selected red jungle fowl (*Gallus gallus*). *Biol Lett*. 2015;11:6-9.
- Buitenhuis AJ, Kjaer JB, Labouriau R, Juul-Madsen HR. Altered circulating levels of serotonin and immunological changes in laying hens divergently selected for feather pecking behaviour. *Poult Sci*. 2006;85:1722-1728.
- Delesalle C, van de Walle GR, Nolten C, et al. Determination of the source of increased serotonin (5-HT) concentrations in blood and peritoneal fluid of colic horses with compromised bowel. *Equine Vet J*. 2008;40:326-331.