

RESEARCH ARTICLE

Open Access



High mobility group box 1 promotes sorafenib resistance in HepG2 cells and in vivo

Yinzong Xiao¹, Lunquan Sun², Yongming Fu¹, Yan Huang¹, Rongrong Zhou¹, Xingwang Hu¹, Pengcheng Zhou¹, Jun Quan¹, Ning Li^{3*} and Xue-Gong Fan^{1*}

Abstract

Background: Primary liver cancer is a lethal malignancy with a high mortality worldwide. Currently, sorafenib is the most effective molecular-targeted drug against hepatocellular carcinoma (HCC). However, the sorafenib resistance rate is high. The molecular mechanism of this resistance has not been fully elucidated. High mobility group box 1 (HMGB1) is a multifaceted protein that plays a key role in the proliferation, apoptosis, metastasis and angiogenesis of HCC cells. In addition, HMGB1 has been suggested to contribute to chemotherapy resistance in tumours, including lung cancer, osteosarcoma, neuroblastoma, leukaemia, and colorectal cancer. This study investigated the association between HMGB1 and sorafenib resistance in HCC.

Methods: HepG2 cells with HMGB1 knockdown or overexpression were generated. The efficacy of sorafenib in these cells was tested using flow cytometry and a cell counting assay. The subcellular localization of HMGB1 in HepG2 cells following sorafenib treatment was measured by western blotting and confocal microscopy. A murine subcutaneous HCC model was generated to examine the association between HMGB1 and the sensitivity of sorafenib treatment.

Results: The HMGB1 knockdown cells exhibited a significantly higher apoptotic level and lower cell viability than the normal HMGB1 expressing cells following the sorafenib treatment. In addition, the cell viability observed in the HMGB1 overexpressing cells was higher than that observed in the control cells following the sorafenib intervention. Sorafenib had a better tumour inhibition effect in the HMGB1 knockdown group in vivo. The amount of mitochondrial HMGB1 decreased, while the amount of cytosolic HMGB1 increased following the exposure to sorafenib. Altogether, HMGB1 translocated from the mitochondria to the cytoplasm outside the mitochondria following the exposure of HepG2 cells to sorafenib.

Conclusions: A novel potential role of HMGB1 in the regulation of sorafenib therapy resistance in HCC was observed. The knockdown of HMGB1 restores sensitivity to sorafenib and enhances HepG2 cell death, while HMGB1 overexpression blunts these effects. The translocation of HMGB1 from the mitochondria to the cytosol following sorafenib treatment provides new insight into sorafenib resistance in HCC.

Keywords: HCC, HMGB1, Sorafenib resistance, Mitochondria

* Correspondence: nxli@hotmail.com; xgfan@hotmail.com

³Department of Blood Transfusion, Xiangya Hospital, Central South University, Changsha 410008, China

¹Hunan Key Laboratory of Viral Hepatitis, Department of Infectious Diseases, Xiangya Hospital, Central South University, Changsha 410008, China

Full list of author information is available at the end of the article



Background

Primary carcinoma of the liver is the second most common cause of death from cancer worldwide. Hepatocellular carcinoma (HCC) accounts for 90% of all liver cancers. HCC has a very poor prognosis, and the overall ratio of mortality to incidence is 0.95 globally [1]. Less than 30% of newly diagnosed HCC patients without surveillance are eligible for curative treatments, such as resection, transplantation, or ablation [2]. In patients with advanced HCC, sorafenib (Nexavar), which is a molecular-targeted therapy, significantly helps prolong the median survival time by approximately 3 months. Sorafenib inhibits B-RAF, vascular endothelial growth factor receptor (VEGFR), and platelet-derived growth factor receptor (PDGFR) [3]. However, only approximately 30% of advanced HCC patients benefit from sorafenib, and acquired resistance often develops within 6 months [4]. The extremely high sorafenib resistance rate has raised great concern worldwide, and the epithelial-mesenchymal transition (EMT), cancer stem cells, and tumour microenvironment may be involved [5]. However, the mechanisms underlying primary and acquired sorafenib resistance in HCC remain unclear. Currently, no other chemotherapeutic agent yields the results obtained with sorafenib; thus, understanding and overcoming sorafenib chemoresistance is critical for improving survival in advanced HCC populations [6].

The high mobility group box 1 (HMGB1) protein has been shown to play pivotal roles in HCC, including tumorigenesis, progression, invasion, metastasis, and prognosis. The mechanisms involved in the context-dependent role of HMGB1 include the regulation of cell proliferation, differentiation, cell death, inflammation and immune function in HCC [7–10].

HMGB1 is a nuclear protein that plays a role in various biological events in the nucleus, including DNA replication, repair, recombination, transcription, and genomic stability [11]. In addition to its significant nuclear role, extracellular HMGB1 is one of the most common damage-associated molecular patterns (DAMPs) with well-defined interactions with the receptor for advanced glycation end products (RAGEs) and Toll-like receptors (TLRs). The binding of HMGB1 to RAGE and TLRs affects HCC invasion, metastasis, and treatment [12, 13]. HMGB1 has also been observed in the cytosol, including the mitochondria [14], but its function in the cytoplasm remains poorly understood. Cytosolic HMGB1 may be involved in different types of cell death, and has been found to be a positive regulator of autophagy through its binding to Beclin-1 [15].

HMGB1 affects tumour growth, metastasis, and prognosis through multiple signalling pathways in cancers, including lung cancer, osteosarcoma, gastric cancer, cervical cancer, and HCC [16–19]. The mechanisms by

which HMGB1 affects these biological processes vary in different malignancies. In addition, the role of HMGB1 in drug resistance in tumours, including lung cancer, osteosarcoma, neuroblastoma, leukaemia, and colorectal cancer, has attracted broad attention [20–23]. Due to its versatile role in cancer, HMGB1 has been proposed as a potential biomarker of survival and a target for cancer therapy [18, 24].

Although the role of HMGB1 in drug resistance in different malignancies has been frequently discussed, its function in anticancer therapy in HCC is not fully understood. Because HMGB1 affects multiple biological events in HCC and contributes to drug resistance in certain cancers [25], this study aims to explore the role of HMGB1 in sorafenib resistance in HCC using the HepG2 cell line and murine models.

Methods

Cell lines

The human hepatocarcinoma cell line HepG2 was purchased from the Cell Bank of Shanghai Academy of Science (Shanghai, China, No. TCHu 72). The cells were cultured in Dulbecco's modified eagle medium (DMEM, Thermo Fisher Scientific Inc., USA) supplemented with 10% foetal bovine serum (FBS, Thermo Fisher Scientific Inc., USA), 100 U/mL penicillin, and 100 mg/mL streptomycin (Sangon, Shanghai, China). All cells were cultured at 37 °C in a humidified incubator with 5% CO₂ concentration.

Establishment of stable HMGB1-overexpressing cells and HMGB1 knockdown cells

shRNAs based on the HMGB1 sequence (NM_001313893.1) were designed as follows: shhmgbl, 5'-GCCCGTTATGAAAGAGAAATTTTTC AAGAGAAAA TTTCTCTTTCATAACGGGTTTTTT-3'. The shHMGB1 shRNA sequences were cloned into PLVX-sh1 vectors (Clontech, Mountain View, CA, USA). Lentiviral vectors encoding the human HMGB1 gene were constructed in the overexpression vector (PLVX-ires-puro) (Clontech) and designated LV-HMGB1. The lentiviral vectors were transfected into HCC cells at a multiplicity of infection (MOI) of 30 to 50 in the presence of polybrene (8 µg/mL). At 72 h after the infection, the infected cells were treated with 1 µg/mL puromycin (Origene) for 2 weeks; then, the HMGB1 expression was detected using western blotting. Pooled populations of knockdown and overexpression cells were obtained 2 weeks after the drug selection without sub-cloning and used for the in vivo experiments.

Reagents

Sorafenib was purchased from Santa Cruz Biotechnology, Inc. (Texas, US No. Sc-220,125). Sorafenib was dissolved

in DMSO (Sigma-Aldrich) and diluted with DMEM or normal saline (NS) to the desired concentration for the in vitro and in vivo studies, respectively. The concentration of the mother liquid was 10 mM/L according to the manufacturer's instructions. The doses chosen for the in vitro (40 mg/kg/d) and in vivo studies were based on those used in other published studies [26–29].

Real-time polymerase chain reaction (RT-PCR)

The total mRNA expression was quantitatively analysed using the SYBR Green fluorescent-based assay (TaKaRa Bio, Otsu, Japan). The RT-PCR assays were performed using glyceraldehyde 3-phosphate dehydrogenase (GAPDH) as an internal control. Independent experiments were repeated three times for each sample, and the relative gene expression levels were analysed using the $2^{-\Delta\Delta CT}$ method. The primers, which were designed by Sangon Biotech (Shanghai), were as follows: For HMGB1: forward 5'-TATGGCAAAGCGGACAAGG-3' and reverse 5'-CTTCGCAACATCACCAATGGA-3'. For GAPDH: forward 5'-GAGAGGGAAATCGTGCGTGAC-3' and reverse 5'-CATCTGCTGGAAGGTGGACA-3'.

Cell viability assay

The cell survival rates and dose-dependent curves of sorafenib were evaluated using the Cell Counting Kit-8 (CCK-8 Kit, Beyotime Biotechnology, China, No. C0038). The cells were plated at a density of 5000 cells per well in 96-well flat-bottomed plates and incubated at 37 °C in a humidified atmosphere of 5% CO₂ for 24 h. The medium was replaced with various concentrations of sorafenib in DMEM and 10% FBS, and the cells were incubated for an additional 24–48 h. The supernatants were removed from the wells, and 10 μL/well of CCK-8 dye were added along with 200 μL/well of fresh DMEM. Absorbance was determined at 450 nm using a plate reader.

Western blotting

The total protein was harvested and mitochondrial proteins were separated using the Mitochondria Extraction Kit (BioVision, NO. K256–25, USA). Nuclear protein was extracted using the Nucleoprotein Extraction Kit (Sangon, NO. C500009, China). For the HepG2 cells, 10 μg protein were loaded to test the whole lysate, while 30 μg protein were loaded to test the cell fractions. For the transfection of the HepG2 cells, 30 μg protein were loaded to examine the level of c-PARP, and 20 μg protein were loaded to examine the level of HMGB1. Following the SDS-PAGE, the proteins were transferred onto PVDF membranes (Sigma-Aldrich), hybridized with specific primary antibodies, and incubated with HRP-conjugated sheep anti-mouse/rabbit IgG (Cell Signaling, USA, 1:2000). The bands were

visualized using an ECL Kit (Merck Millipore, Darmstadt, Germany) per the manufacturer's instruction. HMGB1 (Abcam, USA, 1:1000), c-PARP (Cell Signaling, USA, 1:1000), GAPDH (Cell Signaling, USA, 1:1000), and Actin (Cell Signaling, USA, 1:1000) antibodies were used.

Fluorescence-activated cell sorting (FACS) analysis

Annexin V-PI staining (KeyGen Biotech, China) was performed to measure the phosphatidylserine externalization in HepG2 cells transfected with different siRNAs or plvx vectors for the cell death detection. Briefly, trypsinized cells were collected, washed twice with ice-cold PBS, and resuspended in 200 μL binding buffer containing 5 μL FITC-conjugated Annexin V and 5 μL of propidium iodide. The staining sample was incubated at room temperature for 20 min, and 10,000 cells were immediately analysed using a FACSCalibur flow cytometer (BD Biosciences, USA).

Immunofluorescence analysis and confocal microscopy

The cells were treated with equivalent amounts of sorafenib or DMSO for 24 h. Subsequently, the cells were fixed in 4% paraformaldehyde for 10 min and treated with 100% methanol for membrane permeabilization. The slides were blocked with 2% BSA (Sangon, China) at 37 °C for 30 min, fixed in 4% paraformaldehyde for 10 min, treated with antibodies diluted in 1% BSA overnight at 4 °C, and counterstained with DAPI. The immunofluorescence images were analysed under an Upright Metallurgical Microscope (Leica, Biberach, Germany) using the algorithm settings. The colocalization images were analysed by confocal laser scanning microscopy (CLSM, Leica, Germany). The positive rate (the number of positively stained cells/total cells) was then calculated.

In vivo experiments

Athymic BALB/c nu/nu mice (4–6 weeks old) were purchased from SLAC Laboratory Animal Center (Changsha, China). The mice were weighed and randomly divided into eight groups ($n = 5$). To generate murine subcutaneous tumours, 0.5×10^6 HepG2 cells transfected with control or HMGB1-specific shRNA were injected subcutaneously into the right armpit of the mice. For continuous stimulation in vivo, the mice were maintained under 12-h light/dark cycles with a regular diet of sorafenib (40 mg/kg body weight). The mice were treated for 21 days, and the tumour volumes were measured twice weekly. The volumes were calculated using the formula $V = 1/2a^2b$, where a and b represent the shortest and longest diameters of the tumour, respectively. The mice were weighed twice weekly and sacrificed if they lost >20% of their body weight or

appeared moribund or the tumour reached 1300 to 1500 mm³. The animals were euthanized 21 days post intervention, and the tumour tissues were harvested by resection and fixed in 10% formaldehyde for the immunohistochemistry or stored. The Animal Use Committee of Xiangya Hospital approved all animal treatment protocols, and all mice were treated humanely during the entire study period.

Haematoxylin and eosin (HE) and IHC staining

After embedding in paraffin, the samples were cut into 10- μ m slices and stained with HE. Some samples were cut into 4- μ m slices, incubated with the HMGB1 antibody (Cell Signaling, 1:400), and subsequently incubated with a secondary antibody conjugated with streptavidin–biotin–peroxidase complex (Zhongshan Goldenbridge Biotechnology, Beijing, China). The colour reaction was developed using 3,3-diaminobenzidine tetrahydrochloride (Sigma-Aldrich).

Statistical analysis

The quantitative data are presented as the means \pm s.d of at least three independent experiments. The statistical analyses were performed using SPSS 19.0 software, and Student's t-test was applied to compare the significant differences between two groups. All tests were two-sided, and differences were considered statistically significant at a *p*-value <0.05.

Results

HMGB1 is upregulated in response to sorafenib treatment in HepG2 cells

The effects of sorafenib on the expression of HMGB1 were explored in the HCC cell line HepG2. The HMGB1 mRNA and protein levels were increased following a sorafenib treatment in the clinically relevant range (Fig. 1).

Suppression of HMGB1 increases the sensitivity to sorafenib in HepG2 cells

Sh-HMGB1 HepG2 and sh-control HepG2 cells were generated by transfection with HMGB1 shRNA and a control vector, respectively. The HMGB1 expression was significantly lower in the sh-HMGB1 HepG2 cells than in the control cells (Fig. 2A). According to the cell viability tests, the sh-HMGB1 HepG2 cells treated with sorafenib tended to be more sensitive to the drug than the sh-control cells (Fig. 2B). The apoptotic cells were quantified by determining the percentage of Annexin-V-positive cells 48 h following the sorafenib (5 μ M) treatment using flow cytometry. Significantly higher levels of apoptosis were detected in the sh-HMGB1 cells than in the sh-control cells (Fig. 2C). Consistently, the expression of the apoptosis-related protein c-PARP increased following the sorafenib treatment in the HepG2 HMGB1 knock-down cells (Fig. 2D). Altogether, HMGB1 played a protective role in HepG2 cells following exposure to sorafenib.

Overexpression of HMGB1 increases resistance to sorafenib in HepG2 cells

To further explore the role of HMGB1 in the regulation of sorafenib sensitivity, we overexpressed HMGB1 in HepG2 cells. HMGB1 overexpression vectors were transfected into HepG2 cells, resulting in a significant increase in HMGB1 protein and mRNA in the resultant Plvx-HMGB1 HepG2 cell line (Fig. 3A). Furthermore, according to the cell viability assays, the Plvx-HMGB1 HepG2 cells were less sensitive to the sorafenib treatment than the control cells (Fig. 3B). According to the flow cytometry analysis, compared to the HMGB1 overexpression group, the Plvx-Control group showed a slight but insignificant increase in apoptosis following the sorafenib intervention (Fig. 3C). Compared with the Plvx-HMGB1 HepG2 group, the Plvx-control HepG2 group exhibited a greater increase in the c-PARP levels,




Fig. 1 HMGB1 is upregulated in HepG2 cells following sorafenib treatment (**p* < 0.05). HepG2 cells were treated with 5 μ M sorafenib (Sor) or DMSO, and the HMGB1 expression was measured by (a) qPCR and (b) western blotting




Fig. 2 Suppression of HMGB1 increases the sensitivity to sorafenib in HepG2 cells (* $p < 0.05$; ** $p < 0.01$). **Ai** qPCR and **Aii** western blotting results show that the expression of HMGB1 in HepG2 cells transfected with HMGB1 shRNA (Plvx-sh HMGB1) is significantly lower than that in sh-control HepG2 cells (Plvx-sh Control). **B** The impact of the HMGB1 knockdown on drug sensitivity at different sorafenib doses as determined by a CCK-8 assay. Plvx-sh HMGB1 cells showed an increased sensitivity to sorafenib at 5 μ M. **C** Plvx-sh HMGB1 knockdown and Plvx-sh Control HepG2 cells were treated with 5 μ M sorafenib (treatment) or the same volume of DMSO (control) for 48 h. Annexin V positive cells, consisting of both early and late apoptotic cells, were counted. **D** Expression levels of the apoptosis-related protein c-PARP were examined by western blotting

indicating that the HMGB1 overexpression could reduce sorafenib-induced apoptosis (Fig. 3D).

HMGB1 translocates from the mitochondria to the cytoplasm following sorafenib treatment

To further examine the relationship between HMGB1 and sorafenib resistance, the HMGB1 distribution in HepG2 cells exposed to sorafenib was examined. The total nuclear protein was separated from the cytoplasmic protein, and no significant change in the HMGB1 expression was observed in either fraction following the sorafenib treatment (Fig. 4a). The amount of mitochondrial HMGB1 in the HepG2 cells decreased following the sorafenib exposure, while the amount of cytosolic HMGB1 increased (Fig. 4b). Therefore, HMGB1 translocated from the mitochondria to the cytosol in the HepG2 cells following the sorafenib treatment. In the HepG2 cells, the Tomm20-stained mitochondria and

HMGB1 co-localized (Fig. 4c). The co-localization of the Tomm20-stained mitochondria and HMGB1 significantly decreased following a 48-h sorafenib treatment (Fig. 4d).

Suppression of HMGB1 increases the sensitivity to sorafenib in vivo

To examine the effects of HMGB1 in sorafenib resistance in vivo, BALB/c nu/nu mice were inoculated with HepG2 cells transfected with HMGB1-specific shRNA or control shRNA. Three weeks after the subcutaneous tumour implantation, the tumours reached 100 to 200 mm³ in size, and the athymic mice were treated with 40 mg/kg sorafenib via a p.o. gavage daily. The average tumour volume in the group of mice injected with sh-HMGB1 HepG2 was smaller than that in the group of mice injected with the sh-control HepG2 cells at all time points. Therefore, the




Fig. 3 Overexpression of HMGB1 increases the resistance to sorafenib in HepG2 cells. (* $p < 0.05$; ** $p < 0.01$). **Ai** qPCR and **Aii** western blot analyses show the overexpression of HMGB1 in Plvx-HMGB1 cells compared with that in Plvx-control HepG2 cells. **B** HMGB1 overexpression affects drug sensitivity at different sorafenib doses as determined by a CCK-8 assay. **C** Plvx-HMGB1 cells and Plvx-control HepG2 cells were treated with 5 μ M sorafenib (treatment) or DMSO at same volume (control) for 48 h. The percentage of Annexin V positive cells was measured as an indicator of early and late apoptotic cells. **D** c-PARP was examined to measure the apoptosis levels in HMGB1-overexpressing HepG2 cells and Plvx-Control HepG2 cells

tumour inhibition effect of sorafenib was better in the sh-HMGB1 group (Fig. 5a). The tumours were detached from the mice in each group at the end of the experiment (Fig. 5b). HE staining was performed to examine the tumour, revealing that the abnormal cells were distributed in small nests or cord-like shapes, and some necrotic cells could be observed (Fig. 5c). The HMGB1 immunohistochemical staining (IHC) of the tumour specimens was performed to determine the HMGB1 expression levels in the Plvx-sh HMGB1 and Plvx-HMGB1 groups in vivo (Fig. 5d). Altogether, HMGB1 plays a critical role in the regulation of sorafenib resistance in HepG2 cells in vivo.

Discussion

Sorafenib is a multi-kinase inhibitor with the anticancer effects of suppressing tumour proliferation and angiogenesis and inducing apoptosis. Sorafenib suppresses tumour proliferation by inhibiting the mitogen-activated protein kinase (MAPK) family members Raf-1 and B-Raf. In addition, sorafenib acts by blocking receptors, including VEGFR and PDGFR, leading to the inhibition of angiogenesis [30]. Multiple signalling pathways may contribute to the acquisition of sorafenib resistance. Notably, the epithelial-mesenchymal transition and the balance between apoptosis and autophagy have been shown to play prominent roles in producing resistance




Fig. 4 HMGB1 translocates from the mitochondria to the cytoplasm after sorafenib treatment. HepG2 cells were untreated or treated with 5 μ M sorafenib for 48 h. **a** Nuclear and cytoplasmic proteins were separated, and the expression of HMGB1 in both fractions was examined by western blotting. **b** HMGB1 expression in extracted mitochondria (Mito) and the remaining cytoplasm without the mitochondria (Cyt) was examined. **c** HepG2 cells were untreated (Control) or treated with 5 μ M sorafenib for 48 h (Treatment), and stained for HMGB1 (red), DAPI (blue), and tomm20 (green) to visualize the mitochondria. Cells were examined by confocal microscopy (magnification, $\times 630$). **d** Colocalization of HMGB1 with mitochondria in HepG2 cells was analysed in images from three independent experiments (** $p < 0.01$)

to sorafenib [31–35]. Additionally, the activation of an alteration or escape from the Ras/Raf/MAPK pathway in tumour cells likely contributes to chemoresistance [36]. However, the phosphatidylinositol 3-kinase/Akt (PI3K/Akt) signalling pathway, which is involved in cell survival and death in human malignancies, is suggested to be a major contributing factor [30, 32].

HMGB1 is a versatile protein in tumour biology and cancer therapy [24]. The role of HMGB1 in the intervening chemotherapy response could be associated with its sub-cellular localization and the corresponding

biological events it mediates. Intracellular HMGB1 induces cell proliferation, migration, and invasion via the MAPK and/or PI3K/Akt signalling pathways in malignancies, including myofibroblasts and human cutaneous squamous cell carcinoma [37]. The activation of Akt leads to the reduced phosphorylation of proapoptotic proteins. HMGB1 interacts with MAPK and/or PI3K/Akt to promote cell proliferation and autophagy and plays roles in inflammation. Feng et al. revealed that the activation of the PI3K/Akt signalling pathways mediated HMGB1-induced proliferation in MMC cells [38].




Fig. 5 Suppression of HMGB1 increases the sensitivity to sorafenib in vivo (* $p < 0.05$; ** $p < 0.01$). HepG2 cells stably expressing sh-HMGB1 or sh-control were injected subcutaneously into the flanks of nude mice administered sorafenib (40 mg/kg/day for 21 days). **a** Tumour growth curves in the different groups were recorded. **b** Images of dissected tumours from each group ($n = 5$) at the end of the experiment. **c** HE staining of the tumours. **d** Immunohistochemical analysis and quantitative evaluation of xenograft tumour tissues using HMGB1 antibodies

MAPK positively regulates HMGB1-mediated autophagy [23]. In turn, ERK, which is a class of MAPKs, has been shown to promote HMGB1 release in inflammatory models [39].

Intracellular HMGB1 was thought to be involved in both apoptosis and autophagy. HMGB1 has been suggested to normally have anti-apoptotic effects in cancers, but paradoxical roles in promoting apoptosis were identified in cardiomyocytes and mouse embryonic fibroblasts [16, 40–45]. Furthermore, nuclear HMGB1 indirectly regulates heat shock protein b-1 (HSPB1 or HSP27), which is an important intracellular factor that plays roles in maintaining the quality of mitochondria [46] and may indirectly affect mitochondria-mediated apoptosis. HMGB1-mediated autophagy, resulting in aggravating chemoresistance, has been demonstrated in cancers, including osteosarcoma, lung cancer, leukaemia, and gastric cancer [47–51]. Cytosolic HMGB1 directly binds Beclin1 and subsequently disrupts the Bcl2-Beclin1 interaction, leading to the formation of Beclin-1-mediated autophagosomes [52].

Extracellular HMGB1 may affect chemoresistance through several methods. The classical HMGB1-RAGE signalling axis and HMGB1-TLR4 pathways have been suggested to play important roles in cancer [53]. Because

it is an important DAMP, HMGB1-mediated immunogenic cell death enhances the effects of chemotherapy by promoting tumour cell death. HMGB1 release from cells was demonstrated to promote pancreatic tumour growth by binding RAGE and eventually regulating mitochondrial bioenergetics [47]. Additionally, the interaction between HMGB1 and TLR4 mediates anti-cancer immunity during radio- and/or chemotherapy [54].

Because no previous reports detailed the role of HMGB1 in the HCC chemotherapeutic response, our data suggest for the first time that a knockdown of HMGB1 restores the sensitivity to sorafenib in vitro in HepG2 cells and in vivo. Furthermore, we found that HMGB1 translocates from the mitochondria to the outside cytoplasm in HepG2 cells following treatment with sorafenib. Our findings led to additional questions. Does HMGB1 function in all HCC cell lines or drug-resistant cell lines? Is the phenomenon of mitochondrial HMGB1 translocation associated with the role of HMGB1 in drug sensitivity? What is the function of mitochondrial HMGB1 and cytosolic HMGB1? How does HMGB1 promote sorafenib resistance in HepG2 cells?

While we do not have the answers to these questions, these findings have helped us ask better questions. To provide insight from previous studies, the




Fig. 6 Potential mechanism of HMGB1-induced sorafenib resistance in HCC. Sorafenib targets Raf-1, B-Raf, and eIF4E phosphorylation, which leads to the inhibition of proliferation and induces apoptosis. However, intracellular HMGB1 promotes the downstream MAPK cascade through Erk1/2 phosphorylation. Then, phospho-Erk attenuates the competition between phospho-Bcl-2 and HMGB1 binding to Beclin 1. Nuclear HMGB1 and HSP27 maintain mitophagy and act as co-factors to block apoptosis. Meanwhile, through the HMGB1/RAGE axis, extracellular HMGB1 indirectly promotes tumour growth by accelerating ATP production

roles of HMGB1 in multiple biological events and the functions of sorafenib as a tyrosine kinase inhibitor are summarized in a figure. This overview offers certain insights into the mechanism by which HMGB1 potentially contributes to sorafenib resistance and the potential role of mitochondrial HMGB1 in this process (Fig. 6) [30, 47, 55–57].

This study has certain limitations. The effects were observed in a single HCC cell line, i.e., HepG2, and may be insufficient to prove that HMGB1 plays a role in sorafenib resistance in HCC. Further studies are necessary to determine whether HMGB1 regulates sorafenib resistance in different HCC cell lines. Meanwhile, elucidating the functions of cytosolic HMGB1 and mitochondrial HMGB1 may provide new insights into the different roles of HMGB1.

Conclusions

HMGB1 is involved in sorafenib resistance in HepG2 cells and in vivo. Furthermore, this is the first report to show that HMGB1 downregulation promotes the sensitivity to sorafenib. The translocation of HMGB1 from the mitochondria to the cytoplasm following sorafenib exposure requires further investigation to identify its association with the sensitivity of HepG2 to sorafenib. Our study provides new insights into the relationship between HMGB1 and sorafenib resistance in HepG2 cells. While further studies are necessary, we postulate that HMGB1 provides a new mode for regulating sorafenib resistance and could serve as a novel potential target to stratify patients suitable for sorafenib treatment or contribute to combination drug use in HCC.

Additional file

Additional file 1: FCM of Figs. 2C and 3C. The Excel spreadsheet shows the flow cytometry results of the shHMGB1 HepG2, HMGB1-overexpressing HepG2 cells and corresponding control cells with or without the sorafenib treatment. (CSV 1 kb)

Abbreviations

CCK-8: Cell counting kit-8; DAMPs: Damage-associated molecular patterns; DMEM: Dulbecco's modified eagle medium; EMT: Epithelial mesenchymal transition; FBS: Foetal bovine serum; HCC: Hepatocellular carcinoma; HE: Haematoxylin and eosin; HMGB1: High mobility group box 1; HSPB1: Heat shock protein B-1; IHC: Immunohistochemistry; MAPK: Mitogen-activated protein kinase; PDGFR: Platelet-derived growth factor receptor; PI3K/Akt: Phosphatidylinositol 3-kinase/Akt; RAGEs: Receptor for advanced glycation end products; TLRs: Toll-like receptors; VEGFR: Vascular endothelial growth factor receptor

Acknowledgements

Not applicable.

Funding

This study was supported by grants from the Special National International Technology Cooperation of China (2015DFA31490), the National Natural Science Foundation of China (No.81272253) and the National Major Sciences Research Program of China (973 Program) (No.2013CB910502) received by XGF. XGF supervised the design of the study and critically revised the manuscript. This study was supported by a grant from the National Natural Science Foundation of China (No.81670538) received by NL; NL was involved in the study design and drafting the manuscript.

Availability of data and materials

All data generated or analysed during this study are included in this published article and its Additional file 1.

Authors' contributions

YX, LS, YF, NL and XGF contributed to the experimental design; YX, YF and PZ conducted the cellular and murine experiments; YX, YF and NL contributed to drafting this manuscript; LS, YH, RZ, JQ and XGF were involved in the critical revision of the manuscript. XH and PZ contributed to the experimental quality control and data analysis. All authors have read and approved the manuscript.

Ethics approval and consent to participate

The Animal Use Committee of Xiangya Hospital approved all animal treatment protocols, and all mice were treated humanely during the entire study period.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Author details

¹Hunan Key Laboratory of Viral Hepatitis, Department of Infectious Diseases, Xiangya Hospital, Central South University, Changsha 410008, China. ²Center for Molecular Medicine, Xiangya Hospital, Key Laboratory of Molecular Radiation Oncology of Hunan Province, Central South University, Changsha 410008, China. ³Department of Blood Transfusion, Xiangya Hospital, Central South University, Changsha 410008, China.

Received: 29 June 2017 Accepted: 29 November 2017

Published online: 15 December 2017

References

- Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J, Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin*. 2015;65(2):33–64. 31
- Singal AG, Pillai A, Tiro J. Early detection, curative treatment, and survival rates for hepatocellular carcinoma surveillance in patients with cirrhosis: a meta-analysis. *PLoS Med*. 2014;11(4):e1001624.
- Llovet JM, Villanueva A, Lachenmayer A, Finn RS. Advances in targeted therapies for hepatocellular carcinoma in the genomic era. *Nat Rev Clin Oncol*. 2015;12(7):408–24.
- Cheng AL, Kang YK, Chen Z, Tsao CJ, Qin S. Efficacy and safety of sorafenib in patients in the Asia-Pacific region with advanced hepatocellular carcinoma: a phase III randomised, double-blind, placebo-controlled trial. *Lancet Oncol*. 2009;10(1):25–34.
- Chen J, Jin R, Zhao J, Liu J, Ying H, Yan H, Zhou S, Liang Y, Huang D, Liang X. Potential molecular, cellular and microenvironmental mechanism of sorafenib resistance in hepatocellular carcinoma. *Cancer Lett*. 2015;367(1):1–11.
- Gores GJ. Decade in review-hepatocellular carcinoma: HCC-subtypes, stratification and sorafenib. *Nat Rev Gastro Hepat*. 2014;11(11):645–7.
- Wei J, Wang Z, Li X, Fan X, Duan Y. High-mobility group box 1 is associated with Clinicopathologic features in patients with Hepatocellular carcinoma. *Pathol Oncol Res*. 2012;18(2):293–8.
- Liu F, Zhang Y, Peng Z, Gao H, Li X, Chen M. High expression of high mobility group box 1 (hmgb1) predicts poor prognosis for hepatocellular carcinoma after curative hepatectomy. *J Transl Med*. 2012;10(1):1–10.
- Xiao J, Ding Y, Huang J, Li Q, Liu Y, Ni W, Zhang Y, Zhu Y, Chen L, Chen B. The association of HMGB1 gene with the prognosis of HCC. *PLoS One*. 2014;9(2):e89097.
- Chen R, Hou W, Zhang Q, Kang R, Fan X-G, Tang D. Emerging role of high-mobility group box 1 (HMGB1) in liver diseases. *Mol Med*. 2013;19(1):357.
- Rui K, Chen R, Zhang Q, Wen H, Sha W, Cao L, Jin H, Yan Y, Fan XG, Yan Z. HMGB1 in health and disease. *Mol Asp Med*. 2014;40:1–116.
- Chen RC, Yi PP, Zhou RR, Xiao MF, Huang ZB, Tang DL, Huang Y, Fan XG. The role of HMGB1-RAGE axis in migration and invasion of hepatocellular carcinoma cell lines. *Mol Cell Biochem*. 2014;390(1–2):271–80.
- Ping C, Dai W, Fan W, Jie L, Miao S, Kan C, Li J, Yan Z, Wang C, Jing Y. Ethyl Pyruvate inhibits proliferation and induces apoptosis of Hepatocellular carcinoma via regulation of the HMGB1-RAGE and AKT pathways. *Biochem Bioph Res Co*. 2014;443(4):1162–8.
- Stumbo AC, Cortez E, Rodrigues CA, Henriques MDGMO, Porto LC, Barbosa HS, Carvalho L. Mitochondrial localization of non-histone protein HMGB1 during human endothelial cell–*Toxoplasma gondii* infection. *Cell Biol Int*. 2008;32(2):235–8.
- Tang D, Kang R, Livesey KM, Iijima H, Lotze MT. High mobility group box 1 (HMGB1) activates an Autophagic response to oxidative stress. *Antioxidants Redox S*. 2011;15(8):2185–95.
- Meng Q, Zhao J, Liu H, Zhou G, Zhang W, Xu X, Zheng M. HMGB1 promotes cellular proliferation and invasion, suppresses cellular apoptosis in osteosarcoma. *Tumour Biol*. 2014;35(12):12265–74.
- Pang X, Zhang Y, Wei H, Zhang J, Luo Q, Huang C, Zhang S. Expression and effects of high-mobility group box 1 in cervical cancer. *Int J Mol Sci*. 2014; 15(5):8699–712.
- Wu T, Zhang W, Yang G, Li H, Chen Q, Song R, Zhao L. HMGB1 overexpression as a prognostic factor for survival in cancer: a meta-analysis and systematic review. *Oncotarget*. 2016;7(31):50417–27.
- Lee H, Song M, Shin N, Chang HS, Min BS, Kim HS, Yoo JS, Kim H. Diagnostic significance of serum HMGB1 in colorectal carcinomas. *PLoS One*. 2012;7(4):e34318.
- Zheng H, Chen JN, Yu X, Jiang P, Yuan L, Shen HS, Zhao LH, Chen PF, Yang M. HMGB1 enhances drug resistance and promotes in vivo tumor growth of lung cancer cells. *DNA Cell Biol*. 2016;35(10):622–27.
- Huang J, Liu K, Song D, Ding M, Wang J, Jin Q, Ni J. Krüppel-like factor 4 promotes high-mobility group box 1-induced chemotherapy resistance in osteosarcoma cells. *Cancer Sci*. 2016;107:242.
- Kong Q, LH X, Xu W, Fang JP, HG X. HMGB1 translocation is involved in the transformation of autophagy complexes and promotes chemoresistance in leukaemia. *Int J Oncol*. 2015;47(1):161–70.
- Liu W, Zhang Z, Zhang Y, Chen X, Guo S, Lei Y, Xu Y, Ji C, Bi Z, Wang K. HMGB1-mediated Autophagy modulates sensitivity of colorectal cancer cells to Oxaliplatin via MEK/ERK signaling pathway. *Cancer Biol Ther*. 2015;16(4):511–7.
- Kang R, Zhang Q, Lotze MT, Tang D. HMGB1 in cancer: good, bad, or both? *Clin Cancer Res*. 2013;19(15):4046–57.
- Li N. Potential role of high mobility group box 1 in hepatocellular carcinoma. *Cell Adhes Migr*. 2014;8(5):493–8.
- Rudalska R, Dauch D, Longerich T, McJunkin K, Wuestefeld T, Kang T-W, Hohmeyer A, Pestic M, Leibold J, von Thun A. In vivo RNAi screening identifies a mechanism of sorafenib resistance in liver cancer. *Nat Med*. 2014;20(10):1138–46.
- Newell P, Toffanin S, Villanueva A, Chiang DY, Minguez B, Cabellos L, Savic R, Hoshida Y, Lim KH, Melgar-Lesmes P. Ras pathway activation in hepatocellular carcinoma and anti-tumoral effect of combined sorafenib and rapamycin in vivo. *J Hepatol*. 2009;51(4):725–33.
- Wang Y, Gao J, Zhang D, Zhang J, Ma J, Jiang H. New insights into the antifibrotic effects of sorafenib on hepatic stellate cells and liver fibrosis. *J Hepatol*. 2010;53(1):132–44.
- Sieghart W, Pinter M, Dauser B, Rohr-Udilova N, Piguet A-C, Prager G, Hayden H, Dienes H-P, Dufour J-F, Peck-Radosavljevic M. Erlotinib and sorafenib in an orthotopic rat model of hepatocellular carcinoma. *J Hepatol*. 2012;57(3):592–9.
- Gauthier A, Ho M. Role of sorafenib in the treatment of advanced hepatocellular carcinoma: an update. *Hepatol Res*. 2013;43(2):147–54.
- Malenstein HV, Dekervel J, Verslype C, Cutsem EV, Windmolders P, Nevens F, Pelt JV. Long-term exposure to sorafenib of liver cancer cells induces resistance with epithelial-to-mesenchymal transition, increased invasion and risk of rebound growth. *Cancer Lett*. 2013;329(1):74–83.
- Chen KF, Chen HL, Tai WT, Feng WC, Hsu CH, Chen PJ, Cheng AL. Activation of phosphatidylinositol 3-kinase/Akt signaling pathway mediates acquired resistance to sorafenib in hepatocellular carcinoma cells. *J Pharm Expt Therape*. 2011;337(1):155–61.
- Shimizu S, Takehara T, Hikita H. The let-7 family of microRNAs inhibits Bcl-xL expression and potentiates sorafenib-induced apoptosis in human hepatocellular carcinoma. *J Hepatol*. 2010;52(5):698–704.
- Shimizu S, Takehara T, Hikita H, Kodama T, Tsunematsu H, Miyagi T, Hosui A, Ishida H, Tatsumi T, Kanto T. Inhibition of autophagy potentiates the antitumor effect of the multikinase inhibitor sorafenib in hepatocellular carcinoma. *Int J Cancer*. 2012;131(3):548–57.
- Shi YH, Ding ZB, Zhou J, Hui B, Shi GM, Ke AW, Wang XY, Dai Z, Peng YF, CY G, et al. Targeting autophagy enhances sorafenib lethality for hepatocellular carcinoma via ER stress-related apoptosis. *Autophagy*. 2011;7(10):1159–72.
- Personeni N, Rimassa L, Pressiani T, Destro A, Ligorio C, Tronconi MC, Bozzarelli S, Carnaghi C, Tommaso LD, Giordano L. Molecular determinants of outcome in sorafenib-treated patients with hepatocellular carcinoma. *J Cancer Res Clin*. 2013;139(7):1179–87.
- Sun Y, Tu Y, He Li, Ji C, Cheng BO. High mobility group box 1 regulates tumor metastasis in cutaneous squamous cell carcinoma via the PI3K/AKT and MAPK signaling pathways. *Oncol Lett*. 2016;11(1):59–62.

38. Feng X, Chao W, Min Y, Liu Q, Li H, Liu J, Zhang Y, Hao Y, Lin K, Zhang Y. Role of PI3K/Akt signal pathway on proliferation of mesangial cell induced by HMGB1. *Tissue Cell*. 2016;48(2):121–5.
39. Zhou JR, Zhang LD, Wei HF, Wang X, Ni HL, Yang F, Zhang T, Jiang CL. Neuropeptide Y induces secretion of high-mobility group box 1 protein in mouse macrophage via PKC/ERK dependent pathway. *J Neuroimmunol*. 2013;260(1–2):55–9.
40. Guérin R, Arseneault G, Dumont S, Rokeach LA. Calnexin is involved in apoptosis induced by endoplasmic reticulum stress in the fission yeast. *Mol Biol Cell*. 2008;19(10):4404–20.
41. Brezniceanu ML, Völp K, Bösser S, Solbach C, Lichter P, Joos S, Zörnig M. HMGB1 inhibits cell death in yeast and mammalian cells and is abundantly expressed in human breast carcinoma. *FASEB J*. 2003;17(8):1295–7.
42. Amaravadi RK, Thompson CB. The roles of therapy-induced Autophagy and necrosis in cancer treatment. *Clin Cancer Res*. 2008;13(24):7271–9.
43. Ouyang F, Huang H, Zhang M, Chen M, Huang H, Huang F, Zhou S. HMGB1 induces apoptosis and EMT in association with increased autophagy following H/R injury in cardiomyocytes. *Int J Mol Med*. 2016;27(2–3):237–45.
44. Cui XS, Shen XH, Kim NH. High mobility group box 1 (HMGB1) is implicated in preimplantation embryo development in the mouse. *Mol Reprod Dev*. 2008;75(8):1290–9.
45. Yang H, Antoine DJ, Andersson U, Tracey KJ. The many faces of HMGB1: molecular structure-functional activity in inflammation, apoptosis, and chemotaxis. *J Leukoc Biol*. 2013;93(6):865–73.
46. Narumi T, Shishido T, Otaki Y, Kadowaki S, Honda Y, Funayama A, Honda S, Hasegawa H, Kinoshita D, Yokoyama M. High-mobility group box 1-mediated heat shock protein beta 1 expression attenuates mitochondrial dysfunction and apoptosis. *J Mol Cell Cardiol*. 2015;82(2):1–12.
47. Kang R, Tang D, Schapiro NE, Loux T, Livesey KM, Billiar TR, Wang H, Van HB, Lotze MT, Zeh HJ. The HMGB1/RAGE inflammatory pathway promotes pancreatic tumor growth by regulating mitochondrial bioenergetics. *Oncogene*. 2014;33(5):567–77.
48. Huang J, Ni J, Liu K, Yu Y, Xie M, Kang R, Vernon P, Cao L, Tang D. HMGB1 promotes drug resistance in osteosarcoma. *Cancer Res*. 2011;72(1):230–8.
49. Kang R, Zeh HJ, Lotze MT, Tang D. The Beclin 1 network regulates autophagy and apoptosis. *Cell Death Differ*. 2011;18(4):571–80.
50. Huang J, Liu K, Yu Y, Xie M, Kang R, Vernon PJ, Cao L, Tang D, Ni J. Targeting HMGB1-mediated autophagy as a novel therapeutic strategy for osteosarcoma. *Autophagy*. 2012;8(2):275–7.
51. Liu L, Yang M, Kang R, Wang Z, Zhao Y, Yu Y, Xie M, Yin X, Livesey KM, Lotze MT. HMGB1-induced autophagy promotes chemotherapy resistance in leukemia cells. *Leukemia*. 2010;25(1):23–31.
52. Sun X, Tang D. HMGB1-dependent and -independent autophagy. *Autophagy*. 2014;10(10):1873–6.
53. Sims GP, Rowe DC, Rietdijk ST, Herbst R, Coyle AJ. HMGB1 and RAGE in inflammation and cancer. *Annu Rev Immunol*. 2010;28:367–88.
54. Apetoh L, Ghiringhelli F, Tesniere A, Obeid M, Ortiz C, Criollo A, Mignot G, Maiuri MC, Ullrich E, Saulnier P. Toll-like receptor 4-dependent contribution of the immune system to anticancer chemotherapy and radiotherapy. *Nat Med*. 2007;13(9):1050–9.
55. Soyoung C, Chen Z, Tang LH, Fang Y, Shin SJ, Panarelli NC, Chen YT, Li Y, Jiang X, YCN D. Bcl-xL promotes metastasis independent of its anti-apoptotic activity. *Nat Commun*. 2016;7:10384.
56. Tang D, Kang R, Livesey K, Kroemer G, Billiar T, Vanhouten B, Iii HZ, Lotze M. High-mobility group box 1 is essential for mitochondrial quality control. *Cell Metab*. 2011;13(6):701–11.
57. Tang D, Kang R, Livesey KM, Cheh CW, Farkas A, Loughran P, Hoppe G, Bianchi ME, Tracey KJ, Rd ZH. Endogenous HMGB1 regulates autophagy. *J Cell Biol*. 2010;190(5):881–92.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

