

RESEARCH ARTICLE

Antibodies Directed against Shiga-Toxin Producing *Escherichia coli* Serotype O103 Type III Secreted Proteins Block Adherence of Heterologous STEC Serotypes to HEp-2 Cells

Taseen S. Desin^{1*}, Hugh G. Townsend, Andrew A. Potter

Vaccine & Infectious Disease Organization—International Vaccine Center, University of Saskatchewan, Saskatoon, Saskatchewan, Canada

✉ Current address: Basic Sciences Department, College of Science & Health Professions, King Saud bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

* desint@ksau-hs.edu.sa



OPEN ACCESS

Citation: Desin TS, Townsend HG, Potter AA (2015) Antibodies Directed against Shiga-Toxin Producing *Escherichia coli* Serotype O103 Type III Secreted Proteins Block Adherence of Heterologous STEC Serotypes to HEp-2 Cells. PLoS ONE 10(10): e0139803. doi:10.1371/journal.pone.0139803

Editor: Gunnar F Kaufmann, The Scripps Research Institute and Sorrento Therapeutics, Inc., UNITED STATES

Received: June 22, 2015

Accepted: September 17, 2015

Published: October 9, 2015

Copyright: © 2015 Desin et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: This project was funded by the Saskatchewan Health Research Foundation.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

Shiga toxin-producing *Escherichia coli* (STEC) serotype O103 is a zoonotic pathogen that is capable of causing hemorrhagic colitis and hemolytic uremic syndrome (HUS) in humans. The main animal reservoir for STEC is ruminants and hence reducing the levels of this pathogen in cattle could ultimately lower the risk of STEC infection in humans. During the process of infection, STEC_{O103} uses a Type III Secretion System (T3SS) to secrete effector proteins (T3SPs) that result in the formation of attaching and effacing (A/E) lesions. Vaccination of cattle with STEC serotype O157 T3SPs has previously been shown to be effective in reducing shedding of STEC_{O157} in a serotype-specific manner. In this study, we tested the ability of rabbit polyclonal sera against individual STEC_{O103} T3SPs to block adherence of the organism to HEp-2 cells. Our results demonstrate that pooled sera against EspA, EspB, EspF, NleA and Tir significantly lowered the adherence of STEC_{O103} relative to pre-immune sera. Likewise, pooled anti-STEC_{O103} sera were also able to block adherence by STEC_{O157}. Vaccination of mice with STEC_{O103} recombinant proteins induced strong IgG antibody responses against EspA, EspB, NleA and Tir but not against EspF. However, the vaccine did not affect fecal shedding of STEC_{O103} compared to the PBS vaccinated group over the duration of the experiment. Cross reactivity studies using sera against STEC_{O103} recombinant proteins revealed a high degree of cross reactivity with STEC_{O26} and STEC_{O111} proteins implying that sera against STEC_{O103} proteins could potentially provide neutralization of attachment to epithelial cells by heterologous STEC serotypes.

Introduction

Shiga toxin-producing *Escherichia coli* (STEC) is an enteric pathogen that causes diarrhoeal illness in humans which can lead to hemorrhagic colitis and haemolytic uremic syndrome (HUS), one of the main causes of renal failure in children [1]. Shiga toxins produced by this pathogen play an important role in causing these clinical manifestations. Currently, there is no treatment available for human STEC infections other than supportive care as the administration of antibiotics can exacerbate the disease. STEC O157:H7 is the predominant serotype associated with human infections in North America, while non-O157:H7 serotypes such as O103, O26, O111 are more prevalent in many European countries, South America and parts of Australia [1,2,3]. The main reservoir for STEC is ruminants [4] and therefore intervention strategies aimed at lowering the levels of this pathogen in cattle could ultimately result in improved human health [5].

During the process of infection, STEC uses a Type Three Secretion System (T3SS) to inject virulence factors known as effector proteins directly into host cells, leading to the formation of attaching and effacing lesions (A/E) lesions, which are hallmarks of STEC infections. The major structural components of the STEC T3SS include EspA (filament), EspB and EspD (translocon complex) [6]. The STEC T3SS secretes over 50 effector proteins that are encoded on the LEE Pathogenicity Island or elsewhere on the chromosome (non-LEE effectors) [7]. The translocated intimin receptor, Tir, is an effector protein which enters host cells and forms a receptor that binds to intimin that is expressed on the surface of STEC cells [6]. Many studies have shown that the STEC T3SS is essential for colonization of cattle, implying that this is a major virulence factor employed by this pathogen [8,9,10].

Vaccination of cattle with STEC_{O157} T3SP's has shown to be an effective strategy in reducing the shedding of STEC_{O157} [11,12,13,14,15,16,17,18,19]. However, this protection appears to be serotype specific [20,21]. Therefore, alternative antigens need to be identified that offer protection against non-O157 STEC serotypes. Recently, it has been shown that anti-sera to an extract of STEC_{O157} T3SPs had the highest degree of cross-reactivity with STEC_{O103} recombinant T3SPs [20], suggesting that STEC_{O103} T3SPs may have cross-protective potential. In this study, we tested the effect of sera against STEC_{O103} recombinant proteins on STEC_{O103} and STEC_{O157} adherence to HEp-2 cells. Moreover, we tested the vaccine potential of the recombinant proteins against STEC_{O103} challenge in mice.

Materials and Methods

Bacterial strains and growth conditions

The bacterial strains used in this study comprised of *E. coli* N01-2454 (O103:H2), EDL933 (O157:H7), CL9 (O26:H11) and CL101 (O111:NM) [22,23]. For cloning and protein expression, we used the *E. coli* K-12 lab strains, JM109 (*endA1*, *recA1*, *gyrA96*, *thi*, *hsdR17* (r_k^- , m_k^+), *relA1*, *supE44*, Δ (*lac-proAB*), [F' *traD36*, *proAB*, *laqI^qZAM15*]) and BL21 (F', *dcm*, *ompT*, *hsdS_B* (r_B^- , m_B^-), *gal*, λ (DE3)) obtained from Qiagen and Invitrogen, respectively. The strains were grown in Luria Bertani (LB) medium at 37°C in an orbital shaker (250 rpm), unless otherwise stated. *E. coli* serotypes O103 and O157 were transformed with a green fluorescent protein expressing plasmid, pNR78, for visualization by fluorescence microscopy during the adherence inhibition assays as described [21]. Plasmid pNR78 was constructed in our lab by amplifying the GFP gene from pQBI-25 (Quantum Biotechnologies) which was cloned downstream of the GroEL promoter.

Protein expression and purification

The STEC serotype O103:H2 T3SS genes *escC*, *espA*, *espB*, *espF*, *espG*, *espR1*, *nleA*, *nleE*, *nleF*, *nleG2*, *nleH*, *sepD*, *tccp2* and *tir* were amplified by PCR (Applied Biosystems) based on the

sequence provided by GenBank[®]. Similarly, for cross reactivity studies, *espA*, *espB*, *espF*, *nleA* and *tir* from STEC serotypes O26 and O111 were amplified by PCR. The genes were cloned in either pQE-30 (Qiagen), pET-15b (Novagen), pGEX-5X-1 (GE Healthcare) or pGEX-5X-3, of which the first two are 6x His-tagged protein expression vectors while the latter are Glutathione S-transferase (GST)-fusion expression vectors. The constructs were confirmed by PCR and sequencing (Plant Biotechnology Institute, Saskatoon). Proteins were expressed in *Escherichia coli* K-12 lab strains (JM109 or BL21) and purified using either the method described in the QIAexpressionist[™] manual (Qiagen) for His-tagged proteins or the GST Gene Fusion System Handbook (GE Healthcare) for the GST-fusion protein. Purified protein samples were greater than 90% pure as determined by SDS-PAGE followed by Coomassie blue staining as described previously [20].

Raising polyclonal anti-sera to STEC_{O103} T3SS recombinant proteins

Purified recombinant proteins (100 µg each) were formulated with 30% Emulsigen D (MVP Laboratories) and two New Zealand White rabbits (Charles River) per STEC_{O103} recombinant protein were immunized subcutaneously on day 0, followed by booster injections on days 21 and 42. The rabbits were euthanized on day 56 and sera were collected. Antibody titers against STEC_{O103} recombinant proteins were confirmed using ELISA in duplicate wells as previously described [20]. For antibody titer determinations, the cut-off value was considered to be the average of the blank and two standard deviations. All rabbits used in this study were handled and treated in accordance with the guidelines provided by the Canadian Council on Animal Care (CCAC) as administered by the University Committee on Animal Care and Supply (UCACS), protocol 1994–213. This protocol was approved by the UCACS at the University of Saskatchewan for the present study.

Cell culture

HEp-2 cells (ATCC[®] CCL-23[™], CEDARLANE[®]) were grown in HyClone Dulbecco modified Eagle medium (DMEM; Thermo Scientific) supplemented with 10% fetal bovine serum (FBS; PAA Laboratories) and 1% HEPES Buffer (Invitrogen) at 37°C in a 5% CO₂ incubator. One day prior to the adherence inhibition assays, 10⁵ cells per well were seeded onto eight well chamber slides (Nunc) and allowed to incubate overnight.

Adherence inhibition assays

Adherence of STEC_{O103} and STEC_{O157} to HEp-2 cells was assessed using an assay described elsewhere [21]. Briefly, an overnight culture of STEC grown in LB media was subcultured (1:100) into DMEM containing 10% FBS and 1% HEPES Buffer for 2 hours (until the OD₆₀₀ was 0.2) at 37°C and 5% CO₂ without shaking. For testing the effect of pooled sera against STEC_{O103} T3SPs on adherence, HEp-2 cells were infected with 25 µl of STEC (1.7 x 10⁶ colony forming units), 10 µl of each serum and 225 µl fresh DMEM. The effect of individual anti-serum was tested by infecting HEp-2 cells with 25 µl of STEC (1.7 x 10⁶ CFU), 20 µl of anti-serum and 225 µl fresh DMEM (anti-O103 antibodies were prepared as described previously [21]). The chamber slides were incubated at 37°C and 5% CO₂ for 3 hours (STEC_{O157}) or 3.5 hours (STEC_{O103}). The slides were washed six times with 200 µl Phosphate Buffered Saline (PBS, 0.1M pH 7.2) and fixed with 200 µl PBS containing 3.7% Formaldehyde. This was followed by two washes with PBS after which the slides were allowed to air dry. Coverslips were mounted with Vectashield[®] (Vector) containing DAPI and sealed. The slides were visualized under the fluorescent microscope (Axiovert 200 inverted microscope–Zeiss). Bacteria were observed under FITC, while HEp-2 cells were observed under DAPI. Each experimental group

was first tested using 2 replicate wells in an 8 well chamber slide and 4 random grids were examined per well under the fluorescent microscope as described below. After observing clear differences in STEC adherence to HEp-2 cells between the different treatments, the experiments were repeated independently on a separate occasion using 8 replicates per test group as previously published [21] with 4 random grids per well used for enumerating the number of STEC per HEp-2 cell. The resulting pictures (4 under FITC and 4 under DAPI) per well (total of 8 pictures per well) were used to enumerate the number of STEC and HEp-2 cells per well. The total numbers of STEC per grid were then divided by the total numbers of HEp-2 cells per grid to determine the number of STEC per HEp-2 cell in one grid. This was repeated for the 8 duplicate wells per group, resulting in a total of 64 pictures per test group. Each data point in Figs 1 and 2 represent the average number of STEC per HEp-2 cell from 4 counts (4 random grids) per well. For statistical analysis, the median STEC per HEp-2 cell across the different test groups were compared using a non-parametric analysis as described below.

Immunization of mice with STEC_{O103} T3SPs

Twenty four BALB/C mice were obtained from Charles River Canada. Mice were housed at the VIDO Animal Care Facility (University of Saskatchewan) and handled in accordance with the guidelines provided by the CCAC as administered by the UCACS, protocol 1998–0003. This protocol was approved by the UCACS at the University of Saskatchewan for the present study. Mice were randomly divided into two groups with 12 mice per group. The mice were immunized subcutaneously on Day 0 with 100 µl of either PBS (0.1M, pH 7.2) or a pool of STEC_{O103} recombinant proteins EspA, EspB, EspF, NleA and Tir (1 µg of each protein) followed by a second immunization at Day 21. The vaccines were formulated with 30% Emulsigen D (MVP Laboratories). Sera were collected prior to both immunizations on day 0 and day 21 as well as prior to challenge with STEC_{O103} on day 35. Antibody titers were determined using ELISA in duplicate wells as described previously [20]. For antibody titer determinations, the cut-off value was considered to be the average of the blank and two standard deviations. The mice were challenged as described below.

STEC mouse colonization model

For colonization of mice, we used the streptomycin-treated model as previously described [24,25]. Briefly, mice were given water containing Streptomycin Sulfate (5 g/L) on day 32 for two days. Subsequently, mice were deprived of food and water for 24 hours. On Day 35, mice were orally challenged with 100 µl of 10⁹ cfu of STEC_{O103} Nal^r (resuspended in 20% sucrose). The mice were permitted access to food and water containing Streptomycin for the rest of the experiment. Fecal pellets were collected every 3 days for 21 days post challenge. Shedding of STEC_{O103} was monitored by incubating the fecal samples in 1 ml LB broth for 2 hours at room temperature to allow the pellets to soften. The samples were vortexed, serially diluted in PBS and plated on MacConkey Sorbitol Agar containing Nalidixic Acid (15 µg/ml), Cefixime (5 µg/ml) and Potassium Tellurite (2.5 µg/ml). The plates were incubated overnight at 37°C and STEC colonies were enumerated the following day. Bacterial counts were expressed as cfu per gram of fecal content.

Cross-reactivity of STEC_{O103} T3SS recombinant protein specific sera

Purified STEC_{O26} and STEC_{O111} EspA, EspB, EspF, NleA and Tir recombinant proteins were separated by SDS-PAGE and transferred to a nitrocellulose membrane using a Mini Trans-Blot Electrophoretic Cell (Bio-Rad) as per the manufacturer's instructions. The membranes were probed with either polyclonal sera (1:2500) from mice vaccinated with a pool of the corresponding STEC_{O103} recombinant proteins or with rabbit polyclonal sera (1:2500) raised against




Fig 1. Inhibition of STEC_{O103} adherence to HEp-2 cells by either (A) pooled sera against recombinant STEC_{O103} T3SPs or (B and C) individual serum samples specific for STEC_{O103} recombinant proteins EspB, EspF, EspG or NleA. Anti-O103 refers to antibodies against a secreted fraction of T3SPs from STEC_{O103}. Values are expressed as median bacteria per cell from 8 replicates. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$; ****, $P < 0.0001$.

doi:10.1371/journal.pone.0139803.g001

STEC_{O103} EspA, EspB, EspF, NleA and Tir. Alkaline phosphatase labeled goat anti-mouse or goat anti-rabbit IgG (KPL) antibodies were used as secondary antibodies (1:2000). The membranes were developed using 5-bromo-4-chloro-3-indolyl phosphate (BCIP) and nitroblue tetrazolium (NBT) salt according to the manufacturer's instructions (Sigma).

Statistical analyses

Statistical Analyses were performed using GraphPad Prism 6.02. Adherence inhibition assays were analyzed using a non-parametric analysis (Kruskal-Wallis test) and individual groups were tested using Dunn's multiple comparison test. Mouse antibody titers were presented as medians plus/minus the 25th and 75th percentile ranges. Differences in immune responses between the vaccine and control groups were tested using non-parametric repeated measures ANOVA. A P value of < 0.05 was considered significant.

Results

Immune responses against STEC_{O103} recombinant proteins in rabbits

Polyclonal sera were raised against 14 STEC_{O103} recombinant proteins in rabbits in order to test the adherence inhibition effect of the sera in vitro and for cross reactivity studies. All the




Fig 2. Inhibition of STEC_{O157} adherence to HEp-2 cells by pooled sera against recombinant STEC_{O103} T3SPs. (A) STEC_{O157} adherence to HEp-2 cells was significantly lower in the presence of antibodies against STEC_{O103} EspA, EspB, EspF, NleA and Tir. (B) STEC_{O157} adherence was partially lower in the presence of STEC_{O103} anti-EspA and anti-Tir or anti-EspB, anti-EspF and anti-NleA. Values are expressed as median bacteria per cell from 8 replicates. **, $P < 0.01$; ****, $P < 0.0001$.

doi:10.1371/journal.pone.0139803.g002

recombinant proteins induced a significant IgG specific antibody response as determined by ELISA (Table 1). The mean IgG titer across all the proteins was 911,801, while NleE had the lowest antibody titer (151,399) and EspR1 had the highest antibody titer (2,771,000).

Antibodies against STEC_{O103} T3SP's inhibit adherence of STEC_{O103}

To test the effect of rabbit polyclonal sera against recombinant STEC_{O103} T3SPs on adherence, we used a functional assay where we measured the level of STEC_{O103} adherence to HEp-2 cells. Our results demonstrate that pooled sera against STEC_{O103} recombinant proteins significantly reduced adherence of STEC_{O103} to HEp-2 cells relative to the group incubated with pre-immune sera (Fig 1A). In order to determine which serum samples were involved in this adherence inhibition effect, we tested specific anti-sera to EspA, EspB, EspF, EspG, EscC, EspR1, NleA, NleE, NleF, NleG2, NleH, SepD, Tir and Tccp2 individually in duplicate. We observed that sera against EspB, EspG, EspF and NleA were involved in blocking adherence (data not shown). To confirm this observation, we performed an adherence inhibition assay where sera against EspB, EspG, EspF and NleA were tested individually with 8 replicates. The data clearly suggest that anti-sera to these four proteins were also highly effective in blocking STEC_{O103} adherence to HEp-2 cells compared to the group treated with pre-immune serum (Fig 1B and 1C).

Anti- STEC_{O103} T3SP sera have cross-protective potential

In order to determine if antibodies against STEC_{O103} recombinant proteins can block adherence of other STEC serotypes, we evaluated the effect of pooled sera on STEC_{O157} adherence to HEp-2 cells. Interestingly, our results indicate that incubation of STEC_{O157} with anti-sera to STEC_{O103} EspA, EspB, EspF, NleA and Tir significantly lowered adherence to HEp-2 cells, while anti-sera to the other proteins did not have a major effect (Fig 2A). We further investigated this adherence inhibition effect by testing pooled sera against STEC_{O103} EspA and Tir in one group and sera against EspB, EspF and NleA in another group. Adherence of STEC_{O157} to

Table 1. IgG antibody titers specific for STEC_{O103} recombinant proteins in rabbits.

	Pre-immune	Day 56
Anti-EscC	87.5	241547
Anti-EspA	882.5	838090
Anti-EspB	783.5	1256000
Anti-EspF	711	618333
Anti-EspG	773.5	846100
Anti-EspR1	789.5	2771000
Anti-NleA	2778	1332000
Anti-NleE	555.5	151399
Anti-NleF	2748.5	1036000
Anti-NleG2	652	1358000
Anti-NleH	499.5	614552
Anti-SepD	536.5	483176
Anti-Tccp2	252.5	402873
Anti-Tir	122.5	816244

Antibody titers were determined by setting the cut-off value as the average of the blank and two standard deviations. All serum samples were tested in duplicate wells.

doi:10.1371/journal.pone.0139803.t001

Table 2. Median IgG antibody titers specific for STEC_{O103} EspA, EspB, EspF, NleA or Tir in mice vaccinated with either PBS or a pool of STEC_{O103} recombinant proteins.

Antibody	PBS				Vaccine				P
	Day 0		Day 35		Day 0		Day 35		
Anti-EspA	693	(590–853)	1089	(736–1462)	772	(581–860)	6979	(3109–11618)	0.0007
Anti-EspB	235	(157–304)	600	(353–814)	243	(221–477)	294644	(195617–427747)	<0.0001
Anti-EspF	882	(742–1240)	1232	(973–5538)	1239	(753–1930)	3822	(1067–13092)	0.052
Anti-NleA	481	(161–839)	2734	(422–7809)	552	(250–4314)	1019000	(5364–1385000)	0.0004
Anti-Tir	570	(470–1154)	845	(711–3461)	517	(387–652)	280528	(151906–821988)	<0.0001

Values are expressed as median titers. Numbers in parentheses represent the 25th–75th percentile. Antibody titers were determined by setting the cut-off value as the average of the blank and two standard deviations. All serum samples were tested in duplicate wells.

doi:10.1371/journal.pone.0139803.t002

HEp-2 cells was lower in both groups relative to the control group (Fig 2B) but not to the same level as in the pooled group (Fig 2A). These results suggest that antibodies to STEC_{O103} EspA, EspB, EspF, NleA and Tir proteins have a combined effect on blocking STEC_{O157} adherence and that they have cross-protective potential.

Immunization of mice with T3SP's from STEC_{O103} induces a strong humoral response but does not affect fecal shedding

In order to test the protective capacity of STEC_{O103} effectors, mice were vaccinated subcutaneously with a pool of recombinant proteins and were subsequently infected with STEC_{O103} by oral challenge. Two weeks following the booster immunization, significant EspA-, EspB-, NleA- and Tir-specific IgG titers were detected in the sera relative to the control group (Table 2). In contrast, immunization with EspF elicited a weak IgG specific serum response. To assess the protective capacity of our vaccine, fecal shedding of STEC_{O103} was monitored over 21 days post challenge. The levels of STEC_{O103} were similar in both vaccinates and non-vaccinates throughout the duration of the study, suggesting that antibodies against the antigens used for immunization did not prevent STEC_{O103} from persisting in the intestine (Fig 3), or that the response was not of sufficient magnitude.

STEC_{O26} and STEC_{O111} T3SS proteins display significant cross-reactivity with anti-sera to the corresponding STEC_{O103} proteins

The cross-reactivity of sera against STEC_{O103} T3SS proteins with other STEC serotypes including STEC_{O26} and STEC_{O111}, was first tested by western blotting using rabbit polyclonal sera. Our results indicate that EspB_{O111}, EspF_{O111} and NleA_{O111} reacted strongly with anti-sera to the corresponding STEC_{O103} proteins, while Tir_{O111} displayed a weaker reaction and EspA_{O111} did not react (Fig 4A). The western blot profile for STEC_{O26} proteins was similar with respect to EspB_{O26} and EspF_{O26}. However, EspA_{O26} also reacted strongly, unlike EspA_{O111}, while NleA_{O26} did not react (Fig 4B). Subsequently, sera from mice immunized with a pool of STEC_{O103} recombinant proteins was used to study the cross-reactivity with the equivalent STEC_{O26} and STEC_{O111} proteins. The results indicate that EspB_{O26} reacted strongly with the anti-sera while EspA_{O26}, EspF_{O26} and NleA_{O26} did not (Fig 5A). In contrast, the western blot profile for the STEC_{O111} recombinant proteins showed a significant degree of cross reactivity for EspB_{O111}, NleA_{O111} and Tir_{O111} (Fig 5B). Taken together, the results suggest that EspB_{O103}, NleA_{O103}, and Tir_{O103} possess significant cross-reactive properties with the corresponding




Fig 3. STEC_{O103} shedding in feces following oral challenge in mice. Mice were immunized subcutaneously with either PBS (control) or a pool of STEC_{O103} EspA, EspB, EspF, NleA and Tir followed by a booster immunization three weeks later. Two weeks after the second immunization, mice were orally challenged with 10⁹ cfu of STEC_{O103}. N = 12 for both groups. Values are expressed as median cfu per gram of feces.

doi:10.1371/journal.pone.0139803.g003

proteins from STEC_{O26} and STEC_{O111}. Hence, these proteins may form the basis of a cross-protective vaccine that confers protection against multiple STEC serotypes.

Discussion

Many efforts have been made to develop STEC_{O157} vaccines using the T3SS proteins as targets in order to reduce the levels of the pathogen in cattle [5,12]. However, these vaccination strategies provide limited protection as they are directed only against STEC_{O157} and they are limited in their benefit [20,21]. Non-O157 STEC serotypes are more prevalent in other parts of the world [1] and with the rise in non-O157 STEC infections in humans [26] as well as the increase in the prevalence of these serotypes in cattle [27], a vaccine that can confer protection against multiple serotypes is more desirable. The aim of this study was to determine if STEC_{O103} T3SS proteins could provide protection against STEC_{O103} as well as other heterologous serotypes using adherence-inhibition assays and the streptomycin-treated mouse model.

We used STEC_{O103} T3SS proteins as targets for a potentially cross protective vaccine since the T3SS proteins encoded by this serotype have previously been shown to have the highest degree of cross reactivity with STEC_{O157}, relative to STEC_{O26} and STEC_{O111} [20,21]. Based on this, we over-expressed and purified STEC_{O103} EscC, EspA, EspB, EspF, EspG, EspR1, NleA, NleE, NleF, NleG2, NleH, SepD, Tccp2 and Tir recombinant proteins. In order to test the protective capacity of these proteins, we first examined the effect of rabbit polyclonal sera against the candidate proteins in vitro using a HEp-2 cell adherence inhibition assay which has been successfully used as a functional assay to study STEC adherence [21,28]. This, in turn, may reflect the effect of antibodies on intestinal colonization. Our results demonstrate that pooled sera against STEC_{O103} recombinant proteins significantly inhibited STEC_{O103} adherence. This is in agreement with the findings of Asper *et al*, where anti-sera to all STEC_{O103} secreted proteins blocked adherence by this serotype [21]. We also show for the first time that sera against




Fig 4. Cross-reactivity of STEC_{O103} EspA, EspB, EspF, NleA and Tir specific rabbit polyclonal sera with the corresponding STEC_{O26} and STEC_{O111} recombinant proteins. (A) Western blot using anti-EspA. Lane 1, marker; Lane 2, EspA_{O103} (20.5 kDa); Lane 3, EspA_{O26} (20.5 kDa); Lane 4, EspA_{O111} (20.5 kDa). (B) Western blot using anti-EspB. Lane 1, marker; Lane 2, EspB_{O103} (33.1 kDa); Lane 3, EspB_{O26} (33.2 kDa); Lane 4, EspB_{O111} (32.8 kDa). (C) Western blot using anti-EspF. Lane 1, marker; Lane 2, EspF_{O103} (57 kDa); Lane 3, EspF_{O26} (40 kDa); Lane 4, EspF_{O111} (27 kDa). (D) Western blot using anti-NleA. Lane 1, marker; Lane 2, NleA_{O103} (44 kDa); Lane 3, NleA_{O26} (11.7 kDa); Lane 4, NleA_{O111} (47.5 kDa). (E) Western blot using anti-Tir. Lane 1, marker; Lane 2, Tir_{O103} (56 kDa); Lane 3, Tir_{O111} (56.9 kDa).

doi:10.1371/journal.pone.0139803.g004

individual STEC_{O103} recombinant proteins including, EspB, EspF, EspG and NleA, inhibited adherence of the bacteria to HEp-2 cells. Interestingly, pooled anti-sera to STEC_{O103} recombinant proteins EspA, EspB, EspF, NleA and Tir were able to block adherence of STEC_{O157}, suggesting that these candidate proteins may provide protection against multiple STEC serotypes. However, it appears that the inhibition of STEC_{O157}, unlike that of STEC_{O103}, was due to a combination of STEC_{O103} anti-sera since there was reduced inhibition of STEC_{O157} once the pooled anti-sera were divided into two groups. Taken together, this is the first report which illustrates that sera against STEC_{O103} T3SS recombinant proteins can block adherence of STEC_{O157} to HEp-2 cells.

The streptomycin-treated mouse model [24,29] was used to test the efficacy of the identified candidate STEC_{O103} recombinant proteins as antigens for protection against STEC_{O103}. This model was chosen since it has been widely used by various groups to test their STEC vaccines prior to conducting studies in cattle [25,30,31,32,33]. The mice developed strong serum IgG specific titers against EspA_{O103}, EspB_{O103}, NleA_{O103} and Tir_{O103} following immunization, while the response to the corresponding EspF recombinant protein was weak. The weak response to EspF_{O103} is in line with what was observed for EspF_{O157} in a previous study published by our group [20]. Immunization with STEC_{O103} recombinant proteins did not affect STEC_{O103} fecal shedding over the duration of the experiment relative to the control group.




Fig 5. Cross-reactivity of STEC_{O103} EspA, EspB, EspF, NleA and Tir specific mouse polyclonal sera with the corresponding STEC_{O26} and STEC_{O111} recombinant proteins. Western blot using sera from mice immunized with a combination of STEC_{O103} EspA, EspB, EspF, NleA and Tir to test the cross-reactivity with: (A) STEC_{O26} proteins. Lane 1, marker; Lane 2, EspA_{O26} (20.5 kDa); Lane 3, EspB_{O26} (33.2 kDa); Lane 4, EspF_{O26} (40 kDa); and Lane 5, NleA_{O26} (11.7 kDa). (B) STEC_{O111} proteins. Lane 1, marker; Lane 2, EspA_{O111} (20.5 kDa); Lane 3, EspB_{O111} (32.8 kDa); Lane 4, EspF_{O111} (27 kDa); Lane 5, NleA_{O111} (47.5 kDa); and Lane 6, Tir_{O111} (56.9 kDa).

doi:10.1371/journal.pone.0139803.g005

This was unexpected since similar STEC_{O157} based vaccines have been highly effective in mice [30,33]. In addition, a recent vaccination study by our group illustrated that a combination of nine STEC_{O157} recombinant proteins was highly effective in controlling STEC_{O157} fecal shedding in mice (data not shown). It is possible that our STEC_{O103} vaccine may have been more effective against intestinal colonization had it been administered intranasally. However, both subcutaneous and intranasal immunization of mice with an extract of STEC_{O157} secreted proteins as well as individual recombinant proteins have proven to be highly effective in controlling STEC_{O157} shedding [30]. In addition, the lack of a robust immune response against EspA may have contributed to the persistence of STEC_{O103} in the intestines. Alternatively, since very little work has been done on STEC_{O103} in mice, we speculate that the T3SS may play a different role in STEC_{O103} infection in mice. Therefore, further analysis of the STEC_{O103} T3SS may be required in mice, while a similar vaccine study should be performed in cattle with STEC_{O103}.

The serological cross reactivity of STEC_{O103} recombinant proteins EspA, EspB, EspF, NleA and Tir with the corresponding STEC_{O26} and STEC_{O111} proteins was analyzed by western blotting. Overall our results indicate that there was significant cross reactivity of the STEC_{O26} recombinant proteins, EspA_{O26}, EspB_{O26} and EspF_{O26} when rabbit polyclonal sera were used. These observations are supported by the protein sequence homology of the STEC_{O26} proteins to STEC_{O103}: EspA_{O26} (92%), EspB_{O26} (99%) and EspF_{O26} (91%). The fact that NleA_{O26} did not cross react was not unexpected since the STEC_{O26} genome contains an NleA-like gene which encodes for an 11 kDa protein, while the actual size of NleA_{O103} is 44 kDa. Therefore, sequence homology between NleA_{O103} and NleA_{O26} is expected to be low (58%) with few shared epitopes, if any. We did not show the results for Tir_{O26} since we were unable to express or purify this protein despite numerous attempts. This may be explained by the fact that Tir_{O26} may require co-expression and co-purification with a chaperone [34]. The western blot profile for STEC_{O111} recombinant proteins EspB_{O111}, EspF_{O111}, NleA_{O111} showed a high degree of cross reactivity with the corresponding sera, while there was lower cross reactivity with Tir_{O111}. This is consistent with the observed sequence homologies between the STEC_{O103} and STEC_{O111} proteins: EspB_{O111} (71%), EspF_{O111} (70%), NleA (83%) and Tir (65%). The fact that EspA_{O111} did not react to sera against EspA_{O103} was surprising since EspA_{O111} shares greater

than 81% sequence homology to EspA_{O103}. The serological cross reactivity of the STEC serotypes O26 and O111 recombinant proteins was remarkably lower when mouse polyclonal sera were used. The difference in the results may be due to differences in recognition of epitopes by the mouse and rabbit immune systems. Overall, the data from both cross reactivity studies suggests that EspB_{O103}, NleA_{O103} and Tir_{O103} are highly cross reactive and have the potential to form an efficacious recombinant vaccine that protects cattle not only against STEC_{O103} but other STEC serotypes as well. This finding is supported by two recent studies which demonstrate that EspB_{O157} and Tir_{O157} are immunogenic and protective in cattle against STEC_{O157} [35,36]. Although these studies provide important information about STEC_{O157}, this can be used as a basis for conducting similar studies with STEC_{O103} to test for cross serotype protection.

Vaccination with a commercially available STEC_{O157} T3SS vaccine (Econiche™) is an effective strategy to control STEC shedding in cattle [37]. Many recent studies have proven that this preslaughter intervention does lead to reduced levels of this pathogen in cattle [38,39]. Moreover, Mathews *et al* have recently predicted that vaccination of cattle against STEC_{O157} will have a significant impact on public health by lowering human STEC infections by 85% [40]. Our *in vitro* results are the first steps towards a vaccine that may provide protection against multiple STEC serotypes, which is highly desirable for both North America as well as other regions where non-O157 STEC serotypes are more prevalent. The STEC_{O157} SRP[®] vaccine (contains siderophore and porin proteins) has also shown to be effective in reducing fecal shedding in cattle [41,42]. However, this vaccine also confers limited serotype protection like the Econiche™ vaccine [43,44]. Taken together, the need for an STEC vaccine that provides protection against more than one serotype is required and our *in vitro* results suggest that STEC_{O103} may be a likely candidate, though further testing is required in cattle.

Acknowledgments

The authors would like to thank Neil Rawlyk for technical assistance and members of Dr. Suresh Tikoo's lab for help with the fluorescent microscope. Special thanks to Sherri Tetland and VIDO Animal Care for help with raising anti-sera in rabbits and the mouse experiments. The authors acknowledge the reagents provided by the VIDO Glassware and Media Preparation service. The STEC serotypes were kindly provided by Dr. Mohamed Karmali. Published with permission of the Director of VIDO as journal series number 729.

Author Contributions

Conceived and designed the experiments: TSD HGT AAP. Performed the experiments: TSD. Analyzed the data: TSD HGT AAP. Contributed reagents/materials/analysis tools: TSD HGT AAP. Wrote the paper: TSD HGT AAP.

References

1. Ho NK, Henry AC, Johnson-Henry K, Sherman PM (2013) Pathogenicity, host responses and implications for management of enterohemorrhagic *Escherichia coli* O157:H7 infection. *Can J Gastroenterol* 27: 281–285. PMID: [23712303](#)
2. Beutin L, Kaulfuss S, Herold S, Oswald E, Schmidt H (2005) Genetic analysis of enteropathogenic and enterohemorrhagic *Escherichia coli* serogroup O103 strains by molecular typing of virulence and housekeeping genes and pulsed-field gel electrophoresis. *J Clin Microbiol* 43: 1552–1563. PMID: [15814965](#)
3. Schimmer B, Nygard K, Eriksen HM, Lassen J, Lindstedt BA, Brandal LT, et al. (2008) Outbreak of haemolytic uraemic syndrome in Norway caused by stx2-positive *Escherichia coli* O103:H25 traced to cured mutton sausages. *BMC Infect Dis* 8: 41. doi: [10.1186/1471-2334-8-41](#) PMID: [18387178](#)

4. Gyles CL (2007) Shiga toxin-producing *Escherichia coli*: an overview. *J Anim Sci* 85: E45–62. PMID: [17085726](#)
5. Vande Walle K, Vanrompay D, Cox E (2013) Bovine innate and adaptive immune responses against *Escherichia coli* O157:H7 and vaccination strategies to reduce faecal shedding in ruminants. *Vet Immunol Immunopathol* 152: 109–120. doi: [10.1016/j.vetimm.2012.09.028](#) PMID: [23084625](#)
6. Garmendia J, Frankel G, Crepin VF (2005) Enteropathogenic and enterohemorrhagic *Escherichia coli* infections: translocation, translocation, translocation. *Infect Immun* 73: 2573–2585. PMID: [15845459](#)
7. Tobe T, Beatson SA, Taniguchi H, Abe H, Bailey CM, Fivian A, et al. (2006) An extensive repertoire of type III secretion effectors in *Escherichia coli* O157 and the role of lambdaoid phages in their dissemination. *Proc Natl Acad Sci U S A* 103: 14941–14946. PMID: [16990433](#)
8. Dziva F, van Diemen PM, Stevens MP, Smith AJ, Wallis TS (2004) Identification of *Escherichia coli* O157: H7 genes influencing colonization of the bovine gastrointestinal tract using signature-tagged mutagenesis. *Microbiology* 150: 3631–3645. PMID: [15528651](#)
9. Naylor SW, Roe AJ, Nart P, Spears K, Smith DG, Low JC, et al. (2005) *Escherichia coli* O157: H7 forms attaching and effacing lesions at the terminal rectum of cattle and colonization requires the LEE4 operon. *Microbiology* 151: 2773–2781. PMID: [16079353](#)
10. Sharma VK, Sacco RE, Kunkle RA, Bearson SM, Palmquist DE (2012) Correlating levels of type III secretion and secreted proteins with fecal shedding of *Escherichia coli* O157:H7 in cattle. *Infect Immun* 80: 1333–1342. doi: [10.1128/IAI.05869-11](#) PMID: [2252878](#)
11. Peterson RE, Klopfenstein TJ, Moxley RA, Erickson GE, Hinkley S, Bretschneider G, et al. (2007) Effect of a vaccine product containing type III secreted proteins on the probability of *Escherichia coli* O157:H7 fecal shedding and mucosal colonization in feedlot cattle. *J Food Prot* 70: 2568–2577. PMID: [18044436](#)
12. Potter AA, Klashinsky S, Li Y, Frey E, Townsend H, Rogan D, et al. (2004) Decreased shedding of *Escherichia coli* O157:H7 by cattle following vaccination with type III secreted proteins. *Vaccine* 22: 362–369. PMID: [14670317](#)
13. Van Donkersgoed J, Hancock D, Rogan D, Potter AA (2005) *Escherichia coli* O157:H7 vaccine field trial in 9 feedlots in Alberta and Saskatchewan. *Can Vet J* 46: 724–728. PMID: [16187717](#)
14. Peterson RE, Klopfenstein TJ, Moxley RA, Erickson GE, Hinkley S, Rogan D, et al. (2007) Efficacy of dose regimen and observation of herd immunity from a vaccine against *Escherichia coli* O157:H7 for feedlot cattle. *J Food Prot* 70: 2561–2567. PMID: [18044435](#)
15. Smith DR, Moxley RA, Peterson RE, Klopfenstein TJ, Erickson GE, Clowser SL (2008) A two-dose regimen of a vaccine against *Escherichia coli* O157:H7 type III secreted proteins reduced environmental transmission of the agent in a large-scale commercial beef feedlot clinical trial. *Foodborne Pathog Dis* 5: 589–598. doi: [10.1089/fpd.2008.0080](#) PMID: [18681791](#)
16. Smith DR, Moxley RA, Peterson RE, Klopfenstein TJ, Erickson GE, Bretschneider G, et al. (2009) A two-dose regimen of a vaccine against type III secreted proteins reduced *Escherichia coli* O157:H7 colonization of the terminal rectum in beef cattle in commercial feedlots. *Foodborne Pathog Dis* 6: 155–161. doi: [10.1089/fpd.2008.0136](#) PMID: [19105625](#)
17. Smith DR, Moxley RA, Klopfenstein TJ, Erickson GE (2009) A randomized longitudinal trial to test the effect of regional vaccination within a cattle feedyard on *Escherichia coli* O157:H7 rectal colonization, fecal shedding, and hide contamination. *Foodborne Pathog Dis* 6: 885–892. doi: [10.1089/fpd.2009.0299](#) PMID: [19618995](#)
18. Moxley RA, Smith DR, Luebke M, Erickson GE, Klopfenstein TJ, Rogan D (2009) *Escherichia coli* O157:H7 vaccine dose-effect in feedlot cattle. *Foodborne Pathog Dis* 6: 879–884. doi: [10.1089/fpd.2009.0297](#) PMID: [19737064](#)
19. Allen KJ, Rogan D, Finlay BB, Potter AA, Asper DJ (2011) Vaccination with type III secreted proteins leads to decreased shedding in calves after experimental infection with *Escherichia coli* O157. *Can J Vet Res* 75: 98–105. PMID: [21731179](#)
20. Asper DJ, Karmali MA, Townsend H, Rogan D, Potter AA (2011) Serological response of Shiga toxin-producing *Escherichia coli* type III secreted proteins in sera from vaccinated rabbits, naturally infected cattle, and humans. *Clin Vaccine Immunol* 18: 1052–1057. doi: [10.1128/CVI.00068-11](#) PMID: [21593239](#)
21. Asper DJ, Sekirov I, Finlay BB, Rogan D, Potter AA (2007) Cross reactivity of enterohemorrhagic *Escherichia coli* O157:H7-specific sera with non-O157 serotypes. *Vaccine* 25: 8262–8269. PMID: [17980466](#)
22. Karmali MA, Mascarenhas M, Shen S, Ziebell K, Johnson S, Reid-Smith R, et al. (2003) Association of genomic O island 122 of *Escherichia coli* EDL 933 with verocytotoxin-producing *Escherichia coli* seropathotypes that are linked to epidemic and/or serious disease. *J Clin Microbiol* 41: 4930–4940. PMID: [14605120](#)

23. Tarr PI, Neill MA, Clausen CR, Newland JW, Neill RJ, Moseley SL (1989) Genotypic variation in pathogenic *Escherichia coli* O157:H7 isolated from patients in Washington, 1984–1987. *J Infect Dis* 159: 344–347. PMID: [2644374](#)
24. Wadolkowski EA, Burriss JA, O'Brien AD (1990) Mouse model for colonization and disease caused by enterohemorrhagic *Escherichia coli* O157:H7. *Infect Immun* 58: 2438–2445. PMID: [2196227](#)
25. Zhang XH, He KW, Zhang SX, Lu WC, Zhao PD, Luan XT, et al. (2011) Subcutaneous and intranasal immunization with Stx2B-Tir-Stx1B-Zot reduces colonization and shedding of *Escherichia coli* O157:H7 in mice. *Vaccine* 29: 3923–3929. doi: [10.1016/j.vaccine.2011.02.007](#) PMID: [21338683](#)
26. Smith JL, Fratamico PM, Gunther NWt (2014) Shiga toxin-producing *Escherichia coli*. *Adv Appl Microbiol* 86: 145–197. doi: [10.1016/B978-0-12-800262-9.00003-2](#) PMID: [24377855](#)
27. Gill A, Gill CO (2010) Non-O157 verotoxigenic *Escherichia coli* and beef: a Canadian perspective. *Can J Vet Res* 74: 161–169. PMID: [20885839](#)
28. La Ragione RM, Patel S, Maddison B, Woodward MJ, Best A, Whitlam GC, et al. (2006) Recombinant anti-EspA antibodies block *Escherichia coli* O157:H7-induced attaching and effacing lesions in vitro. *Microbes Infect* 8: 426–433. PMID: [16298154](#)
29. Mohawk KL, O'Brien AD (2011) Mouse models of *Escherichia coli* O157:H7 infection and shiga toxin injection. *J Biomed Biotechnol* 2011: 258185. doi: [10.1155/2011/258185](#) PMID: [21274267](#)
30. Babiuk S, Asper DJ, Rogan D, Mutwiri GK, Potter AA (2008) Subcutaneous and intranasal immunization with type III secreted proteins can prevent colonization and shedding of *Escherichia coli* O157:H7 in mice. *Microb Pathog* 45: 7–11. doi: [10.1016/j.micpath.2008.01.005](#) PMID: [18487034](#)
31. Gao X, Cai K, Li T, Wang Q, Hou X, Tian R, et al. (2011) Novel fusion protein protects against adherence and toxicity of enterohemorrhagic *Escherichia coli* O157:H7 in mice. *Vaccine* 29: 6656–6663. doi: [10.1016/j.vaccine.2011.06.106](#) PMID: [21742003](#)
32. Cai K, Gao X, Li T, Wang Q, Hou X, Tu W, et al. (2011) Enhanced immunogenicity of a novel Stx2Am-Stx1B fusion protein in a mice model of enterohemorrhagic *Escherichia coli* O157:H7 infection. *Vaccine* 29: 946–952. doi: [10.1016/j.vaccine.2010.11.035](#) PMID: [21134452](#)
33. Amani J, Salmanian AH, Rafati S, Mousavi SL (2010) Immunogenic properties of chimeric protein from espA, eae and tir genes of *Escherichia coli* O157:H7. *Vaccine* 28: 6923–6929. doi: [10.1016/j.vaccine.2010.07.061](#) PMID: [20709010](#)
34. Abe A, de Grado M, Pfuetzner RA, Sanchez-Sanmartin C, Devinney R, Puente JL, et al. (1999) Enteropathogenic *Escherichia coli* translocated intimin receptor, Tir, requires a specific chaperone for stable secretion. *Mol Microbiol* 33: 1162–1175. PMID: [10510231](#)
35. McNeilly TN, Mitchell MC, Rosser T, McAteer S, Low JC, Smith DG, et al. (2010) Immunization of cattle with a combination of purified intimin-531, EspA and Tir significantly reduces shedding of *Escherichia coli* O157:H7 following oral challenge. *Vaccine* 28: 1422–1428. doi: [10.1016/j.vaccine.2009.10.076](#) PMID: [19903545](#)
36. Vilte DA, Larzabal M, Garbaccio S, Gammella M, Rabinovitz BC, Elizondo AM, et al. (2011) Reduced faecal shedding of *Escherichia coli* O157:H7 in cattle following systemic vaccination with gamma-intimin C(2)(8)(0) and EspB proteins. *Vaccine* 29: 3962–3968. doi: [10.1016/j.vaccine.2011.03.079](#) PMID: [21477674](#)
37. Vogstad AR, Moxley RA, Erickson GE, Klopfenstein TJ, Smith DR (2014) Stochastic simulation model comparing distributions of STEC O157 faecal shedding prevalence between cattle vaccinated with type III secreted protein vaccines and non-vaccinated cattle. *Zoonoses Public Health* 61: 283–289. doi: [10.1111/zph.12069](#) PMID: [23826923](#)
38. Vogstad AR, Moxley RA, Erickson GE, Klopfenstein TJ, Smith DR (2013) Assessment of heterogeneity of efficacy of a three-dose regimen of a type III secreted protein vaccine for reducing STEC O157 in feces of feedlot cattle. *Foodborne Pathog Dis* 10: 678–683. doi: [10.1089/fpd.2012.1374](#) PMID: [23692077](#)
39. Boland KG, Hayles AN, Miller CB, Kerr T, Brown WC, Lahmers KK (2013) Regional immune response to immunization with *Escherichia coli* O157:H7-derived intimin in cattle. *Clin Vaccine Immunol* 20: 562–571. doi: [10.1128/CVI.00743-12](#) PMID: [23408521](#)
40. Matthews L, Reeve R, Gally DL, Low JC, Woolhouse ME, McAteer SP, et al. (2013) Predicting the public health benefit of vaccinating cattle against *Escherichia coli* O157. *Proc Natl Acad Sci U S A* 110: 16265–16270. doi: [10.1073/pnas.1304978110](#) PMID: [24043803](#)
41. Fox JT, Thomson DU, Drouillard JS, Thornton AB, Burkhardt DT, Emery DA, et al. (2009) Efficacy of *Escherichia coli* O157:H7 siderophore receptor/porin proteins-based vaccine in feedlot cattle naturally shedding *E. coli* O157. *Foodborne Pathog Dis* 6: 893–899. doi: [10.1089/fpd.2009.0336](#) PMID: [19737065](#)

42. Thomson DU, Loneragan GH, Thornton AB, Lechtenberg KF, Emery DA, Burkhardt DT, et al. (2009) Use of a siderophore receptor and porin proteins-based vaccine to control the burden of *Escherichia coli* O157:H7 in feedlot cattle. *Foodborne Pathog Dis* 6: 871–877. doi: [10.1089/fpd.2009.0290](https://doi.org/10.1089/fpd.2009.0290) PMID: [19737063](https://pubmed.ncbi.nlm.nih.gov/19737063/)
43. Cernicchiaro N, Renter DG, Cull CA, Paddock ZD, Shi X, Nagaraja TG (2014) Fecal shedding of non-O157 serogroups of Shiga toxin-producing *Escherichia coli* in feedlot cattle vaccinated with an *Escherichia coli* O157:H7 SRP vaccine or fed a *Lactobacillus*-based direct-fed microbial. *J Food Prot* 77: 732–737. doi: [10.4315/0362-028X.JFP-13-358](https://doi.org/10.4315/0362-028X.JFP-13-358) PMID: [24780326](https://pubmed.ncbi.nlm.nih.gov/24780326/)
44. Paddock ZD, Renter DG, Cull CA, Shi X, Bai J, Nagaraja TG (2014) *Escherichia coli* O26 in feedlot cattle: fecal prevalence, isolation, characterization, and effects of an *E. coli* O157 vaccine and a direct-fed microbial. *Foodborne Pathog Dis* 11: 186–193. doi: [10.1089/fpd.2013.1659](https://doi.org/10.1089/fpd.2013.1659) PMID: [24286301](https://pubmed.ncbi.nlm.nih.gov/24286301/)