

Impact of Severe Acute Respiratory Syndrome Coronavirus 2 Viral Load on Risk of Intubation and Mortality Among Hospitalized Patients With Coronavirus Disease 2019

Reed Magleby,^{1,a} Lars F. Westblade,^{1,2,3,a} Alex Trzebucki,¹ Matthew S. Simon,^{1,2} Mangala Rajan,² Joel Park,^{1,4} Parag Goyal,^{1,2} Monika M. Safford,^{1,2} and Michael J. Satlin^{1,2}

¹NewYork–Presbyterian Hospital/Weill Cornell Medical Center, New York, New York, USA, ²Department of Medicine, Weill Cornell Medicine, New York, New York, USA, ³Department of Pathology and Laboratory Medicine, Weill Cornell Medicine, New York, New York, USA, and ⁴Department of Emergency Medicine, Weill Cornell Medicine, New York, New York, USA

(See the Editorial Commentary by Mathers on pages e4206–7.)

Background. Patients hospitalized with coronavirus disease 2019 (COVID-19) frequently require mechanical ventilation and have high mortality rates. However, the impact of viral burden on these outcomes is unknown.

Methods. We conducted a retrospective cohort study of patients hospitalized with COVID-19 from 30 March 2020 to 30 April 2020 at 2 hospitals in New York City. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) viral load was assessed using cycle threshold (Ct) values from a reverse transcription-polymerase chain reaction assay applied to nasopharyngeal swab samples. We compared characteristics and outcomes of patients with high, medium, and low admission viral loads and assessed whether viral load was independently associated with intubation and in-hospital mortality.

Results. We evaluated 678 patients with COVID-19. Higher viral load was associated with increased age, comorbidities, smoking status, and recent chemotherapy. In-hospital mortality was 35.0% (Ct <25; n = 220), 17.6% (Ct 25–30; n = 216), and 6.2% (Ct >30; n = 242) with high, medium, and low viral loads, respectively ($P < .001$). The risk of intubation was also higher in patients with a high viral load (29.1%) compared with those with a medium (20.8%) or low viral load (14.9%; $P < .001$). High viral load was independently associated with mortality (adjusted odds ratio [aOR], 6.05; 95% confidence interval [CI], 2.92–12.52) and intubation (aOR, 2.73; 95% CI, 1.68–4.44).

Conclusions. Admission SARS-CoV-2 viral load among hospitalized patients with COVID-19 independently correlates with the risk of intubation and in-hospital mortality. Providing this information to clinicians could potentially be used to guide patient care.

Keywords. SARS-CoV-2; coronavirus disease 2019; viral load; hospitalized patients; outcomes.

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a novel pathogen that has rapidly caused a devastating pandemic of coronavirus disease 2019 (COVID-19). As of 10 June 2020, SARS-CoV-2 had infected more than 7 million people and killed more than 400 000 people throughout the world [1]. Although the majority of patients who develop COVID-19 have mild presentations [2], 18%–33% of patients who are hospitalized require mechanical ventilation and up to 20% of hospitalized patients die [3–7]. Investigations of risk factors for intubation and mortality with COVID-19 in hospitalized patients have largely focused on patient characteristics, such as older age, obesity, and comorbidities, as well as presenting symptoms and laboratory parameters [5, 7–9]. In

contrast, the impact of SARS-CoV-2 viral load on clinical outcomes in hospitalized patients has not been thoroughly investigated. In 2 studies of hospitalized patients in China, those with severe presentations of COVID-19 had higher viral loads than those with mild presentations. However, the impact of SARS-CoV-2 viral load on the risk of intubation or death was not evaluated [10, 11].

The current standard-of-care test to diagnose COVID-19 is to collect a nasopharyngeal (NP) swab and use a reverse transcription-polymerase chain reaction (RT-PCR) assay to detect SARS-CoV-2 RNA [12]. These RT-PCR assays provide clinicians only with information on whether SARS-CoV-2 is detected or not detected. However, these assays also contain quantitative information on cycle threshold (Ct) values that are inversely correlated with viral load and are not reported clinically. We hypothesized that assessing SARS-CoV-2 viral load by analyzing Ct values from an initial NP swab sample could be a clinically valuable tool for identifying patients at highest risk of intubation and death and provide insights into the pathogenesis of COVID-19. We therefore conducted this retrospective analysis of SARS-CoV-2 viral loads on admission, clinical

Received 26 May 2020; editorial decision 16 June 2020; accepted 26 June 2020; published online June 30, 2020.

^aR. M. and L. F. W. contributed equally to this work.

Correspondence: M. J. Satlin, Weill Cornell Medical College, 1300 York Avenue, A-421, New York, NY 10021 (mjs9012@med.cornell.edu).

Clinical Infectious Diseases® 2021;73(11):e4197–205

© The Author(s) 2020. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com.
DOI: 10.1093/cid/ciaa851

presentations, and outcomes at 2 affiliated New York City hospitals using a high-throughput RT-PCR assay.

METHODS

Study Population and Setting

This retrospective, observational study consisted of all patients who were hospitalized at NewYork–Presbyterian Hospital/Weill Cornell Medical Center and affiliated Lower Manhattan Hospital and had an NP swab sample collected and analyzed for SARS-CoV-2 using the cobas 6800 RT-PCR System (Roche Molecular Systems, Inc., Branchburg, NJ) between 30 March 2020 and 30 April 2020. The predominant NP swab collection and transport kits used were the BD Universal Viral Transport System (Becton, Dickinson and Company, Franklin Lakes, NJ) and the Universal Transport Medium (Hardy Diagnostics, Santa Maria, CA). Patients who did not have an NP swab sample collected and analyzed within 1 day of hospital admission or whose sample was analyzed on a different diagnostic platform or at a different institution were excluded. The policy during the study period was to only perform SARS-CoV-2 tests in patients who were thought to require hospital admission. However, some patients who were tested were subsequently discharged from the emergency department (ED) without hospital admission.

Viral Load Assessment

The US Food and Drug Administration provided emergency use authorization approval for the cobas SARS-CoV-2 RT-PCR test, and the test was performed according to the manufacturer's instructions [13]. This assay amplifies 2 targets within the SARS-CoV-2 genome: ORF1ab, a SARS-CoV-2–specific target, and the E gene, a pan-Sarbecovirus target that is present in SARS-CoV-2 and SARS-CoV but not in seasonal coronaviruses or Middle East respiratory syndrome CoV. For routine clinical care, results are classified as detected if either the ORF1ab or E gene is detected or as not detected if neither target is detected. However, the instrument also generates a Ct value for each target that correlates inversely with quantitative viral load and is not released to clinicians. The Ct value represents the number of replication cycles required for sufficient gene amplification to produce a fluorescent signal that crosses a predefined threshold.

For this study, we reviewed Ct values for both gene targets for all initial SARS-CoV-2 tests that were performed on NP swab samples that were collected from study participants for routine clinical care within 1 day of hospital admission. We separated the Ct values for the SARS-CoV-2–specific target (ORF1ab) into terciles based on the quantitative values. We then designated high viral load samples as the lowest Ct tercile, medium viral load samples as the middle tercile, and low viral load samples as the highest tercile. Specimens that were designated positive for SARS-CoV-2 but for which only the E gene was detected were designated low viral load samples.

Data Collection

Data were retrospectively abstracted manually from the electronic medical record using a quality-controlled protocol and entered into a REDCap database [14]. All data collectors were trained, and a random resampling of data previously showed high interrater reliability (mean Cohen's kappa of 0.92) [5]. Data included demographics, comorbidities, social characteristics, selected outpatient medications on admission, presenting symptoms on arrival to the hospital, oxygen supplementation required within 3 hours of presentation, laboratory parameters, chest radiograph findings, concurrent bloodstream infections, in-hospital complications, and in-hospital mortality. Clinical data after hospital discharge were not consistently available, thus only outcomes that occurred during the hospital admission were analyzed. The Weill Cornell Medicine Institutional Review Board approved the study with a waiver of informed consent.

Statistical Analyses

We compared baseline characteristics and outcomes of hospitalized patients with COVID-19 who had high, medium, and low initial viral loads using the nonparametric nptrend command in Stata (StataCorp, College Station, TX) that tests for trend across ordered groups. Continuous variables were represented with medians and interquartile ranges and categorical variables were represented as proportions. A 2-sided *P* value of $\leq .05$ was used to designate statistical significance. The risk of in-hospital intubation and death was also compared across 8 numerical Ct value ranges. We also constructed Cox proportional hazards models to compare the cumulative risks of intubation and death during the inpatient admission among patients with high, medium, and low viral loads. We then identified baseline factors that were associated with in-hospital mortality and intubation using univariate logistic regression models. All variables that were statistically significantly associated with each outcome were then entered into separate multivariate logistic regression models. Adjusted odds ratios of mortality and intubation were calculated for each of these variables with 95% confidence intervals (CIs). Analyses were conducted using Stata, version 15.0.

RESULTS

Cycle Threshold Values and Establishment of Viral Load Categories

A total of 678 NP swab samples were available for analysis from unique hospitalized patients who met the study inclusion criteria (Figure 1). Ct values for the ORF1ab locus ranged from 14.3 to 36.4 (Supplementary Figure), and 10 samples were considered positive for SARS-CoV-2 based on detection of the E gene even though ORF1ab was not detected. The median Ct value was 27.9. The lowest third of Ct values were <25.2 , the middle third were between 25.2 and 30.3, and the highest third were >30.3 . For simplicity, we designated high viral load samples to have Ct values <25 ($n = 220$), medium viral load samples




Figure 1. Study flow diagram. Abbreviations: ED, emergency department; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

to have Ct values between 25 and 30 ($n = 216$), and low viral load samples to have Ct values >30 or for which only the E gene was detected ($n = 242$). Of the 49 patients who had a positive SARS-CoV-2 test but were discharged from the ED and not admitted, the median Ct value was 29.1.

Patient Characteristics and Presentations Stratified by Viral Load

The median age of patients with high, medium, and low viral loads was 72, 69, and 63 years, respectively ($P < .001$; [Table 1](#)). In addition to older age, patients with higher viral loads were more likely to have coronary artery disease, congestive heart failure, cerebrovascular disease, hypertension, chronic obstructive pulmonary disease (COPD), chronic kidney disease, and active cancer. They were also more likely to be a former or current smoker or have received recent chemotherapy. Patients with high viral loads had a median of 7 days from symptom onset until hospital admission compared with 8 and 10 days for patients with medium and low viral loads, respectively ($P < .001$). Patients with higher viral loads were also more likely to require oxygen by a nonrebreather, high-flow nasal cannula or mechanical ventilation within 3 hours of presentation to the ED but were less likely to present with fever, nausea, or vomiting. Lymphopenia, anemia, and thrombocytopenia were more common among patients with higher viral loads, whereas alanine aminotransferase elevations were less common. There were no differences in chest X-ray findings among patients with high, medium, or low viral loads. There were also no differences in viral loads among different racial or ethnic categories or between patients who did and did not use angiotensin-converting

enzyme inhibitors (ACEIs), angiotensin II receptor blockers (ARBs), or hydroxychloroquine.

Outcomes Stratified by Viral Load

The last day of study follow-up was 8 June 2020. By that day, 19.2% of patients had died during their admission, 75.8% had been discharged alive, 1.6% had been transferred to another hospital, and 3.4% were still hospitalized. The risk of intubation and death increased with higher viral loads. In-hospital mortality was 35.0% in patients with a high viral load compared with 17.6% in patients with a medium viral load and 6.2% in patients with a low viral load ($P < .001$; [Table 1](#)). The risk of intubation was 29.1% in patients with a high viral load compared with 20.8% and 14.9% in patients with a medium or low viral load, respectively ($P < .001$; [Table 1](#)). These associations were also observed in time-based analyses ([Figure 2](#)) in which, compared with a low viral load, a high viral load was associated with a hazard ratio (HR) of 5.06 (95% CI, 2.86–8.96; $P < .001$) for in-hospital mortality and an HR of 2.15 (95% CI, 1.31–3.53; $P = .003$) for intubation, whereas a medium viral load was associated with mortality (HR, 2.52; 95% CI, 1.36–4.67; $P = .003$) but not with risk of intubation (HR, 1.53; 95% CI, .91–2.60; $P = .11$). When viral load was assessed by continuous Ct values instead of being grouped into high, medium, and low viral load categories, the risk of death increased with decreasing Ct values and the risk of intubation was greater with Ct values <27 compared with >27 ([Figure 3](#)).

In a multivariate model that adjusted for age, race, coronary artery disease, congestive heart failure, cerebrovascular

Table 1. Baseline Characteristics of Hospitalized Patients With Coronavirus 2019 Stratified by Admission Viral Load

Variable	High Viral Load (Ct <25), n = 220	Medium Viral Load (Ct 25–30), n = 216	Low Viral Load (Ct >30), n = 242	P Value ^a
Age, years	72 (60–81)	69 (58–79)	63 (50–73)	<.001
Female gender	81 (36.8)	84 (38.9)	99 (40.9)	.37
Race ^b (n = 585)				
White	89 (45.1)	79 (43.4)	91 (44.2)	.54
Black	25 (12.7)	27 (14.8)	30 (14.6)	.74
Asian	42 (21.3)	32 (17.6)	36 (17.5)	.23
Hispanic ethnicity ^b (n = 628)	48 (23.9)	54 (27.1)	60 (26.3)	.58
Comorbidities				
Obesity: body mass index >30 ^b (n = 663)	60 (27.6)	70 (33.5)	79 (33.3)	.20
Coronary artery disease	44 (20.0)	45 (21.8)	31 (12.8)	.039
Congestive heart failure	28 (12.7)	15 (6.9)	13 (5.4)	.004
Cerebrovascular disease	27 (12.3)	15 (6.9)	13 (5.4)	.007
Diabetes	82 (37.3)	71 (32.9)	77 (31.8)	.22
Hypertension	136 (61.8)	144 (66.7)	121 (50.0)	.008
Chronic pulmonary disease	49 (22.3)	40 (18.5)	38 (15.7)	.07
Chronic obstructive pulmonary disease	21 (9.6)	11 (5.1)	9 (3.7)	.009
Asthma	20 (9.1)	22 (10.1)	20 (8.3)	.75
Chronic kidney disease	32 (14.6)	22 (10.2)	20 (8.3)	.032
End-stage renal disease	22 (10.0)	13 (6.0)	12 (5.0)	.035
Human immunodeficiency virus infection	4 (1.8)	3 (1.4)	5 (2.1)	.83
Active cancer	22 (10.0)	14 (6.5)	6 (2.5)	.001
Solid tumor	13 (5.9)	8 (3.7)	5 (2.1)	.032
Hematologic malignancy	9 (4.1)	6 (2.8)	2 (0.8)	.025
Transplant recipient	15 (6.8)	8 (3.7)	9 (3.7)	.12
Rheumatologic disease	12 (5.5)	13 (6.0)	12 (5.0)	.81
Social characteristics				
Former or current smoker	78 (35.5)	54 (25.0)	62 (25.6)	.022
Known sick contacts	41 (18.6)	41 (19.0)	34 (14.1)	.18
Healthcare worker	7 (3.2)	10 (4.6)	9 (3.7)	.80
Home medications				
Angiotensin-converting enzyme inhibitor/angiotensin II receptor blocker	68 (30.9)	74 (34.3)	68 (28.1)	.49
Hydroxychloroquine	5 (2.3)	1 (0.5)	9 (3.7)	.27
Immunosuppressive medications	27 (12.3)	21 (9.7)	19 (7.9)	.11
Oral steroids	13 (5.9)	11 (5.1)	9 (3.7)	.19
Calcineurin inhibitor	11 (5.0)	7 (3.2)	9 (3.7)	.49
Mycophenolate	12 (5.5)	7 (3.2)	9 (3.7)	.36
Chemotherapy within the previous 6 months	9 (4.1)	3 (1.4)	1 (0.4)	.004
Clinical presentation				
Days of symptoms prior to admission^b (n = 611)	7 (3–9)	8 (5–13)	10 (6–14)	<.001
Symptoms				
Fever	138 (62.7)	152 (70.4)	173 (71.5)	.046
Cough	160 (72.7)	148 (68.5)	166 (68.6)	.34
Dyspnea	156 (70.9)	136 (63.0)	169 (69.8)	.85
Sore throat	18 (8.2)	17 (7.9)	11 (4.6)	.12
Headache	16 (7.3)	15 (6.9)	25 (10.3)	.23
Myalgias	38 (17.3)	41 (19.0)	54 (22.3)	.17
Nausea or vomiting	30 (13.6)	31 (14.3)	51 (21.1)	.03
Diarrhea	49 (22.3)	50 (23.2)	68 (28.1)	.14
Altered mental status	38 (17.3)	41 (19.0)	29 (12.0)	.11
Anosmia	7 (3.2)	12 (5.6)	14 (5.8)	.20
Ageusia	12 (5.5)	20 (9.3)	20 (8.3)	.27
Highest oxygen requirement within 3 hours of arrival to the emergency department				
No supplemental oxygen required	78 (35.5)	84 (38.9)	92 (38.0)	.58
Oxygen by low-flow nasal cannula	81 (36.8)	86 (39.8)	113 (46.7)	.03

Table 1. Continued

Variable	High Viral Load (Ct <25), n = 220	Medium Viral Load (Ct 25–30), n = 216	Low Viral Load (Ct >30), n = 242	P Value ^a
Oxygen by nonrebreather mask, high-flow nasal cannula, or noninvasive mechanical ventilation	50 (22.7)	38 (17.6)	33 (13.6)	.011
Mechanical ventilation	11 (5.0)	8 (3.7)	4 (1.7)	.046
Laboratory values				
Leukocytosis: white blood cell count >11 × 10 ⁹ cells/L ^b (n = 602)	41 (22.2)	39 (19.8)	40 (18.6)	.46
Lymphopenia: absolute lymphocyte count < 1 × 10⁹ cells/L^b (n = 594)	135 (71.8)	140 (72.5)	127 (59.6)	.008
Anemia: hemoglobin <12 g/dL^b (n = 602)	52 (27.4)	45 (22.8)	41 (19.1)	.048
Thrombocytopenia: platelet count < 150 × 10⁹/L^b (n = 602)	46 (24.2)	45 (22.8)	26 (12.1)	.002
Aspartate aminotransferase elevation ^{b,c} (n = 580)	118 (64.5)	144 (75.4)	142 (68.9)	.38
Alanine aminotransferase elevation^{b,d} (n = 589)	46 (24.7)	44 (22.8)	79 (37.6)	.004
Troponin I >0.5 ng/mL ^b (n = 229)	17 (14.5)	9 (8.4)	5 (6.8)	.07
Inflammatory makers				
Procalcitonin, ng/mL ^b (n = 504)	0.26 (0.12–0.64)	0.22 (0.1–0.56)	0.2 (0.11–0.45)	.12
C-reactive protein, mg/dL ^b (n = 373)	12.9 (7.8–20.4)	11.1 (6.3–19.0)	11.4 (6.7–20.2)	.57
Ferritin, ng/mL ^b (n = 521)	849 (409–1417)	842 (409–1542)	821 (340–1361)	.42
D-dimer, ng/mL ^b (n = 405)	535 (309–990)	472 (300–980)	547 (354–1716)	.17
Chest X-ray findings				
Clear	22 (10.0)	13 (6.0)	17 (7.0)	.24
Unilateral infiltrates	27 (12.3)	23 (10.7)	25 (10.3)	.51
Bilateral infiltrates	166 (75.5)	169 (78.2)	194 (80.2)	.22
Concurrent bloodstream infection	8 (3.6)	5 (2.3)	3 (1.2)	.09
Outcomes				
Intubation	64 (29.1)	45 (20.8)	36 (14.9)	<.001
Days until intubation	2 (0–3)	2 (0–4)	2 (0–5)	.66
In-hospital mortality	77 (35.0)	36 (17.6)	14 (6.2)	<.001
Days until death	7 (4–14)	8 (3–15)	10 (3–32)	.32
Other complications				
Myocardial infarction	16 (7.3)	10 (4.6)	5 (2.1)	.007
Congestive heart failure	14 (6.4)	6 (2.8)	6 (2.5)	.032
Arrhythmia	29 (13.2)	18 (8.3)	20 (8.3)	.08
Acute kidney injury requiring hemodialysis	33 (15.0)	18 (8.3)	7 (2.9)	<.001

Variables are expressed as number (%) or median (interquartile range). Bolded values indicated variables with statistically significant associations.

Abbreviation: Ct, cycle threshold.

^aP values were calculated using the nonparametric nptrend command in Stata, version 15.0, that tests for trend across ordered groups.

^bThis variable was not assessed in all participants. The denominator is listed next to the variable.

^cAspartate aminotransferase elevation indicates a value >34 units/L.

^dAlanine aminotransferase elevation indicates a value >55 units/L.

disease, hypertension, COPD, days of symptoms prior to admission, symptoms upon presentation, initial chest X-ray findings, and level of oxygen support within 3 hours of arrival to the ED (Table 2), having a high viral load was independently associated with increased risk of in-hospital mortality (adjusted odds ratio [aOR], 6.05; 95% CI, 2.92–12.52; $P < .001$) compared with having a low viral load. The risk of in-hospital mortality was also higher in patients with a medium viral load compared with a low viral load, but this association was not statistically significant (aOR, 2.06; 95% CI, .98–4.34; $P = .058$). Compared with those with a low viral load, having a high viral load was also independently associated with increased risk of intubation (aOR, 2.73; 95% CI, 1.68–4.44; $P < .001$), whereas the risk of intubation associated with a medium viral load did not reach statistical significance (aOR, 1.59; 95% CI, .96–2.63; $P = .07$).

Patients with higher viral loads were also more likely to develop myocardial infarction, congestive heart failure, and acute kidney injury requiring hemodialysis (Table 1).

DISCUSSION

In this study, we demonstrate that patients who were admitted to the hospital with high SARS-CoV-2 viral loads, as assessed by Ct values of NP swab samples, were more likely to be intubated or die during their hospitalization. This association persisted even when we adjusted for age, comorbidities, presenting symptoms, chest radiography findings, and degree of presenting hypoxia.

Prior studies have indicated that viral load correlates with severity of COVID-19 presentation [10, 11], and our study of a larger cohort of hospitalized patients adds to this knowledge




Figure 2. Probability of in-hospital survival (A) and freedom from intubation (B) during COVID-19 hospitalization among patients with high, medium, and low viral loads. A medium viral load was associated with a hazard ratio (HR) of in-hospital mortality of 2.52 (95% confidence interval [CI], 1.36–4.67; $P = .003$) and a HR of intubation of 1.53 (95% CI, .91–2.60; $P = .11$) compared with a low viral load. A high viral load was associated with a HR of in-hospital mortality of 5.06 (95% CI, 2.86–8.96; $P < .001$) and a HR of intubation of 2.15 (95% CI, 1.31–3.53; $P = .003$) compared with a low viral load.

base by identifying that admission viral load has important prognostic implications. Reporting SARS-CoV-2 viral load based on Ct values from admission NP swab samples could therefore help identify patients who are at highest risk of adverse outcomes and who therefore may benefit from more intensive monitoring. Identifying patients with high viral loads could also be helpful for allocating scarce therapeutic interventions

such as antiviral agents (eg, remdesivir) [15]. Our findings also suggest that stratification or adjustment for baseline viral load would benefit the design of clinical trials of antiviral agents for COVID-19. It is also possible that viral load could be used along with other factors, such as age, comorbidities, and severity of symptoms and hypoxia, to decide on the need for hospital admission. However, additional studies that evaluate viral loads




Figure 3. Cycle threshold values from nasopharyngeal swab samples on admission and risk of intubation and death during the hospitalization.

and clinical outcomes among all patients who present to the ED are warranted prior to pursuing this strategy clinically.

Older age and the presence of comorbidities such as hypertension, coronary artery disease, congestive heart failure, COPD, and cancer are known to be associated with worse outcomes in COVID-19 [2, 7, 16, 17]. These patients may have decreased cardiopulmonary reserve and thus are less likely to tolerate the physiologic insults caused by COVID-19. Our findings suggest that these patients also have higher SARS-CoV-2 viral loads when they present to the hospital, which may contribute to the worse outcomes observed in these patients. Reasons for higher viral loads specifically in these populations are not well understood and warrant further investigation.

Given that SARS-CoV-2 uses the angiotensin-converting enzyme 2 receptor (ACE2) for entry into host cells [18], there have been concerns that use of ACEIs and ARBs may upregulate ACE2 expression and lead to increased viral proliferation into host cells [19]. Although patients with hypertension and congestive heart failure were more likely to have higher viral loads, use of ACEIs and ARBs was not associated with higher viral load. Our findings are consistent with those from observational studies that have not demonstrated worse outcomes in patients who use ACEIs or ARBs [20–22] and support the recommendations of professional societies of not discontinuing these medications in the setting of COVID-19 [23].

Another notable finding from this study is that there were no differences in admission SARS-CoV-2 viral loads or outcomes among different racial or ethnic groups. In the United States, Hispanic and black communities have been disproportionately affected by COVID-19, with a greater proportion of deaths among these patients than what would be expected

based on their population proportions [24–26]. Our finding that admission viral loads were not different among race and ethnicity groups suggests that these disparities are not related to viral load but instead may be related to comorbid illnesses and nonbiological factors such as social determinants of health. This further underscores the importance of studies that examine the impact of social determinants of health on outcomes during the COVID-19 pandemic.

We also found that patients with higher viral loads were more likely to develop myocardial infarction, congestive heart failure, and acute kidney injury. It is unclear whether these associations were from chance, were related to increased hypoxia in heart and kidney tissue, or were related to increased viral infection of these organs. A recent autopsy study demonstrated that SARS-CoV-2 frequently directly infects both the heart and kidney [27] and that kidney injury and myocardial injury are commonly reported complications of severe COVID-19 [28, 29]. Additional studies are warranted to assess the relationships between viral loads in NP swab samples, disease burden in the heart and kidney, and clinical outcomes.

Our study has limitations. We evaluated only the viral load of a single NP swab sample per patient at the time of hospital admission and thus could not assess viral load dynamics over time or the infectious burden at the time of infection onset. However, we found that this single sample on admission had important prognostic implications among hospitalized patients. In order to maintain consistency, we only analyzed Ct values from a single diagnostic platform, and thus our findings may not apply to all COVID-19 diagnostic assays. However, other studies have demonstrated that the Panther Fusion SARS-CoV-2 test (Hologic, Inc) and the Xpert Xpress SARS-CoV-2 test (Cepheid,

Table 2. Multivariate Logistic Regression Models of Factors Associated With Intubation and In-hospital Mortality

Variable	Intubation		Mortality	
	Adjusted OR (95% CI)	PValue	Adjusted OR (95% CI)	PValue
Age, per year increase			1.10 (1.07–1.13)	<.001
White race			1.00 (.58–1.72)	.99
Obesity	1.41 (.93–2.12)	.10		
Coronary artery disease			1.48 (.79–2.78)	.22
Congestive heart failure			2.10 (.89–4.93)	.09
Cerebrovascular disease			1.24 (.54–2.86)	.61
Hypertension			0.74 (.41–1.33)	.31
Chronic obstructive pulmonary disease			0.65 (.23–1.82)	.42
Use of oral steroids as an outpatient	1.86 (.84–4.12)	.13		
Days of symptoms prior to admission, per day increase			0.97 (.92–1.02)	.23
Symptoms on admission				
Fever	1.37 (.87–2.17)	.18	1.38 (.77–2.46)	.28
Cough			0.64 (.36–1.13)	.12
Dyspnea	1.99 (1.20–3.30)	.008		
Headache			0.56 (.16–1.97)	.37
Myalgias			1.24 (.60–2.58)	.56
Nausea or vomiting			0.50 (.21–1.19)	.12
Altered mental status			1.29 (.65–2.55)	.46
Ageusia			0.88 (.26–3.00)	.84
Highest level of supplemental oxygen within 3 hours of arrival to the emergency department^a				
None			Reference	
Oxygen by nasal cannula			3.79 (1.86–7.73)	<.001
Oxygen by nonbreather mask, high-flow nasal cannula, or noninvasive mechanical ventilation			5.58 (2.50–12.46)	<.001
Mechanical ventilation			23.34 (6.29–86.51)	<.001
Chest X-ray findings				
No infiltrates	Reference		Reference	
Unilateral infiltrates	1.72 (.30–9.84)	.54	4.36 (1.09–17.45)	.037
Bilateral infiltrates	9.94 (2.37–41.74)	.002	4.98 (1.47–16.94)	.01
Viral load by nasopharyngeal swab				
Low viral load (Ct >30)	Reference		Reference	
Medium viral load (Ct 25–30)	1.59 (.96–2.63)	.07	2.06 (.98–4.34)	.058
High viral load (Ct <25)	2.73 (1.68–4.44)	<.001	6.05 (2.92–12.52)	<.001

Only variables that had a significant association with intubation or mortality in a univariate logistic regression model were included in the corresponding multivariate model. Empty cells indicate that the variable was not associated with the corresponding outcome in the univariate logistic regression model. Variables with statistically significant associations are in bold.

Abbreviations: CI, confidence interval; OR, odds ratio.

^aThis variable was analyzed as a risk factor for mortality but was not analyzed as a factor associated with intubation because 1 of the oxygen supplementation categories was mechanical ventilation.

Inc), 2 commonly used RT-PCR assays, yield nearly identical Ct values as the cobas 6800 assay used in this study [30, 31]. Thus, we suspect that our findings may also be applicable to other diagnostic platforms. We encourage others to evaluate the relationship between clinical outcomes and Ct values using other diagnostic platforms and other patient populations. Another potential role for reporting SARS-CoV-2 viral loads through Ct values is to guide the use of isolation precautions, given that viral load correlates with infectivity [32–34]. Our study did not assess this potential use of Ct values, but we believe this is an important area for future investigation. Another limitation is that our study was retrospective and relied on data that were documented in the electronic medical record and thus could have misclassified patient characteristics or outcomes. However, our data abstraction process used a standardized protocol and our

queries identified high interrater reliability for data collection. Last, we focused on in-hospital mortality and did not capture deaths that occurred after discharge from the hospital.

In conclusion, we found that admission SARS-CoV-2 viral loads, as determined by Ct values that are generated with standard-of-care diagnostic assays, are independently associated with intubation and death among hospitalized patients with COVID-19. These findings highlight the critical role of viral load in SARS-CoV-2 pathogenesis and suggest that Ct values should be reported to assist clinicians in identifying patients at high risk for adverse COVID-19-related outcomes.

Supplementary Data

Supplementary materials are available at *Clinical Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility

of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. The authors acknowledge the work of our data abstraction team at Weill Cornell Medicine who are highlighted by name in our supplementary document. We also acknowledge Melissa Cushing, MD, Massimo Loda, MD, and the clinical laboratory scientists who performed the severe acute respiratory syndrome coronavirus 2 laboratory testing.

Financial support. This work was partially supported by the National Center for Advancing Translational Science (UL1 TR002384 to Julianne Imperato-McGinley) at the National Institutes of Health.

Potential conflicts of interest. L. F. W. reports receipt of consulting fees from Roche Molecular Systems, Inc and receipt of grants from BioFire Diagnostics, LLC outside the submitted work. M. S. S. reports receipt of consulting fees from Roche Diagnostics and receipt of grants from Amgen outside the submitted work. M. J. S. reports receipt of grants from BioFire Diagnostics outside the submitted work. All other authors report no potential conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. World Health Organization. Coronavirus disease 2019 (COVID-19): situation report—142. Available at: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200610-covid-19-sitrep-142.pdf?sfvrsn=180898cd_6. Accessed 11 June 2020.
2. Wu Z, McGoogan JM. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-19) outbreak in China: summary of a report of 72314 cases from the Chinese Center for Disease Control and Prevention. *JAMA* 2020 [Epub ahead of print].
3. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA* 2020; 323:2052–9.
4. Gold JAW, Wong KK, Szablewski CM, et al. Characteristics and clinical outcomes of adult patients hospitalized with COVID-19—Georgia, March 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69:545–50.
5. Goyal P, Choi JJ, Pinheiro LC, et al. Clinical characteristics of Covid-19 in New York City. *N Engl J Med* 2020; 382:2372–4.
6. Cummings MJ, Baldwin MR, Abrams D, et al. Epidemiology, clinical course, and outcomes of critically ill adults with COVID-19 in New York City: a prospective cohort study. *Lancet* 2020; 395:1763–70.
7. Argenziano MG, Bruce SL, Slater CL, et al. Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: retrospective case series. *BMJ* 2020; 369:m1996.
8. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA* 2020; 323:1061–9.
9. Liang W, Liang H, Ou L, et al. Development and validation of a clinical risk score to predict the occurrence of critical illness in hospitalized patients with COVID-19. *JAMA Intern Med* 2020 [Epub ahead of print].
10. Zheng S, Fan J, Yu F, et al. Viral load dynamics and disease severity in patients infected with SARS-CoV-2 in Zhejiang province, China, January–March 2020: retrospective cohort study. *BMJ* 2020; 369:m1443.
11. Liu Y, Yan LM, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *Lancet Infect Dis* 2020; S1473-3099:30232-2.
12. Sethuraman N, Jeremiah SS, Ryo A. Interpreting diagnostics tests for SARS-CoV-2. *JAMA* 2020 [Epub ahead of print].
13. cobas SARS-CoV-2 EUA assay. Roche Diagnostics. package insert. v1.0 March 2020.
14. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data capture (REDCap)—a metadata-driven methodology and workflow process for providing translational research informatics support. *J Biomed Inform* 2009; 42:377–81.
15. Ison MG, Wolfe C, Boucher HW. Emergency use authorization of remdesivir: the need for a transparent distribution process. *JAMA* 2020 [Epub ahead of print].
16. Wu C, Chen X, Cai Y, et al. Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020; 180:1–11.
17. Liang W, Guan W, Chen R, et al. Cancer patients in SARS-CoV-2 infection: a nationwide analysis in China. *Lancet Oncol* 2020; 21:335–7.
18. Lan J, Ge J, Yu J, et al. Structure of the SARS-CoV-2 spike receptor-binding domain bound to the ACE2 receptor. *Nature* 2020; 581:215–20.
19. Ishiyama Y, Gallagher PE, Averill DB, Tallant EA, Brosnihan KB, Ferrario CM. Upregulation of angiotensin-converting enzyme 2 after myocardial infarction by blockade of angiotensin II receptors. *Hypertension* 2004; 43:970–6.
20. Li J, Wang X, Chen J, Zhang H, Deng A. Association of renin-angiotensin system inhibitors with severity or risk of death in patients with hypertension hospitalized for coronavirus disease 2019 (COVID-19) infection in Wuhan, China. *JAMA Cardiol* 2020 [Epub ahead of print].
21. Mancia G, Rea F, Ludergnani M, Apolone G, Corrao G. Renin-angiotensin-aldosterone system blockers and the risk of Covid-19. *N Engl J Med* 2020; 382:2431–40.
22. Mehta N, Kalra A, Nowacki AS, et al. Association of use of angiotensin-converting enzyme inhibitors and angiotensin II receptor blockers with testing positive for coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020 [Epub ahead of print].
23. American College of Cardiology. HFSA/AAC/AHA statement addresses concerns re: using RAAS antagonists in COVID-19. Available at: <https://www.acc.org/latest-in-cardiology/articles/2020/03/17/08/59/hfsa-acc-aha-statement-addresses-concerns-re-using-raas-antagonists-in-covid-19>. Accessed 16 May 2020.
24. New York State Department of Health. COVID-19 fatalities. Updated May 16, 2020. Available at: <https://covid19tracker.health.ny.gov/views/NYS-COVID19-Tracker/NYSDOHCOVID-19Tracker-Fatalities?%3Aembed=yes&%3Atoolbar=no&%3Atabs=n>. Accessed 17 May 2020.
25. Reyes C, Husain N, Gutowski C, St Clair S, Pratt G. Chicago's coronavirus disparity: black Chicagoans are dying at nearly six times the rate of white residents, data show. *Chicago Tribune*. Published April 7, 2020. Available at: <https://www.chicagotribune.com/coronavirus/ct-coronavirus-chicago-coronavirus-deaths-demographics-lightfoot-20200406-77nlyhiavjz2wa4ckivh7mu-story.html>. Accessed 16 May 2020.
26. Garg S, Kim L, Whitaker M, et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019—COVID-NET, 14 States, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep* 2020; 69:458–64.
27. Puelles VG, Lütgehetmann M, Lindenmeyer MT, et al. Multiorgan and renal tropism of SARS-CoV-2. *N Engl J Med* 2020 [Epub ahead of print].
28. Shi S, Qin M, Shen B, et al. Association of cardiac injury and mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol* 2020 [Epub ahead of print].
29. Pei G, Zhang Z, Peng J, et al. Renal involvement and early prognosis in patients with COVID-19 pneumonia. *J Am Soc Nephrol* 2020; 31:1157–65.
30. Craney AR, Velu P, Satlin MJ, et al. Comparison of two high-throughput reverse transcription-polymerase chain reaction systems for detection of severe acute respiratory syndrome coronavirus 2. *J Clin Microbiol* 2020 [Epub ahead of print].
31. Smithgall MC, Scherbergova I, Whittier S, Green DA. Comparison of Cepheid Xpert Xpress and Abbott ID now to Roche Cobas for the rapid detection of SARS-CoV-2. *J Clin Virol* 2020; 128:104428.
32. He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat Med* 2020; 26:672–5.
33. Bullard J, Dust K, Funk D, et al. Predicting infectious SARS-CoV-2 from diagnostic samples. *Clin Infect Dis* 2020 [Epub ahead of print].
34. La Scola B, Le Bideau M, Andreati J, et al. Viral RNA load as determined by cell culture as a management tool for discharge of SARS-CoV-2 patients from infectious disease wards. *Eur J Clin Microbiol Infect Dis* 2020; 39:1059–61.