



Published in final edited form as:

*Obesity (Silver Spring)*. 2018 June ; 26(6): 1088–1095. doi:10.1002/oby.22192.

## Impact of Early-Life Weight Status on Cognitive Abilities in Children

Nan Li<sup>1</sup>, Kimberly Yolton<sup>2</sup>, Bruce P. Lanphear<sup>3</sup>, Aimin Chen<sup>4</sup>, Heidi J. Kalkwarf<sup>5</sup>, and Joseph M. Braun<sup>1</sup>

<sup>1</sup>Department of Epidemiology, Brown University, Providence, Rhode Island, United States

<sup>2</sup>Department of Pediatrics, Division of General and Community Pediatrics, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, United States

<sup>3</sup>Child and Family Research Institute, BC Children's Hospital and the Faculty of Health Sciences, Simon Fraser University, Burnaby, British Columbia, Canada

<sup>4</sup>Division of Epidemiology, Department of Environmental Health, University of Cincinnati College of Medicine, Cincinnati, Ohio, United States

<sup>5</sup>Department of Pediatrics, Division of Gastroenterology, Hepatology and Nutrition, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, United States

### Abstract

**Objective**—Whether obesity is associated with childhood cognition is unknown. Given the sensitivity of the developing brain to environmental factors, we examined whether early-life weight status was associated with children's cognition.

**Methods**—Using data from mother-child pairs enrolled in the HOME Study (2003–2006), we assessed children's early-life weight status using weight-for-length/height standard deviation (SD) scores. We administered a battery of neuropsychological tests to assess cognition, executive function, and visual-spatial abilities at ages 5 and 8 years. Using linear mixed models, we estimated associations between early-life weight status and cognition.

**Results**—Among 233 children, 167 were lean ( $< 1SD$ ) and 48 were non-lean ( $> 1SD$ ). After covariate adjustment, the results suggest that full-scale Intelligence Quotient (IQ) scores decreased with 1-unit increase in weight-for-height SD score ( $\beta = -1.4$ , CI:  $-3.0, 0.1$ ). For individual component scores, with 1-unit increase in weight-for-height SD score, perceptual reasoning ( $\beta = -1.7$ , CI:  $-3.3, 0.0$ ) and working memory ( $\beta = -2.4$ ; CI:  $-4.4, -0.4$ ) scores decreased. Weight status was generally not associated with other cognition measures.

---

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: [http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

Corresponding Author: Joseph M. Braun, Brown University, Box G-S121(2), Providence, RI 02912, Telephone: 401-863-5397, joseph\_braun\_1@brown.edu.

**Competing financial interests:** The authors declared no conflict of interest.

**Conclusions**—Within this cohort of typically developing children, early-life weight status was inversely associated with children’s perceptual reasoning and working memory scores, and possibly full-scale IQ scores.

### Keywords

children; adiposity; cognitive function

---

## Introduction

Childhood obesity is an important public-health issue.<sup>1</sup> The global prevalence of childhood overweight and obesity has increased substantially since the 1990s, and approximately 60 million children are expected to be obese in 2020.<sup>2</sup> In the United States, approximately 17% of children and adolescents were obese and another 15% were overweight during 2009–2010.<sup>3</sup> Young children with excess adiposity are more susceptible to becoming overweight or obese in later childhood,<sup>4</sup> and at increased risk for adverse health consequences and obesity in adulthood.<sup>5</sup>

Obesity is associated with lower cognition in adults; these associations may be mediated by pro-inflammatory cytokines, leptin, and C-reactive protein.<sup>6,7</sup> In addition, obesity-induced dysregulation of appetite hormones, such as ghrelin and glucagon-like peptide 1 (GLP-1), may have detrimental effects on cognition since these hormones act in multiple brain regions that are relevant to cognitive abilities.<sup>7–10</sup> In rodents, both obesity and high fat diets can adversely affect hippocampal-dependent learning.<sup>11,12</sup>

The association between obesity and cognition in children is less well understood. Previous studies examining childhood weight status and cognition using cross-sectional designs were not able to specify the directionality of this relationship.<sup>13,14</sup> Results from prospective studies are inconsistent, possibly due to discrepancies in age and weight distribution, assessment of weight status or cognitive abilities, and confounder adjustment.<sup>15–21</sup> Furthermore, only two studies have examined the impact of early-life obesity on childhood cognition, however, both studies focused on early growth. Given that cognition develops rapidly in the first few years of life, it is important to investigate whether weight status in early life has an impact on cognitive function in children during this critical period of neurodevelopment.<sup>22</sup>

To address these gaps in the current literature, we investigated whether weight status in the first two years of life was associated with an array of cognitive abilities in school-age children using data from a prospective cohort study. Our analysis was guided by the hypothesis that adipose tissue produces adipocytokines or inflammatory molecules that may adversely affect children’s neurodevelopment.

## Methods

### Study Participants

We used data from mother-child pairs participating in the Health Outcomes and Measures of the Environment (HOME) Study, a longitudinal cohort that enrolled pregnant women in the

Cincinnati, OH metropolitan area from 2003 to 2006 and conducted follow-up visits with their children through age 8 years. Inclusion criteria and recruitment for the HOME Study has been described previously.<sup>23</sup> Of 1,263 eligible women, 468 (37%) enrolled in our study. The current analysis was restricted to 233 children who had early-life weight and length/height measurements, covariates, and at least one measurement of cognitive abilities at age 5 or 8 years.

The HOME Study protocols were approved by the institutional review boards at Cincinnati Children's Hospital Medical Center and cooperating delivery hospitals. All mothers provided written informed consent for themselves and their children.

### Early-Life Weight Status Measures

Study staff measured weight and length/height when children were ages 1 and 2 years. We used an infant scale to measure weight at age 1 year and a pediatric scale at age 2 years (ScaleTronix Inc., White Plains, NY). We measured length with a length board at age 1 year and standing height with a stadiometer at age 2 years. All measures were done in triplicate and the average was used.

According to World Health Organization (WHO) guidelines, weight-for-length/height z-score (WHZ) should be used to assess weight status for children under age 5 years, with  $WHZ > 2SD$  considered overweight and  $WHZ > 3SD$  considered obese.<sup>24</sup> We calculated WHZ using WHO sex-specific standard data.<sup>25</sup> Because only a small number of children in this study had a  $WHZ > 2SD$  ( $N=18$ ), the threshold for being overweight or obese, we grouped children into two categories: lean ( $< 1SD$ ) and non-lean ( $> 1SD$ ). We used WHZ at 2 years to determine children's weight status for 85% of children ( $n=199$ ), while WHZ at age 1 year was used for 15% ( $n=34$ ) of children with no weight and height data collected at 2 years (we used the phrase "early-life" to represent ages 1–2 years). We calculated Pearson correlation coefficients to examine the correlation of WHZs at different ages, including 1-month, 1-year, 2-year, 3-year, and 4-year. We focused on weight at age 1–2 years rather than other ages or WHZs trajectories because our primary interest was in the impact of adiposity in the first two years of life, a period of development when cognition develops rapidly.<sup>22</sup>

### Cognitive Outcomes

Trained examiners assessed several domains of children's cognitive abilities at ages 5 and 8 years using valid and reliable neuropsychological instruments that have been used extensively in epidemiological studies. These included tests of general cognitive abilities, attention, impulsivity, working memory, and reference memory.

We administered the Wechsler Preschool and Primary Scales of Intelligence-III (WPPSI-III) and Wechsler Intelligence Scale for Children-IV (WISC-IV) at ages 5 and 8 years, respectively.<sup>26,27</sup> Both Wechsler instruments measure children's overall intellectual abilities (Full-Scale Intelligence Quotient [FSIQ]), verbal abilities (Verbal IQ [VIQ], WPPSI-III and Verbal Comprehension Index [VCI], WISC-IV), perceptual reasoning and organization skills (Performance IQ [PIQ], WPPSI-III and Perceptual Reasoning Index [PRI], WISC-IV), and speed of mental and graphomotor processing (Processing Speed Quotient [PSQ], WPPSI-III and Processing Speed Index [PSI], WISC-IV). Additionally, digit span and letter-number

sequencing (Working Memory Index [WMI]) was measured using the WISC-IV. Standardized scores on these tests have a mean = 100 and standard deviation (SD) = 15, and lower scores indicate lower cognitive abilities.

We administered the Conners' Kiddie Continuous Performance Test and Conners' Continuous Performance Test-II (K-CPT/CPT-II) at ages 5 and 8 years, respectively.<sup>28,29</sup> These two computerized tasks assess children's attention, impulsivity, and executive control. During the test, children were told to press the spacebar on a computer whenever they saw any picture except for the "soccer ball" picture (K-CPT) or letter except for the letter "X" (CPT-II). Both CPTs generate T-scores for the following measures: commission errors that measures response inhibition (i.e., hitting spacebar when the non-target stimulus is presented); omission errors that assesses sustained attention (i.e., failing to hit spacebar when a target stimulus is presented); and hit reaction time (RT) that measures speed of processing and responding. Tau is an ex-Gaussian indicator of RTs, and higher values of tau indicate instances of longer RTs in the RT stream. Higher T-scores on the K-CPT/CPT-II tests indicate poorer performance.

Finally, we administered the Virtual Morris Water Maze (VMWM), a computerized version of the rodent Morris Water Maze, to test children's learning and spatial reference memory at age 8 years.<sup>30</sup> Children were instructed to find an escape platform as quickly as possible in a virtual pool using a joystick. Children first completed four practice trials where the platform was visible in a room with no visual cues. Subsequently, children completed four blocks of four trials with visual landmarks on the walls and the platform hidden in a fixed location. Performance was measured by the average latency to find the platform and distance from the start location to the platform across the four blocks. Children also completed a 30-s probe trial in the same room with the platform removed; performance was measured by the percentage of time spent in the correct quadrant, which assesses children's visual-spatial reference memory.

### Covariate Assessment

We used directed acyclic graphs to identify potential confounders in our exposure-outcome association (Figure S1).<sup>31</sup> Maternal age, education, marital status, length of breastfeeding, child's race, and interest in playing video games were assessed using questionnaires administered by trained interviewers. Child's sex and birthweight were abstracted from hospital medical charts. We calculated maternal body mass index (BMI) using weight and height measurements collected at the 16-week prenatal clinic visit. We measured mothers' FSIQ using the Wechsler Abbreviated Scale of Intelligence.<sup>32</sup> We assessed prenatal tobacco smoke exposure by measuring the concentration of cotinine - a sensitive and specific biomarker of secondhand and active tobacco smoke exposure - in maternal serum samples collected during 16 and 26 weeks of pregnancy. We used the mean of the log<sub>10</sub>-transformed serum cotinine concentrations in our analyses.

### Statistical Analyses

We analyzed the association between early-life WHZ and cognitive abilities (measured repeatedly at ages 5 and 8 years) using linear mixed models. Outcomes of FSIQ score, each



–0.4) were statistically significant (Table 2). We observed linear associations of WHZ with FSIQ, VCI, PRI, and WMI scores (all non-linearity  $P$ -values  $> 0.20$ ) (Figure 1 and Figure S2). There was some evidence of a non-linear association between WHZ and PSI scores, where PSI scores increased and reached a peak at a WHZ of  $\sim 0.5$  SD, and then decreased with increasing WHZ (non-linearity  $P$ -value=0.05). When WHZ was modelled as a binary variable, children who were non-lean had lower FSIQ scores compared to lean children ( $\beta = -4.3$ , 95% confidence interval (CI):  $-8.4, -0.2$ ) (Table S2). For individual component scores, children who were non-lean had lower working memory ( $\beta = -7.7$ , CI:  $-12, -3.0$ ) scores than lean children. Children who were non-lean also had lower scores of perceptual reasoning and processing speed compared to lean children; however, these associations were not statistically significant.

The K-CPT/CPT-II was administered to 225 children, with a total of 340 repeated measurements at ages 5 and 8 years. The RT T-score increased with 1-unit increase in WHZ ( $\beta = 1.1$ , CI= $-0.2, 2.5$ ), and the RT T-score was higher among non-lean children compared to lean children ( $\beta = 3.6$ , CI= $0.7, 6.6$ ). RT T-score increased monotonically with WHZ (non-linearity  $p = 0.70$ ) (Figure 2). WHZ was not associated with commission or omission T-scores or tau.

Visual-spatial abilities were measured among 185 children at age 8 years. Latency (seconds) and distance (pool units) to find the hidden platform was not associated with weight status (Table 2 and Table S2). In the probe trial, the portion of children who found the correct quadrant was not associated with weight status.

Generally, the association between WHZ and cognitive abilities did not differ by child sex. However, WHZ (modeled as either continuous or binary variable) was inversely associated with WMI scores and VMWM distance among boys (sex  $\times$  WHZ  $P$ -values  $< 0.05$ ), but not girls (Table S3–S4). Other WHZ-cognition associations were consistent across boys and girls. When we examined the associations with IQ subtests, 1-unit increase in early-life WHZ was inversely associated with Block Design ( $\beta = -0.4$ , CI:  $-0.7, -0.1$ ), but was not associated with other PRI subtests (Table S5). Results from sensitivity analyses using WHO BMIZ were similar to our main findings (Table S6).

## Discussion

In the present study, we investigated the associations between early-life weight status and children's cognitive abilities in a longitudinal cohort of children from Cincinnati, Ohio. Our findings suggest that early-life WHZ may be inversely associated with FSIQ, PRI, and WMI scores after adjusting for potential confounders. Early-life WHZ was also suggestively associated with higher RT T-scores on the K-CPT/CPT-II. Other measures of cognitive abilities were not associated with early-life WHZ.

Prospective studies examining the association between weight status and cognitive abilities in children have yielded inconsistent results.<sup>15–21</sup> Two studies have examined the impact of early-life weight status on cognitive abilities; however, both studies focused on change in BMI or weight over the first few years of life.<sup>15,16</sup> In a study of Finnish children, both small

and large body size were associated with decreased visual-motor integration at age 56 months.<sup>15</sup> This inverted U-shape relationship is similar to the pattern we observed when examining WHZ and children's PSI, a measure of visual perception and organization, as well as motor skills. In a large population-based study of British children, rapid weight gain between birth and age 25 months (defined as an increase of  $>0.67$  in the SD score of weight) was not associated with cognitive abilities at ages 49 months or 8 years, measured using the WPPSI and WISC-III, respectively.<sup>16</sup> Unlike our study, this British study focused on weight gain in the first two years of life, without accounting for children's height and obesity status. This could result in misclassification of early life adiposity because children with constant overweight/obesity during the first two years may not be considered as having rapid growth, in contrast, lean children who grew, but did not become overweight or obese were still classified as having rapid growth.

Other prospective studies have examined the impact of obesity in childhood on global measures of cognition, with varied results.<sup>17–21</sup> Consistent with one of our findings, two studies reported that obesity at ages 3–5 years was associated with lower cognitive abilities in boys only at ages 5 or 8–12 years.<sup>17,18</sup> In contrast, two studies reported that children who were underweight had lower cognitive abilities.<sup>19,20</sup> Very few children were underweight in our study, and thus we were not able to precisely examine this association. Finally, a study of Dutch children did not observe associations between BMI at age 4 years and cognitive abilities at age 7 years assessed using the Kaufman Assessment Battery for Children.<sup>21</sup> However, this study had a small number of children with overweight (N=18), limiting their statistical power. In sum, discrepancies in the findings above may be explained by differences in the age of exposure and outcome measurements, measures of weight status, weight status distribution of the study population, different lengths of follow-up, measures of cognitive abilities, and adjustment for confounding factors.<sup>15–21</sup>

Executive function is a set of self-regulatory cognitive processes that aid in managing thoughts, emotions, and goal-directed behaviors. It includes domains like inhibitory control, working memory, reward sensitivity, and attention.<sup>14</sup> Executive function is associated with academic success in children and is critical for physical health and success throughout life.<sup>35</sup> Previous studies reported a consistent inverse association between obesity and executive function in children; however, most studies were cross-sectional in design and thus, the directionality of the association cannot be determined.<sup>14</sup> Some speculate that children with lower executive function have lower self-regulation of caloric intake and physical activity, which ultimately leads to overweight or obesity, whereas others argue that obesity may decrease executive function through the action of adipocytokines or inflammatory molecules.<sup>7,14,36</sup> Moreover, some have proposed a bidirectional relation where low executive function promotes excess adiposity and excess adiposity exacerbates executive function decrements.<sup>36</sup>

Our results suggest that non-lean children had higher RT T-scores (indicating lower speed of processing and responding) compared to lean children. Some cross-sectional studies included in a systematic review by Reinert et al. reported shorter reaction time and more commission errors among children with higher BMI.<sup>14</sup> While the reasons for the inconsistency with our own results is unclear, it is possible that the shorter reaction time in

those studies may reflect high impulsivity, which may have led to over-consumption of food and higher BMI. Given the cross-sectional nature of prior studies, it is not possible to delineate the temporal relation between BMI and these executive functions. In contrast, the longitudinal design of our study ensured that we could determine the directionality of the relationship (from adiposity to executive function) and ensure that it was not influenced by bidirectionality, since weight status was measured years prior to executive function assessment and at an age when eating behavior was more affected by parental influence than children's self-regulation.<sup>37</sup> A population-based study in Germany investigated the 1-year longitudinal associations and found that BMI was not associated with executive function.<sup>38</sup> However, the 1-year follow-up time might be too short to allow overweight-associated pathophysiological processes to affect executive function, and they did not adjust for mothers' BMI in the analysis. The CPT tests utilized in our study assess only the attention and inhibitory control aspects of executive functioning, therefore, additional longitudinal studies with repeated and comprehensive measures of executive function are needed.

In our study, weight status in general was not associated with visual-spatial abilities measured by VMWM, except that weight-for-height may be inversely associated with distance in the VMWM hidden platform trials among boys (Table S3–S4). We found that early-life WHZ was inversely associated with performance on Block Design test, a PRI subtest that also measures visuospatial organization. Consistent with our findings, BMI was inversely associated with block design test performance among school-age US children and adolescents in a cross-sectional study.<sup>13</sup> Some other studies examined the relationship between obesity and visual-spatial abilities in children. In a UK cohort study, obesity at age 3 was associated with worse visuospatial skills at age 5 in boys.<sup>18</sup> A randomized controlled trial conducted among school-age Danish children found that visuospatial construction skills were improved after obesity intervention among children with overweight or obesity, indicating that obesity may play a detrimental role in visual-spatial abilities.<sup>39</sup> The underlying mechanisms for these sex-specific effects are unclear. Sex hormones or psychosocial factors may partly explain the differences. Given the sparsity of the current literature, additional prospective studies are needed to confirm these findings and examine whether the associations differ by sex.

There are several biological mechanisms by which early life adiposity could affect neurodevelopment. Adipose tissue produces pro-inflammatory cytokines that activate inflammatory pathways in children and adults.<sup>7</sup> Systematic inflammation may affect multiple brain regions relevant to cognitive abilities, and was shown to adversely affect spatial learning and memory in rodents.<sup>7</sup> In addition, the dysregulation of appetite-regulating hormones among children with excess adiposity may adversely affect cognition.<sup>7</sup> Ghrelin, a hunger hormone, can cross the blood-brain-barrier and activate hippocampal regions to improve memory in rodents.<sup>9</sup> GLP-1, which promotes satiety, also acts on multiple brain regions that are involved in cognitive abilities, such as the hypothalamus and the prefrontal cortex.<sup>8</sup> In rodents deficiencies in ghrelin and GLP-1 caused deficits in learning and memory.<sup>7,10</sup>

Our study had some limitations. This study had limited statistical power to detect small effects and precisely estimate associations, especially for children who were overweight/



obese, because of our modest sample size. Thus, we were not able to examine children with obesity and overweight separately. However, by using different categories of weight status, we were able to examine the group of children who were at the high-end of normal WHZ range ( $>1SD$ ) compared to lean children ( $1SD$ ), which brings attention to this group of children who are at risk of being overweight or obese later in childhood and adolescence.<sup>40</sup> Furthermore, while WHZ at age 2 years was used in most children as an exposure, WHZ at age 1 year was used in some children when 2-year data were not available. However, 2-year WHZ was highly correlated with 1-year, 3-year, and 4-year WHZ in this study, which reduced the possibility of exposure misclassification. The correlation pattern (and the one observed for repeated BMI z-scores, Table S7) indicates that 2-year WHZ tends to predict weight status in childhood, which is consistent with previous findings.<sup>4</sup> Another limitation is that we did not use direct measurement of adiposity, such as dual energy X-ray absorptiometry, which can provide a more accurate assessment of body composition.<sup>41</sup> However, given that young children may not be cooperative for this measurement, WHZ is a more practical measurement of excess weight in early childhood and is endorsed by WHO to assess body composition for children under 5 years. Finally, we made several comparisons and it is possible that some of our results may be chance findings. However, adjusting for multiple comparisons may undeservedly reduce statistical power<sup>42</sup>, and the general pattern of our results suggests an inverse association between early life adiposity and cognitive abilities.

Our study had several strengths. Our prospective data enabled us to investigate whether weight status in the critical period of brain development - the first two years of life - impacts cognitive abilities in childhood. Compared to prior cross-sectional studies examining the association between childhood weight status and cognition, the prospective design helped us determine the directionality of the association. Furthermore, we repeatedly administered valid and reliable neuropsychological tests to assess an array of cognitive abilities. In addition, weight and length/height were measured by trained study staff, rather than self-reported by parents, which provided more accurate measurements and reduced potential exposure misclassification. Finally, we collected detailed information on covariates in this study, which enabled us to adjust for multiple potential confounding factors including sociodemographic factors, perinatal factors, and maternal IQ.

## Conclusion

Excess early-life weight-for-height may be inversely associated with full-scale IQ, perceptual reasoning scores, and working memory scores (boys only), and suggestively associated with longer reaction time among school-aged children. Future prospective cohort studies should confirm these findings and investigate if early-life weight-for-height is associated with school performance, attention-deficit/hyperactivity disorder diagnosis, learning disabilities, or special education service use.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

**Sources of Financial Support:** This work was supported by NIEHS grants R01 ES024381, R01 ES025214, R01 ES020349, R00 ES020346, P01 ES011261, and R01 ES014575.

## References

1. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002; 360:473–482. [PubMed: 12241736]
2. de Onis M, Blossner M, Borghi E. Global prevalence and trends of overweight and obesity among preschool children. *The American journal of clinical nutrition*. 2010; 92:1257–1264. [PubMed: 20861173]
3. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity and trends in body mass index among US children and adolescents, 1999–2010. *JAMA*. 2012; 307:483–490. [PubMed: 22253364]
4. Glavin K, Roelants M, Strand BH, et al. Important periods of weight development in childhood: a population-based longitudinal study. *BMC public health*. 2014; 14:160. [PubMed: 24524269]
5. Freedman DS, Mei Z, Srinivasan SR, Berenson GS, Dietz WH. Cardiovascular risk factors and excess adiposity among overweight children and adolescents: the Bogalusa Heart Study. *J Pediatr*. 2007; 150:12–17e12. [PubMed: 17188605]
6. Elias MF, Elias PK, Sullivan LM, Wolf PA, D'Agostino RB. Lower cognitive function in the presence of obesity and hypertension: the Framingham heart study. *Int J Obes Relat Metab Disord*. 2003; 27:260–268. [PubMed: 12587008]
7. Miller AL, Lee HJ, Lumeng JC. Obesity-associated biomarkers and executive function in children. *Pediatr Res*. 2015; 77:143–147. [PubMed: 25310758]
8. Pannacciulli N, Le DS, Salbe AD, et al. Postprandial glucagon-like peptide-1 (GLP-1) response is positively associated with changes in neuronal activity of brain areas implicated in satiety and food intake regulation in humans. *Neuroimage*. 2007; 35:511–517. [PubMed: 17317222]
9. Carlini VP, Martini AC, Schiöth HB, Ruiz RD, Fiol de Cuneo M, de Barioglio SR. Decreased memory for novel object recognition in chronically food-restricted mice is reversed by acute ghrelin administration. *Neuroscience*. 2008; 153:929–934. [PubMed: 18434026]
10. During MJ, Cao L, Zuzga DS, et al. Glucagon-like peptide-1 receptor is involved in learning and neuroprotection. *Nat Med*. 2003; 9:1173–1179. [PubMed: 12925848]
11. Wu A, Molteni R, Ying Z, Gomez-Pinilla F. A saturated-fat diet aggravates the outcome of traumatic brain injury on hippocampal plasticity and cognitive function by reducing brain-derived neurotrophic factor. *Neuroscience*. 2003; 119:365–375. [PubMed: 12770552]
12. Bocarsly ME, Fasolino M, Kane GA, et al. Obesity diminishes synaptic markers, alters microglial morphology, and impairs cognitive function. *Proc Natl Acad Sci U S A*. 2015; 112:15731–15736. [PubMed: 26644559]
13. Li Y, Dai Q, Jackson JC, Zhang J. Overweight is associated with decreased cognitive functioning among school-age children and adolescents. *Obesity (Silver Spring)*. 2008; 16:1809–1815. [PubMed: 18551126]
14. Reinert KR, Po'e EK, Barkin SL. The relationship between executive function and obesity in children and adolescents: a systematic literature review. *J Obes*. 2013; 2013:820956. [PubMed: 23533726]
15. Heinonen K, Raikonen K, Pesonen AK, et al. Prenatal and postnatal growth and cognitive abilities at 56 months of age: a longitudinal study of infants born at term. *Pediatrics*. 2008; 121:e1325–1333. [PubMed: 18450875]
16. Beyerlein A, Ness AR, Streuling I, Hadders-Algra M, von Kries R. Early rapid growth: no association with later cognitive functions in children born not small for gestational age. *Am J Clin Nutr*. 2010; 92:585–593. [PubMed: 20592132]
17. Black N, Johnston DW, Peeters A. Childhood Obesity and Cognitive Achievement. *Health Econ*. 2015; 24:1082–1100. [PubMed: 26123250]
18. Martin A, Booth JN, Young D, et al. Associations between obesity and cognition in the pre-school years. *Obesity (Silver Spring)*. 2016; 24:207–214. [PubMed: 26638123]

19. Wisniewski SL. Childhood obesity among the poor in Peru: Are there implications for cognitive outcomes? *Econ Hum Biol.* 2017; 26:51–60. [PubMed: 28284174]
20. Bisset S, Fournier M, Fournier M, Pagani L, Janosz M. Predicting academic and cognitive outcomes from weight status trajectories during childhood. *Int J Obes (Lond).* 2013; 37:154–159. [PubMed: 22828944]
21. Veldwijk J, Scholtens S, Hornstra G, Bemelmans WJ. Body mass index and cognitive ability of young children. *Obes Facts.* 2011; 4:264–269. [PubMed: 21921648]
22. Garon N, Bryson SE, Smith IM. Executive function in preschoolers: a review using an integrative framework. *Psychol Bull.* 2008; 134:31–60. [PubMed: 18193994]
23. Braun JM, Kalloo G, Chen A, et al. Cohort Profile: The Health Outcomes and Measures of the Environment (HOME) study. *Int J Epidemiol.* 2017; 46:24. [PubMed: 27006352]
24. World Health Organization. [Accessed August 22, 2017] Obesity and overweight fact sheet [Web page]. <http://www.who.int/mediacentre/factsheets/fs311/en/>
25. WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards: Length/height-for-age, weight-for-age, weight-for-length, weight-for-height and body mass index-for-age: Methods and development. Geneva: World Health Organization; 2006.
26. Wechsler, D. Wechsler Preschool and Primary Scale of Intelligence. 3. San Antonio: The Psychological Corporation; 2002.
27. Wechsler, D. Wechsler Intelligence Scale for Children-IV. 3. San Antonio: The Psychological Corporation; 2003.
28. Conners, C. Conners' Kiddie Continuous Performance Test. North Tonawanda: Multi-Health Systems; 2006.
29. Conners, C. Conners' Continuous Performance Test. 2. Toronto: Multi-Health Systems; 2004.
30. Astur RS, Ortiz ML, Sutherland RJ. A characterization of performance by men and women in a virtual Morris water task: a large and reliable sex difference. *Behavioural brain research.* 1998; 93:185–190. [PubMed: 9659999]
31. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology.* 1999; 10:37–48. [PubMed: 9888278]
32. Wechsler, D. Wechsler Abbreviated Scale of Intelligence. New York: The Psychological Corporation: Harcourt Brace & Company; 1999.
33. Braun JM, Daniels JL, Poole C, et al. A prospective cohort study of biomarkers of prenatal tobacco smoke exposure: the correlation between serum and meconium and their association with infant birth weight. *Environ Health.* 2010; 9:53. [PubMed: 20799929]
34. Desquilbet L, Mariotti F. Dose-response analyses using restricted cubic spline functions in public health research. *Statistics in medicine.* 2010; 29:1037–1057. [PubMed: 20087875]
35. Diamond A, Lee K. Interventions shown to aid executive function development in children 4 to 12 years old. *Science.* 2011; 333:959–964. [PubMed: 21852486]
36. Hayes JF, Eichen DM, Barch DM, Wilfley DE. Executive function in childhood obesity: Promising intervention strategies to optimize treatment outcomes. *Appetite.* 2017
37. Parsons TJ, Power C, Logan S, Summerbell CD. Childhood predictors of adult obesity: a systematic review. *Int J Obes Relat Metab Disord.* 1999; 23(Suppl 8):S1–107.
38. Groppe K, Elsner B. Executive function and weight status in children: A one-year longitudinal perspective. *Child Neuropsychol.* 2017; 23:129–147. [PubMed: 26416504]
39. Huang T, Larsen KT, Jepsen JR, et al. Effects of an obesity intervention program on cognitive function in children: A randomized controlled trial. *Obesity (Silver Spring).* 2015; 23:2101–2108. [PubMed: 26337394]
40. Cunningham SA, Kramer MR, Narayan KM. Incidence of childhood obesity in the United States. *The New England journal of medicine.* 2014; 370:403–411. [PubMed: 24476431]
41. Wells JC. A critique of the expression of paediatric body composition data. *Arch Dis Child.* 2001; 85:67–72. [PubMed: 11420208]
42. Rothman KJ. No adjustments are needed for multiple comparisons. *Epidemiology.* 1990; 1:43–46. [PubMed: 2081237]


### Study Importance Questions

#### What is already known about this subject?

- Obesity is associated with lower cognition in adults
- The association between obesity and cognition in children is less well understood


#### What does your study add?

- Our study found that early-life weight status was inversely associated with children's perceptual reasoning and working memory scores, and possibly full-scale IQ scores.



**Figure 1.** Adjusted Restricted Cubic Polynomial Spline of Weight-For-Height Z-Scores and Children's Full-Scale IQ, Perceptual Reasoning Index, Working Memory Index, and Processing Speed Index at 5 or 8 Years of Age <sup>a</sup>

<sup>a</sup>Adjusted for continuous variables maternal age, BMI, IQ, child's birthweight, age at outcome measurement, and maternal cotinine concentration (mean of log<sub>10</sub>-transformed cotinine concentrations during 16 and 26 weeks of pregnancy); and categorical variables maternal education (high school graduate or less, tech school or some college, college graduate or above), marital status (married, unmarried), length of breastfeeding (0 month, 6 months, >6 months), child's sex (Male, Female), race (non-Hispanic White, non-Hispanic Black, other).



**Figure 2.** Adjusted Restricted Cubic Polynomial Spline of Weight-For-Height Z-Scores and Children's Hit Reaction Time T-score at 5 or 8 Years of Age <sup>a</sup>

<sup>a</sup>Adjusted for continuous variables maternal age, BMI, IQ, child's birthweight, age at outcome measurement, and maternal cotinine concentration (mean of log<sub>10</sub>-transformed cotinine concentrations during 16 and 26 weeks of pregnancy); and categorical variables maternal education (high school graduate or less, tech school or some college, college graduate or above), marital status (married, unmarried), length of breastfeeding (0 month, 6 months, >6 months), child's sex (Male, Female), race (non-Hispanic White, non-Hispanic Black, other).

**Table 1**  
 Descriptive Statistics of Weight-For-Height Z-Score at Ages 1 or 2 Years and Full-Scale Intelligence Quotient (IQ) Scores at Ages 5 or 8 Years among the Health Outcomes and Measures of the Environment Study Women and Children (N=233)

	Weight-For-Height Z-Score (N=233)		Full-Scale IQ (N=191)		Full-Scale IQ (n=204)	
	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)
<b>Child Sex</b>						
Male	105	0.5 (1.1)	84	101 (15)	90	103 (15)
Female	128	0.5 (1.1)	107	104 (15)	114	102 (16)
<b>Child Race</b>						
Non-Hispanic White	141	0.5 (1.0)	117	108 (13)	123	108 (12)
Non-Hispanic Black	75	0.5 (1.2)	60	91 (13)	68	91 (15)
Other	17	0.5 (1.2)	14	105 (12)	13	110 (16)
<b>Child Birthweight</b>						
Small for gestational age (<10%)	23	-0.2 (1.0)	19	96 (16)	21	96 (18)
Appropriate for gestational age	177	0.4 (1.0)	144	103 (15)	155	103 (16)
Large for gestational age (>90%)	33	1.5 (1.0)	28	102 (14)	28	103 (13)
<b>Breastfeeding Duration</b>						
0 month	42	0.5 (1.3)	33	95 (16)	38	96 (17)
>0-6 months	88	0.4 (1.1)	67	101 (16)	76	101 (16)
>6 months	103	0.5 (1.0)	91	106 (14)	90	107 (14)
<b>Maternal Age at Delivery (years)</b>						
18-25	52	0.4 (1.2)	40	93 (14)	48	94 (16)
>25-35	148	0.5 (1.0)	122	104 (15)	126	104 (15)
>35	33	0.4 (1.1)	29	110 (14)	30	110 (13)
<b>Annual Household Income</b>						
>\$80K	60	0.4 (0.8)	49	112 (11)	53	112 (12)

	Weight-For-Height Z-Score (N=233)		5y		Full-Scale IQ (N=191)		8y		Full-Scale IQ (n=204)	
	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)	N	Mean (SD)
\$40–80K	83	0.6 (1.1)	75	105 (13)	70	105 (13)	70	105 (13)	70	105 (13)
<\$40K	90	0.4 (1.2)	67	93 (15)	81	93 (15)	81	94 (16)	81	94 (16)
Maternal Education										
College graduate	116	0.5 (0.9)	96	110 (12)	98	110 (12)	98	110 (12)	98	110 (12)
Tech school/Some College	66	0.2 (1.1)	56	98 (14)	58	98 (14)	58	101 (14)	58	101 (14)
High school graduate	51	0.7 (1.4)	39	90 (14)	48	90 (14)	48	90 (16)	48	90 (16)
Marital Status										
Married	154	0.5 (1.0)	131	107 (13)	132	107 (13)	132	108 (13)	132	108 (13)
Unmarried	79	0.5 (1.2)	60	92 (14)	72	92 (14)	72	93 (16)	72	93 (16)
Gestational Smoking Status										
Unexposed (<0.015ng/ml)	77	0.4 (1.0)	64	107 (13)	65	107 (13)	65	107 (13)	65	107 (13)
Secondhand Smoke (<3.0 ng/ml)	133	0.5 (1.1)	113	100 (15)	118	100 (15)	118	101 (16)	118	101 (16)
Active Smoking (>3.0 ng/ml)	23	0.7 (1.3)	14	97 (21)	21	97 (21)	21	96 (17)	21	96 (17)
Maternal Body Mass Index (kg/m <sup>2</sup> )										
< 25 (lean)	93	0.2 (1.0)	73	105 (16)	80	105 (16)	80	104 (16)	80	104 (16)
25–30 (overweight)	79	0.5 (1.0)	65	104 (14)	72	104 (14)	72	105 (14)	72	105 (14)
>30 (obese)	61	0.9 (1.3)	53	97 (15)	52	97 (15)	52	97 (16)	52	97 (16)
Maternal Full-scale IQ										
1 <sup>st</sup> Tercile 58–99	78	0.4 (1.3)	63	92 (14)	71	92 (14)	71	92 (15)	71	92 (15)
2 <sup>nd</sup> Tercile >99–114	80	0.3 (0.9)	64	106 (15)	67	106 (15)	67	107 (13)	67	107 (13)
3 <sup>rd</sup> Tercile >114	75	0.7 (1.0)	64	109 (12)	66	109 (12)	66	109 (14)	66	109 (14)

SD: standard deviation



Unadjusted and Adjusted Difference or Relative Risk in Children's Cognitive Abilities by 1-unit change in Early Life Weight-For-Height Z-Score (WHZ)<sup>a</sup>

Table 2

	N	Unadjusted Difference or Relative Risk (95% CI)	P-value	Adjusted Difference or Relative Risk <sup>b</sup> (95% CI)	P-value
WPPSI-III/WISC-IV Full-Scale IQ	233	-0.7 (-2.4, 1.0)	0.41	-1.4 (-3.0, 0.1)	0.07
WPPSI-III/WISC-IV Verbal	233	-0.7 (-2.4, 0.9)	0.39	-0.8 (-2.2, 0.7)	0.30
WPPSI-III/WISC-IV Perceptual Reasoning	233	-0.3 (-2.0, 1.3)	0.70	<b>-1.7 (-3.3, 0.0)</b>	<b>0.047</b>
WPPSI-III/WISC-IV Processing Speed	224	-0.3 (-1.7, 1.2)	0.74	-0.6 (-2.4, 1.2)	0.51
WISC-IV Working Memory	204	-1.3 (-3.2, 0.5)	0.16	<b>-2.4 (-4.4, -0.4)</b>	<b>0.02</b>
VMWM Latency (Seconds)	185	0.2 (-1.0, 1.5)	0.84	0.4 (-1.0, 1.8)	0.60
VMWM Distance (Pool Units)	185	-0.6 (-1.9, 0.6)	0.49	-0.3 (-1.7, 1.1)	0.69
VMWM Time in Correct Quadrant	185	1.0 (0.9, 1.2)	0.99	1.0 (0.8, 1.2)	0.66
K-CPT/CPT-II Commissions T-score	225	0.3 (-0.6, 1.2)	0.53	0.4 (-0.6, 1.4)	0.46
K-CPT/CPT-II Omissions T-score	225	0.0 (-1.8, 1.8)	0.99	1.2 (-0.8, 3.2)	0.25
K-CPT/CPT-II Reaction Time T-score	225	1.1 (-0.2, 2.4)	0.10	1.1 (-0.2, 2.5)	0.10
K-CPT/CPT-II Tau	188	1.0 (0.97, 1.1)	0.55	1.0 (0.98, 1.1)	0.16

<sup>a</sup>Relative risks were calculated for VMWM time spent in correct quadrant (0 vs. >0) and K-CPT/CPT-II Tau; differences were calculated for other outcomes.

<sup>b</sup> Adjusted for continuous variables maternal age, BMI, IQ, child's birthweight, age at outcome measurement, and maternal serum cotinine concentration (mean of log10-transformed cotinine concentrations during 16 and 26 weeks of pregnancy); and categorical variables maternal education (high school graduate or less, tech school or some college, college graduate or above), marital status (married, unmarried), length of breastfeeding (0 month, 6 months, >6 months), child's sex (Male, Female), race (non-Hispanic White, non-Hispanic Black, other). Interest in playing video games (a lot, some, a little) was adjusted in models with VMWM outcomes.

WPPSI-III: Wechsler Preschool and Primary Scales of Intelligence-III.

WISC-IV: Wechsler Intelligence Scales for Children-fourth version. VMWM: Virtual Morris Water Maze. K-CPT: Kiddie Conner's Continuous Performance Task. CPT-II: Conner's Continuous Performance Task-II