

## Article

# Acute Administration of Bioavailable Curcumin Alongside Ferrous Sulphate Supplements Does Not Impair Iron Absorption in Healthy Adults in a Randomised Trial

Helena Tiekou Lorinczova <sup>1</sup>, Gulshanara Begum <sup>1</sup>, Derek Renshaw <sup>2</sup> and Mohammed Gulrez Zariwala <sup>1,\*</sup>

<sup>1</sup> Centre for Nutraceuticals, School of Life Sciences, University of Westminster, 115 New Cavendish Street, London W1W 6UW, UK; w1505041@my.westminster.ac.uk (H.T.L.); begumru@westminster.ac.uk (G.B.)

<sup>2</sup> Centre for Sport, Exercise and Life Sciences, Institute for Health and Wellbeing, Coventry University, Priory St, Coventry CV1 5FB, UK; derek.renshaw@coventry.ac.uk

\* Correspondence: zariwam@wmin.ac.uk; Tel.: +44-20-7911-5000 (ext. 65086)

**Abstract:** Ferrous sulphate (FS) is a cost effective, readily available iron supplement for iron deficiency (ID). The pro-oxidant effect of oral ferrous iron is known to induce inflammation, causing gastric side-effects and resulting in poor compliance. Curcumin is a potent antioxidant and has also been shown to exhibit iron chelation in-vitro, although it is not established whether these effects are retained in-vivo. The aim of this study was therefore to assess the influence of a formulated bioavailable form of curcumin (HydroCurc<sup>TM</sup>; 500 mg) on acute iron absorption and status in a double blind, placebo-controlled randomized trial recruiting 155 healthy participants (79 males; 26.42 years  $\pm$  0.55 and 76 females; 25.82 years  $\pm$  0.54). Participants were randomly allocated to five different treatment groups: iron and curcumin placebo (FS0\_Plac), low dose (18 mg) iron and curcumin placebo (FS18\_Plac), low dose iron and curcumin (FS18\_Curc), high dose (65 mg) iron and curcumin placebo (FS65\_Plac), and high dose iron and curcumin (FS65\_Curc). Participants were provided with the supplements according to their relevant treatment groups at baseline (0 min), and blood collection was carried out at 0 min and at 180 min following supplementation. In the treatment groups, significant difference was observed in mean serum iron between baseline (0 min) and at end-point (180 min) ( $F(1, 144) = 331.9, p < 0.0001$ ) with statistically significant intra-group increases after 180 min ( $p < 0.0001$ ) in the FS18\_Plac (8.79  $\mu\text{mol/L}$ ), FS18\_Curc (11.41  $\mu\text{mol/L}$ ), FS65\_Plac (19.09  $\mu\text{mol/L}$ ), and FS65\_Curc (16.39  $\mu\text{mol/L}$ ) groups. A significant difference was also observed between the two time points in serum TIBC levels and in whole blood haemoglobin (HGB) in the treatment groups, with a significant increase (1.55%/2.04 g/L) in HGB levels from baseline to end-point observed in the FS65\_Curc group ( $p < 0.05$ ). All groups receiving iron demonstrated an increase in transferrin saturation (TS%) in a dose-related manner, demonstrating that increases in serum iron are translated into increases in physiological iron transportation. This study demonstrates, for the first time, that regardless of ferrous dose, formulated curcumin in the form of HydroCurc<sup>TM</sup> does not negatively influence acute iron absorption in healthy humans.

**Keywords:** curcumin; iron; ferrous sulphate; supplementation; nutrition; antioxidant



check for updates

**Citation:** Lorinczova, H.T.; Begum, G.; Renshaw, D.; Zariwala, M.G. Acute Administration of Bioavailable Curcumin Alongside Ferrous Sulphate Supplements Does Not Impair Iron Absorption in Healthy Adults in a Randomised Trial. *Nutrients* **2021**, *13*, 2300. <https://doi.org/10.3390/nu13072300>

Academic Editors: Teresa Partearroyo and Gregorio Varela-Moreiras

Received: 28 May 2021

Accepted: 30 June 2021

Published: 3 July 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

Iron is an essential mineral nutrient that plays a key role in most biological processes necessary to sustain life [1,2]. Iron deficiency (ID) is ranked as the most prevalent nutritional deficiency [3] affecting around 30% of the global population, equating to over 2 billion people [4,5]. ID is caused by inadequate dietary iron intake, as well as uptake by the body to replenish iron that is lost daily mainly due to erythrocyte turnover in addition to skin and gut cell sloughing. ID has been attributed to more than 60% of all anaemia cases worldwide and is associated with reduced life expectancy, altered immune response, developmental delays, and adverse pregnancy outcomes [3,6]. As iron is fundamental

for growth and metabolism, ID even without anaemia can lead to impaired cognitive and physical development in children, compromise physical and cognitive performance in adults [3,7–9], and has been linked with fatigue [10], impaired quality of life [11], and reduced mood [3,9].

Ferrous sulphate is considered the global ‘gold standard’ iron supplement due to its extensive history of safe usage, documented effectiveness, and low cost [12,13]. However, non-compliance with ferrous sulphate is as high as 50% due to negative gastrointestinal (GI) effects attributed to iron intake [14–16]. The scope for ID treatment success is thus limited often by poor compliance, particularly due to adverse GI effects such as constipation, diarrhoea, and flatulence [3]. These GI symptoms are often accompanied by increased inflammation as a consequence of excessive iron levels in the gut, facilitating the production of detrimental reactive oxygen species (ROS) and consequent damage to the gut mucosa [17].

Antioxidants can counteract the effects of ROS and have therefore generated interest as molecules that could potentially ameliorate the pro-oxidant side-effects of iron [18,19]. Curcumin, a non-flavonoid polyphenol, is the most biologically active antioxidant component in the rhizomatous spice *Curcuma longa* Linnaeus or turmeric [20]. Curcumin is comprised of three curcuminoids: curcumin (80–75%), demethoxycurcumin (DMC; 20%), and bisdemethoxycurcumin (BDMC; 2–5%) [21]. Curcumin has been extensively studied for its ROS scavenging properties [22–25]. It acts as a free-radical chain breaker, capable of donating hydrogen to ROS due to the presence of a hydroxyl group in its structure [26]. Curcumin also suppresses inflammation by mechanisms including downregulation of NF- $\kappa$ B (nuclear factor kappa-light-chain-enhancer of activated B cells), thereby demonstrating potent anti-inflammatory activity [27]. In addition to its antioxidant and anti-inflammatory action, curcumin has also been shown to have neuroprotective properties, particularly in relation to memory and cognition [28]. A meta-analysis of randomised controlled trials confirmed a significant positive impact of curcumin supplementation on levels of brain-derived neurotrophic factor (BDNF), a factor that affects cognitive function, learning, and memory [29]. Serum BDNF levels were also shown to increase significantly in healthy adults following co-administration of iron and curcumin supplements recently by Tiekou Lorinczova and co-workers [30]. The ability of curcumin to potentially counter iron-mediated ROS generation and inflammation in the GI tract, in concert with its ability to have a beneficial central effect, raises the intriguing possibility of co-administering iron and curcumin supplements to limit the oxidative stress associated with iron supplementation while synergising the beneficial effects of both molecules.

However, a body of research has shown whole turmeric or curcumin adversely affects iron absorption and iron status in animal models [31–33]. Whole turmeric is known to inhibit iron absorption by 20–90% in humans, reducing iron absorption in a dose-dependent manner [34]. Curcumin and its analogue curcuminoids, DMC and BDMC, can exist in two different forms depending on their environment, keto and enol [35]. In their enol form, they are capable of accepting as well as donating hydrogen and have metal chelation characteristics [36]. Curcumin binds ferric iron (Fe<sup>3+</sup>) to form a ferric–curcumin complex that is dose-dependent and Fe<sup>3+</sup> specific [37].

Despite these observations, there remains considerable ambiguity relating to the nature of interactions between iron and curcumin in the context of absorption kinetics and physiological effects in humans. The aim of this research study was therefore to assess the influence of a formulated bioavailable form of curcumin (HydroCurc™) on acute iron absorption in humans. To the best knowledge of the authors, curcuminoid constituents have not been assessed previously in humans in relation to oral iron supplementation and in relation to circulatory iron levels.

## 2. Materials and Methods

### 2.1. Study Design

Briefly, 155 healthy participants were recruited in a double blind, placebo-controlled randomized trial. Blocked, gender balanced randomisation was carried out using Study Randomizer software [38], and sample size was calculated using G\*Power 3.1.9.2 software [39,40].

### 2.2. Ethics Approval and Study Registration

The study protocol was approved by the Faculty of Science and Technology Ethics Committee, University of Westminster under the application identification: ETH1718-0907. The study and the study protocol are registered with ClinicalTrials.gov (NCT04465851) and public since 7 June 2020.

### 2.3. Inclusion and Exclusion Criteria

All participants enrolled in the study aligned with the inclusion criteria of being healthy, aged 18–40 years, with ferritin levels in the normal range for the United Kingdom (UK; 15–300 µg/L for men and 15–200 µg/L for women) according to Dooley and Worwood, and Fitzsimons and colleagues [41,42]. Any prospective participants with deficient haemoglobin (HGB) levels (<130 g/L for men and <120 g/L for women) were not eligible for inclusion in the study [5].


Participants were excluded from the trial if they had any diagnosis of medical condition or comorbidities as per the exclusion criteria (such as, currently trying to conceive, pregnant, lactating, experiencing any chronic menstrual disorders, or reported undergoing any menopausal changes). Additionally, participants were excluded if they had any issues related to oral supplement ingestion; were on any medication or supplementation; if their alcohol consumption exceeded 21 units/week; or if they experienced any chronic GI symptoms, eating disorders, psychological conditions, or any hypo/hypertensive blood pressure (BP) measurements [43,44].

### 2.4. Safety Screening

All potential study participants were screened during a medical history interview for health conditions that may preclude participation in the study, conducted in 2018–2019, with respect to the inclusion/exclusion criteria. Interviews were supported with the use of the General Symptoms Questionnaire (GSQ-65). All participants were interviewed for a duration of 60 min and included the use of open-ended questions to gain comprehensive understanding of their health status. Furthermore, a fasting (12 h, overnight) blood sample was collected (for ferritin and full blood count analysis) and BP was measured using Omron M6 Comfort blood pressure monitor, (Omron, Hoofddorp, The Netherlands). Participants were not enrolled if anomalous results or findings from the safety screening were established.

### 2.5. Participants

Study participants were randomly allocated to one of five different treatment groups using blocked, gender balanced randomisation (with 31 participants in each group) using the Study Randomizer software [38]. The five different treatment groups were ferrous sulphate placebo + curcumin placebo (FS0\_Plac), ferrous sulphate–18 mg elemental iron + placebo (FS18\_Plac), ferrous sulphate–18 mg elemental iron + 500 mg curcumin (FS18\_Curc), ferrous sulphate–65 mg elemental iron + placebo (FS65\_Plac), and ferrous sulphate–65 mg elemental iron +500 mg curcumin (FS65\_Curc) (Figure 1).



**Figure 1.** Study design. Random allocation to treatment groups post safety screening ( $n = 31$ /group). At baseline (0 min), both body measurements and blood samples were collected. At end-point (180 min), only blood samples were collected.

## 2.6. Supplements

Participants were provided with two different doses of ferrous sulphate iron supplements depending on the group they were randomized to. The higher dose of ferrous sulphate was 200 mg with 65 mg elemental iron and is based upon the traditional first line oral iron therapy for treatment and prophylaxis of ID and ID anaemia generally used worldwide [45]. The lower dose was 55 mg ferrous sulphate containing 18 mg elemental iron, based upon the recommended Daily Value (DV) of iron as per the US Food and Drug Administration (FDA) [45,46]. Formulated curcumin supplement (HydroCurc™, Pharmako Biotechnologies Pty Ltd. New South Wales, Australia) included 500 mg/day comprising of 80% curcumin, 17% DMC, and 3% BDMC (~85% total curcuminoids), entrapped in a patented delivery system (LipiSpurse®) [47]. For the placebo, microcrystalline cellulose was used. This cellulose was also used as an inert bulking agent in the capsules of active ingredients. The ferrous sulphate and curcumin supplements were packed into white-opaque hydroxypropyl methylcellulose (HPMC) capsules (sizes used were #1 and #00). Participants were provided with one ferrous sulphate iron at 0 min and with one curcumin supplement 60 min after the iron supplement on the day of the trial, in the fasted state, by trained research staff.

## 2.7. Physical Examination

Blood pressure was measured by a trained research associate or research assistant as described in the safety screening section. Anthropometric measurements were collected on the day of the study (baseline measures) by trained research staff. Height was measured to the nearest millimeter (without shoes) using a Seca 247 stadiometer (Seca GmbH & Co. KG, Hamburg, Germany). Weight, body mass index (BMI), and body fat % were measured using Seca mBCA 515 medical Body Composition Analyzer (Seca GmbH & Co. KG, Hamburg, Germany). Care was taken to instruct participants to void bladder of urine, empty pockets, and remove any excessive clothing and any jewellery prior to all measurements. Waist circumference was obtained using a soft tape measure at the narrowest point between the lower rib and the top of the hip bone.

### 2.8. Blood Sample Collection and Processing

Blood tests were carried out for each participant, following an overnight fast (12-h fast). Venous blood samples were collected (by certified phlebotomists) by venipuncture (from the antecubital fossa) using a 21G needle, at time 0 min and 180 min after iron supplementation. Approximately 10 mL of blood was collected from each participant using a Becton Dickinson (BD) Vacutainer<sup>®</sup> serum-separating tube (SST) and a BD Vacutainer<sup>®</sup> ethylenediamine tetra-acetic acid (EDTA) tube (BD, Oxford, UK). Blood in the SSTs were left to coagulate at room temperature for 45 min and then centrifuged (Hettich 340r, Hettich GmbH & Co. KG, Tuttlingen, Germany). To render samples acellular, centrifugation was performed for 10 min at 5000 rpm/3857 × g at 4 °C. Post centrifugation, aliquots of serum supernatant were stored at −80 °C in 1.5 mL microcentrifuge tubes. The EDTA tubes were kept on ice prior to whole blood analysis (on the day of collection) and then centrifuged for 10 min at 3857 × g at 4 °C. Similarly, post centrifugation, aliquots of plasma supernatant were stored at −80 °C immediately.

### 2.9. Analysis of Serum Ferritin Concentration

Serum ferritin samples were defrosted and analysed using a Horiba ABX Pentra 400 (Horiba Ltd., Kyoto, Japan) multiparametric medical bench top chemistry analyser and the Horiba ABX Pentra CRP CP High Sensitivity and ABX Ferritin 2 CP reagents. This chemistry analyser is compliant with the National Committee for Clinical Laboratory Standards (NCCLS) [48]. In accordance with the manufactures processes, ferritin levels were determined by latex-enhanced immunoturbidimetric assay [49].

### 2.10. Analysis of Serum Iron Profile

Serum iron, total iron binding capacity (TIBC), transferrin saturation (TS), and unsaturated iron-binding capacity (UIBC) were assessed externally (using necessary standards and controls for each analyte) at Health Services laboratories (HSL Analytics LLP, London, UK); Project No. P197 as primary outcome measures of this study. Serum iron and UIBC were measured by direct colorimetric method, and subsequently, serum TIBC and TS were calculated using the equations from Elsayed et al. [50]:

$$\text{TIBC} = \text{serum iron} + \text{UIBC}$$

$$\text{TS} = \text{serum iron} / (\text{serum iron} + \text{UIBC}) \times 100$$

### 2.11. Analysis of Whole Blood Haemoglobin (HGB) Concentration

HGB, a secondary outcome measure, was analysed using the Sysmex XP-300 (Sysmex Corporation, Kobe, Japan) automated haematology analyser. The Sysmex XP-300 is compliant with International Standards Organisation (ISO)/International Electrotechnical Commission (IEC) 17043:2010, via the Sysmex Network Communication Service. The Sysmex instrument aspirated 50 µL of each sample. HGB was analysed using a non-cyanide haemoglobin detection method [51]. The XP-300 was calibrated by Sysmex Corporation according to the manufacturer's specifications, and quality was guaranteed using Sysmex internal quality control.

### 2.12. Analysis of Serum Thiobarbituric Acid Reactive Substances (TBARS) Concentration

Serum TBARS were determined by TBARS parameter assay kit (R&D Systems, Minneapolis, MN, USA), with a sensitivity of 0.007–0.055 µM minimum detectable dose range as another secondary outcome measure of the study. The assay procedure was performed according to the manufacturer's instructions and plates were read using a microplate reader (SPECTROstar<sup>®</sup> Nano, BMG Labtech GmbH, Ortenberg, Germany) at the absorbance of 530 nm.

### 2.13. Statistical Analysis

Values are expressed as mean  $\pm$  Standard Error of Mean (SEM). All parameters were checked for normality using Shapiro–Wilk (S-W) tests, and results were statistically analysed using a two-way, repeated measures analysis of variance (ANOVA) or mixed effects model [52–54]. Post-hoc tests (Sidak's and Tukey's) were applied to assess differences between and within treatment groups with multiplicity adjusted P values reported for each comparison. Data analysis was carried out using PRISM software package (Version 9, Graphpad Software Inc., San Diego, CA, USA) and IBM Statistical Package for the Social Sciences (SPSS, Version 25.0, IBM Corp., New York, NY, USA).

## 3. Results

### 3.1. Participant Characteristics

Of the 155 participants that started the study, there were 79 males (26.42 years  $\pm$  0.55) and 76 females (25.82 years  $\pm$  0.54). Participant 105 was excluded from further analysis (Figure S1) due to BMI of 49.65 kg/m<sup>2</sup>, the only BMI that fell in class 3 obese category ( $\geq 40$  kg/m<sup>2</sup>) [55,56] and the only value detected as 3\*IQR or an extreme outlier in SPSS. The age of participants ranged from 19 to 39 years of age. Overall, the mean age of the included 154 participants was 26.12 years ( $\pm$  0.39). In addition, mean height was 1.71 m ( $\pm$ 0.01), and mean weight was 68.93 kg ( $\pm$  1.13). Anthropometric characteristics of participants within each of the five treatment groups are presented in Table 1.

**Table 1.** Anthropometric characteristics of participants split by gender.

Characteristics	FS0_Plac		FS18_Plac		FS18_Curc		FS65_Plac		FS65_Curc		Total	
	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female	Male	Female
Age (years)	25.87 $\pm$ 1.18	26.73 $\pm$ 1.31	26.11 $\pm$ 1.27	25.08 $\pm$ 1.47	25.60 $\pm$ 1.13	23.53 $\pm$ 1.24	26.92 $\pm$ 1.25	27.06 $\pm$ 1.14	27.12 $\pm$ 1.35	26.36 $\pm$ 1.08	26.32 $\pm$ 0.55	25.81 $\pm$ 0.56
	(n = 15)	(n = 16)	(n = 18)	(n = 13)	(n = 15)	(n = 16)	(n = 13)	(n = 17)	(n = 17)	(n = 14)	(n = 78)	(n = 76)
Height (m)	1.77 $\pm$ 0.02	1.67 $\pm$ 0.02	1.76 $\pm$ 0.01	1.64 $\pm$ 0.01	1.77 $\pm$ 0.02	1.59 $\pm$ 0.02	1.78 $\pm$ 0.01	1.63 $\pm$ 0.02	1.78 $\pm$ 0.2	1.65 $\pm$ 0.02	1.77 $\pm$ 0.01	1.64 $\pm$ 0.01
	(n = 15)	(n = 15)	(n = 18)	(n = 12)	(n = 15)	(n = 15)	(n = 13)	(n = 17)	(n = 17)	(n = 14)	(n = 78)	(n = 73)
Weight (kg)	75.22 $\pm$ 3.15	66.36 $\pm$ 3.25	80.70 $\pm$ 3.70	60.08 $\pm$ 3.07	74.58 $\pm$ 2.78	57.76 $\pm$ 2.64	77.13 $\pm$ 2.27	60.27 $\pm$ 2.89	75.24 $\pm$ 2.26	58.54 $\pm$ 1.91	76.69 $\pm$ 1.32	60.64 $\pm$ 1.28
	(n = 15)	(n = 15)	(n = 18)	(n = 12)	(n = 15)	(n = 15)	(n = 13)	(n = 17)	(n = 17)	(n = 14)	(n = 78)	(n = 73)
BMI (kg/m <sup>2</sup> )	24.06 $\pm$ 0.83	23.72 $\pm$ 0.85	25.94 $\pm$ 1.06	22.38 $\pm$ 1.21	23.71 $\pm$ 0.79	22.93 $\pm$ 1.03	24.32 $\pm$ 0.56	22.46 $\pm$ 0.82	23.95 $\pm$ 0.77	21.48 $\pm$ 0.65	24.44 $\pm$ 0.38	22.61 $\pm$ 0.41
	(n = 15)	(n = 15)	(n = 18)	(n = 12)	(n = 15)	(n = 15)	(n = 13)	(n = 17)	(n = 17)	(n = 14)	(n = 78)	(n = 73)
Body Fat (%)	18.37 $\pm$ 1.68	32.16 $\pm$ 1.57	21.97 $\pm$ 2.02	29.45 $\pm$ 2.44	18.35 $\pm$ 1.86	30.29 $\pm$ 2.07	19.13 $\pm$ 1.51	27.95 $\pm$ 1.52	19.44 $\pm$ 1.85	28.18 $\pm$ 1.25	19.56 $\pm$ 0.82	29.59 $\pm$ 0.79
	(n = 15)	(n = 15)	(n = 18)	(n = 12)	(n = 15)	(n = 15)	(n = 13)	(n = 17)	(n = 17)	(n = 14)	(n = 78)	(n = 73)


Values presented in mean  $\pm$  standard error of mean (SEM).

### 3.2. Baseline Levels of Assessed Biomarkers

Baseline comparisons between each of the supplementation groups showed no statistical difference in any of the measured biological marker levels.

### 3.3. Effects of Acute Supplementation on Serum Iron Levels

When assessing the effects of each treatment within the groups, there was a significant difference observed over time ( $F(1, 144) = 331.9, p < 0.0001$ ) with statistically significant intra-group increases in mean serum iron after 180 min ( $p < 0.0001$ ) in the FS18\_Plac, FS18\_Curc, FS65\_Plac, and FS65\_Curc groups. The lower dose groups, FS18\_Curc and the corresponding curcumin placebo group FS18\_Plac, after 180 min, had a 11.41  $\mu\text{mol/L}$  and 8.79  $\mu\text{mol/L}$  increase in serum iron, respectively. The higher dose groups, FS65\_Curc and the corresponding iron and placebo group FS65\_Plac, after 180 min, had 16.39  $\mu\text{mol/L}$  and 19.09  $\mu\text{mol/L}$  increase in serum iron, respectively. There was no significant increase observed in the FS0\_Plac group (Figure 2).



**Figure 2.** Serum iron levels at baseline and 180 min following supplementation. Results are presented as mean  $\pm$  SEM Serum iron levels ( $\mu\text{mol/L}$ ). *ns* indicates no significant difference ( $p > 0.05$ ); \* represents significance values when comparing time points within the same the conditions (\*\*\*\*  $p < 0.0001$ ).

Inter-group analysis of serum iron levels at 180 min showed significant between group difference ( $F(4, 149) = 10.73, p \leq 0.0001$ ). Statistically significant differences were observed between each treatment group except FS18\_Plac vs. FS18\_Curc, FS18\_Curc vs. FS65\_Curc and FS65\_Plac vs. FS65\_Curc (Table 2).


**Table 2.** Inter-group differences in serum iron levels ( $\mu\text{mol/L}$ ) at 180 min.

Groups		MD $\pm$ SED	Sig. *	p-Value
FS0_Plac	vs. FS18_Plac	$-7.18 \pm 2.18$	**	0.0096
FS0_Plac	vs. FS18_Curc	$-10.02 \pm 2.16$	****	<0.0001
FS0_Plac	vs. FS65_Plac	$-19.88 \pm 2.17$	****	<0.0001
FS0_Plac	vs. FS65_Curc	$-15.63 \pm 2.16$	****	<0.0001
FS18_Plac	vs. FS18_Curc	$-2.84 \pm 2.19$	NS	0.6917
FS18_Plac	vs. FS65_Plac	$-12.71 \pm 2.19$	****	<0.0001
FS18_Plac	vs. FS65_Curc	$-8.45 \pm 2.19$	**	0.0013
FS18_Curc	vs. FS65_Plac	$-9.86 \pm 2.18$	****	<0.0001
FS18_Curc	vs. FS65_Curc	$-5.61 \pm 2.18$	NS	0.0771
FS65_Plac	vs. FS65_Curc	$4.25 \pm 2.18$	NS	0.2935

Results presented as mean difference (MD)  $\pm$  standard error of difference (SED). Sig.\* represents significance values when comparing groups at the same timepoint (\*\*  $p < 0.01$ , \*\*\*\*  $p < 0.0001$ , NS  $p > 0.05$ ).

### 3.4. Effects of Acute Supplementation on Serum TIBC Levels

When evaluating the effect of treatment group on serum TIBC levels after 180 min of supplementation, a significant difference was observed between the two time points ( $F(1, 137) = 19.76, p < 0.0001$ ). A significant increase of  $2.36 \mu\text{mol/L}$  ( $p = 0.0102$ ) and  $2.11 \mu\text{mol/L}$  ( $p = 0.0406$ ) in TIBC levels from baseline to end-point was observed in the treatment groups FS0\_Plac and FS18\_Plac, respectively. There was no significant difference observed within the rest of treatment groups (FS18\_Curc, FS65\_Plac and FS65\_Curc) over time (Figure 3).




**Figure 3.** Effect of 180 min supplementation on mean serum total iron binding capacity (TIBC,  $\mu\text{mol/L}$ ) levels (mean, SEM). \* represents significance values when comparing time points within the same conditions ( $p < 0.05$ ); ns indicates no significant difference ( $p > 0.05$ ).

There were no significant differences observed in TIBC levels between groups 180 min after supplementation ( $F(4, 148) = 1.143, p = 0.3387$ ).

### 3.5. Effects of Acute Supplementation on Serum UIBC Levels

When assessing the effects of each treatment within the groups, there was a significant difference observed over time ( $F(1, 137) = 210.8, p < 0.0001$ ) with statistically significant intra-group decreases in mean serum UIBC levels in the FS18\_Plac ( $6.61 \mu\text{mol/L}, p = 0.0002$ ), FS18\_Curc ( $9.69 \mu\text{mol/L}, p < 0.0001$ ), FS65\_Plac ( $18.06 \mu\text{mol/L}, p < 0.0001$ ), and FS65\_Curc ( $15.69 \mu\text{mol/L}, p < 0.0001$ ) groups over time. There was no significant decrease observed in the FS0\_Plac group (Figure 4).



**Figure 4.** Effect of 180 min supplementation on mean serum unsaturated iron-binding capacity (UIBC,  $\mu\text{mol/L}$ ) levels (mean, SEM). \* represents significance values when comparing time points within the same the conditions (\*\*\*  $p = 0.0002$ , \*\*\*\*  $p < 0.0001$ ); ns indicates no significant difference ( $p > 0.05$ ).



When assessing the effects of each treatment on UIBC levels between groups at 180 min, a significant difference ( $F(4, 148) = 7.226, p < 0.0001$ ) was detected. Statistically significant differences were observed between each treatment group except FS0\_Plac vs FS18\_Curc, FS18\_Plac vs FS18\_Curc, and FS65\_Plac vs FS65\_Curc (Table 3).


**Table 3.** Inter-group differences in serum unsaturated iron-binding capacity (UIBC) levels ( $\mu\text{mol/L}$ ) at 180 min.

Groups		MD $\pm$ SED		Sig. *	p-Value
FS0_Plac	vs.	FS18_Plac	4.57 $\pm$ 2.80	NS	0.4782
FS0_Plac	vs.	FS18_Curc	9.56 $\pm$ 2.78	**	0.0062
FS0_Plac	vs.	FS65_Plac	19.43 $\pm$ 2.83	****	<0.0001
FS0_Plac	vs.	FS65_Curc	18.10 $\pm$ 2.81	****	<0.0001
FS18_Plac	vs.	FS18_Curc	4.99 $\pm$ 2.83	NS	0.3944
FS18_Plac	vs.	FS65_Plac	7.00 $\pm$ 2.86	****	<0.0001
FS18_Plac	vs.	FS65_Curc	13.53 $\pm$ 2.85	****	<0.0001
FS18_Curc	vs.	FS65_Plac	9.87 $\pm$ 2.85	**	0.0055
FS18_Curc	vs.	FS65_Curc	8.54 $\pm$ 2.84	*	0.0235
FS65_Plac	vs.	FS65_Curc	-1.33 $\pm$ 2.87	NS	0.9906

Results presented as mean difference (MD)  $\pm$  standard error of difference (SED). Sig.\* represents significance values when comparing groups at the same timepoint (\*  $p < 0.05$ , \*\*  $p < 0.01$ , \*\*\*\*  $p < 0.0001$ , NS  $p > 0.05$ ).

### 3.6. Effects of Acute Supplementation on Serum Transferrin Saturation (TS) %

When assessing the effects of each treatment on serum TS after 180 min of supplementation, a within group statistically significant difference was observed ( $F(1, 137) = 308.4, p < 0.0001$ ). There were statistically significant ( $p < 0.0001$ ) increases of 12.77%, 17.24%, 30.00%, and 27.96% in TS observed in the FS18\_Plac, FS18\_Curc, FS65\_Plac, and FS65\_Curc, respectively. There was no significant increase observed in the FS0\_Plac group over time (Figure 5).



**Figure 5.** Effect of 180 min supplementation on mean serum transferrin saturation (TS) % (mean, SEM). \* represents significance values when comparing time points within the same the conditions (\*\*\*\*  $p < 0.0001$ ); ns represents no significant difference ( $p > 0.05$ ).

When assessing the inter-group effects of each treatment on TS at 180 min, a significant difference ( $F(4, 148) = 10.39, p < 0.0001$ ) was detected. Statistically significant differences were observed between each treatment groups except FS0\_Plac vs. FS18\_Curc, FS18\_Plac vs. FS18\_Curc, and FS65\_Plac vs. FS65\_Curc (Table 4).


**Table 4.** Inter-group differences in transferrin saturation (TS %) at 180 min.

Groups		MD ± SED	Sig. *	p-Value	
FS0_Plac	vs.	FS18_Plac	−10.06 ± 3.78	NS	0.0627
FS0_Plac	vs.	FS18_Curc	−16.30 ± 3.76	***	0.0002
FS0_Plac	vs.	FS65_Plac	−32.20 ± 3.82	****	<0.0001
FS0_Plac	vs.	FS65_Curc	−29.21 ± 3.80	****	<0.0001
FS18_Plac	vs.	FS18_Curc	−6.24 ± 3.82	NS	0.4772
FS18_Plac	vs.	FS65_Plac	−22.14 ± 3.88	****	<0.0001
FS18_Plac	vs.	FS65_Curc	−19.14 ± 3.85	****	<0.0001
FS18_Curc	vs.	FS65_Plac	−15.90 ± 3.86	***	0.0005
FS18_Curc	vs.	FS65_Curc	−12.90 ± 3.83	**	0.0077
FS65_Plac	vs.	FS65_Curc	2.99 ± 3.89	NS	0.9388

Results presented as mean difference (MD) ± standard error of difference (SED). Sig.\* represents significance values when comparing groups at the same timepoint (\*\*  $p < 0.01$ , \*\*\*  $p < 0.001$ , \*\*\*\*  $p < 0.0001$ , NS  $p > 0.05$ ).

### 3.7. Effects of Acute Supplementation on Whole Blood HGB Levels

When evaluating the effect of treatment group on whole blood HGB levels after 180 min of supplementation, a significant difference was observed between the two time points ( $F(1, 147) = 22.35, p < 0.0001$ ). A significant increase of 1.55%/2.04 g/L in HGB levels from baseline to end-point (180 min) was observed in participants taking FS65\_Curc (\*  $p < 0.05$ ) (Figure 6 and Figure S2).



**Figure 6.** Haemoglobin (HGB) levels are expressed as percentage difference compared to baseline levels (SEM). \* represents significant difference over time (\*  $p < 0.05$ ).

There was no significant difference observed between treatment groups after 180 min of supplementation ( $F(4, 149) = 0.1042, p = 0.9809$ ).

### 3.8. Effects of Acute Supplementation on Serum TBARS Levels

There were no significant intra- ( $F(1, 144) = 0.1889, p = 0.6645$ ) or inter-group ( $F(4, 149) = 0.9583, p = 0.4323$ ) differences observed in TBARS levels (level of oxidative stress) in this acute study as a result of iron and curcumin co-administration after 180 min.

## 4. Discussion

The polyphenol curcumin has been purported to have many health benefits as a result of its potential to target numerous signaling avenues and its impact on cellular level activity [57]. Despite its significant therapeutic potential, the pharmacological use

of curcumin has been limited due to its poor bioavailability, limited bio-distribution, poor stability, and short activity half-life [58]. Formulation science strategies have shown that carrier-based delivery systems such as liposomes and micelles may address these limitations and enhance curcumin's therapeutic potential [59–61]. This study utilised curcumin supplement in the form of HydroCurc™, a formulation consisting of 85% total curcuminoids entrapped in a proprietary delivery system (LipiSpense®) that has been shown to enhance bioavailability in humans [47].

Studies assessing the effects of curcumin on iron levels have provided mixed results. In murine models, it was shown that curcumin decreases systemic iron levels when administered consistently over several months [31,62,63]. Curcumin was shown to moderately chelate iron-forming insoluble complexes that may influence iron absorption in the gut, although this has largely been shown in in-vitro studies [64].

In the current study, co-administration of ferrous iron and curcumin in the form of HydroCurc™ does not negatively affect iron absorption compared to ferrous iron alone. Mean increases in serum iron after 180 min, in both FS65\_Plac and FS65\_Curc, are in line with expected increases in the literature [65–68]. Furthermore, HGB levels were also significantly elevated at 180 min compared to baseline in the group receiving the high iron dose of 65 mg ferrous sulphate with HydroCurc™. HGB increases can result from dehydration or a reduction in the plasma vascular fraction [69]. However, this is unlikely to explain the resulting increase in HGB levels as all participants across the study were given free access to drinking water ad libitum.

Current understanding of iron homeostasis indicates that iron absorbed via the GI tract would first need to pass through the liver and bone marrow, before being incorporated into HGB during erythropoiesis [70]. However, this seems unlikely to explain the result in the current study given the short time frame of 180 min (3 h). Longitudinal research into iron supplementation, lasting several weeks (rather than hours), demonstrated significantly increased HGB levels in mildly anaemic populations [71]. However, the authors of the current study did not find any studies reporting acute changes (within 3 h) in HGB levels in a healthy population with normal serum levels. Although the researchers cannot rule out the possibility that this acute change is due to a physiological process, it cannot be explained based on current understanding of iron incorporation into HGB during erythropoiesis [70]. This outcome presents an intriguing avenue for in-depth exploration in further investigations.

The current study used serum iron analysis to evaluate iron absorption, as it increases in response to changes in dietary intakes of iron [3]. Dietary non-haem iron, typically found in the ferric ( $\text{Fe}^{3+}$ ) iron form, is oxidised into absorbable soluble ferrous ( $\text{Fe}^{2+}$ ) iron in the gastric and intestinal pH environment [72,73]. Given reports of diurnal changes in serum iron concentrations [74–76] and previous recommendations to standardize sampling [50], all participants' baseline measures of serum iron were assessed in the morning following an overnight fast (12 h fast). Following randomization, all participants were observed to take oral ferrous sulphate on an empty stomach with water only (to circumvent any confounding effect of dietary components, which could influence iron absorption in an acute absorption test) [68]. To assess acute iron absorption from ferrous sulphate, it is advised to administer a dose yielding 65 mg of elemental iron and assess serum iron at 2–3 h post supplementation [67,68]. This method is used to evaluate gut iron absorption in humans [68,77,78]. If gut iron absorption is adequate, there is approximately a 17.9  $\mu\text{mol/L}$  increase in serum iron levels as suggested by Alleyne et al. [68] or an increase of two to three times above baseline levels [67].

The current study also demonstrates that the increases in serum iron are translated into an increase in iron transportation via transferrin saturation levels. All groups receiving iron demonstrated an increase in transferrin saturation (TS%) and in a dose-related manner. Clinically, this measure is used as an indicator of iron-related diseases such as iron overload e.g., haemochromatosis or iron insufficiency e.g., iron deficiency anaemia [50]. However, in the current study, transferrin saturation (%) is an indicator of the saturation in iron

transport capacity between different treatments and iron doses. Interestingly, there were no differences in transferrin saturation in the presence of HydroCurc™ compared to iron alone at either dose of iron. This finding indicates that HydroCurc™ has no effect on iron uptake or bioavailability at low or high doses of iron. Additionally, the dose response differences in transferrin saturation (%) as well as unsaturated iron binding capacity (UIBC) between the 18 mg and 65 mg groups indicates additional reserve capacity in the iron metabolism pathways in healthy participants to mobilise different doses of iron.

## 5. Conclusions

In the current study, regardless of ferrous dose (18 mg or 65 mg iron), formulated curcumin in the form of HydroCurc™ does not inhibit ferrous iron absorption following ingestion. Furthermore, the significant increase in HGB levels observed in the FS65\_Plac group is retained in the FS65\_Curc group, demonstrating that the serum iron levels result in a measurable physiological effect that is not diminished by the addition of formulated curcumin. To the best knowledge of the authors, this study has demonstrated for the first time that formulated curcumin does not appear to negatively influence acute iron absorption in healthy humans.

**Supplementary Materials:** The following are available online at <https://www.mdpi.com/article/10.3390/nu13072300/s1>, Figure S1: The study compliance after 155 participants were enrolled and randomised equally into 5 treatment groups; Figure S2: Effect of 180 min supplementation on mean Haemoglobin (HGB) levels.

**Author Contributions:** Conceptualisation and methodology: M.G.Z., D.R. and G.B. Supervision: M.G.Z., D.R. and G.B. Investigation and data curation: H.T.L. Data analysis: H.T.L. Writing—original draft preparation: H.T.L., D.R. and M.G.Z. Writing—review and editing: M.G.Z., D.R., G.B. and H.T.L. All authors have read and agreed to the published version of the manuscript.

**Funding:** The study was supported by Knowledge Exchange funding from Gencor Pacific Ltd. (UT722720).

**Institutional Review Board Statement:** The study was conducted according to the guidelines of the Declaration of Helsinki, and approved by the Faculty of Science and Technology Ethics Committee, University of Westminster (ID: ETH1718-0907 22 February 2018).

**Informed Consent Statement:** Informed consent was obtained from all subjects involved in the study.

**Data Availability Statement:** The data presented in this study are available on request from the corresponding author. The data are not publicly available due to ethical, legal and privacy issues.

**Acknowledgments:** We gratefully acknowledge the time and dedication of all participants who participated in this study. We would like to acknowledge all colleagues from The University of Westminster who contributed support and expertise to this study. We thank Owen Fitzsimons for assistance with participant enrolment and randomization, phlebotomy and investigation, Lina Temouri for assistance with administrative duties, phlebotomy and laboratory work, Chrystalla Ferrier for assisting with the operation of the Ilab Aries instrument and Yvoni Kyriakidou for assistance with phlebotomy. We thank Sonita Chea, Hashum Sum and Ruchita Venkatesh for their laboratory and administration assistance and Gloria Koncz for her administrative assistance. The authors would also like to thank the Research Interns and Visiting Researchers who assisted over the course of this project.

**Conflicts of Interest:** The funder was involved in the study design process but had no involvement in the data collection or analysis. The authors declare no other conflict of interest.

## References

1. Aisen, P.; Enns, C.; Wessling-Resnick, M. Chemistry and Biology of Eukaryotic Iron Metabolism. *Int. J. Biochem. Cell Biol.* **2001**, *33*, 940–959. [\[CrossRef\]](#)
2. Lieu, P.T.; Heiskala, M.; Peterson, P.A.; Yang, Y. The Roles of Iron in Health and Disease. *Mol. Asp. Med.* **2001**, *22*, 1–87. [\[CrossRef\]](#)
3. Lopez, A.; Cacoub, P.; Macdougall, I.C.; Peyrin-Biroulet, L. Iron Deficiency Anaemia. *Lancet* **2016**, *387*, 907–916. [\[CrossRef\]](#)

4. Gakidou, E.; Afshin, A.; Abajobir, A.A.; Abate, K.H.; Abbafati, C.; Abbas, K.M.; Abd-Allah, F.; Abdulle, A.M.; Abera, S.F.; Aboyans, V.; et al. Global, Regional, and National Comparative Risk Assessment of 84 Behavioural, Environmental and Occupational, and Metabolic Risks or Clusters of Risks, 1990–2016: A Systematic Analysis for the Global Burden of Disease Study 2016. *Lancet* **2017**, *390*, 1345–1422. [[CrossRef](#)]
5. World Health Organization. *Nutritional Anaemias: Tools for Effective Prevention and Control*; WHO: Geneva, Switzerland, 2017.
6. Roberts, D.J.; Weatherall, D.J. Introduction: The Complexity and Challenge of Preventing, Treating, and Managing Blood Diseases in the Developing Countries. *Hematol. Oncol. Clin. North Am.* **2016**, *30*, 233–246. [[CrossRef](#)] [[PubMed](#)]
7. Haas, J.D.; Brownlie IV, T. Iron Deficiency and Reduced Work Capacity: A Critical Review of the Research to Determine a Causal Relationship. *J. Nutr.* **2001**, *131*, 676S–690S. [[CrossRef](#)] [[PubMed](#)]
8. Brownlie IV, T.; Utermohlen, V.; Hinton, P.S.; Giordano, C.; Haas, J.D. Marginal Iron Deficiency without Anemia Impairs Aerobic Adaptation among Previously Untrained Women. *Am. J. Clin. Nutr.* **2002**, *75*, 734–742. [[CrossRef](#)]
9. Jáuregui-Lobera, I. Iron Deficiency and Cognitive Functions. *Neuropsychiatr. Dis. Treat.* **2014**, *10*, 2087–2095. [[CrossRef](#)]
10. Patterson, A.J.; Brown, W.J.; Powers, J.R.; Roberts, D.C.K. Iron Deficiency, General Health and Fatigue: Results from the Australian Longitudinal Study on Women’s Health. *Qual. Life Res.* **2000**, *9*, 491–497. [[CrossRef](#)]
11. Enjuanes, C.; Klip, I.T.; Bruguera, J.; Cladellas, M.; Ponikowski, P.; Banasiak, W.; Van Veldhuisen, D.J.; Van Der Meer, P.; Jankowska, E.A.; Comín-Colet, J. Iron Deficiency and Health-Related Quality of Life in Chronic Heart Failure: Results from a Multicenter European Study. *Int. J. Cardiol.* **2014**, *174*, 268–275. [[CrossRef](#)]
12. NICE Anaemia—Iron Deficiency. Available online: <https://cks.nice.org.uk/topics/anaemia-iron-deficiency/> (accessed on 12 May 2021).
13. Tolkien, Z.; Stecher, L.; Mander, A.P.; Pereira, D.I.A.; Powell, J.J. Ferrous Sulfate Supplementation Causes Significant Gastrointestinal Side-Effects in Adults: A Systematic Review and Meta-Analysis. *PLoS ONE* **2015**, *10*, e0117383. [[CrossRef](#)]
14. Gastarena, M.A.I.; Gil, A.G.; Azqueta, A.; Coronel, M.P.; Gimeno, M. A Comparative Study on the Gastrointestinal Tolerance of Different Antianaemic Preparations. *Hum. Exp. Toxicol.* **2003**, *22*, 137–141. [[CrossRef](#)] [[PubMed](#)]
15. De Souza, A.I.; Filho, M.B.; Bresani, C.C.; Ferreira, L.O.C.; Figueiroa, J.N. Adherence and Side Effects of Three Ferrous Sulfate Treatment Regimens on Anemic Pregnant Women in Clinical Trials. *Cad. Saude Publica* **2009**, *25*, 1225–1233. [[CrossRef](#)] [[PubMed](#)]
16. DeLoughery, T.G. Safety of Oral and Intravenous Iron. *Acta Haematol.* **2019**, *142*, 8–12. [[CrossRef](#)] [[PubMed](#)]
17. Wessling-Resnick, M. Excess Iron: Considerations Related to Development and Early Growth. *Am. J. Clin. Nutr.* **2017**, *106*, 1600S–1605S. [[CrossRef](#)] [[PubMed](#)]
18. Assunção, M.; Santos-Marques, M.J.; Carvalho, F.; Andrade, J.P. Green Tea Averts Age-Dependent Decline of Hippocampal Signaling Systems Related to Antioxidant Defenses and Survival. *Free Radic. Biol. Med.* **2010**, *48*, 831–838. [[CrossRef](#)] [[PubMed](#)]
19. Fahnestock, M.; Marchese, M.; Head, E.; Pop, V.; Michalski, B.; Milgram, W.N.; Cotman, C.W. BDNF Increases with Behavioral Enrichment and an Antioxidant Diet in the Aged Dog. *Neurobiol. Aging* **2012**, *33*, 546–554. [[CrossRef](#)]
20. Kulkarni, R.; Deshpande, A.; Saxena, K.; Varma, M.; Sinha, A.R. Ginger Supplementary Therapy for Iron Absorption in Iron Deficiency Anemia. *Indian J. Tradit. Knowl.* **2012**, *11*, 78–80.
21. Ramezani, M.; Hatampour, M.; Sahebkar, A. Promising Anti-Tumor Properties of Bisdemethoxycurcumin: A Naturally Occurring Curcumin Analogue The Present Review Discusses the Findings on the Anti-Tumor Effects of BDMC, Underlying Mechanisms, and the Relevance of Finding for Translational Studies in Human. *Wiley Online Libr.* **2017**, *233*, 880–887. [[CrossRef](#)]
22. Pulla Reddy, A.C.; Lokesh, B.R. Studies on Spice Principles as Antioxidants in the Inhibition of Lipid Peroxidation of Rat Liver Mitochondria. *Mol. Cell. Biochem.* **1992**, *111*, 117–124. [[CrossRef](#)]
23. Hjorth Tønnesen, H.; Smistad, G.; Ågren, T.; Karlsen, J. Studies on Curcumin and Curcuminoids. XXIII: Effects of Curcumin on Liposomal Lipid Peroxidation. *Int. J. Pharm.* **1993**, *90*, 221–228. [[CrossRef](#)]
24. Borsari, M.; Ferrari, E.; Grandi, R.; Saladini, M. Curcuminoids as Potential New Iron-Chelating Agents: Spectroscopic, Polarographic and Potentiometric Study on Their Fe(III) Complexing Ability. *Inorg. Chim. Acta* **2002**, *328*, 61–68. [[CrossRef](#)]
25. Bernabé-Pineda, M.; Ramírez-Silva, M.T.; Romero-Romo, M.A.; González-Vergara, E.; Rojas-Hernández, A. Spectrophotometric and Electrochemical Determination of the Formation Constants of the Complexes Curcumin-Fe(III)-Water and Curcumin-Fe(II)-Water. *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.* **2004**, *60*, 1105–1113. [[CrossRef](#)]
26. Salem, M.; Rohani, S.; Gillies, E.R. Curcumin, a Promising Anti-Cancer Therapeutic: A Review of Its Chemical Properties, Bioactivity and Approaches to Cancer Cell Delivery. *RSC Adv.* **2014**, *4*, 10815–10829. [[CrossRef](#)]
27. Buhmann, C.; Mobasher, A.; Busch, F.; Aldinger, C.; Stahlmann, R.; Montaseri, A.; Shakibaei, M. Curcumin Modulates Nuclear Factor KB (Nf-KB)-Mediated Inflammation in Human Tenocytes in Vitro: Role of the Phosphatidylinositol 3-Kinase/Akt Pathway. *J. Biol. Chem.* **2011**, *286*, 28556–28566. [[CrossRef](#)]
28. Mishra, S.; Palanivelu, K. The Effect of Curcumin (Turmeric) on Alzheimer’s Disease: An Overview. *Ann. Indian Acad. Neurol.* **2008**, *11*, 13–19. [[CrossRef](#)] [[PubMed](#)]
29. Sarraf, P.; Parohan, M.; Javanbakht, M.H.; Ranji-Burachaloo, S.; Djalali, M. Short-Term Curcumin Supplementation Enhances Serum Brain-Derived Neurotrophic Factor in Adult Men and Women: A Systematic Review and Dose-Response Meta-Analysis of Randomized Controlled Trials. *Nutr. Res.* **2019**, *69*, 1–8. [[CrossRef](#)]
30. Tiekou Lorinczova, H.; Fitzsimons, O.; Mursaleen, L.; Renshaw, D.; Begum, G.; Zariwala, M.G. Co-Administration of Iron and a Bioavailable Curcumin Supplement Increases Serum BDNF Levels in Healthy Adults. *Antioxidants* **2020**, *9*, 645. [[CrossRef](#)] [[PubMed](#)]

31. Chin, D.; Huebbe, P.; Frank, J.; Rimbach, G.; Pallauf, K. Curcumin May Impair Iron Status When Fed to Mice for Six Months. *Redox Biol.* **2014**, *2*, 563–569. [CrossRef]
32. Cory, H.; Passarelli, S.; Szeto, J.; Tamez, M.; Mattei, J. The Role of Polyphenols in Human Health and Food Systems: A Mini-Review. *Front. Nutr.* **2018**, *5*, 87. [CrossRef] [PubMed]
33. Samba-Mondonga, M.; Constante, M.; Fragoso, G.; Calvé, A.; Santos, M.M. Curcumin Induces Mild Anemia in a DSS-Induced Colitis Mouse Model Maintained on an Iron-Sufficient Diet. *PLoS ONE* **2019**, *14*, e0208677. [CrossRef] [PubMed]
34. Tuntipopipat, S.; Zeder, C.; Siriprapa, P.; Charoenkiatkul, S. Inhibitory Effects of Spices and Herbs on Iron Availability. *Int. J. Food Sci. Nutr.* **2009**, *60*, 43–55. [CrossRef] [PubMed]
35. Bich, V.T.; Thuy, N.T.; Binh, N.T.; Huong, N.T.M.; Yen, P.N.D.; Luong, T.T. Structural and Spectral Properties of Curcumin and Metal- Curcumin Complex Derived from Turmeric (*Curcuma Longa*). In *Physics and Engineering of New Materials*; Springer Proceedings in Physics; Springer Science and Business Media, LLC: Berlin, Germany, 2009; Volume 127, pp. 271–278.
36. Ferrari, E.; Benassi, R.; Sacchi, S.; Pignedoli, F.; Asti, M.; Saladini, M. Curcumin Derivatives as Metal-Chelating Agents with Potential Multifunctional Activity for Pharmaceutical Applications. *J. Inorg. Biochem.* **2014**, *139*, 38–48. [CrossRef]
37. Srichairatanakool, S.; Thephinlap, C.; Phisalaphong, C.; Porter, J.; Fucharoen, S. Curcumin Contributes to In Vitro Removal of Non-Transferrin Bound Iron by Deferiprone and Desferrioxamine in Thalassaemic Plasma. *Med. Chem.* **2007**, *3*, 469–474. [CrossRef] [PubMed]
38. Study Randomizer App.Studyrandomizer.Com. Available online: <https://app.studyrandomizer.com> (accessed on 12 May 2021).
39. Faul, F.; Erdfelder, E.; Lang, A.G.; Buchner, A. G\*Power 3: A Flexible Statistical Power Analysis Program for the Social, Behavioral, and Biomedical Sciences. *Behav. Res. Methods* **2007**, *39*, 175–191. [CrossRef]
40. Erdfelder, E.; Faul, F.; Buchner, A.; Lang, A.G. Statistical Power Analyses Using G\*Power 3.1: Tests for Correlation and Regression Analyses. *Behav. Res. Methods* **2009**, *41*, 1149–1160. [CrossRef]
41. Dooley, J.; Worwood, M. *Guidelines on Diagnosis and Therapy: Genetic Haemochromatosis*; British Committee for Standards in Haematology: London, UK, 2000; pp. 1–33.
42. Fitzsimons, E.J.; Cullis, J.O.; Thomas, D.W.; Tsochatzis, E.; Griffiths, W.J.H. Diagnosis and Therapy of Genetic Haemochromatosis (Review and 2017 Update). *Br. J. Haematol.* **2018**, *181*, 293–303. [CrossRef]
43. Blood Pressure UK. Available online: <http://www.bloodpressureuk.org/your-blood-pressure/understanding-your-blood-pressure/what-do-the-numbers-mean/> (accessed on 12 May 2021).
44. Williams, B.; Mancia, G.; Spiering, W.; Rosei, E.A.; Azizi, M.; Burnier, M.; Clement, D.L.; Coca, A.; De Simone, G.; Dominiczak, A.; et al. 2018 ESC/ESH Guidelines for the Management of Arterial Hypertension. The Task Force for the Management of Arterial Hypertension of the European Society of Cardiology (ESC) and the European Society of Hypertension (ESH). *J. Hypertens* **2018**, *36*, 1953–2041. [CrossRef] [PubMed]
45. Goddard, A.F.; James, M.W.; McIntyre, A.S.; Scott, B.B. Guidelines for the Management of Iron Deficiency Anaemia. *Gut* **2011**, *60*, 1309–1316. [CrossRef]
46. Food and Drug Administration (FDA) Center for Food Safety and Applied Nutrition. *Guidance for Industry: A Food Labeling Guide*; Food and Drug Administration: College Park, MD, USA, 2013.
47. Briskey, D.; Sax, A.; Mallard, A.R.; Rao, A. Increased Bioavailability of Curcumin Using a Novel Dispersion Technology System (LipiSpense®). *Eur. J. Nutr.* **2019**, *58*, 2087–2097. [CrossRef]
48. Coudène, P.; Marson, B.; Badiou, S.; Flavier, S.; Anelli, S.; Cristol, J.P.; Dupuy, A.M. Evaluation of the ABX Pentra 400: A Newly Available Clinical Chemistry Analyser. *Clin. Chem. Lab. Med.* **2005**, *43*, 782–792. [CrossRef] [PubMed]
49. Simó, J.M.; Joven, J.; Clivillé, X.; Sans, T. Automated Latex Agglutination Immunoassay of Serum Ferritin with a Centrifugal Analyser. *Clin. Chem.* **1994**, *40*, 625–629. [CrossRef]
50. Elsayed, M.E.; Sharif, M.U.; Stack, A.G. Transferrin Saturation: A Body Iron Biomarker. In *Advances in Clinical Chemistry*; Academic Press Inc.: Cambridge, MA, USA, 2016; Volume 75, pp. 71–97.
51. Van Dievoet, M.A.; Louagie, H.; Ghys, T. Performance Evaluation of the Sysmex®XP-300 in an Oncology Setting: Evaluation and Comparison of Hematological Parameters with the Sysmex®XN-3000. *Int. J. Lab. Hematol.* **2016**, *38*, 490–496. [CrossRef] [PubMed]
52. Glass, G.V.; Peckham, P.D.; Sanders, J.R. Consequences of Failure to Meet Assumptions Underlying the Fixed Effects Analyses of Variance and Covariance. *Rev. Educ. Res.* **1972**, *42*, 237–288. [CrossRef]
53. Harwell, M.R.; Rubinstein, E.N.; Hayes, W.S.; Olds, C.C. Summarizing Monte Carlo Results in Methodological Research: The One- and Two-Factor Fixed Effects ANOVA Cases. *J. Educ. Stat.* **1992**, *17*, 315–339. [CrossRef]
54. Lix, L.M.; Keselman, J.C.; Keselman, H.J. Consequences of Assumption Violations Revisited: A Quantitative Review of Alternatives to the One-Way Analysis of Variance F Test. *Rev. Educ. Res.* **1996**, *66*, 579–619. [CrossRef]
55. WHO. Physical Status: The Use and Interpretation of Anthropometry, Report of a World Health Organization (WHO) Expert Committee. In *World Health Organization Technical Report Series*; WHO: Geneva, Switzerland, 1995; Volume 854, p. 452.
56. WHO. Obesity: Preventing and Managing the Global Epidemic. Report of a WHO Consultation. In *World Health Organization Technical Report Series*; World Health Organization: Geneva, Switzerland, 2000; Volume 894.
57. Hewlings, S.; Kalman, D. Curcumin: A Review of Its Effects on Human Health. *Foods* **2017**, *6*, 92. [CrossRef]
58. Prasad, S.; Gupta, S.C.; Tyagi, A.K.; Aggarwal, B.B. Curcumin, a Component of Golden Spice: From Bedside to Bench and Back. *Biotechnol. Adv.* **2014**, *32*, 1053–1064. [CrossRef]

59. Zhongfa, L.; Chiu, M.; Wang, J.; Chen, W.; Yen, W.; Fan-Havard, P.; Yee, L.D.; Chan, K.K.; Zhongfa, L.; Chiu, Á.M.; et al. Enhancement of Curcumin Oral Absorption and Pharmacokinetics of Curcuminoids and Curcumin Metabolites in Mice. *Cancer Chemother Pharm.* **2012**, *69*, 679–689. [[CrossRef](#)]
60. Zupančič, Š.; Kocbek, P.; Zariwala, M.G.; Renshaw, D.; Gul, M.O.; Elsaid, Z.; Taylor, K.M.G.; Somavarapu, S. Design and Development of Novel Mitochondrial Targeted Nanocarriers, Dqasomes for Curcumin Inhalation. *Mol. Pharm.* **2014**, *11*, 2334–2345. [[CrossRef](#)]
61. Mursaleen, L.; Somavarapu, S.; Zariwala, M.G. Deferoxamine and Curcumin Loaded Nanocarriers Protect Against Rotenone-Induced Neurotoxicity. *J. Parkinson's Dis.* **2020**, *10*, 99–111. [[CrossRef](#)] [[PubMed](#)]
62. Jiao, Y.; Wilkinson IV, J.; Di, X.; Wang, W.; Hatcher, H.; Kock, N.D.; D'Agostino, R.; Knovich, M.A.; Torti, F.M.; Torti, S.V. Curcumin, a Cancer Chemopreventive and Chemotherapeutic Agent, Is a Biologically Active Iron Chelator. *Blood* **2009**, *113*, 462–469. [[CrossRef](#)] [[PubMed](#)]
63. Badria, F.A.; Ibrahim, A.S.; Badria, A.F.; Elmarakby, A.A. Curcumin Attenuates Iron Accumulation and Oxidative Stress in the Liver and Spleen of Chronic Iron-Overloaded Rats. *PLoS ONE* **2015**, *10*, e0134156. [[CrossRef](#)] [[PubMed](#)]
64. Ak, T.; Gülçin, I. Antioxidant and Radical Scavenging Properties of Curcumin. *Chem. Biol. Interact.* **2008**, *174*, 27–37. [[CrossRef](#)] [[PubMed](#)]
65. Höglund, S. Iron Absorption in Apparently Healthy Men and Women: III. Studies in Iron Absorption. *Acta Med. Scand.* **1969**, *186*, 487–491. [[CrossRef](#)]
66. Ekenved, G.; Norrby, A.; Sölvell, L. Serum Iron Increase as a Measure of Iron Absorption—Studies on the Correlation with Total Absorption. *Scand. J. Haematol.* **1976**, *17*, 31–49. [[CrossRef](#)]
67. Wick, M.; Pinggera, W.; Lehmann, P. *Iron Metabolism: Diagnosis and Therapy of Anemias*; Springer Science & Business Media: Wien, Austria, 1996; ISBN 9783709136881.
68. Alleyne, M.; Horne, M.K.; Miller, J.L. Individualized Treatment for Iron-Deficiency Anemia in Adults. *Am. J. Med.* **2008**, *121*, 943–948. [[CrossRef](#)]
69. Pstras, L.; Debowska, M.; Wojcik-Zaluska, A.; Zaluska, W.; Waniewski, J. Hemodialysis-Induced Changes in Hematocrit, Hemoglobin and Total Protein: Implications for Relative Blood Volume Monitoring. *PLoS ONE* **2019**, *14*, e0220764. [[CrossRef](#)]
70. Anderson, G.J.; Frazer, D.M. Current Understanding of Iron Homeostasis. *Am. J. Clin. Nutr.* **2017**, *106*, 1559S–1566S. [[CrossRef](#)]
71. Schoorl, M.; Schoorl, M.; van der Gaag, D.; Bartels, P.C.M. Effects of Iron Supplementation on Red Blood Cell Hemoglobin Content in Pregnancy. *Hematol. Rep.* **2012**, *4*, 91–94. [[CrossRef](#)]
72. Porter, R.; FitzSimons, D.W. *Iron Metabolism*; Elsevier/Experta Medica: Amsterdam, The Netherlands, 1977; Volume 51, ISBN 9021940574.
73. Santiago, P. Ferrous versus Ferric Oral Iron Formulations for the Treatment of Iron Deficiency: A Clinical Overview. *Sci. World J.* **2012**, *2012*, 846824. [[CrossRef](#)] [[PubMed](#)]
74. Stengle, J.M.; Schade, A.L. Diurnal-Nocturnal Variations of Certain Blood Constituents in Normal Human Subjects: Plasma Iron, Siderophilin, Bilirubin, Copper, Total Serum Protein and Albumin, Haemoglobin and Haematocrit. *Br. J. Haematol.* **1957**, *3*, 117–124. [[CrossRef](#)]
75. Sinniah, R.; Doggart, J.R.; Neill, D.W. Diurnal Variations of the Serum Iron in Normal Subjects and in Patients with Haemochromatosis. *Br. J. Haematol.* **1969**, *17*, 351–358. [[CrossRef](#)]
76. Casale, G.; de Nicola, P. Circadian Rhythms in the Aged: A Review. *Arch. Gerontol. Geriatr.* **1984**, *3*, 267–284. [[CrossRef](#)]
77. Andersen, S.L.; Gyruup, C.; Handberg, A.; Nielsen, G.L. Oral Iron Absorption Test Should Not Be Performed with Iron Drops Containing Ferric Iron. *Dan. Med J.* **2015**, *62*, A5116.
78. Ginanjar, E.; Indrawati, L.; Setianingsih, I.; Atmakusumah, D.; Harahap, A.; Timan, I.S.; Marx, J.J.M. Iron Absorption in Iron-Deficient Women, Who Received 65 Mg Fe with an Indonesian Breakfast, Is Much Better from NaFe(III)EDTA than from Fe(II)SO<sub>4</sub>, with an Acceptable Increase of Plasma NTBI. A Randomized Clinical Trial. *Pharmaceuticals* **2018**, *11*, 85. [[CrossRef](#)] [[PubMed](#)]