

# Antimicrobial Peptides in Innate Immunity against Mycobacteria

Dong-Min Shin and Eun-Kyeong Jo\*

Department of Microbiology and Infection Signaling Network Research Center, Chungnam National University, School of Medicine, Daejeon 301-747, Korea

Antimicrobial peptides/proteins are ancient and naturally-occurring antibiotics in innate immune responses in a variety of organisms. Additionally, these peptides have been recognized as important signaling molecules in regulation of both innate and adaptive immunity. During mycobacterial infection, antimicrobial peptides including cathelicidin, defensin, and hepcidin have antimicrobial activities against mycobacteria, making them promising candidates for future drug development. Additionally, antimicrobial peptides act as immunomodulators in infectious and inflammatory conditions. Multiple crucial functions of cathelicidins in antimycobacterial immune defense have been characterized not only in terms of direct killing of mycobacteria but also as innate immune regulators, i.e., in secretion of cytokines and chemokines, and mediating autophagy activation. Defensin families are also important during mycobacterial infection and contribute to antimycobacterial defense and inhibition of mycobacterial growth both *in vitro* and *in vivo*. Hepcidin, although its role in mycobacterial infection has not yet been characterized, exerts antimycobacterial effects in activated macrophages. The present review focuses on recent efforts to elucidate the roles of host defense peptides in innate immunity to mycobacteria.

[Immune Network 2011;11(5):245-252]

## INTRODUCTION

Tuberculosis remains one of the most serious infectious diseases globally. In 2009, it had an estimated incidence of 9.4 million cases and 1.7 million people died of tuberculosis globally. Generally, one third of the world's population is

thought to be latently infected (1). Upon infection, *Mycobacterium tuberculosis* invades and successfully replicate inside host macrophages. Although infected host cells can harbor viable mycobacteria, only ~10% of infected people develop tuberculosis (2). Therefore, the interaction between bacterial pathogenesis and the magnitude of the host immune response determines the outcome of the disease (3).

Earlier studies demonstrated the potential therapeutic roles of antimicrobial peptides in a variety of human diseases, including atopic dermatitis, cystic fibrosis and Crohn's disease (4). Recent studies have emphasized the roles of cathelicidin LL-37 in antimycobacterial immune defense, especially in human monocytes/macrophages (5,6). Vitamin D was found to be important in the regulation of LL-37 expression in monocytes, macrophages, and respiratory epithelial cells (6,7). Defensins have been widely studied as an antimicrobial peptide family present in airway fluid and reported to possess antimicrobial activities, including those against mycobacterial infection (8,9). Additionally, hepcidin, an antimicrobial peptide that regulates iron homeostasis, inhibits *M. tuberculosis* growth *in vitro* and inflicts structural damage on this notorious pathogen (10). Moreover, these antimicrobial peptide molecules influence a variety of physiological processes and also function as crucial signaling mediators in host defense and inflammation (11).


Despite these advances in research of the role of antimicrobial peptides in mycobacterial infection, the regulatory mechanisms of these antimicrobial peptides and their exact roles in inflammation during mycobacterial infection remain

Received on September 5, 2011. Revised on September 16, 2011. Accepted on September 23, 2011.

© This is an open access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

\*Corresponding Author. Tel: 82-42-580-8243; Fax: 82-42-585-3686; E-mail: hayoungj@cnu.ac.kr

Keywords: Antimicrobial peptides, Innate Immunity, Mycobacteria



**Figure 1.** A schematic diagram for the role of antimicrobial peptides such as cathelicidins and defensins in host immune system. Left diagram shows biological effects of cathelicidin (hCAP-18/LL-37) in immunity. hCAP-18/LL-37 is synthesized and released from epithelial cells in response to microbial infection or physical injury. hCAP-18/LL-37 participates in the recruitment of neutrophils and other circulating cells including monocytes/macrophages at sites of infection by chemotaxis through secretion of several cytokines/chemokines. Release of hCAP-18/LL-37 from keratinocytes results in induced wound healing. Also, they contribute to direct killing activity against invading pathogens and to indirect antimicrobial activity by promoting autophagy activation/maturation in monocytes/macrophages. Right diagram shows immunological functions of defensins in various immune cells. Defensins induced by various physiological stimuli including TLRs or infection.  $\alpha$ -defensins are synthesized and released from monocytes/macrophages or neutrophils/eosinophils, whereas  $\beta$ -defensins are synthesized and released from not only their cells but also DCs, airway epithelium or skin. Released peptides have direct antimicrobial killing effects and they also have indirect killing effects by interacting with various target cells and tissues to promote secondary responses that may be crucial for modulating inflammation, the recruitment of immune cells, and activation/maturation of several type of immune cell.

to be clarified. Thus, understanding the molecular mechanisms of expression of antimicrobial peptides and their role as immune modulators during the host innate response to mycobacteria will help in the design of new therapies against tuberculosis. Due to the increasing global incidence of multi-drug-resistant tuberculosis, peptide-derived microbicides represent promising candidate therapeutics in the struggle against resistant mycobacteria.

## GENERAL OVERVIEW OF ANTIMICROBIAL PEPTIDES/PROTEINS

Antimicrobial defense peptides/proteins can be produced by

activated macrophages and assist in elimination of ingested bacteria (4,5). Antimicrobial peptides such as cathelicidins and defensins play a crucial role in biological processes, including antimicrobial activities and immunomodulatory functions (summarized in Fig. 1). Cathelicidins are bipartite molecules consisting of an N-terminal cathelin domain and a C-terminal domain, which has antimicrobial activity (12,13). The N-terminal cathelin domain is known as a hallmark of the intracellular storage part of cathelicidins (12,13). Cathelicidins show constitutive and/or inducible expression in various cells and tissues (12,13). Their tissue/cell-specific expression is regulated by several stimuli including infection of microbes, inflammatory cytokines (13-15). Many studies reported that

hCAP-18/LL-37 contributes to elimination of bacteria (14), systemic protection against microbial invasion (13), chemotaxis-attraction through secretion of several cytokines/chemokines (16), wound healing (16) and autophagy activation/maturation (17).

Defensins are antimicrobial/cytotoxic peptides which contain 29~35 amino acid residues, including 6 invariant cysteine residues (18). Defensins are expressed by various physiological/biological stimuli including TLRs or infection (18). The antimicrobial spectrum of defensins include not only gram positive microbes but also gram negative bacteria, including mycobacteria, *Treponema pallidum*, fungi, and some viruses (18-20). Defensins exert nonspecific antimicrobial/cytotoxic activity against mammalian target cells and microorganisms (18). In addition to their antimicrobial/cytotoxic properties, some defensins act as opsonins (18), contribute to selective chemo-attractants for monocytes (18), DCs and T cells (21), and promote to wound healing (8,22,23) and regulation of inflammation (24,25).

### CATHELICIDIN hCAP-18/LL-37

The cathelicidin family, a key member of host defense peptide families, is derived from leukocytes and epithelial cells and has an important role in elimination of pathogenic microbes (13-15,26). Various inflammatory or infectious stimuli can induce cathelicidin LL-37, which then exhibits antimicrobial activity against a number of bacteria and fungi (12, 14,22). hCAP-18/LL-37 is currently the only identified human cathelicidin, and profoundly affects multiple biological and pathological conditions (14, 26). Cathelicidins contain a conserved N-terminal cathelin domain and a variable C-terminal cationic antimicrobial domain that becomes active and has antimicrobial activity. The mature peptide LL-37 comprises the C-terminal portion and is expressed in various cell and tissue types, including neutrophils, monocytes, keratinocytes, lymphocytes, and epithelial cells of the skin, testis, and the gastrointestinal and respiratory tracts (12,15).

Accumulating evidence supports an early defensive role for LL-37 at various sites. During mycobacterial infection, the highest expression of LL-37 was observed in alveolar macrophages (27). Other studies showed the importance of neutrophils in host defense against mycobacteria. When the overall immunity of blood cells to mycobacterial infection was evaluated in tuberculosis contactors, neutrophil counts were associated with a high risk of tuberculosis infection and re-

striction of mycobacterial growth (28). Additionally, the neutrophil peptides, cathelicidin LL-37 and lipocalin2 (Lcn2, also known as neutrophil gelatinase-associated lipocalin [NGAL]) contributed to inhibition of mycobacterial growth and immune defense against tuberculosis (28). Earlier studies showed that the synthetic peptide LL-37 had broad antimicrobial activity in airway epithelial cells of the lung (29) and in bronchoalveolar lavage fluids (30). Recently, we reported that *M. ulcerans* infection significantly induces antimicrobial peptide LL-37 in human primary keratinocytes via TLR2- and Dectin-1-dependent pathways (31). These reports emphasize a role for LL-37 in the early innate response to mycobacteria.

In mycobacterial infection of human mononuclear cells, LL-37 is induced in a vitamin D-dependent manner, and plays an important role in inhibition of intracellular mycobacteria through NADPH oxidase 2-dependent mechanisms (6,32). Similarly, *M. bovis* bacillus Calmette-Guérin (BCG)-induced up-regulation of the antimicrobial peptide cathelicidin LL-37 in human epithelial cells was dependent on NADPH/ROS signaling pathways (33). Of note, vitamin D is crucial for the regulation of LL-37 induction, which can be expressed in human monocytes and respiratory epithelial cells through conversion of vitamin D into its active metabolites (6,7).

Recently, emerging roles of autophagy in the regulation of innate immune functions have been reported (34). Importantly, the autophagy pathway has been known to be a key defensive mechanism to eliminate *M. tuberculosis* through phagosomal maturation (35). Our recent data have shown that vitamin D<sub>3</sub> actively induces autophagy in human monocytes and inhibits intracellular mycobacterial growth through increased autophagosomal maturation (17). In this study, we found that LL-37 plays an important role in the induction and maturation of autophagy pathways activated by vitamin D<sub>3</sub> in human monocytes (17). LL-37 regulated the transcriptional expression of the autophagy-related genes *beclin-1* and *atg-5* via C/EBP- $\beta$  and MAPK activation (17). In addition, both defensin- $\beta$ 4 and cathelicidin are induced by distinct pathways in human monocytes, but cooperate to activate the TLR2/1-mediated antimycobacterial activity (9). Furthermore, recent studies have shown that the mycobacterial lipoprotein LpQH actively induces autophagy through functional vitamin D receptor signaling and following induction of LL-37-dependent pathways (36).

Besides autophagy regulation, additional novel functions of LL-37 have been reported: regulation of chemoattraction, inhibition of apoptosis, wound healing, angiogenesis, and re-

lease of cytokines/chemokines (15,16). LL-37 can also function as an immune regulator, mediating *M. tuberculosis*-induced ROS release and production of pro-inflammatory cytokines and chemokines (32). In mice, the only cathelicidin CRAMP was reported and structurally similar, but shorter, than human LL-37 (12). Both human LL-37 and murine CRAMP has a similar pattern of tissue distribution and biological function (15). For example, both LL-37 and CRAMP have been shown to be chemotactic for various immune cells, including neutrophils, monocytes, macrophages, and T cells (37,38). Further, increasing evidence indicates that various cytokines and signals affect induction of cathelicidin expression (16). The Th1 cytokine IFN- $\gamma$  up-regulates, whereas the Th2 cytokine IL-4 down-regulates, TLR2/1-mediated induction of cathelicidin (39). In this way, cell-mediated immune responses can cross-talk with innate immune pathways via cathelicidin and other AMPs (39). Collectively, these data suggest that cathelicidin LL-37 exerts not only direct antimicrobial activity but is also an important immune modulator of autophagy regulation during mycobacterial infection.

## DEFENSINS

Human defensins constitute a large portion of the pulmonary innate host defense system. Earlier studies showed that defensins are present in high concentrations on respiratory epithelia and selectively target microbial structures (40). Additionally, defensins function as signaling molecules, which link the adaptive immune system to invader microorganisms (40). Similar to cathelicidins, precursors of defensins that contain a characteristic  $\beta$ -sheet-rich fold and a framework of six disulfide-linked cysteines, require proteolytic cleavage for antimicrobial activity (41). The small (3~5 kDa) human cationic defensins are a delineated family of effectors of host defense, inflammation, and cytotoxicity (18). Defensins are divided into three subfamilies:  $\alpha$ -,  $\beta$ -, and  $\theta$ -defensins (25). Six  $\alpha$ -defensins, four human  $\beta$ -defensins (HBD1~4) (reviewed in Ref. 25), and additional  $\beta$ -defensin gene clusters have been identified by computational analysis (42).

High-throughput studies using microarray analyses of gene expression profiles of PBMCs from patients with tuberculosis and *M. tuberculosis*-infected healthy donors found that the effector molecules  $\alpha$ -defensin 1, 3, and 4 are upregulated in patients with tuberculosis (43). Human defensins show synergy with antituberculous drugs, thus suggesting that they may be a promising adjunct to antituberculous chemotherapy

(44). Moreover, a protective role for  $\alpha$ -defensin against mycobacterial infection has been reported in human eosinophils (19).  $\alpha$ -defensin is induced in eosinophils upon stimulation with *M. bovis* BCG and lipomannan and shows a synergistic effect with eosinophil cationic protein on mycobacterial growth inhibition (19).

The human  $\beta$ -defensins HBD-1 and HBD-2 are predominantly expressed at epithelial sites, and less well defined than  $\alpha$ -defensin family (23,25). Both HBD-1 and HBD-2 has bactericidal activity against both Gram-positive and Gram-negative bacteria (23,25). At least six HBD-1 isoforms (range in length from 36 to 47 amino acids) have been identified in urine, whereas a single isoform of HBD-2 (41 amino acids in length) has been isolated from respiratory epithelial secretions and saliva (23). While HBD-1 is constitutively expressed, HBD-2 is upregulated during bacterial infection or in response to endogenous inflammatory cytokines (23,45,46), suggesting a role for HBD-2 in regulation of antimicrobial and inflammatory responses. During mycobacterial infection, HBD-2 can be transported into mycobacteria-containing macrophage phagosomes to exert mycobactericidal and mycobacteristatic activity (20). In airway epithelial cells, HBD-2 mRNA is induced by *M. bovis* BCG infection and is upregulated by TNF- $\alpha$  produced by *M. bovis* BCG-infected cells (46). HBD-2 expression is also triggered by bacterial LPS/TLR4 stimulation through a CD14-dependent mechanism and ultimately results in activation of NF- $\kappa$ B (24).

Recent studies have emphasized the role of HBD4 in the innate immune defense against mycobacteria (9,35). TLR2/1-mediated IL-1 $\beta$  is required for up-regulation of HBD4 expression, which has antimicrobial activity against intracellular mycobacterial infection (9). Moreover, intratracheal administration of l-isoleucine into mice infected with the antibiotic-sensitive strain H37Rv and a multidrug-resistant clinical isolate significantly up-regulated  $\beta$ -defensins 3 and 4 and inhibited bacillary loads (47). *M. bovis* BCG-mediated expression of HBD2 was found to be regulated by the protein kinase C (PKC), JNK and PI3K/Akt pathways in airway epithelial cells (46). Of interest, more highly virulent *M. bovis* strains exhibit lower murine defensin- $\beta$ 4 expression, and *vice versa*, during early infection (48). In experimental tuberculosis, initial expression of murine defensin- $\beta$ 3 and defensin- $\beta$ 4 in airway epithelial cells was correlated with temporary control of mycobacterial growth (49). Additionally, high and stable production of mouse  $\beta$ -defensins, mBD3 and mBD4 during latent infection is associated with long-term control of myco-

**Table I.** The roles of AMPs in immune system

AMPs	Expressed cells organs	Functions	References
<i>hCAP-18/LL-37</i>	Neutrophils, monocytes/macrophages, keratinocytes, lymphocytes, epithelial cells and bronchoalveolar lavage fluids	Elimination of pathogenic microbes (e.g. Mycobacteria, fungi) Induction and maturation of autophagy Inhibition of apoptosis Regulation of chemoattraction Release of cytokines/chemokines Wound healing, angiogenesis Up-regulation of Th1 cytokine (e.g. IFN- $\gamma$ )/down-regulation of Th2 cytokine (e.g. IL-4)	Ref. 14,26 Ref. 17 Ref. 16 Ref. 16 Ref. 16 Ref. 16 Ref. 37
<i>Defensin</i>	Eosinophils, PBMCs, DCs, T-cells, monocytes/macrophages, airway epithelial cells		
<i><math>\alpha</math>-defensin</i>		Up-regulation in TB patients Protective effects against mycobacteria in eosinophils	Ref. 41 Ref. 19
<i><math>\beta</math>-defensin</i>		Mycobactericidal and Mycobacteristatic activity Regulation of Inflammation (e.g. TNF- $\alpha$ ) Activation of NF $\kappa$ B Chemotactic effects in immature DCs and T cells	Ref. 20 Ref. 44 Ref. 24 Ref. 21
<i><math>\theta</math>-defensin</i>		Still unknown yet	
<i>Hepcidin</i>	hepatocytes, macrophages	Homeostatic regulation of iron absorption, iron recycling and iron mobilization Inhibition of invasion of microbes and tumor cells Direct antimicrobial activity	Ref. 55,56 Ref. 59 Ref. 10

bacterial proliferation (49).

Similar to hCAP-18/LL-37, HBD-2 plays roles other than direct antimicrobial action. These include chemotactic roles for immature dendritic cells and memory T cells through a chemokine receptor CCR6-dependent mechanism. This mechanism promotes adaptive immune responses by recruiting immune cells to sites of microbial invasion (21). Unlike  $\alpha$ - and  $\beta$ -defensins,  $\theta$ -defensins are found in some non-human primates, but not in humans, gorillas, bonobos, or chimpanzees (50). Tang *et al.* first isolated a trisulfide-containing antimicrobial peptide, termed rhesus theta defensin 1 (RTD-1), from granules of neutrophils and monocytes of the rhesus macaque (51). Although  $\theta$ -defensins have antimicrobial activity against diverse pathogens (51-53), especially viruses (54-56), there is, as yet, no evidence that  $\theta$ -defensins are involved in defense against mycobacterial infection. The roles of antimicrobial peptides in mycobacterial infection are summarized in Table I. Future studies will reveal the multiple roles of various human defensins in the regulation of immune responses and host defense against mycobacterial infection.

## HEPCIDIN

Hepcidin is a cationic amphipathic bactericidal peptide primarily produced in the liver, and acts as a homeostatic regulator of iron absorption, recycling, and mobilization. Its expression is markedly induced during infectious and inflammatory conditions (57,58). Hepcidin is synthesized by iron loading and cytokine IL-6, and decreased by anemia and hypoxia. The major mechanism of hepcidin function is thought to be related to regulation of transmembrane iron transport through binding with ferroportin, an iron exporter expressed in hepatocytes and macrophages (59,60). The resulting decrease in extracellular iron concentrations probably makes less available for invading microorganisms and tumor cells, thereby contributing to host defense and controlling chronic diseases (57,58,61). As a novel mediator of innate immunity, hepcidin and related therapeutics are promising candidates for the treatment of various diseases, such as hemochromatosis and anemia from chronic inflammation (57,58,61).

Hepcidin is expressed in macrophages after infection with the intracellular pathogens *M. avium* and *M. tuberculosis*.

Stimulation of macrophages with mycobacteria and IFN- $\gamma$  synergistically induced hepcidin mRNA and protein, which localized to the mycobacteria-containing phagosomes (10). Additionally, hepcidin possesses direct antimicrobial activity and causes damage to *M. tuberculosis* (10). Further investigation of the signaling mechanisms responsible for hepcidin mRNA expression showed that STAT1 and NF- $\kappa$ B activation and induction of C/EBP $\beta$  were involved in IFN- $\gamma$  and *M. tuberculosis*-induced hepcidin expression by macrophages (62). These data strongly suggest that *M. tuberculosis*-induced hepcidin expression by activated effector cells may contribute to host defense against mycobacteria. However, future studies are needed to clarify the exact roles and mechanisms of hepcidin expression in innate immune cells during mycobacterial infection.

## CONCLUDING REMARKS

The antimicrobial peptides can contribute to antimycobacterial innate immunity through direct (killing) and indirect (immune modulation) activities. During mycobacterial infection, the cathelicidin, defensin, and hepcidin peptide families have been reported to exhibit antimicrobial activities and immunomodulatory functions. These peptides are produced in different types of innate immune cells, such as macrophages, neutrophils, keratinocytes and epithelial cells. As a bridge between the innate and adaptive immune responses, cathelicidin may contribute to host antibacterial defenses and dampen harmful inflammation. Furthermore, antimycobacterial immune defense is linked to cathelicidin expression and its role in mediating autophagy. IL-1 $\beta$ , a crucial cytokine in antimycobacterial defense, is required for defensin expression, which is critical for innate immunity to mycobacteria. The accumulating data will enable development of therapeutic options and innovative antibiotics derived from host antimicrobial peptides.

## ACKNOWLEDGEMENTS

We thank coworkers and students for many fruitful discussions. This work was supported by the Korea Science & Engineering Foundation through the Infection Signaling Network Research Center (R13-2007-020-01000-0) at Chungnam National University. We apologize to colleagues whose work and publications could not be referenced owing to space constraints.

## CONFLICTS OF INTEREST

The author have no financial conflict of interest.

## REFERENCES

1. WHO: Global Tuberculosis Control 2010. Geneva: WHO; 2010. 7.
2. Pieters J: Mycobacterium tuberculosis and the macrophage: maintaining a balance. Cell Host Microbe 3:399-407, 2008.
3. Flynn JL, Chan J: Tuberculosis: latency and reactivation. Infect Immun 69:4195-4201, 2001.
4. Zaiou M: Multifunctional antimicrobial peptides: therapeutic targets in several human diseases. J Mol Med 85: 317-329, 2007.
5. Liu PT, Modlin RL: Human macrophage host defense against Mycobacterium tuberculosis. Curr Opin Immunol 20:371-376, 2008.
6. Liu PT, Stenger S, Li H, Wenzel L, Tan BH, Krutzik SR, Ochoa MT, Schaubert J, Wu K, Meinken C, Kamen DL, Wagner M, Bals R, Steinmeyer A, Zügel U, Gallo RL, Eisenberg D, Hewison M, Hollis BW, Adams JS, Bloom BR, Modlin RL: Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. Science 311: 1770-1773, 2006.
7. Hansdotir S, Monick MM, Hinde SL, Lovan N, Look DC, Hunninghake GW: Respiratory epithelial cells convert inactive vitamin D to its active form: potential effects on host defense. J Immunol 181:7090-7099, 2008.
8. Cole AM, Waring AJ: The role of defensins in lung biology and therapy. Am J Respir Med 1:249-259, 2002.
9. Liu PT, Schenk M, Walker VP, Dempsey PW, Kanchanapoomi M, Wheelwright M, Vazirnia A, Zhang X, Steinmeyer A, Zügel U, Hollis BW, Cheng G, Modlin RL: Convergence of IL-1 $\beta$  and VDR activation pathways in human TLR2/1-induced antimicrobial responses. PLoS One 4:e5-810, 2009.
10. Sow FB, Florence WC, Satoskar AR, Schlesinger LS, Zwilling BS, Lafuse WP: Expression and localization of hepcidin in macrophages: a role in host defense against tuberculosis. J Leukoc Biol 82:934-945, 2007.
11. Beisswenger C, Bals R: Functions of antimicrobial peptides in host defense and immunity. Curr Protein Pept Sci 6: 255-264, 2005.
12. Lehrer RI, Ganz T: Cathelicidins: a family of endogenous antimicrobial peptides. Curr Opin Hematol 9:18-22, 2002.
13. Zanetti M: The role of cathelicidins in the innate host defenses of mammals. Curr Issues Mol Biol 7:179-196, 2005.
14. Fahy RJ, Wewers MD: Pulmonary defense and the human cathelicidin hCAP-18/LL-37. Immunol Res 31:75-89, 2005.
15. Zanetti M: Cathelicidins, multifunctional peptides of the innate immunity. J Leukoc Biol 75:39-48, 2004.
16. Lai Y, Gallo RL: AMPed up immunity: how antimicrobial peptides have multiple roles in immune defense. Trends Immunol 30:131-141, 2009.
17. Yuk JM, Shin DM, Lee HM, Yang CS, Jin HS, Kim KK, Lee ZW, Lee SH, Kim JM, Jo EK: Vitamin D $\beta$  induces autophagy

- in human monocytes/macrophages via cathelicidin. *Cell Host Microbe* 6;231-243, 2009.
18. Lehrer RI, Lichtenstein AK, Ganz T: Defensins: antimicrobial and cytotoxic peptides of mammalian cells. *Annu Rev Immunol* 11;105-128, 1993.
  19. Driss V, Legrand F, Hermann E, Loiseau S, Guerardel Y, Kremer L, Adam E, Woerly G, Dombrowicz D, Capron M: TLR2-dependent eosinophil interactions with mycobacteria: role of alpha-defensins. *Blood* 113;3235-3244, 2009.
  20. Kisich KO, Heifets L, Higgins M, Diamond G: Antimycobacterial agent based on mRNA encoding human beta-defensin 2 enables primary macrophages to restrict growth of *Mycobacterium tuberculosis*. *Infect Immun* 69;2692-2699, 2001.
  21. Yang D, Chertov O, Bykovskaia SN, Chen Q, Buffo MJ, Shogan J, Anderson M, Schröder JM, Wang JM, Howard OM, Oppenheim JJ: Beta-defensins: linking innate and adaptive immunity through dendritic and T cell CCR6. *Science* 286;525-528, 1999.
  22. De Smet K, Contreras R: Human antimicrobial peptides: defensins, cathelicidins and histatins. *Biotechnol Lett* 27;1337-1347, 2005.
  23. O'Neil DA, Porter EM, Elewaut D, Anderson GM, Eckmann L, Ganz T, Kagnoff MF: Expression and regulation of the human beta-defensins hBD-1 and hBD-2 in intestinal epithelium. *J Immunol* 163;6718-6724, 1999.
  24. Becker MN, Diamond G, Verghese MW, Randell SH: CD14-dependent lipopolysaccharide-induced beta-defensin-2 expression in human tracheobronchial epithelium. *J Biol Chem* 275;29731-29736, 2000.
  25. Oppenheim JJ, Biragyn A, Kwak LW, Yang D: Roles of antimicrobial peptides such as defensins in innate and adaptive immunity. *Ann Rheum Dis* 62(Suppl 2);ii17-21, 2003.
  26. Kai-Larsen Y, Agerberth B: The role of the multifunctional peptide LL-37 in host defense. *Front Biosci* 13;3760-3767, 2008.
  27. Rivas-Santiago B, Hernandez-Pando R, Carranza C, Juarez E, Contreras JL, Aguilar-Leon D, Torres M, Sada E: Expression of cathelicidin LL-37 during *Mycobacterium tuberculosis* infection in human alveolar macrophages, monocytes, neutrophils, and epithelial cells. *Infect Immun* 76;935-941, 2008.
  28. Martineau AR, Newton SM, Wilkinson KA, Kampmann B, Hall BM, Nawroly N, Packe GE, Davidson RN, Griffiths CJ, Wilkinson RJ: Neutrophil-mediated innate immune resistance to mycobacteria. *J Clin Invest* 117;1988-1994, 2007.
  29. Bals R, Wang X, Zasloff M, Wilson JM: The peptide antibiotic LL-37/hCAP-18 is expressed in epithelia of the human lung where it has broad antimicrobial activity at the airway surface. *Proc Natl Acad Sci U S A* 95;9541-9546, 1998.
  30. Agerberth B, Grunewald J, Castañón-Velez E, Olsson B, Jörnvall H, Wigzell H, Eklund A, Gudmundsson GH: Antibacterial components in bronchoalveolar lavage fluid from healthy individuals and sarcoidosis patients. *Am J Respir Crit Care Med* 160;283-290, 1999.
  31. Lee HM, Shin DM, Choi DK, Lee ZW, Kim KH, Yuk JM, Kim CD, Lee JH, Jo EK: Innate immune responses to *Mycobacterium ulcerans* via toll-like receptors and dectin-1 in human keratinocytes. *Cell Microbiol* 11;678-692, 2009.
  32. Yang CS, Shin DM, Kim KH, Lee ZW, Lee CH, Park SG, Bae YS, Jo EK: NADPH oxidase 2 interaction with TLR2 is required for efficient innate immune responses to mycobacteria via cathelicidin expression. *J Immunol* 182;3696-3705, 2009.
  33. Méndez-Samperio P, Pérez A, Torres L: Role of reactive oxygen species (ROS) in *Mycobacterium bovis* bacillus Calmette Guérin-mediated up-regulation of the human cathelicidin LL-37 in A549 cells. *Microb Pathog* 47;252-257, 2009.
  34. Jo EK: Innate immunity to mycobacteria: vitamin D and autophagy. *Cell Microbiol* 12;1026-1035, 2010.
  35. Gutierrez MG, Master SS, Singh SB, Taylor GA, Colombo MI, Deretic V: Autophagy is a defense mechanism inhibiting BCG and *Mycobacterium tuberculosis* survival in infected macrophages. *Cell* 119;753-766, 2004.
  36. Shin DM, Yuk JM, Lee HM, Lee SH, Son JW, Harding CV, Kim JM, Modlin RL, Jo EK: Mycobacterial lipoprotein activates autophagy via TLR2/1/CD14 and a functional vitamin D receptor signalling. *Cell Microbiol* 12;1648-1665, 2010.
  37. De Yang, Chen Q, Schmidt AP, Anderson GM, Wang JM, Wooters J, Oppenheim JJ, Chertov O: LL-37, the neutrophil granule- and epithelial cell-derived cathelicidin, utilizes formyl peptide receptor-like 1 (FPR1) as a receptor to chemoattract human peripheral blood neutrophils, monocytes, and T cells. *J Exp Med* 192;1069-1074, 2000.
  38. Kurosaka K, Chen Q, Yarovinsky F, Oppenheim JJ, Yang D: Mouse cathelin-related antimicrobial peptide chemoattracts leukocytes using formyl peptide receptor-like 1/mouse formyl peptide receptor-like 2 as the receptor and acts as an immune adjuvant. *J Immunol* 174;6257-6265, 2005.
  39. Edfeldt K, Liu PT, Chun R, Fabri M, Schenk M, Wheelwright M, Keegan C, Krutzik SR, Adams JS, Hewison M, Modlin RL: T-cell cytokines differentially control human monocyte antimicrobial responses by regulating vitamin D metabolism. *Proc Natl Acad Sci U S A* 107;22593-22598, 2010.
  40. Ganz T: Antimicrobial polypeptides in host defense of the respiratory tract. *J Clin Invest* 109;693-697, 2002.
  41. Wilson CL, Ouellette AJ, Satchell DP, Ayabe T, López-Boado YS, Stratman JL, Hultgren SJ, Matrisian LM, Parks WC: Regulation of intestinal alpha-defensin activation by the metalloproteinase matrilysin in innate host defense. *Science* 286;113-117, 1999.
  42. Schutte BC, Mitros JP, Bartlett JA, Walters JD, Jia HP, Welsh MJ, Casavant TL, McCray PB Jr: Discovery of five conserved beta-defensin gene clusters using a computational search strategy. *Proc Natl Acad Sci U S A* 99;2129-2133, 2002.
  43. Jacobsen M, Reppilber D, Gutschmidt A, Neher A, Feldmann K, Mollenkopf HJ, Ziegler A, Kaufmann SH: Candidate biomarkers for discrimination between infection and disease caused by *Mycobacterium tuberculosis*. *J Mol Med (Berl)* 85;613-621, 2007.
  44. Kalita A, Verma I, Khuller GK: Role of human neutrophil peptide-1 as a possible adjunct to antituberculosis chemotherapy. *J Infect Dis* 190;1476-1480, 2004.
  45. Harder J, Meyer-Hoffert U, Teran LM, Schwichtenberg L, Bartels J, Maune S, Schröder JM: Mucoid *Pseudomonas aer-*

- uginosa, TNF-alpha, and IL-1beta, but not IL-6, induce human beta-defensin-2 in respiratory epithelia. *Am J Respir Cell Mol Biol* 22;714-721, 2000.
46. Méndez-Samperio P, Miranda E, Trejo A: Mycobacterium bovis Bacillus Calmette-Guérin (BCG) stimulates human beta-defensin-2 gene transcription in human epithelial cells. *Cell Immunol* 239;61-66, 2006.
  47. Rivas-Santiago CE, Rivas-Santiago B, León DA, Castañeda-Delgado J, Hernández Pando R: Induction of  $\beta$ -defensins by l-isoleucine as novel immunotherapy in experimental murine tuberculosis. *Clin Exp Immunol* 164;80-89, 2011.
  48. Aguilar León D, Zumárraga MJ, Jiménez Oropeza R, Gioffré AK, Bernardelli A, Orozco Estévez H, Cataldi AA, Hernández Pando R: Mycobacterium bovis with different genotypes and from different hosts induce dissimilar immunopathological lesions in a mouse model of tuberculosis. *Clin Exp Immunol* 157;139-147, 2009.
  49. Rivas-Santiago B, Sada E, Tsutsumi V, Aguilar-Leon D, Contreras JL, Hernandez-Pando R: beta-Defensin gene expression during the course of experimental tuberculosis infection. *J Infect Dis* 194;697-701, 2006.
  50. Cole AM, Wang W, Waring AJ, Lehrer RI: Retrocyclins: using past as prologue. *Curr Protein Pept Sci* 5;373-381, 2004.
  51. Tang YQ, Yuan J, Osapay G, Osapay K, Tran D, Miller CJ, Ouellette AJ, Selsted ME: A cyclic antimicrobial peptide produced in primate leukocytes by the ligation of two truncated alpha-defensins. *Science* 286;498-502, 1999.
  52. Leonova L, Kokryakov VN, Aleshina G, Hong T, Nguyen T, Zhao C, Waring AJ, Lehrer RI: Circular minidefensins and posttranslational generation of molecular diversity. *J Leukoc Biol* 70;461-464, 2001.
  53. Tran D, Tran PA, Tang YQ, Yuan J, Cole T, Selsted ME: Homodimeric theta-defensins from rhesus macaque leukocytes: isolation, synthesis, antimicrobial activities, and bacterial binding properties of the cyclic peptides. *J Biol Chem* 277;3079-3084, 2002.
  54. Brandt CR, Akkarawongsa R, Altmann S, Jose G, Kolb AW, Waring AJ, Lehrer RI: Evaluation of a theta-defensin in a Murine model of herpes simplex virus type 1 keratitis. *Invest Ophthalmol Vis Sci* 48;5118-5124, 2007.
  55. Wang W, Cole AM, Hong T, Waring AJ, Lehrer RI: Retrocyclin, an antiretroviral theta-defensin, is a lectin. *J Immunol* 170;4708-4716, 2003.
  56. Yang C, Boone L, Nguyen TX, Rudolph D, Limpakarnjanarat K, Mastro TD, Tappero J, Cole AM, Lal RB: Theta-Defensin pseudogenes in HIV-1-exposed, persistently seronegative female sex-workers from Thailand. *Infect Genet Evol* 5;11-15, 2005.
  57. Ganz T: Hepcidin and its role in regulating systemic iron metabolism. *Hematology Am Soc Hematol Educ Program* 29-35, 2006.
  58. Ganz T: Hepcidin—a peptide hormone at the interface of innate immunity and iron metabolism. *Curr Top Microbiol Immunol* 306;183-198, 2006.
  59. Ganz T, Nemeth E: Iron sequestration and anemia of inflammation. *Semin Hematol* 46;387-393, 2009.
  60. Nemeth E, Ganz T: The role of hepcidin in iron metabolism. *Acta Haematol* 122;78-86, 2009.
  61. Atanasiu V, Manolescu B, Stoian I: Hepcidin the link between inflammation and anemia in chronic renal failure. *Rom J Intern Med* 44;25-33, 2006.
  62. Sow FB, Alvarez GR, Gross RP, Satoskar AR, Schlesinger LS, Zwilling BS, Lafuse WP: Role of STAT1, NF-kappaB, and C/EBPbeta in the macrophage transcriptional regulation of hepcidin by mycobacterial infection and IFN-gamma. *J Leukoc Biol* 86;1247-258, 2009.