



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Cardiothoracic Imaging

Pulmonary infarcts in COVID-19

Mary M. Salvatore*

Department of Radiology, Columbia University Medical Center, New York, NY, United States of America



ARTICLE INFO

Keywords:
COVID-1

SARS-CoV-2 binds to the angiotensin-converting enzyme 2 (ACE2) receptors found on vascular endothelial cells among others.¹ Goshua et al. reported that von Willebrand factor is released by activated vascular endothelial cells and modulates platelet adhesiveness and aggregation; it was most elevated in sickest COVID-19 patients.² Thus, the endothelial cell infection leads to a hypercoagulable state providing a rationale for the extent of microthrombi seen in those who succumbed to the disease.

Chest computer tomography (CT) allows the noninvasive evaluation of pulmonary emboli and sequela of endothelial injury. Pulmonary emboli and Hampton's humps (Fig. 1) are known CT findings associated with vascular clotting and parenchymal ischemia and/or infarction.³ Larger pulmonary emboli are relatively rare compared to the microthrombotic disease reported in COVID-19.¹ A recent study found pulmonary emboli in 20% of 51 COVID-19 positive patients and 19% of COVID-19 negative patients.⁴ However, microthrombi were identified in 90% of autopsies performed on patients with COVID-19.⁵

Hampton's humps representing ischemic or infarcted lung are found in less than 36% of patients with pulmonary emboli who do not have COVID-19 infection.⁶ The lung is spared in many patients due to its dual blood supply with a bronchial circulation that connects to the pulmonary arterial circulation and responds to ischemia by increasing its volume 300% following pulmonary embolus.⁷ In contrast, Lax et al. found infarcts at autopsy in 8 of 11(81%) of patients with COVID-19 related pulmonary thrombosis.⁸

The reason for increased infarcts in patients with COVID-19 is likely multifactorial. First, there is a larger clot burden overall increasing the likelihood for infarct or ischemia.¹ The micro-thrombotic disease of COVID-19 effects vessels smaller than 3 mm, which are more often associated with infarction than large central embolic disease.⁷ Embolic disease spares the bronchial circulation allowing it to compensate for diminished pulmonary arterial supply. In COVID-19, the bronchial

endothelial cells are also vulnerable to infection and may be compromised.

Autopsies of COVID-19 patients demonstrate a heavy clot burden with significant infarct and ischemia. PE's are not diagnosed more frequently in COVID-19 likely because the micro thrombi are too small. Increased pulmonary infarcts and ischemia were identified at autopsy and should be visible on CT as Hampton's humps. Zhao et al. demonstrated peripheral opacities in 87 of 101 patients (86%) who were infected with SARS-CoV-2.⁹ VQ scan and Dual Energy CT could




Fig. 1. COVID-19 positive patient with Hampton's hump in the left lower lobe.

* Department of Radiology, Columbia University Irving Medical Center, 630 West 168th Street, New York, NY, United States of America
E-mail address: ms5680@cumc.columbia.edu.



Fig. 2. COVID-19 positive patient with extensive peripheral consolidation.

potentially prove if the peripheral opacities of COVID-19 (Fig. 2) represent ischemia suspected with micro thrombotic disease.^{10–14}

Declaration of competing interest

Mary Salvatore received lecture fees from Genentech and Boehringer Ingelheim and serves as a consultant for Genentech and Boehringer Ingelheim and has received a research grant from Genentech and Boehringer Ingelheim.

References

- Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel A, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet* 2020 May 2;395 (10234):1417–8.
- Goshua G, Pine AB, Meizlish ML, Chang CH, Zhang H, Bahel P, Baluha A, Bar N, Bona RD, Burns AJ, Dela Cruz CS, Dumont A, Halene S, Hwa J, Koff J, Menninger H, Neparidze N, Price C, Siner JM, Tormey C, Rinder HM, Chun HJ, Lee AI. Endotheliopathy in COVID-19-associated coagulopathy: evidence from a single-centre, cross-sectional study. *Lancet Haematol* 2020 Aug;7(8):e575–82.
- Machin A, Gill G, Pappachan JM. Hampton's hump. *BMJ Case Rep* 2013 Nov 6;2013:bcr2013201789.
- Capaccione KM, Li G, Salvatore MM. Pulmonary embolism rate in patients infected with SARS-CoV-2. *Blood Res* 2020 Dec 31;55(4):275–8.
- De Michele S, Sun Y, Yilmaz MM, Katsyv I, Salvatore M, Dzierba AL, Marboe CC, Brodie D, Patel NM, Garcia CK, Saqi A. Forty Postmortem Examinations in COVID-19 Patients. *Am J Clin Pathol* 2020 Nov 4;154(6):748–60.
- Patel UB, Ward TJ, Kadoch MA, et al. Radiographic features of pulmonary embolism: Hampton's hump. *Postgrad Med J* 2014;90:420–1.
- AA Frazier JR Galvin TJ Franks et al From the archives of the AFIP: pulmonary vasculature: hypertension and infarction. *Radiographics*. 20 (2): 491–5.
- Lax SF, Skok K, Zechner P, Kessler HH, Kaufmann N, Koelblinger C, Vander K, Bargfrieder U, Trauner M. Pulmonary arterial thrombosis in COVID-19 with fatal outcome: results from a prospective, single-center, clinicopathologic case series. *Ann Intern Med* 2020 Sep 1;173(5):350–61.
- Zhao W, Zhong Z, Xie X, Yu Q, Liu J. Relation between chest CT findings and clinical conditions of coronavirus disease (COVID-19) pneumonia: a multicenter study. *Am J Roentgenol* 2020;214(5):1072–7.
- Lang M, Som A, Carey D, et al. Pulmonary vascular manifestations of COVID-19 pneumonia. *Radiol Cardiothorac Imaging* 2020;2(3):e200277.
- Khanduri S, Chawla H, Khan A, Ali I, Krishnam A, Malik S, Khan N, Patel YD, Surbhi Rehman M. Spectrum of chest dual-energy computed tomography findings in COVID patients in North India. *Cureus* 2021 Jan 4;13(1):e12489.
- Ganti S, Kok SSX, Aftab S, Halagatti SV, Wong SBS, Uppaluri AS. Spectrum of lung perfusion changes on dual-energy CT in COVID-19: incremental benefit to conventional CT. *Emerg Radiol* 2021 Jan 8:1–5.
- Dhawan RT, Gopalan D, Howard L, Vicente A, Park M, Manalan K, Wallner I, Marsden P, Dave S, Branley H, Russell G, Dharmarajah N, Kon OM. Beyond the clot: perfusion imaging of the pulmonary vasculature after COVID-19. *Lancet Respir Med* 2021 Jan;9(1):107–16.
19. *Clin Nucl Med* 2020 Dec;45(12):e523–4.