

Bis(trimethylammonium) naphthalene-1,5-disulfonate

Yu Jin

Ordered Matter Science Research Center, Southeast University, Nanjing 211189, People's Republic of China
Correspondence e-mail: jinyunihao@yahoo.cn


Received 2 December 2011; accepted 7 December 2011

Key indicators: single-crystal X-ray study; $T = 293\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.041; wR factor = 0.109; data-to-parameter ratio = 19.0.

The asymmetric unit of the title compound, $2\text{C}_3\text{H}_{10}\text{N}^+ \cdot \text{C}_{10}\text{H}_6\text{S}_2\text{O}_6^{2-}$, contains a half-anion, which is completed by inversion symmetry, and one cation. The cations and anions are associated via strong N–H···O(sulfonate) hydrogen-bonding interactions, forming cation–anion–cation groups. Secondary interactions such as C–H(ammonium)···O(sulfonate) and van der Waals interactions link the cations and anions together in a three-dimensional crystal structure, with zigzag rows of cations lying between layers of anions.

Related literature

The title compound was investigated as part of our search for simple ferroelectric compounds. For general background to ferroelectric metal-organic frameworks, see: Ye *et al.* (2006); Zhang *et al.* (2008, 2009, 2010); Fu *et al.* (2009). For a related structure, see: Wang & Yang (2011).



Experimental

Crystal data

$2\text{C}_3\text{H}_{10}\text{N}^+ \cdot \text{C}_{10}\text{H}_6\text{O}_6\text{S}_2^{2-}$
 $M_r = 406.51$

Monoclinic, $P2_1/c$
 $a = 8.3428(17)\text{ \AA}$

$b = 10.502(2)\text{ \AA}$
 $c = 11.742(2)\text{ \AA}$
 $\beta = 105.81(3)^\circ$
 $V = 989.8(3)\text{ \AA}^3$
 $Z = 2$

Mo $K\alpha$ radiation
 $\mu = 0.30\text{ mm}^{-1}$
 $T = 293\text{ K}$
 $0.3 \times 0.3 \times 0.2\text{ mm}$

Data collection

Rigaku Mercury CCD diffractometer
Absorption correction: multi-scan (*CrystalClear*; Rigaku, 2005)
 $R_{\text{int}} = 0.036$
 $T_{\min} = 0.489$, $T_{\max} = 1.000$

10031 measured reflections
2265 independent reflections
2016 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.036$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.041$
 $wR(F^2) = 0.109$
 $S = 1.10$
2265 reflections

119 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.36\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H} \cdots A$	$D-\text{H}$	$\text{H} \cdots A$	$D \cdots A$	$D-\text{H} \cdots A$
N1–H1E···O2 ⁱ	0.91	1.81	2.718 (2)	173
C1–H1C···O1	0.96	2.43	3.372 (3)	166
C2–H2B···O3 ⁱⁱ	0.96	2.31	3.232 (4)	162

Symmetry codes: (i) $x + 1, y, z$; (ii) $x + 1, -y + \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *CrystalClear* (Rigaku, 2005); cell refinement: *CrystalClear*; data reduction: *CrystalClear*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The author thanks the Ordered Matter Science Research Center, Southeast University, for its excellent experimental conditions and its generous financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BH2404).

References

- Fu, D.-W., Ge, J.-Z., Dai, J., Ye, H.-Y. & Qu, Z.-R. (2009). *Inorg. Chem. Commun.* **12**, 994–997.
- Rigaku (2005). *CrystalClear*. Rigaku Corporation, Tokyo, Japan.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Wang, C. & Yang, S. L. (2011). *Acta Cryst. E* **67**, o1847.
- Ye, Q., Song, Y.-M., Wang, G.-X., Chen, K., Fu, D.-W., Hong Chan, P. W., Zhu, J.-S., Huang, S. D. & Xiong, R.-G. (2006). *J. Am. Chem. Soc.* **128**, 6554–6555.
- Zhang, W., Chen, L.-Z., Xiong, R.-G., Nakamura, T. & Huang, S. D. (2009). *J. Am. Chem. Soc.* **131**, 12544–12545.
- Zhang, W., Xiong, R.-G. & Huang, S. D. (2008). *J. Am. Chem. Soc.* **130**, 10468–10469.
- Zhang, W., Ye, H.-Y., Cai, H.-L., Ge, J.-Z., Xiong, R.-G. & Huang, S. D. (2010). *J. Am. Chem. Soc.* **132**, 7300–7302.

supplementary materials

Acta Cryst. (2012). E68, o90 [doi:10.1107/S1600536811052718]

Bis(trimethylammonium) naphthalene-1,5-disulfonate

Y. Jin

Comment

Ferroelectric compounds have displayed a variety of technical applications, such as ferroelectric random access memories, infrared detectors, piezoelectric sensors, nonlinear optical devices, as a result of their excellent ferroelectric, piezoelectric, pyroelectric, and optical properties. Most of the new ferroelectric metal-organic compounds consistent with the necessary requirements for ferroelectric properties have been explored. However, the conditions required in these systems, such as a phase transition, a good electric hysteresis loop and electric domain, and a dielectric anomaly, often disappear (Zhang *et al.*, 2009). Hence, pure organic compounds can be of great potential and can probably make up for the drawbacks found in ferroelectric metal-organic compounds. Reversible structural phase transition remains one of the prominent properties for ferroelectrics. For a small part of these compounds, the components can be arranged in a disordered fashion at a relative high temperature and in an ordered fashion at a relative low temperature. The transition from the disordered arrangement to the ordered one gives rise to sharp change in the physical properties of the compound (Fu *et al.*, 2009; Zhang *et al.*, 2008, 2010; Ye *et al.*, 2006). As part of our search for simple ferroelectric compounds, we have investigated the title compound and report now its room temperature crystal structure.

The centrosymmetric anion and one cation are shown in Fig. 1 with the hydrogen bonds listed in Table 1. The existence of numerous hydrogen-bonding interactions helps to make the substance more stable, and thus forms a three-dimensional layered structure. The interactions tie the cations and anions together in sheets with zigzag rows of cations lying between layers of anions (Fig. 2). There are only van der Waals interactions between layers. The reported structure is similar to that of a related naphthalene-1,5-disulfonate salt (Wang & Yang, 2011).

Experimental

$(C_3H_{10}N^+)_2(C_{10}H_6S_2O_6^{2-})$ was formed from a mixture of $N(CH_3)_3$ (8 mL), $C_{10}H_8O_6S_2$ (288.28 mg, 1.00 mmol), and distilled water (10 ml), which was stirred for few minutes at room temperature, giving a clear transparent solution. After evaporation over few days, block-shaped colorless crystals suitable for X-ray diffraction were obtained in about 82% yield, filtered and washed with distilled water.

Refinement

H atoms bound to carbon and nitrogen were placed at idealized positions [$C-H = 0.93-0.96 \text{ \AA}$, $N-H = 0.91 \text{ \AA}$] and allowed to ride on their parent atoms with U_{iso} fixed at 1.2 $U_{\text{eq}}(C,N)$.

supplementary materials

Figures




Fig. 1. The molecular structure of the title compound with displacement ellipsoids drawn at the 30% probability level.




Fig. 2. The crystal structure of the title compound viewed along the c axis. Intermolecular interactions are shown as dashed lines.

Bis(trimethylammonium) naphthalene-1,5-disulfonate

Crystal data

$2\text{C}_3\text{H}_{10}\text{N}^+\cdot\text{C}_{10}\text{H}_6\text{O}_6\text{S}_2^{2-}$	$F(000) = 432$
$M_r = 406.51$	$D_x = 1.364 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -P 2ybc	Cell parameters from 3450 reflections
$a = 8.3428 (17) \text{ \AA}$	$\theta = 6.2\text{--}55.3^\circ$
$b = 10.502 (2) \text{ \AA}$	$\mu = 0.30 \text{ mm}^{-1}$
$c = 11.742 (2) \text{ \AA}$	$T = 293 \text{ K}$
$\beta = 105.81 (3)^\circ$	Block, colourless
$V = 989.8 (3) \text{ \AA}^3$	$0.3 \times 0.3 \times 0.2 \text{ mm}$
$Z = 2$	

Data collection

Rigaku Mercury CCD diffractometer	2265 independent reflections
Radiation source: fine-focus sealed tube graphite	2016 reflections with $I > 2\sigma(I)$
ω scans	$R_{\text{int}} = 0.036$
Absorption correction: multi-scan (<i>CrystalClear</i> ; Rigaku, 2005)	$\theta_{\text{max}} = 27.5^\circ, \theta_{\text{min}} = 3.2^\circ$
$T_{\text{min}} = 0.489, T_{\text{max}} = 1.000$	$h = -10 \rightarrow 10$
10031 measured reflections	$k = -13 \rightarrow 13$
	$l = -15 \rightarrow 15$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites

$R[F^2 > 2\sigma(F^2)] = 0.041$	H-atom parameters constrained
$wR(F^2) = 0.109$	$w = 1/[\sigma^2(F_o^2) + (0.0505P)^2 + 0.3652P]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.10$	$(\Delta/\sigma)_{\text{max}} < 0.001$
2265 reflections	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
119 parameters	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$
0 restraints	Extinction correction: <i>SHELXL97</i> (Sheldrick, 2008), $F_c^* = kF_c[1 + 0.001x F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
0 constraints	Extinction coefficient: 0.055 (4)
Primary atom site location: structure-invariant direct methods	

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.7732 (3)	0.1761 (3)	0.4624 (2)	0.0783 (8)
H1B	0.7949	0.2389	0.4090	0.117*
H1C	0.6593	0.1826	0.4647	0.117*
H1D	0.7929	0.0927	0.4357	0.117*
C2	0.8549 (4)	0.1033 (3)	0.6651 (3)	0.0926 (10)
H2B	0.9280	0.1194	0.7423	0.139*
H2C	0.8760	0.0197	0.6395	0.139*
H2D	0.7413	0.1087	0.6683	0.139*
C3	0.8748 (4)	0.3266 (3)	0.6265 (3)	0.0824 (9)
H3B	0.9491	0.3339	0.7046	0.124*
H3C	0.7629	0.3436	0.6293	0.124*
H3D	0.9061	0.3868	0.5749	0.124*
C4	0.4055 (2)	0.39604 (16)	0.66437 (14)	0.0336 (4)
H4A	0.3567	0.3452	0.7105	0.040*
C5	0.37869 (18)	0.37009 (14)	0.54673 (13)	0.0279 (3)
C6	0.44909 (18)	0.44781 (14)	0.47321 (13)	0.0258 (3)
C7	0.4218 (2)	0.42584 (15)	0.35041 (14)	0.0315 (4)
H7A	0.3535	0.3589	0.3146	0.038*
C8	0.4940 (2)	0.50113 (16)	0.28418 (14)	0.0353 (4)
H8A	0.4764	0.4842	0.2040	0.042*
N1	0.88425 (18)	0.19790 (16)	0.58184 (15)	0.0444 (4)
H1E	0.9904	0.1864	0.5772	0.053*
O1	0.3624 (2)	0.15499 (14)	0.43979 (16)	0.0635 (5)
O2	0.21016 (15)	0.17580 (12)	0.58538 (12)	0.0406 (3)
O3	0.11286 (19)	0.28226 (17)	0.39993 (14)	0.0646 (5)
S1	0.25686 (5)	0.23499 (4)	0.48711 (4)	0.03500 (17)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0441 (12)	0.127 (3)	0.0569 (15)	-0.0063 (14)	0.0017 (11)	-0.0125 (16)
C2	0.112 (2)	0.086 (2)	0.0729 (18)	-0.0519 (19)	0.0134 (17)	0.0023 (15)

supplementary materials

C3	0.0762 (18)	0.0563 (15)	0.104 (2)	0.0190 (13)	0.0066 (16)	-0.0161 (14)
C4	0.0363 (8)	0.0363 (8)	0.0301 (8)	-0.0014 (7)	0.0123 (7)	0.0050 (6)
C5	0.0247 (7)	0.0274 (7)	0.0310 (8)	0.0002 (6)	0.0068 (6)	0.0011 (6)
C6	0.0256 (7)	0.0249 (7)	0.0265 (7)	0.0029 (6)	0.0064 (6)	0.0005 (6)
C7	0.0347 (8)	0.0298 (8)	0.0285 (8)	-0.0014 (6)	0.0061 (6)	-0.0041 (6)
C8	0.0423 (9)	0.0385 (9)	0.0254 (7)	-0.0014 (7)	0.0100 (7)	-0.0011 (6)
N1	0.0267 (7)	0.0521 (10)	0.0534 (10)	-0.0035 (6)	0.0092 (7)	-0.0075 (8)
O1	0.0751 (11)	0.0409 (8)	0.0925 (12)	-0.0193 (7)	0.0537 (10)	-0.0242 (8)
O2	0.0332 (6)	0.0420 (7)	0.0471 (7)	-0.0069 (5)	0.0118 (5)	0.0085 (5)
O3	0.0455 (8)	0.0811 (11)	0.0533 (9)	-0.0226 (8)	-0.0101 (7)	0.0189 (8)
S1	0.0322 (2)	0.0350 (3)	0.0379 (3)	-0.00957 (16)	0.00984 (17)	-0.00121 (16)

Geometric parameters (\AA , $^\circ$)

C1—N1	1.473 (3)	C4—H4A	0.9300
C1—H1B	0.9600	C5—C6	1.426 (2)
C1—H1C	0.9600	C5—S1	1.7744 (16)
C1—H1D	0.9600	C6—C7	1.416 (2)
C2—N1	1.460 (3)	C6—C6 ⁱ	1.424 (3)
C2—H2B	0.9600	C7—C8	1.359 (2)
C2—H2C	0.9600	C7—H7A	0.9300
C2—H2D	0.9600	C8—C4 ⁱ	1.400 (2)
C3—N1	1.459 (3)	C8—H8A	0.9300
C3—H3B	0.9600	N1—H1E	0.9100
C3—H3C	0.9600	O1—S1	1.4338 (15)
C3—H3D	0.9600	O2—S1	1.4541 (13)
C4—C5	1.365 (2)	O3—S1	1.4379 (16)
C4—C8 ⁱ	1.400 (2)		
N1—C1—H1B	109.5	C6—C5—S1	120.33 (11)
N1—C1—H1C	109.5	C7—C6—C6 ⁱ	119.11 (17)
H1B—C1—H1C	109.5	C7—C6—C5	123.01 (14)
N1—C1—H1D	109.5	C6 ⁱ —C6—C5	117.88 (16)
H1B—C1—H1D	109.5	C8—C7—C6	120.95 (15)
H1C—C1—H1D	109.5	C8—C7—H7A	119.5
N1—C2—H2B	109.5	C6—C7—H7A	119.5
N1—C2—H2C	109.5	C7—C8—C4 ⁱ	120.58 (15)
H2B—C2—H2C	109.5	C7—C8—H8A	119.7
N1—C2—H2D	109.5	C4 ⁱ —C8—H8A	119.7
H2B—C2—H2D	109.5	C3—N1—C2	110.7 (2)
H2C—C2—H2D	109.5	C3—N1—C1	113.8 (2)
N1—C3—H3B	109.5	C2—N1—C1	110.8 (2)
N1—C3—H3C	109.5	C3—N1—H1E	107.0
H3B—C3—H3C	109.5	C2—N1—H1E	107.0
N1—C3—H3D	109.5	C1—N1—H1E	107.0
H3B—C3—H3D	109.5	O1—S1—O3	114.11 (11)
H3C—C3—H3D	109.5	O1—S1—O2	112.49 (9)
C5—C4—C8 ⁱ	120.29 (15)	O3—S1—O2	111.10 (8)

C5—C4—H4A	119.9	O1—S1—C5	105.98 (8)
C8 ⁱ —C4—H4A	119.9	O3—S1—C5	106.41 (9)
C4—C5—C6	121.16 (14)	O2—S1—C5	106.10 (8)
C4—C5—S1	118.50 (12)		

Symmetry codes: (i) $-x+1, -y+1, -z+1$.

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1E···O2 ⁱⁱ	0.91	1.81	2.718 (2)	173.
C1—H1C···O1	0.96	2.43	3.372 (3)	166
C2—H2B···O3 ⁱⁱⁱ	0.96	2.31	3.232 (4)	162
N1—H1E···S1 ⁱⁱ	0.91	2.76	3.5967 (18)	154.

Symmetry codes: (ii) $x+1, y, z$; (iii) $x+1, -y+1/2, z+1/2$.

supplementary materials

Fig. 1




Fig. 2

