

Findings: From January 2016 to November 2018, 60 patients were randomly assigned (30 participants per group). In the whole study population, the average volume of nodules was 25 ml. The two groups were similar in terms of basal nodule volume, thyroid function, histology, symptoms/cosmetic score and procedure time. At six months, participants in the RFA group showed a reduction volume of 64.3% (95% CI: 57.5% - 71.2%) compared to 53.2% (95% CI: 47.2% - 59.2%) in the LA group ($p=0.015$) and this difference was also confirmed in a linear regression model adjusted for age, baseline volume and proportion of cellular component (Laser vs. RFA percent change $\Delta = -12.8$, $P=0.018$). We have not recorded any significant difference in terms of successful rate at six months after treatment between the two groups (86.7% in the RFA vs 66.7% in the LA, $p=0.127$). At six months, both symptoms and cosmetic scores improved (compressive symptom score: 2.13 vs 3.9 for RFA, $p < 0.001$; 2.4 vs. 3.87 for LA, $p < 0.001$; cosmetic score: 1.65 vs 2.2 for RFA $p < 0.001$, 1.85 vs 2.2 for LA $p < 0.001$) without any statistically significant difference between the two groups. No statistical difference between the two groups was detected at six months as regards the TSH level. High rate of cellularity negatively affects the volume reduction in RFA group (r coefficient -0.41, $p=0.034$) while histological features did not affect the efficacy of the LA. The adverse event rates were 37% and 43% for RFA and LA, respectively, with no requirement for hospitalization.

Interpretation: Both techniques are very effective in reducing the volume of thyroid nodules. RFA appears to be more effective than LA, but both techniques showed no difference in terms of success rate six months after treatment. The safety of the two techniques is very satisfactory.

Reproductive Endocrinology

OVARIAN FUNCTION — FROM OLIGOMENORRHEA TO AMENORRHEA

Diagnostic Criteria for Polycystic Ovary Syndrome in Adolescents: Impact on Prevalence and Longitudinal Body Mass Index Trajectories

Chau Thien Tay, MBBS, FRACP¹, Helena Jane Teede, MBBS, PHD, FRACP², Roger Hart, MD, PHD³, Arul Earnest, PHD¹, Dorota Doherty, PHD³, Martha Hickey, PHD⁴, Lisa J. Moran, BSc (Hons), BND, G Cert Pub Health, PHD, APD, MBBS, FRACP, PhD¹, Anju Elizabeth Joham, MBBS, FRACP, PhD¹.

¹Monash Centre for Health Research and Implementation, Clayton, Australia, ²Monash Centre for Health Research and Implementation, Clayton VIC, Australia, ³University of Western Australia, Subiaco, Australia, ⁴University of Melbourne, Melbourne, Australia.

OR31-02

Background: Polycystic ovary syndrome (PCOS) is characterised by oligo-anovulation (OA), hyperandrogenism (HA) and polycystic ovary morphology (PCO). While the Rotterdam criteria (defined as 2 out of 3 features) are the most widely used criteria in adults, controversy remains for the diagnostic criteria in adolescents as many PCOS features overlap with normal pubertal physiological changes. The 2018 international evidence-based PCOS guideline recommends modified Rotterdam criteria (OA

and HA) in adolescents based on expert consensus. We aimed to 1) compare the prevalence of PCOS using original and modified Rotterdam criteria in an unselected adolescent cohort and 2) explore the association between diagnostic phenotypes and long-term body mass index (BMI) trajectories. **Methods:** 227 adolescent females of the Western Australian Pregnancy Cohort (Raine) Study undertook detailed PCOS assessment at the mean age of 15.3 years (mean age of menarche 12.4 years). Detailed anthropometric measurements were collected from birth until age 22 years. T-test was used for group BMI comparisons and longitudinal BMI was analysed using Generalised Estimating Equations with PCOS by time and PCOS phenotypes by time as interaction terms. **Results:** PCOS was diagnosed in 66 (29.1%) participants using original Rotterdam criteria versus 37 (16.3%) participants using modified Rotterdam criteria. Using modified Rotterdam criteria, participants with PCOS had higher mean group BMI than participants without PCOS from age 5 years onwards. Significant interaction was detected between PCOS and time ($p < 0.001$) on longitudinal BMI gain where higher BMI gain was observed in participants with PCOS from age 14 years onwards. Only the modified criteria phenotype was significantly associated with long-term BMI gain whereas other PCOS phenotypes had similar BMI trajectories as participants without PCOS ($p < 0.001$). **Conclusions:** Our findings validate the PCOS guideline recommendation as modified Rotterdam criteria reduce over-diagnosis of PCOS in adolescents and accurately identify the phenotype at risk of long-term weight gain. The BMI trajectories of females with and without PCOS diverge from early childhood suggesting that metabolic dysfunction in PCOS commences early in the pre-pubertal period. **Disclosures:** Nothing to disclose. **Funding:** PCOS CRE scholarship and Research Training Program Scholarship awarded to CT; NHMRC Medical Research Future fund awarded to HT; National Heart Foundation Future Leader Fellowship awarded to LM; NHMRC early career fellowship awarded to AJ.

Neuroendocrinology and Pituitary

CASE REPORTS IN CLASSICAL AND UNUSUAL CAUSES OF HYPOPITUITARISM

Congenital Nephrogenic Diabetes Insipidus with First Presentation as an Adult: A Case Report

Yuan-yuan Liu, MD, Peter Sargious, MD, Gregory A. Kline, MD, Alexander A. Leung, MD MPH.

Department of Medicine, University of Calgary, Calgary, AB, Canada.

SAT-238

Congenital Nephrogenic Diabetes Insipidus with First Presentation as an Adult: A Case Report

Background:

Congenital nephrogenic diabetes insipidus (NDI) is a rare inherited condition, usually presenting during the first year of life. It is characterized by a renal insensitivity to arginine vasopressin. About 90% of patients are males with X-linked NDI who have mutations in the vasopressin V2 receptor (AVPR2) gene encoding the vasopressin V2 receptor. Females are typically asymptomatic. Here, we report