

mRNA and protein expression for novel GABA_A receptors θ and $\rho 2$ are altered in schizophrenia and mood disorders; relevance to FMRP-mGluR5 signaling pathway

SH Fatemi^{1,2,3}, TD Folsom¹, RJ Rooney⁴ and PD Thuras⁵

Fragile X mental retardation protein (FMRP) is an RNA-binding protein that targets ~5% of all mRNAs expressed in the brain. Previous work by our laboratory demonstrated significantly lower protein levels for FMRP in lateral cerebella of subjects with schizophrenia, bipolar disorder and major depression when compared with controls. Absence of FMRP expression in animal models of fragile X syndrome (FXS) has been shown to reduce expression of gamma-aminobutyric acid A (GABA_A) receptor mRNAs. Previous work by our laboratory has found reduced expression of FMRP, as well as multiple GABA_A and GABA_B receptor subunits in subjects with autism. Less is known about levels for GABA_A subunit protein expression in brains of subjects with schizophrenia and mood disorders. In the current study, we have expanded our previous studies to examine the protein and mRNA expression of two novel GABA_A receptors, theta (GABR θ) and rho 2 (GABR $\rho 2$) as well as FMRP, and metabotropic glutamate receptor 5 (mGluR5) in lateral cerebella of subjects with schizophrenia, bipolar disorder, major depression and healthy controls, and in superior frontal cortex (Brodmann Area 9 (BA9)) of subjects with schizophrenia, bipolar disorder and healthy controls. We observed multiple statistically significant mRNA and protein changes in levels of GABR θ , GABR $\rho 2$, mGluR5 and FMRP molecules including concordant reductions in mRNA and proteins for GABR θ and mGluR5 in lateral cerebella of subjects with schizophrenia; for increased mRNA and protein for GABR $\rho 2$ in lateral cerebella of subjects with bipolar disorder; and for reduced mRNA and protein for mGluR5 in BA9 of subjects with bipolar disorder. There were no significant effects of confounds on any of the results.

Translational Psychiatry (2013) 3, e271; doi:10.1038/tp.2013.46; published online 18 June 2013

Introduction

Impairment of the gamma-aminobutyric acid (GABA) signaling system is believed to partially account for behavioral and cognitive deficits associated with schizophrenia and mood disorders.^{1,2} Reduction of GABA_A mediated signal transmission has also been associated with anxiety, panic, impaired learning and memory.^{3–5} GABA_A receptors are responsible for mediating the fast inhibitory action of GABA,⁶ and are important sites for clinical action of a number of drugs including benzodiazepines, barbiturates and anesthetics. Recent work has suggested that proper GABAergic neurotransmission is required for network oscillations that facilitate the processing of information both in and between various brain regions and that this may be required for normal cognition.¹ Altered expression of GABA_A receptor subunits could impair these oscillations and result in improper cognitive function. Little is currently known about GABA_A receptor subunit expression in schizophrenia and mood disorders, although it is likely that changes in GABA_A receptor expression would result in reduced GABAergic transmission.

Recent evidence^{7–9} provides a linkage between GABA neurotransmission and fragile X mental retardation protein (FMRP). FMRP is an RNA-binding protein that has been estimated to regulate translation of 842 transcripts in the brain.¹⁰ In animal models of fragile X syndrome (FXS), the absence of FMRP is accompanied by reduced mRNA expression of GABA_A receptor subunits including alpha 1 ($\alpha 1$), $\alpha 3$, $\alpha 4$, beta 1 ($\beta 1$), $\beta 2$, delta (δ), gamma 1 ($\gamma 1$) and $\gamma 2$ in frontal cortex, whereas there was no change in the cerebellum.^{7–9} A functional consequence of this reduced expression has been observed in fragile X mental retardation 1 (Fmr1)-knocked out mice that display impaired GABAergic signaling in striatal neurons, as measured by increased frequency of spontaneous and miniature inhibitory postsynaptic currents and reduced paired pulse ratio of inhibitory postsynaptic currents.¹¹

FMRP normally represses metabotropic glutamate receptor 5 (mGluR5) signaling, whereas the absence of FMRP has been hypothesized to lead to unregulated mGluR5 signaling, and ultimately result in the various abnormal phenotypes

¹Division of Neuroscience Research, Department of Psychiatry, University of Minnesota Medical School, Minneapolis, MN, USA; ²Department of Pharmacology, University of Minnesota Medical School, Minneapolis, MN, USA; ³Department of Neuroscience, University of Minnesota Medical School, Minneapolis, MN, USA; ⁴Genome Explorations, Memphis, TN, USA and ⁵VA Medical Center, Department of Psychiatry, Minneapolis, MN, USA

Correspondence: Dr SH Fatemi, Division of Neuroscience Research, Department of Psychiatry, University of Minnesota Medical School, 420 Delaware St SE, MMC 392, Minneapolis, MN 55455, USA.

E-mail: fatem002@umn.edu

Keywords: bipolar disorder; FMRP; GABR $\rho 2$; GABR θ ; mGluR5; schizophrenia

Received 27 February 2013; revised 11 April 2013; accepted 22 April 2013

associated with FXS.¹² Animal studies using antagonists of the mGluR5 receptor have rescued learning and behavioral deficits associated with FXS, and reduced seizures in FMR1-knock out mice.^{13–17}

Recently, we reported on reduced protein expression of FMRP in lateral cerebellum from subjects with schizophrenia, bipolar disorder and major depression.¹⁸ These results are novel, as gene association studies have not identified fragile X mental retardation 1 (*FMR1*), the gene that codes FMRP, as a candidate gene for schizophrenia.^{19–20} However, a recent study verified our earlier study, finding reduced FMRP expression in peripheral blood lymphocytes from subjects with schizophrenia.²¹ Kovács *et al.*²¹ found that age of onset and IQ predicted FMRP levels, but chlorpromazine-equivalent antipsychotic dose did not. Importantly, none of the study subjects showed the CGG triplet expansion that normally causes silencing of the *FMR1* gene in subjects with FXS.²¹ Combined with our findings of reduced FMRP expression in the cerebellar vermis and prefrontal cortex of subjects with autism,^{22–23} who were not comorbid for FXS, reduction of FMRP expression may be a hallmark of multiple psychiatric disorders.

Based on the evidence from animal models that decrease in FMRP expression results in reduced expression of GABA_A receptor subunit mRNA and our finding of significantly reduced FMRP in cerebella of subjects with schizophrenia and mood disorders,¹⁸ we hypothesized that we would observe reduced expression of the GABA_A receptor subunits in lateral cerebella from the same diagnostic groups. An initial screen of several GABA receptor subunits in lateral cerebella of subjects with schizophrenia, bipolar disorder and major depression found reductions in GABA_B receptor subunits one and two (GABBR1 and GABBR2).²⁴ Here, we report novel findings regarding alterations in levels of mRNA and protein for GABA_A receptor theta (GABRθ) and GABA_A receptor rho 2 (GABRρ2), as well as mGluR5 and FMRP levels in the lateral cerebellum and Brodmann Area 9 (BA9) of subjects with schizophrenia and mood disorders. These results demonstrate the disruption of the GABAergic and FMRP-mGluR5 signaling systems in subjects with schizophrenia and mood disorders.

Materials and methods

Brain procurement. The Institutional Review Board of the University of Minnesota—School of Medicine has approved this study. Post-mortem lateral cerebella were obtained from the Stanley Foundation Neuropathology Consortium under approved ethical guidelines. Post-mortem superior frontal cortex (BA9) was obtained from the McLean 74 Cohort, Harvard Brain and Tissue Resource Center. DSM-IV diagnoses were established prior to death by psychiatrists using information from all available medical records and family interviews. Details regarding the subject selection, demographics, diagnostic process and tissue processing were collected by the Stanley Medical Research Foundation, and the Harvard Brain and Tissue Resource Center. The Stanley collection consisted of 15 subjects with schizophrenia, 15 with bipolar disorder, 14 with major depression without psychotic features and 14 normal controls (Table 1). The McLean 74 Cohort consists of 20 subjects with schizophrenia, 19 with bipolar disorder and 29 normal controls (Table 2). All groups were matched for age, sex, race, post-mortem interval and hemispheric side.

Sodium dodecyl sulfate polyacrylamide gel electrophoresis and western blotting. Brain tissue was prepared as previously described.^{18,23–28} For lateral cerebellum, 60 μg of tissue was used, whereas for BA9, 30 μg of tissue was used. For mGluR5 and FMRP, we used 6% resolving gels, whereas for GABRθ, GABRρ2, neuronal-specific enolase (NSE) and β-actin we used 10% resolving gels. We minimized interblot variability by including samples from subjects of each group (control, schizophrenia, bipolar disorder and major depression) on each gel. Samples were run in duplicate. Samples were electrophoresed for 15 min at 75 V, followed by 55 min at 150 V. Samples were then electroblotted onto nitrocellulose membranes for 2 h at 300 mA at 4 °C. Blots were blocked with 0.2% I-Block (Tropix, Bedford, MA, USA) in phosphate-buffered saline with 0.3% Tween 20 for 1 h at room temperature (RT), followed by an overnight incubation in primary antibodies at 4 °C. The primary antibodies used were anti-GABRθ

Table 1 Demographic information for the four diagnostic groups from Stanley Medical Research Institute

	Bipolar	Control	Schizophrenia	Depression	F or χ^2	P
Age	42.33 (11.72)	46.64 (9.46)	44.53 (13.11)	46.57 (9.66)	0.49 ^a	0.69
Sex	6F, 9M	5F, 9M	6F, 9M	6F, 8M	0.15 ^b	0.99
Race	14W, 1B	13W, 1B	12W, 3A	14W	14.12 ^b	0.12
PMI	32.53 (16.12)	24.5 (9.85)	33.67 (14.62)	27.57 (11.13)	1.51 ^a	0.22
pH	6.18 (0.23)	6.26 (0.25)	6.16 (0.26)	6.18 (0.23)	0.51 ^a	0.68
Side of brain	7L, 8R	7L, 7R	9L, 6R	9L, 5R	1.20 ^b	0.75
Brain weight	1441.2 (171.5)	1511 (165.4)	1471.7 (108.2)	1443.57 (127.56)	0.71 ^a	0.55
Family hx	0.93 (0.8)	0.13 (0.52)	1.13 (0.83)	0.73 (0.46)	27.19 ^b	0.0001
Suicidal death	9 (5 Violent)	0	6 (2 Violent)	9 (4 Violent)	14.04 ^b	0.029
Drug/Alc hx	0.8 (0.77)	0.36 (0.74)	0.53 (0.74)	0.43 (0.64)	5.08 ^b	0.45
Age of onset	21.47 (8.35)	—	23.2 (7.96)	33.36 (13.68)	3.63 ^a	0.001
Duration of illness	20.13 (9.67)	—	21.67 (11.24)	12.29 (11.37)	2.47 ^a	0.018
Severity of substance abuse	1.93 (1.98)	0.14 (0.54)	1.20 (1.86)	1.15 (2.04)	22.21 ^b	0.10
Severity of alcohol abuse	2.27 (1.98)	1.14 (1.03)	1.47 (1.59)	1.93 (2.02)	9.52 ^b	0.85
Fluphenazine (lifetime)	20 826.67 (24 015.96)	—	52 266.67 (62 061.57)	—	6.21 ^a	0.019

^aANOVA. ^b χ^2 test.

Bold values indicate significant ($P < 0.05$) values.

Table 2 Demographic information for the three diagnostic groups from the McLean 74 Cohort

	Bipolar	Control	Schizophrenia	F, t or χ^2	P
Age	61.75 (19.20)	58.59 (15.12)	60.71 (12.07)	0.24 ^a	0.79
Sex	4M:15F	17M:12F	14M:6F	10.38 ^b	0.006
PMI	22.25 (5.39)	21.76 (3.85)	23.86 (7.20)	0.92 ^a	0.41
pH	6.45 (0.76)	6.45 (0.17)	6.49 (0.30)	0.07 ^a	0.93
Side of brain	10L:9R	14L:15R	10L:10R	0.087 ^b	0.96
Suicidal death	4 (1 violent)	0	0	10.96 ^b	0.027
Drug/Alc hx	0.42 (0.51)	0.69 (0.66)	0.30 (0.47)	6.84 ^b	0.15
Severity of substance abuse	0.58 (1.39)	0.41 (0.95)	0.70 (1.45)	0.33 ^b	0.72
Severity of alcohol abuse	0.95 (1.43)	0.86 (1.73)	0.60 (1.23)	0.39 ^b	0.75
Age of onset	23.53 (7.67)	—	20.52 (3.52)	1.45 ^c	0.16
Duration of illness	39.69 (18.27)	—	40.18 (12.09)	0.091 ^c	0.93
Use of MS	9	—	1	9.17 ^b	0.002

Abbreviation: MS, mood stabilizer.

^aANOVA. ^b χ^2 test. ^ct-test for bipolar versus schizophrenia.
Bold values indicate significant ($P < 0.05$) values.

(ab49188, Abcam (Cambridge, MA, USA) 1:1000), anti-GABRP2 (ab83223, Abcam, 1:500), anti-FMRP (MAB2160, Millipore (Temecula, CA, USA), 1:500), anti-mGluR5 (ab53090, Abcam, 1:300), anti-NSE (1:2000; Abcam) and anti- β actin (A5441, Sigma Aldrich (St Louis, MO, USA), 1:5000). Blots were washed for 30 min in phosphate-buffered saline supplemented with 0.3% Tween 20 (PBST) for 30 min at RT, and were subsequently incubated in the proper secondary antibodies. Secondary antibodies were goat anti-mouse IgG (A9044, Sigma Aldrich, 1:80000) and goat anti-rabbit IgG (A9169, Sigma Aldrich, 1:80000). Blots were washed twice in PBST for 15 min each. Following the second wash, bands were visualized using the ECL-plus detection system (GE Healthcare, Buckinghamshire, UK) and exposed to CL-Xposure film (Thermo Scientific, Rockford, IL, USA). The molecular weights of ~224 (dimer) and 112 kDa (monomer) for mGluR5; 73 kDa (FMRP); 70 kDa (GABR θ); 54 kDa (GABRP2); 46 kDa (NSE) and 42 kDa (β -actin) immunoreactive bands were quantified with background subtraction using a Bio-Rad GS-800 Calibrated Densitometer (Bio-Rad, Hercules, CA, USA) and Quantity One 1-D Analysis software (Bio-Rad). Sample densities were analyzed blind to nature of diagnosis. Results obtained are based on at least two independent experiments.

Quantitative real-time polymerase chain reaction (qRT-PCR). We performed qRT-PCR as previously described.²⁸ Raw data were analyzed as previously described²⁸ using the Sequence Detection Software RQ Manager (ABI, Foster City, CA, USA), whereas relative quantitation using the comparative threshold cycle (C_T method) was performed in Bioconductor using the ABqPCR package in Microsoft Excel (ABI Technote no. 2: Relative Gene Expression Quantitation). Calculations were done assuming that 1 delta C_T equals a two-fold difference in expression. Significance values were determined using unpaired *t*-tests. The probe IDs used were: (1) GABA_A receptor theta (GABRQ): Hs00610921_m1; (2) GABA_A receptor rho 2 (GABRR2): Hs00266703_m1; (3) fragile X mental retardation 1 (FMR1): Hs00924547_m1; (4) metabotropic glutamate 5 (GRM5): Hs00168275_m1; (5) beta actin: Hs99999903_m1 and (6) glyceraldehyde 3-phosphate dehydrogenase (GAPDH): Hs99999905_m1.

Statistical analysis. All protein measurements for each group were normalized against β -actin and NSE, and expressed as ratios of GABR θ / β -actin, GABRP2/ β -actin, FMRP/ β -actin, mGluR5/ β -actin, GABR θ /NSE, GABRP2/NSE, FMRP/NSE and mGluR5/NSE. Statistical analysis was performed as previously described,^{18,23,25} with $P < 0.05$ considered significant. Group comparisons were conducted using analysis of variance (ANOVA). Follow-up independent *t*-tests were then conducted if the results were significant. Group differences on possible confounding factors were explored using χ^2 tests for categorical variables, and ANOVA for continuous variables. Where group differences were found, analysis of covariance was used to explore these effects on group differences for continuous variables, and factorial ANOVA with interaction terms for categorical variables. All analyses were conducted using SPSS v.17 (SPSS, Chicago, IL, USA).

Results

Western blotting results for GABR θ , GABRP2, FMRP and mGluR5 in the lateral cerebellum. All protein measurements were normalized against β -actin or NSE. In lateral cerebella, ANOVA identified group differences for GABR θ / β -actin ($F(3,48) = 5.49, P < 0.003$), GABR θ /NSE ($F(3,48) = 5.61, P < 0.002$), GABRP2/ β -actin ($F(3,40) = 2.90, P < 0.047$), GABRP2/NSE ($F(3,40) = 3.05, P < 0.039$), mGluR5 monomer/ β -actin ($F(3,46) = 4.15, P < 0.011$) and mGluR5 monomer/NSE ($F(3,46) = 5.18, P < 0.004$) (Figure 1; Table 3). There was a group difference for FMRP/NSE ($F(3,50) = 4.93, P < 0.004$) (Figure 1; Table 3).

Follow-up *t*-tests found significant reductions in protein for GABR θ / β -actin ($P < 0.001$), GABR θ /NSE ($P < 0.001$), mGluR5 monomer/ β -actin ($P < 0.050$), mGluR5 monomer/NSE ($P < 0.030$) and FMRP/NSE ($P < 0.001$) in subjects with schizophrenia (Table 3; Figures 2 and 3). In lateral cerebella from subjects with bipolar disorder, there were significant reductions in GABR θ / β -actin ($P < 0.012$), GABR θ /NSE ($P < 0.005$), mGluR5 monomer/ β -actin ($P < 0.001$), mGluR5 monomer/NSE ($P < 0.004$) and FMRP/NSE ($P < 0.003$), and significant increased expression of GABRP2/ β -actin ($P < 0.0044$) and GABRP2/NSE ($P < 0.009$) (Table 3;




Figure 1 Representative bands for mGluR5, FMRP, GABR θ , GABR ρ 2, NSE and β -actin in lateral cerebellum (a) and BA9 (b) of subjects with schizophrenia and mood disorders. B, bipolar disorder; C, control; D, major depression; S, schizophrenia. FMRP and β -actin images for lateral cerebellum reprinted from Fatemi *et al.*,¹⁸ with permission from Elsevier.

Table 3 Western blotting results for FMRP, GABR θ , mGluR5 and GABR ρ 2 values expressed as ratios to β -actin and neuronal-specific enolase (NSE) in lateral cerebellum

	ANOVA		Control		Schizophrenia		Bipolar disorder		Major depression	
	F	P	Protein	P	Protein	P	Protein	P	Protein	P
GABR θ / β -actin	5.49	0.003	0.425 \pm 0.105	RG	0.231 \pm 0.136	0.001	0.297 \pm 0.090	0.012	0.301 \pm 0.138	0.014
GABR ρ 2/ β -actin	2.90	0.047	0.023 \pm 0.013	RG	0.048 \pm 0.042	NC	0.047 \pm 0.014	0.0044	0.058 \pm 0.038	0.0085
mGluR5 dimer/ β -actin	2.14	NC	0.176 \pm 0.111	RG	0.125 \pm 0.10	NC	0.127 \pm 0.101	NC	0.211 \pm 0.064	NC
mGluR5 monomer/ β -actin	4.15	0.011	0.038 \pm 0.030	RG	0.015 \pm 0.023	0.05	0.0086 \pm 0.0098	0.001	0.019 \pm 0.020	NC
FMRP/ β -actin ^a	6.22	0.001	0.070 \pm 0.052	RG	0.017 \pm 0.029	0.039	0.029 \pm 0.033	0.014	0.023 \pm 0.021	0.005
β -actin	0.83	NC	25.8 \pm 2.07	RG	25.2 \pm 2.07	NC	26.9 \pm 2.91	NC	26.1 \pm 4.38	NC
GABR θ /NSE	5.61	0.002	0.87 \pm 0.18	RG	0.48 \pm 0.30	0.001	0.58 \pm 0.20	0.005	0.62 \pm 0.24	0.012
GABR ρ 2/NSE	3.05	0.039	0.045 \pm 0.025	RG	0.1 \pm 0.094	NC	0.096 \pm 0.052	0.009	0.13 \pm 0.085	0.006
mGluR5 dimer/NSE	1.99	NC	0.32 \pm 0.20	RG	0.26 \pm 0.23	NC	0.26 \pm 0.22	NC	0.44 \pm 0.12	NC
mGluR5 monomer/NSE	5.18	0.004	0.08 \pm 0.48	RG	0.032 \pm 0.049	0.03	0.017 \pm 0.021	0.004	0.041 \pm 0.041	0.047
FMRP/NSE	4.93	0.004	0.092 \pm 0.07	RG	0.023 \pm 0.04	0.001	0.042 \pm 0.05	0.003	0.033 \pm 0.033	0.001
NSE	0.93	NC	20.2 \pm 3.34	RG	20.1 \pm 1.67	NC	20.2 \pm 3.24	NC	21.6 \pm 2.87	NC

Abbreviations: ANOVA, analysis of variance; NC, no change; RG, reference group.

^aFMRP data reprinted from Schizophrenia Research, 124(1–3):246–247, Fatemi, S.H., Kneeland, R.E., Liesch, S.B., Folsom, T.D., Fragile X mental retardation protein levels are decreased in major psychiatric disorders, page 246, Copyright (2010), with permission from Elsevier.

Bold values indicate significant ($P < 0.05$) values.

Figures 2 and 3). In subjects with major depression, follow-up *t*-tests found significant reductions in GABR θ / β -actin ($P < 0.014$), GABR θ /NSE ($P < 0.012$), mGluR5 monomer/NSE ($P < 0.047$) and FMRP/NSE ($P < 0.001$), and significantly increased expression of GABR ρ 2/ β -actin ($P < 0.0085$) and GABR ρ 2/NSE ($P < 0.006$) (Table 3; Figures 2 and 3). There were no significant changes in protein levels of mGluR5 dimer in lateral cerebella.

Western blotting results for GABR θ , GABR ρ 2, FMRP and mGluR5 in BA9. In BA9, ANOVA identified group differences for GABR θ / β -actin ($F(2,64) = 4.04$, $P < 0.022$),

GABR θ /NSE ($F(2,62) = 4.54$, $P < 0.014$), mGluR5 monomer/ β -actin ($F(2,63) = 10.72$, $P < 0.001$), mGluR5 monomer/NSE ($F(2,63) = 8.14$, $P < 0.001$), FMRP/ β -actin ($F(2,59) = 3.85$, $P < 0.027$) and FMRP/NSE ($F(2,60) = 4.26$, $P < 0.019$) (Figure 1; Table 4). Follow-up *t*-tests found significant reductions of GABR θ / β -actin, mGluR5 monomer/ β -actin and FMRP/ β -actin ($P < 0.017$, $P < 0.001$ and $P < 0.018$, respectively), and GABR θ /NSE, mGluR5 monomer/NSE and FMRP/NSE ($P < 0.019$, $P < 0.003$ and $P < 0.029$, respectively) in BA9 of subjects with schizophrenia (Table 4, Figures 4 and 5). In subjects with bipolar disorder, follow-up *t*-tests found significant reductions in GABR θ / β -actin,




Figure 2 Expression of GABRA6/β-actin (a), GABRP2/β-actin (b), mGluR5 dimer/β-actin (c), mGluR5 monomer/β-actin (d), FMRP/β-actin (e), and β-actin (f) in lateral cerebella of healthy controls versus subjects with bipolar disorder, major depressive disorder and schizophrenia. Histogram bars shown as mean ± s.e., * $P < 0.05$. FMRP and β-actin data reprinted from Fatemi *et al.*,¹⁸ with permission from Elsevier.

mGluR5 monomer/β-actin and FMRP/β-actin ($P < 0.024$, $P < 0.001$ and $P < 0.030$, respectively), and GABRA6/NSE, mGluR5 monomer/NSE and FMRP/NSE ($P < 0.011$, $P < 0.001$ and $P < 0.011$, respectively) (Table 4; Figures 4 and 5). There were no significant changes in protein levels for GABRP2 or mGluR5 dimer in BA9.

Analysis of confounds for protein data in lateral cerebellum and BA9. In the analysis of protein data from lateral cerebella, no significant differences were found between groups on hemisphere side, ethnicity, gender, history of substance abuse, severity of alcohol abuse or substance abuse, post-mortem interval, age, pH or brain weight (Table 1). We also compared the groups on family history and suicide, and found significant differences ($P < 0.0001$ and $P < 0.029$, respectively), but further analysis revealed that these factors had no significant impact on any of the results. We did find that subjects with schizophrenia and bipolar disorder had significantly longer duration of illness than did those with depression

($t(47) = 2.47$, $P < 0.018$). Age of onset was significantly later for subjects with major depression compared to subjects with schizophrenia or bipolar disorder ($t(41) = 3.63$, $P < 0.001$). ANOVA controlling for age of onset and duration of illness did find that subjects with depression displayed significantly higher mGluR5 dimer/NSE ($F(2,31) = 4.31$, $P < 0.02$) and mGluR5 dimer/β-actin ($F(2,31) = 4.31$, $P < 0.02$) than subjects with bipolar disorder or schizophrenia while controlling for duration. However, as mGluR5 dimer values did not change significantly between the groups, this finding is not meaningful.

For protein data from BA9, no significant differences were found between diagnostic groups on hemisphere side, history of substance abuse, severity of alcohol abuse or substance abuse, post-mortem interval, age or pH (Table 2). Nor did we find significant differences on use of barbiturates, opiates, amphetamines, cocaine or propoxyphene. We also compared subjects with schizophrenia versus subjects with bipolar disorder on disease duration, age of onset, use of antipsychotic, antidepressant and anticonvulsant medications,




Figure 3 Expression of GABRθ/NSE (a), GABRρ2/NSE (b), mGluR5 dimer/NSE (c), mGluR5 monomer/NSE (d), FMRP/NSE (e), and NSE (f) in lateral cerebella of healthy controls versus subjects with bipolar disorder, major depressive disorder and schizophrenia. Histogram bars shown as mean ± s.e., **P* < 0.05.

Table 4 Western blotting results for FMRP, GABRθ, mGluR5 and GABRρ2 values expressed as ratios to β-actin and neuronal-specific enolase (NSE) in BA9

	ANOVA		Control		Schizophrenia		Bipolar disorder	
	<i>F</i>	<i>P</i>	Protein	<i>P</i>	Protein	<i>P</i>	Protein	<i>P</i>
GABRθ/β-actin	4.04	0.022	1.34 ± 0.40	RG	1.06 ± 0.32	0.017	1.07 ± 0.44	0.024
GABRρ2/β-actin	0.38	NC	0.23 ± 0.17	RG	0.21 ± 0.14	NC	0.26 ± 0.23	NC
mGluR5 dimer/β-actin	1.39	NC	0.78 ± 0.38	RG	0.57 ± 0.42	NC	0.70 ± 0.49	NC
mGluR5 monomer/β-actin	10.72	0.001	0.19 ± 0.13	RG	0.089 ± 0.058	0.001	0.071 ± 0.042	0.001
FMRP/β-actin	3.85	0.027	0.81 ± 0.40	RG	0.54 ± 0.40	0.018	0.56 ± 0.29	0.030
β-actin	0.19	NC	7.06 ± 1.68	RG	7.31 ± 1.23	NC	7.29 ± 1.82	NC
GABRθ/NSE	4.54	0.014	1.51 ± 0.40	RG	1.23 ± 0.40	0.019	1.20 ± 0.36	0.011
GABRρ2/NSE	0.54	NC	0.26 ± 0.20	RG	0.23 ± 0.14	NC	0.30 ± 0.27	NC
mGluR5 dimer/NSE	0.66	NC	0.90 ± 0.59	RG	0.72 ± 0.67	NC	0.74 ± 0.50	NC
mGluR5 monomer/NSE	8.14	0.001	0.22 ± 0.17	RG	0.11 ± 0.07	0.003	0.086 ± 0.052	0.001
FMRP/NSE	4.26	0.019	1.05 ± 0.55	RG	0.71 ± 0.50	0.029	0.64 ± 0.36	0.011
NSE	0.01	NC	6.48 ± 1.79	RG	6.54 ± 2.27	NC	6.44 ± 2.61	NC

Abbreviations: ANOVA, analysis of variance; NC, no change; RG, reference group. Bold values indicate significant (*P* < 0.05) values.

and found no significant differences. We did find that 21.1% of bipolar patients died by suicide versus none for the other diagnostic groups ($\chi^2(2) = 10.96, P < 0.027$). We also found

that there were significantly more female subjects ($\chi^2(2) = 10.38, P < 0.006$) in the bipolar group (78.9%) than in either normal controls (41.4%) or subjects with




Figure 4 Expression of GABR0/β-actin (a), GABRp2/β-actin (b), mGluR5 dimer/β-actin (c), mGluR5 monomer/β-actin (d), FMRP/β-actin (e) and β-actin (f) in BA9 of healthy controls versus subjects with bipolar disorder and schizophrenia. Histogram bars shown as mean ± s.e., * $P < 0.05$.

schizophrenia (30%). We also found higher rates of mood stabilizer use in patients with bipolar disorder (47.4%) than in patients with schizophrenia (5%) ($\chi^2(1) = 9.17$, $P < 0.002$). Further analyses controlling for gender, mood stabilizer use and suicide found the initial differences on outcome measures as a function of diagnostic groups to be unchanged.

qRT-PCR results for GABR0, GABRp2, FMRP and mGluR5 in lateral cerebellum and BA9. For qRT-PCR experiments, all values were normalized against both β-actin and GAPDH, and these values were averaged. In the lateral cerebella, ANOVA identified group differences for *GABRQ* (GABR0; $P < 0.046$), *GABRR2* (GABRp2; $P < 0.017$) and *GRM5* (mGluR5; $P < 0.034$) (Table 5). There were significantly reduced mRNA values for *GABRQ* ($P < 0.016$) and *GRM5* ($P < 0.039$) in the lateral cerebella of subjects with schizophrenia (Table 5), similar to protein changes in the same region. *GABRR2* mRNA was significantly increased

($P < 0.019$) in the lateral cerebella in subjects with bipolar disorder, mirroring similar changes in protein levels, and *GRM5* mRNA expression was significantly reduced ($P < 0.009$) in subjects with major depression (Table 5). In BA9, ANOVA identified group differences for *GABRR2* ($P < 0.003$) and *GRM5* ($P < 0.048$) (Table 5). In BA9 of subjects with schizophrenia, there was significantly increased mRNA for *GABRQ* ($P < 0.03$). In BA9 of subjects with bipolar disorder, there was significantly increased mRNA for *GABRR2* ($P < 0.0001$) and significantly reduced mRNA for *GRM5* ($P < 0.04$), similar to changes in mGluR5 protein levels in the same region (Table 5). FMR1 mRNA values did not show significant changes in either of the brain areas (Table 5).

Discussion

The current studies demonstrate abnormal processing of mRNA and protein expression for two novel GABA_A




Figure 5 Expression of GABRθ/NSE (a), GABRρ2/NSE (b), mGluR5 dimer/NSE (c), mGluR5 monomer/NSE (d), FMRP/NSE (e) and NSE (f) in BA9 of healthy controls versus subjects with bipolar disorder and schizophrenia. Histogram bars shown as mean \pm s.e., * $P < 0.05$.

receptors, θ and $\rho 2$, as well as FMRP and mGluR5 in lateral cerebella and BA9 of subjects with schizophrenia and mood disorders. The most salient results included: (1) FMRP protein levels were significantly decreased in all the brain sites in schizophrenia, bipolar disorder and major depression; (2) mGluR5 protein levels were significantly reduced in all the brain sites in schizophrenia and bipolar disorder; (3) mRNA levels for mGluR5 were significantly reduced in lateral cerebellum of subjects with schizophrenia and major depression, and BA9 of subjects with bipolar disorder; (4) Protein levels for GABRθ were reduced significantly in all the brain sites in schizophrenia, bipolar disorder and major depression; (5) mRNA levels for GABRθ were elevated significantly in BA9 of subjects with schizophrenia, in contrast mRNA for the same receptor was decreased significantly in lateral cerebellum of subjects with schizophrenia; (6) Protein levels for GABRρ2 were increased significantly in lateral cerebellum of subjects with bipolar disorder and major depression; simultaneously,

mRNA for the same receptor was also increased significantly in all the brain sites in subjects with bipolar disorder.

The GABRθ gene (*GABRQ*) is clustered with GABA_A receptor epsilon (*GABRE*) and GABA_A receptor alpha 3 (*GABRA3*) at Xq28.²⁹ In rat, GABRθ mRNA has been shown embryonically (E17/E19) to localize to the hypothalamus, tegmentum, pontine nuclei and medulla, suggesting a possible role in midbrain development.³⁰ GABRθ mRNA is expressed in multiple brain regions in human including amygdala, dorsal raphe, hippocampus, hypothalamus, locus coeruleus and substantia nigra.³¹ The locus coeruleus is relevant to psychiatric disorders, as it is the largest noradrenergic nucleus and has important roles in the regulation of anxiety states, vigilance, attention and memory functions.³² GABRθ forms a functional receptor when coexpressed with alpha, beta and gamma subunits.³¹ The functional properties of GABA_A receptors that include the θ subunit have not been well characterized.³⁰

Table 5 qRT-PCR results for GABRQ, GABRR2, GRM5 and FMR1 in lateral cerebella, and BA9 of subjects with schizophrenia and mood disorders

Lateral cerebellum		Schizophrenia		Bipolar disorder		Major depression	
Gene	ANOVA	Fold change	P	Fold change	P	Fold change	P
GABRQ	0.046	0.64	0.016	1.25	0.47	0.730	0.11
GABRR2	0.017	1.33	0.11	1.58	0.019	1.019	0.91
GRM5	0.034	0.49	0.039	0.80	0.39	0.530	0.009
FMR1	0.099	0.65	0.16	1.06	0.85	0.606	0.052

BA9		Schizophrenia		Bipolar disorder		Major depression	
Gene	ANOVA	Fold change	P	Fold change	P		
GABRQ	0.075	1.43	0.03	1.03	0.78	NTA	
GABRR2	0.003	1.21	0.32	2.10	0.0001		
GRM5	0.048	1.05	0.68	0.77	0.04		
FMR1	0.705	0.88	0.31	0.96	0.68		

Abbreviations: ANOVA, analysis of variance; FMR1, fragile X mental retardation 1; NTA, no tissue available; qRT-PCR, quantitative real-time polymerase chain reaction.

Note: for lateral cerebella, ANOVA is based on six comparisons: C versus S, C versus B, C versus D, S versus B, S versus D and B versus D.

For BA9, ANOVA based on three comparisons: C versus S, C versus B and S versus B.

Bold values indicate significant ($P < 0.05$) values.

Recent studies have investigated possible associations of the gene that codes for GABR θ (*GABRQ*) with multiple disorders.^{33–36} However, single-nucleotide polymorphisms of GABR θ were not associated with susceptibility to bipolar disorder,^{33–34} migraine³⁵ or essential tremor.³⁶ However, the GABRQ-478F allele showed an association with the improvement of tremor with ethanol use among men.³⁶

The levels of GABR θ receptor protein are reduced significantly in both BA9 and lateral cerebellum of the subjects with schizophrenia (Figure 6). In contrast, mRNA for GABR θ receptor is downregulated in the lateral cerebella, whereas in BA9 its mRNA is upregulated (Figure 6). As both mRNA and protein are concordantly downregulated in the lateral cerebellum, a severe chronic receptor deficit may be responsible for our observed results; while in BA9, increased mRNA expression may be a compensatory response to chronic receptor downregulation, suggesting that different mechanisms may be at work (Figure 6).

In subjects with bipolar disorder, GABR θ receptor protein is reduced in the lateral cerebellum while its mRNA is upregulated (Figure 7). By the same token, protein for this receptor is downregulated in BA9, with its mRNA level unchanged (Figure 7). Here, the mechanisms for these alterations may again be different in the two brain sites. In lateral cerebellum, chronic GABR θ protein downregulation could lead to upregulation of its mRNA in a feedback loop. In BA9, normal mRNA levels with decreased protein levels indicate a defective step either in processing of protein in rough endoplasmic reticulum or subsequent cell compartments (such as Golgi or secretory granules), leading to reduced protein production (Figure 7).

In subjects with major depression, although protein levels for GABR θ are reduced significantly in lateral cerebellum, its mRNA levels remain normal (Figure 8). This scenario again indicates that the deficit lies at rough endoplasmic reticulum or a subsequent cellular compartment causing the chronic receptor protein downregulation (Figure 8). In the absence of available BA9 tissue, the fate of GABR θ in major depression will await future determinations.




Figure 6 Summary of mRNA and protein expression for GABR θ , GABRp2, mGluR5 and FMRP in lateral cerebella and BA9 of subjects with schizophrenia. Concordant results for mRNA and protein were obtained for GABR θ and mGluR5 in lateral cerebellum. Decreased expression of GABR θ protein in BA9 may lead to a positive feedback loop, increasing mRNA expression. Protein levels for mGluR5 and FMRP were reduced significantly in both brain sets. \uparrow , increased expression; \downarrow , reduced expression, --, no change.

The gene that codes for GABRp2 (*GABRR2*) is localized to 6q15.³⁷ GABRp2 mRNA is widely distributed in the brain, including prefrontal cortex, hippocampus and cerebellum.^{38,39} In adult rat cerebellum, GABRp2 has been localized to

Purkinje cells and basket-like cells only.⁴⁰ GABR ρ 2 has been shown to associate with α 1 and γ 2 receptor subunits.⁴¹ In cerebellum, GABA_A receptors that include the ρ 2 subunit help mediate a component of phasic inhibitory GABAergic transmission at interneuron–Purkinje cell synapses.⁴²




Figure 7 Summary of mRNA and protein expression for GABR θ , GABR ρ 2, mGluR5 and FMRP in lateral cerebella and BA9 of subjects with bipolar disorder. Concordant results for mRNA and protein were obtained for GABR ρ 2 in lateral cerebellum. Decreased expression of GABR θ protein in lateral cerebellum may lead to a positive feedback loop, increasing mRNA expression. Protein levels for GABR θ , mGluR5 and FMRP were decreased significantly in both brain sites. \uparrow , increased expression; \downarrow , reduced expression, --, no change.




Figure 8 Summary of mRNA and protein expression for GABR θ , GABR ρ 2, mGluR5 and FMRP in lateral cerebella of subjects with major depression. There were no concordant results in subjects with major depression. However, protein levels for GABR θ and FMRP were reduced significantly, whereas it increased for GABR ρ 2 in major depression. \uparrow , increased expression; \downarrow , reduced expression, --, no change.

A recent study has demonstrated an association between a single-nucleotide polymorphism of *GABRR2* (GABR ρ 2) (rs1570932) and a component of the bipolar phenotype, namely bipolar patients with psychotic symptoms, similar to those experienced by subjects with schizophrenia.³³ A second study found an single-nucleotide polymorphism (rs12201676) associated with bipolar disorder that is flanked by *GABRR1* (15 kb away) and *GABRR2* (17 kb away) genes.⁴³ *GABRR2* has also been associated with alcoholism.⁴⁴

Levels of GABR ρ 2 mRNA and protein did not change in lateral cerebellum or BA9 of subjects with schizophrenia (Figure 6). However, in subjects with bipolar disorder, a concordant and significant increase was observed in mRNA and protein levels of GABR ρ 2 in lateral cerebellum, indicating chronic upregulation in gene and protein product in this disorder (Figure 7). Interestingly, in BA9 of bipolar subjects, GABR ρ 2 mRNA levels were also elevated significantly, but no protein change was observed (Figure 7). In major depression, GABR ρ 2 protein levels were also elevated but without any change in mRNA, indicating abnormalities in processing GABR ρ 2 protein in rough endoplasmic reticulum compartment or subsequent cellular stations (Figure 8). Thus, GABR ρ 2 changes were confined to brains of subjects with mood disorders and were not seen in schizophrenia and, at least in the case of bipolar disorder, reflect upregulation of GABR ρ 2 mRNA/protein.

The *FMR1* gene is located at Xq27.3. FMRP has been shown to localize in multiple regions of neurons, including the soma, dendrites, synaptic spines and the axon.^{11,45,46} FMRP controls multiple post-transcriptional events, including splicing, nuclear export and translation.^{46,47} FMRP protein is significantly downregulated in schizophrenia, bipolar disorder and major depression in lateral cerebellum,¹⁸ and in BA9 for subjects with schizophrenia and bipolar disorder in the absence of any mRNA abnormalities (Figures 6–8). This




Figure 9 Summary of relationships between GABR θ , GABR ρ 2, mGluR5 and FMRP in three major psychiatric disorders: schizophrenia, bipolar disorder and major depression. Although there are clear biochemical connections between FMRP, mGluR5 and GABR ρ 2, no direct relationship can be established between GABR θ and FMRP. \uparrow , increased expression; \downarrow , reduced expression, --, no change.

picture is similar to what we have described in idiopathic cases of autism without evidence of any effects in the gene for FMRP, and thus replicative of the post-transcriptional abnormalities affecting protein synthesis (Figure 9). Changes in mRNA expression do not always correlate with similar changes in protein expression, including expression of FMRP. A recent study found that in subjects with the FMR1 premutation (expanded 5' CGG repeat, but without full symptoms of FXS), there were both significantly increased FMR1 mRNA and significantly reduced levels of FMRP.⁴⁸ Similarly, the recent findings of Kovács *et al.*²¹ demonstrated the downregulation of FMRP protein levels in the absence of any change in FMR1 mRNA or expansion of the 5' CGG triplet repeat in peripheral blood lymphocytes of subjects with schizophrenia. As reduced FMRP expression has been identified in four major psychiatric disorders, identifying the post-transcriptional abnormalities that may be responsible for this reduction would have a major impact on the etiology and treatment of these disorders. Additionally, verification of reduced FMRP protein levels in peripheral blood lymphocytes of subjects with schizophrenia confirm our data at least in schizophrenia, and validate our additional new findings of reduced FMRP in BA9 of subjects with schizophrenia.

The gene for mGluR5 is located at 11q14.2-q14.3. mGluR5, like other metabotropic glutamate receptors contains seven membrane-spanning domains and a large extracellular N-terminus,⁴⁹ and are G-protein coupled. Metabotropic glutamate receptors are found throughout the CNS, with high concentrations in cerebral cortex, hippocampus, striatum, hypothalamus, midbrain, cerebellum, medulla and pons.⁵⁰ There were concordant and significant reductions in levels of mRNA and protein for mGluR5 in lateral cerebellum of subjects with schizophrenia (Figure 6). In BA9, despite significant reductions in protein levels, mRNA levels were normal (Figure 6). Thus, protein levels for mGluR5 were downregulated in both brain sites in schizophrenia. In a similar vein, mGluR5 protein levels were reduced significantly in both lateral cerebellum and BA9 in subjects with bipolar disorder despite normal mRNA levels, indicating post-transcriptional abnormalities in the pathway for mGluR5 protein synthesis (Figure 7). Interestingly, in lateral cerebellum of subjects with major depression, despite downregulated mRNA for mGluR5, the protein levels were normal (Figure 8). It is possible that unknown mechanisms affecting transit for protein rescue the product for release, despite low turnover for its mRNA, in major depression.

Recently, Matosin *et al.*⁵¹ showed no significant alteration in mGluR5-binding density or mGluR5 protein levels in dorsolateral prefrontal cortex of subjects with schizophrenia. However, close inspection of their western blotting data showed highly oversaturated bands for the monomeric mGluR5 protein levels for both control and subjects with schizophrenia, potentially masking any differences between the two groups. Although several other reports did not show any change in mGluR5 protein^{52,53} or mRNA^{54–56} in prefrontal cortex of subjects with schizophrenia, these results could be due to the use of different analytic techniques or brain regional effects. However, other reports have reported the decreases in mGluR5 mRNA in prefrontal cortex of subjects with schizoaffective disorder⁵⁶ and in those with major

depression,⁵⁷ supporting our current data showing significant decreases in mGluR5 protein levels in lateral cerebellum and BA9 of subjects with schizophrenia and bipolar disorder, and decreases in mRNA levels in lateral cerebella of subjects with schizophrenia and major depression, and BA9 of subjects with bipolar disorder. Additionally, we have previously observed increased expression of mGluR5 protein in BA9 and cerebellar vermis of children with autism (Figure 9).^{22,23} However, although there is a great deal of overlap in the symptomologies of autism and FXS, there is less overlap between FXS and schizophrenia, and mood disorders.

Interactions between the aforementioned four proteins may alter GABAergic transmission. The cytoplasmic domains of GABRP1 and GABRP2 interact with MAP1B (Figure 9).⁵⁸ Disruption of ρ -MAP1B interactions leads to a doubling of the inward current of GABA_C receptors from bipolar cells in retinal slices in the presence of low levels of GABA.⁵⁸ MAP1B mRNA is targeted by FMRP¹⁰ (Figure 9), and in FMR1-knock out mice there is an abnormal upregulation of MAP1B.⁵⁹ With reduced expression of FMRP, one might speculate that there would be increased expression of MAP1B in subjects with schizophrenia and mood disorders. However, a preliminary study involving anterior cingulate cortex found reduced expression of MAP1B protein in subjects with bipolar disorder, with no change in subjects with schizophrenia or major depression.⁶⁰ Further experiments involving multiple brain sites would be required to see if this is a regional difference or if there is a global reduction. Altered expression of MAP1B could in turn cause changes in GABAergic neurotransmission through GABA receptors that contain ρ subunits. Currently, we do not know of a link between FMRP and GABR θ (Figure 9). No changes in GABR θ mRNA were identified among the GABA_A receptor subunits that show reduced expression in animal models of FXS.^{7–9}

In conclusion, FMRP is significantly downregulated in the lateral cerebella¹⁸ and BA9 from subjects with schizophrenia, bipolar disorder and major depression, potentially causing GABA receptor changes and altered expression of mGluR5 in the three disorders in the absence of any FMR1 chromosomal abnormalities. Additionally, we have identified selective abnormalities in mRNA and protein levels of two novel GABA_A receptors, namely GABR θ and GABRP2, in subjects with schizophrenia and mood disorders. These changes could potentially explain changes in GABAergic transmission and consequent deficits associated with these disorders including anxiety, panic, and impaired learning and memory. Our results also identify potential novel targets for future pharmacologic intervention. Lastly, despite significance and novelty of these results, the study should be considered exploratory, requiring further future confirmation in other brain sites.

Conflict of interest

The authors declare no conflicts of interest. Dr Fatemi has patents related to Reelin as a marker of psychiatric disorders. He derives no income as yet from these patents.

Acknowledgements. Grant support by the National Institutes of Mental Health (Grant no. 1R01MH086000-01A2) to SHF is gratefully acknowledged. SHF is also supported by the Bernstein Endowed Chair in Adult Psychiatry. Tissue samples

from the Stanley Medical Research Institute and assistance with demographic information from Dr Edwin Fuller-Torrey and Dr Maree J Webster to SHF is gratefully acknowledged. Tissue samples from the Harvard Brain Tissue Resource Center, which is supported in part by PHS grant number R24 MH068855 is gratefully acknowledged.

Disclaimer. NIMH had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report and in the decision to submit the paper for publication.

- Gonzalez-Burgos G, Fish KN, Lewis DA. GABA neuron alterations, cortical circuit dysfunction and cognitive deficits in schizophrenia. *Neural Plast* 2011; **2011**: 723184.
- Luscher B, Shen Q, Sahr N. The GABAergic deficit hypothesis of major depressive disorder. *Mol Psychiatry* 2011; **16**: 383–406.
- Heulens I, D'Hulst C, Braat S, Rooms L, Kooy RF. Involvement and therapeutic potential of the GABAergic system in the fragile X syndrome. *ScientificWorldJournal* 2010; **10**: 2198–2206.
- Pinna G, Costa E, Guidotti A. Fluoxetine and norfluoxetine stereospecifically and selectively increase brain neurosteroid content at doses that are inactive on 5-HT reuptake. *Psychopharmacology (Berl)* 2006; **186**: 362–372.
- Pinna G, Costa E, Guidotti A. SSRIs act as selective brain steroidogenic stimulants (SBSs) at low doses that are inactive on 5-HT reuptake. *Curr Opin Pharmacol* 2009; **9**: 24–30.
- Brandon NJ, Smart TG, Moss SJ. Regulation of GABAA Receptors by protein phosphorylation. In: Martin DL, Olsen RW (eds) *GABA in the Nervous System: The View at Fifty Years*. Lippincott, Williams and Wilkins: Philadelphia, PA, USA, 2000 pp191–206.
- D'Hulst C, De Geest N, Reeve SP, Van Dam D, De Deyn PP, Hassan BA et al. Decreased expression of the GABA_A receptor in fragile X syndrome. *Brain Res* 2006; **1121**: 238–245.
- El Idrissi A, Ding XH, Scalia J, Trenkner E, Brown WT, Dobkin C. Decreased GABA_A receptor expression in the seizure-prone fragile X mouse. *Neurosci Lett* 2005; **377**: 141–146.
- Gantois I, Vandescampele J, Speleman F, Reyniers E, D'Hooge R, Severijnen LA et al. Expression profiling suggests underexpression of the GABAA receptor subunit delta in the fragile X knockout mouse model. *Neurobiol Dis* 2006; **21**: 346–357.
- Darnell JC, Van Driesche SJ, Zhang C, Hung KY, Mele A, Fraser CE et al. FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism. *Cell* 2011; **146**: 247–261.
- Centonze D, Rossi S, Mercaudo V, Napoli I, Ciotti MT, De Chiara V et al. Abnormal striatal GABA transmission in the mouse model for fragile X syndrome. *Biol Psychiatry* 2008; **63**: 963–973.
- Bear MF, Huber KM, Warren ST. The mGluR theory of fragile X mental retardation. *Trends Neurosci* 2004; **27**: 370–377.
- de Vrij FM, Levenson J, van der Linde HC, Koekoek SK, De Zeeuw CI, Nelson DL et al. Rescue of behavioral phenotype and neuronal protrusion morphology in Fmr1 KO mice. *Neurobiol Dis* 2008; **31**: 127–132.
- Dölen G, Osterweil E, Shankaranarayana Rao BS, Smith GB, Auerbach D, Chattarji S et al. Correction of fragile X syndrome in mice. *Neuron* 2007; **56**: 955–962.
- Westmark CJ, Westmark PR, Malter JS. MPEP reduces seizure severity in Fmr-1 KO mice over expressing human Beta. *Int J Clin Exp Pathol* 2009; **3**: 56–68.
- Yan QJ, Rammal M, Tranfaglia M, Bauchwitz RP. Suppression of two major fragile X syndrome mouse model phenotypes by the mGluR5 antagonist MPEP. *Neuropharmacology* 2005; **49**: 1053–1066.
- Yuskaitis CJ, Mines MA, King MK, Sweatt JD, Miller CA, Jope RS. Lithium ameliorates altered glycogen synthase kinase-3 and behavior in a mouse model of fragile X syndrome. *Biochem Pharmacol* 2010; **79**: 632–646.
- Fatemi SH, Kneeland RE, Liesch SB, Folsom TD. Fragile X mental retardation protein levels are decreased in major psychiatric disorders (letter). *Schizophr Res* 2010a; **124**: 246–247.
- Jönsson E, Björck E, Wahlström J, Gustavsson P, Sedvall G. Screening for CGG trinucleotide repeat expansion in the fragile X mental retardation 1 gene in schizophrenic patients. *Psychiatr Genet* 2005; **5**: 157–160.
- Ashworth A, Abusaad I, Walsh C, Nanko S, Murray RM, Asherson P et al. Linkage analysis of the Fragile X gene FMR-1 and schizophrenia: no evidence for linkage but report of a family with schizophrenia and an unstable triplet repeat. *Psychiatr Genet* 1996; **6**: 81–86.
- Kovács T, Keleman O, Kéri S. Decreased fragile X mental retardation protein (FMRP) is associated with lower IQ and earlier illness onset in patients with schizophrenia. *Psychiatry Res* 2013; **S0165-1781**: 00845–1.
- Fatemi SH, Folsom TD. Dysregulation of fragile X mental retardation protein and metabotropic glutamate receptor 5 in superior frontal cortex of individuals with autism: a postmortem brain study. *Mol Autism* 2011; **2**: 6–16.
- Fatemi SH, Folsom TD, Kneeland RE, Liesch SB. Metabotropic glutamate receptor 5 upregulation in children with autism is associated with underexpression of both fragile X mental retardation protein and GABAA receptor beta 3 in adults with autism. *Anat Rec* 2011a; **294**: 1635–1645.
- Fatemi SH, Folsom TD, Thuras PD. Deficits in GABA(B) receptor system in schizophrenia and mood disorders: a postmortem study. *Schizophr Res* 2011b; **128**: 37–43.
- Fatemi SH, King DP, Reutiman TJ, Folsom TD, Laurence JA, Lee S et al. PDE4B polymorphisms and decreased PDE4B expression are associated with schizophrenia. *Schizophr Res* 2008; **101**: 36–49.
- Fatemi SH, Reutiman TJ, Folsom TD, Thuras PD. GABA(A) receptor downregulation in brains of subjects with autism. *J Autism Dev Disord* 2009a; **39**: 233–230.
- Fatemi SH, Folsom TD, Reutiman TJ, Thuras PD. Expression of GABA(B) receptors is altered in brains of subjects with autism. *Cerebellum* 2009b; **8**: 64–69.
- Fatemi SH, Reutiman TJ, Folsom TD, Rooney RJ, Patel DH, Thuras PD. mRNA and protein levels for GABAAalpha4, alpha5, beta1, and GABABR1 receptors are altered in brains of subjects with autism. *J Autism Dev Disord* 2010b; **40**: 743–750.
- Korpi ER, Gründer G, Lüddens H. Drug interactions at GABA(A) receptors. *Prog Neurobiol* 2002; **67**: 113–159.
- Pape JR, Bertrand SS, Lafon P, Odessa MF, Chaigniau M, Stiles JK et al. Expression of GABA(A) receptor alpha3-, theta-, and epsilon-subunit mRNAs during rat CNS development and immunolocalization of the epsilon subunit in developing postnatal spinal cord. *Neuroscience* 2009; **160**: 85–96.
- Bonnert TP, McKernan RM, Farrar S, le Bourdellès B, Heavens RP, Smith DW et al. Theta, a novel gamma-aminobutyric acid type A receptor subunit. *Proc Natl Acad Sci USA* 1999; **96**: 9891–9896.
- Aston-Jones G, Chiang C, Alexinsky T. Discharge of noradrenergic locus coeruleus neurons in behaving rats and monkeys suggests a role in vigilance. *Prog Brain Res* 1991; **88**: 501–520.
- Breuer R, Hamshere ML, Strohmaier J, Mattheisen M, Degenhardt F, Meier S et al. Independent evidence for the selective influence of GABA(A) receptors on one component of the bipolar disorder phenotype. *Mol Psychiatry* 2011; **16**: 587–589.
- Craddock N, Jones L, Jones IR, Kirov G, Green EK, Grozeva D et al. Strong genetic evidence for a selective influence of GABAA receptors on a component of the bipolar disorder phenotype. *Mol Psychiatry* 2010; **15**: 146–153.
- Fernandez F, Esposito T, Lea RA, Colson NJ, Ciccociocola A, Gianfrancesco F et al. Investigation of gamma-aminobutyric acid (GABA) A receptors genes and migraine susceptibility. *BMC Med Genet* 2008; **9**: 109.
- García-Martín E, Martínez C, Alonso-Navarro H, Benito-León J, Lorenzo-Betancor O, Pastor P et al. Gamma-aminobutyric acid GABRA4, GABRE, and GABRQ receptor polymorphisms and risk for essential tremor. *Pharmacogenet Genomics* 2011; **21**: 436–439.
- Ma DQ, Whitehead PL, Menold MM, Martin ER, Ashley-Koch AE, Mei H et al. Identification of significant association and gene-gene interaction of GABA receptor subunit genes in autism. *Am J Hum Genet* 2005; **77**: 377–388.
- Alakujala A, Palgi M, Wegelius K, Schmidt M, Enz R, Paulin L et al. GABA receptor rho subunit expression in the developing rat brain. *Brain Res Dev Brain Res* 2005; **154**: 15–23.
- Boue-Grabot E, Roudbaraki M, Bascles L, Tramu G, Bloch B, Garret M. Expression of GABA receptor rho subunits in rat brain. *J Neurochem* 1998; **70**: 899–907.
- Rozzo A, Armellini M, Franzot J, Chiaruttini C, Nistri A, Tongiorgi E. Expression and dendritic mRNA localization of GABAC receptor rho1 and rho2 subunits in developing rat brain and spinal cord. *Eur J Neurosci* 2002; **15**: 1747–1758.
- Milligan CJ, Buckley NJ, Garret M, Deuchars J, Deuchars SA. Evidence for inhibition mediated by coassembly of GABAA and GABAC receptor subunits in native central neurons. *J Neurosci* 2004; **24**: 7241–7250.
- Harvey VL, Duguid IC, Krasel C, Stephens GJ. Evidence that GABA rho subunits contribute to functional ionotropic GABA receptors in mouse cerebellar Purkinje cells. *J Physiol* 2006; **577**: 127–139.
- Wang KS, Liu XF, Aragom N. A genome-wide meta-analysis identifies novel loci associated with schizophrenia and bipolar disorder. *Schizophr Res* 2010; **124**: 192–199.
- Xuei X, Flury-Wetherill L, Dick D, Goate A, Tischfield J, Nurnberger J Jr et al. GABRR1 and GABRR2, encoding the GABA-A receptor subunits rho1 and rho2, are associated with alcohol dependence. *Am J Med Genet B Neuropsychiatr Genet* 2010; **153B**: 418–427.
- Antar LN, Afroz R, Dichtenberg JB, Carroll RC, Bassell GJ. Metabotropic glutamate receptor activation regulates fragile X mental retardation protein and FMR1 mRNA localization differentially in dendrites and at synapses. *J Neurosci* 2004; **24**: 2648–2655.
- De Rubeis S, Bagni C. Fragile X mental retardation protein control of neuronal mRNA metabolism: insights into mRNA stability. *Mol Cell Neurosci* 2010; **43**: 43–50.
- Keene JD. RNA regulons: coordination of post-translational events. *Nat Rev Genet* 2007; **8**: 533–543.
- Hessl D, Wang JM, Schneider A, Koldewyn K, Le L, Iwahashi C et al. Decreased fragile X mental retardation protein expression underlies amygdala dysfunction in carriers of the fragile X premutation. *Biol Psychiatry* 2011; **70**: 859–865.
- Minakami R, Katsuki F, Yamamoto T, Nakamura K, Sugiyama H. Molecular cloning and the functional expression of two isoforms of human metabotropic glutamate receptor subtype 5. *Biochem Biophys Res Commun* 1994; **199**: 1136–1143.
- Hinoi E, Ogita K, Takeuchi Y, Ohashi H, Maruyama T, Yoneda Y. Characterization with [3H]quisqualate of group I metabotropic glutamate receptor subtype in rat central and peripheral excitable tissues. *Neurochem Int* 2001; **38**: 277–285.

51. Matosin N, Frank E, Deng C, Huang X-F, Newell KA. Metabotropic glutamate receptor 5 binding and protein expression in schizophrenia and following antipsychotic drug treatment. *Schizophr Res* 2013; **146**: 170–176.
52. Gupta DS, McCullumsmith RE, Beneyto M, Haroutunian V, Davis KL, Meador-Woodruff JH. Metabotropic glutamate receptor protein expression in the prefrontal cortex and striatum in schizophrenia. *Synapse* 2005; **57**: 123–131.
53. Corti C, Xuereb JH, Crepaldi L, Corsi M, Michielin F, Ferraguti F. Altered levels of glutamatergic receptors and Na⁺/K⁺ ATPase- α 1 in the prefrontal cortex of subjects with schizophrenia. *Schizophr Res* 2011; **128**: 7–14.
54. Ohnuma T, Tessler S, Arai H, Faull RL, McKenna PJ, Emson PC. Gene expression of metabotropic glutamate receptor 5 and excitatory amino acid transporter 2 in the schizophrenic hippocampus. *Brain Res Mol Brain Res* 2000; **85**: 24–31.
55. Richardson-Burns SM, Haroutunian V, Davis KL, Watson SJ, Meador-Woodruff JH. Metabotropic glutamate receptor mRNA expression in the schizophrenic thalamus. *Biol Psychiatry* 2000; **47**: 22–28.
56. Volk DW, Eggan SM, Lewis DA. Alterations in metabotropic glutamate receptor 1 α and regulator of G protein signaling 4 in the prefrontal cortex in schizophrenia. *Am J Psychiatry* 2010; **167**: 1489–1498.
57. Deschwanden A, Karolewicz B, Feyissa AM, Treyer V, Ametamey SM, Johayem A *et al*. Reduced metabotropic glutamate receptor 5 density in major depression determined by ((11)C)ABP688 PET and postmortem study. *Am J Psychiatry* 2011; **168**: 724–734.
58. Billups D, Hanley JG, Orme M, Attwell D, Moss SJ. GABAC receptor sensitivity is modulated by interaction with MAP1B. *J Neurosci* 2001; **20**: 8643–8650.
59. Lu R, Wang H, Liang Z, Ku L, O'donnell WT, Li W *et al*. The fragile X protein controls microtubule-associated protein 1B translation and microtubule stability in brain neuron development. *Proc Natl Acad Sci USA* 2004; **101**: 15201–15206.
60. Bouras C, Kövari E, Hof PR, Riederer BM, Giannakopoulos P. Anterior cingulate cortex pathology in schizophrenia and bipolar disorder. *Acta Neuropathol* 2001; **102**: 373–379.



Translational Psychiatry is an open-access journal published by Nature Publishing Group. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivs 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-nd/3.0/>