

RESEARCH ARTICLE

Thermal imaging ruled out as a supplementary assessment in patients with fibromyalgia: A cross-sectional study

Nuria Sempere-Rubio^{1,2}, Marta Aguilar-Rodríguez^{1,2}, Marta Inglés¹, Ruth Izquierdo-Alventosa¹, Pilar Serra-Añó^{1,2*}

1 Department of Physiotherapy, University of Valencia, Valencia, Spain, **2** Faculty of Physiotherapy, Department of Physiotherapy, UBIC Research Group, University of Valencia, Valencia, Spain

 These authors contributed equally to this work.

* pilar.serra@uv.es



Abstract

Background

The diagnosis of fibromyalgia syndrome (FMS) syndrome is often complicated and relies on diagnostic criteria based mostly on the symptoms reported by patients. Implementing objective complementary tests would be desirable to better characterize this population.

Objective

The purpose of this cross-sectional study was to compare the skin temperature at rest using thermography in women with FMS and healthy women.

Methods

Eighty-six women with FMS and 92 healthy controls volunteered to participate. The temperature of all participants was measured by infra-red thermography, registering the skin surface temperature (minimum, maximum and average) at rest in different areas: neck, upper and lower back, chest, knees and elbows. In order to analyze the differences in the skin temperature between groups, inferential analyses of the data were performed using Mann-Whitney U test.

Results

The results showed no significant difference in skin temperature between groups in the neck, upper back, chest and elbows ($p > 0.05$). The lower back and knees areas showed significant differences between groups ($p < 0.05$), although these differences did not reach a minimum of clinically detectable change.

Conclusions

Women with fibromyalgia presented no clinically meaningful reduction or difference in skin temperature at rest when compared with a group of healthy women. The infra-red

OPEN ACCESS

Citation: Sempere-Rubio N, Aguilar-Rodríguez M, Inglés M, Izquierdo-Alventosa R, Serra-Añó P (2021) Thermal imaging ruled out as a supplementary assessment in patients with fibromyalgia: A cross-sectional study. PLoS ONE 16(6): e0253281. <https://doi.org/10.1371/journal.pone.0253281>

Editor: Claudia Sommer, University of Würzburg, GERMANY

Received: January 8, 2021

Accepted: May 29, 2021

Published: June 16, 2021

Copyright: © 2021 Sempere-Rubio et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its [Supporting information files](#).

Funding: This work was supported by Conselleria d'Educació, Ciència i Esports [Valencian Regional Department of Education, Science and Sports], València, Spain [Grant GV2016/140]. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: No authors have competing interests.

thermography is not an effective supplementary assessment tool in women with fibromyalgia.

Introduction

Fibromyalgia (FM) is a common chronic pain syndrome accompanied by other symptoms such as fatigue, headache, sleep disturbances and anxiety [1,2]. This population show increased inflammation expression [3], due to a mitochondrial dysfunction-dependent event implicated in their pathophysiology [4]. Indeed, previous studies have reported high levels of the pro-inflammatory cytokines interleukin (IL)-6, IL-8 in peripheral blood [5,6].

Further, some previous studies have demonstrated that women with FM may have a distorted activity of the autonomous nervous system (ANS), showing alterations in the heart rate variability [7], sleeps disorders [8] and dysfunction of the microcirculation, affecting sweating functions and also body temperature [9,10].

FM is diagnosed according to the classification criteria established by the American College of Rheumatology (ACR) [11]. The diagnosis is not easy and may be frequently unattended [12]. The use of objective assessments to help with the diagnosis is essential to better characterize the syndrome and plan the therapeutic approach of this population [13].

Thermal imaging has been proved as an effective tool to study diseases in which skin temperature can reflect the presence of inflammation in underlying tissues due to a clinical abnormality in chronic pain conditions, such as in rheumatoid arthritis [14], osteoarthritis [15], frozen shoulder [16], and tendinitis [17]. This is because infrared radiation emitted by the human skin, reflects the real-time microcirculatory dynamics of the cutaneous surface of patients [18], being conditioned by the underneath metabolic activity, skin microcirculation and ANS activity [19]. The latter is well correlated with skin temperature [20] because of its role in human thermoregulation [21] and is also related with the exacerbation of the pain in FM population [22].

Pain is detected by primary afferent C-fibers that release substance P into the spinal cord, causing hyperalgesia and cooling. These findings suggest a relationship between the measurement of skin temperature and pain assessment. Therefore, this may be a good option in FM population because thermography is conducted with a portable camera that requires no contact with the individual [23], is non-ionizing [24] and non-invasive [25].

Nevertheless, there is no consensus about the usefulness of thermography. Some authors reported a greater number of hot spots in FM compared with healthy individuals [26,27] whilst others reported that the heat distribution patterns of people with FM are similar to those of the healthy population [28].

We hypothesized that the skin temperature in FM population is lower than in healthy population because of ANS impairment. Therefore, the goal of this study was to compare the skin temperature at rest between women with FM and healthy women.

Materials and methods

Participants and setting

A cross-sectional design was carried out. For the recruitment of the participants, a non-probabilistic sampling was performed. Specifically, a combination of an incidental sampling and a snowball-type sampling. First, women with FMS and their age-matched controls who met their respective inclusion criteria were contacted. Women with FM were recruited from

several associations of FM and specialized medical units while healthy women were recruited through advertisements in local associations and on social networks. Further, once these participants were included as potential participants, recruitment continued through snowball sampling (i.e., word of mouth). They were recruited over a period of one year and six months. Of the total number of volunteers (i.e. 95 women with FM and 95 healthy controls), we discarded those who were not at the postmenopausal stage because of the well-known influence of the menstrual cycle on the temperature [29]. Therefore, a total of 86 women with fibromyalgia syndrome (FM) from different Fibromyalgia associations, and who met the diagnostic criteria of the American College of Rheumatology (ACR, 2010), as assessed by rheumatologists, formed the Fibromyalgia group (FMG) [11]. Ninety-two women, age-matched and without FM symptoms formed the control group (CG). Exclusion criteria for both groups were: suffering any rheumatologic, acute or terminal disease and failing to accomplish the requirements of the thermography measurement protocol recommended by Delphi study conducted by Moreira et al [23]. Further, none of them should present any symptom of inflammation or any sign of hot flashes due to the menopause.

All the assessments were carried out in the biomechanics laboratory of the Department of Physiotherapy at the University of Valencia.

This study complied with the ethical principles of the Declaration of Helsinki, adopted by the World Medical Association on medical research in humans and its revision in 2013. The project was approved by the Ethics Committee on Human Research of our institution. All enrolled participants provided informed written consent prior to the study.

Patients were not involved in the design, conduct, reporting, or dissemination of our research.

Measurements

The participants' skin temperature measurements were taken at the research laboratory, always at between 3 and 6 pm in an 8m² room with black walls. The temperature was kept at 24° C and relative humidity at 44%, tested by a digital hygrometer (ThermoPro TP65), without any direct air source in the exploration area. The participants removed their clothes and remained standing in their underwear to acclimate to the room temperature. Body hair was removed from the exploration area. After 15 minutes, the skin temperature was measured [30] by a physiotherapist widely experienced in the analysis of skin surface temperature with a thermographic camera, using the standard protocol recommended by Gomes Moreira et al. in their Delphi study [23]. The participants were instructed to avoid alcohol and caffeine beverages, smoking, using cosmetics and showering for four hours before the assessment. They should have further avoided performing physical activity, receiving massage, physical therapy, saunas, electro diagnostic tests or taking steroids, sympathetic blockers, vasoactive drugs, opioids and transdermal patches, in the 24 hours prior to measurement. Participants were reminded of these recommendations by phone 48 h before the tests started.

The thermography camera was a FLIR E60BX Thermal Imaging Camera, [FLIR Systems, Inc., Oregon, USA; 76800 Pixels (320 x 240)], featuring a 320 x 240 60Hz infrared detector with a 0.045° C thermal sensitivity and a -20 to 120° C (-4 to 248° F) temperature range. The camera was turned on 30 minutes prior to the test to allow sensor stabilization following the manufacturer's guidelines.

The camera was perpendicularly pointed to the explored region using a tripod and at a distance of 1 m. The images were taken with an emissivity of 0.98 ϵ [23,30]. The selected areas for our study were modified with regard to the Regions of Interest (ROI) described by Ammer et al. [31], in order to select the largest skin area possible. A total of six regions that are




Fig 1. Regions of interest used in the study. A: Neck; B: Upper back; C: Lower back; D: Chest; E: Medial knee; F: Lateral elbow. Regions in which significant differences between groups were obtained are represented by bold lines.

<https://doi.org/10.1371/journal.pone.0253281.g001>

described below were assessed: neck, upper back, lower back, chest, knees and elbows (Fig 1). We discarded the assessment of more distal regions to avoid that the results could be influenced by other autonomic syndromes that have been described in people suffering from fibromyalgia, like Raynaud Syndrome [32] or peripheral sensory-motor polyneuropathy [33].

Neck. A rectangle was drawn on the back of the neck. The upper edge was adjacent to the hairline, while the lower edge corresponded to a horizontal line at the height of the spinous process of the first thoracic vertebrae. The lateral edges were the vertical outline of the neck.

Upper back. A polygon with six angles was drawn. The upper edge was a horizontal line at the height of the spinous process of the first thoracic vertebrae. From either end, two lines ran bilaterally to the upper lateral angle at the acromioclavicular joint. From this point, two lines ran bilaterally to the axillar fold. The lower edge was formed by the horizontal line joining these two latter points.

Lower back. A trapezium was drawn in which the upper edge was a horizontal line on the iliac crest. The lower edge was placed above the intergluteal cleft, adjacent to the natal cleft. The lateral edges were the outline of the skin.

Chest. A trapezium was drawn having an upper edge formed by a horizontal line connecting the two acromia. From either end two lines ran bilaterally to the axillar fold. The lower edge was formed by the horizontal line joining these two points. **Medial knee.** A circumference

was drawn for each knee. The edge of the circumference was adjacent to the patella and the popliteal fold for its internal face. The average of the temperature values of the two knees were obtained for subsequent analyses.

Lateral elbow. A circumference was drawn for each elbow. The edge of the circumference was adjacent to the cubital fold and the lower edges of the elbow. The average of the temperature values of the two elbows were obtained for subsequent analyses.

The processing of the images was carried out using the *FLIR Research IR Professional Analyzing Software* (Version 1.2, Wilsonville, OR). A palette of 85–100 colors, with a thermal window of 0.15 uC for each color, was used. Thermal sensitivity of 0.51 uC was utilized per each shade, based on a rainbow-colored colorimetric scale, in which the colors ranged from the hottest (red) to the coldest (blue) [34]. The data obtained from each area were: the maximum, minimum and average temperature.

Sample size calculation

Sample size was calculated taking into consideration a medium effect size of the mean difference of temperature between groups, a type I error of 5% and a type II error of 10%. This calculation resulted in 70 patients in each group. 86 women were included in the FMG and 92 in the CG. Ultimately, the temperature of the elbows and knees of 1 woman of the FMG could not be analyzed, including 85 women for the analysis of these regions. G-Power[®] version 3.1. was used for sample size estimation (Institute for Experimental Psychology, University of Düsseldorf, Düsseldorf, Germany).

Statistics

Statistical analysis was performed using SPSS software Version 22 (SPSS Inc., Chicago, IL, USA). Standard statistical methods were used to obtain the mean and standard deviation (SD). The normality assumption was checked using Shapiro-Wilk. Inferential analyses of the data were performed using Mann-Whitney U test with one inter-subject factor having two categories (FMG and CG).

As results of the Mann-Whitney U test, we provide the U statistic calculated as:

$$U = n_1 n_2 + \frac{n_1(n_1 + 1)}{2} - R_1$$

Where n_1 is the sample size for the FMG and n_2 is the sample size of CG and R_1 is the sum of ranks for group with FM.

Further, we provide the Z-score to convert the statistic U in a standardize value. The effect size, when significant differences were obtained, was calculated using $r (z/\sqrt{N})$.

Furthermore, to explore the relationship between suffering the syndrome and the different categories of the demographic data (i.e. civil and employment status, studies level, smoking habits and alcohol intake), we conducted Chi Square tests (χ^2). The type I error for the statistical tests was set at 5% ($p \leq 0.05$).

Results

Participants

The mean (SD) age for CG was 54.89 (6.31) years whilst for FMG it was 54.91 (7.00) years. There was no significant difference in age between groups ($p > 0.05$). The mean (SD) body mass index (BMI) for CG was 27.83 (4.75) whilst for FMG it was 25.97 (4.00). There was a

Table 1. Demographic data of the participants.

	FM	C	p-value of the χ^2
Civil Status			
Married/in union	68	66	$\chi^2(3) = 3.10, p = 0.38$
Single	6	11	
Widowed	1	4	
Separate/Divorced	11	11	
Employment status			
Active	44	60	$\chi^2(3) = 4.31, p = 0.23$
Unemployed	18	14	
Pensioner	21	14	
Housekeeper	3	4	
Studies level			
Illiterate	1	2	$\chi^2(3) = 15.01, p < 0.05^*$
Primary	37	28	
Secondary	36	26	
University	12	36	
Smoking habits			
Daily	19	14	$\chi^2(3) = 3.04, p = 0.39$
Not daily	3	7	
Not at the present	31	30	
Never	33	41	
Alcohol intake			
Daily	6	5	$\chi^2(4) = 3.26, p = 0.52$
Some days per week	25	34	
Some days per month	15	21	
Some days in the last year	16	14	
Not in the last year	24	18	

FM: Fibromyalgia group; C: Control Group;

*: Statistical relationship among the categories ($p < 0.05$).

<https://doi.org/10.1371/journal.pone.0253281.t001>

significant difference in BMI between groups ($p < 0.05$) being both values included in the 'overweight' category [35]. Table 1 shows demographic data of the participants.

Thermography outcomes

Table 2 shows the results of the thermographic assessment. Our results showed that there were no significant differences in the skin temperature between groups in the neck, upper back and chest ($p > 0.05$). There were significant differences between groups in the lower back for minimum temperature ($U = 3210.00, Z = -2.17, p < 0.05$), maximum temperature ($U = 3270.00, Z = -2.00, p < 0.05$) and average temperature ($U = 3161.00, Z = -2.32, p < 0.05$). Further, there were significant differences between groups in the knees for minimum temperature ($U = 3038.50, Z = -2.56, p < 0.05$), maximum temperature ($U = 3124.00, Z = -2.31, p < 0.05$) and average temperature ($U = 2738.50, Z = -3.44, p < 0.05$). There were not significant differences between groups in the elbows assessment for minimum temperature, maximum temperature and average temperature ($p > 0.05$). The significant differences obtained did not exceed 0.5°C which is considered the minimum clinical finding (44) Fig 1 show the regions in which significant differences were obtained in bold lines.

Table 2. Skin temperature of the study participants.

	FM		C		Size effect (r)
	Mean (SD)	Median (range)	Mean (SD)	Median (range)	
Neck_min	31.58 (1.28)	31.60 (27.45–34.05)	31.89 (1.06)	31.93 (26.75–34.20)	
Neck_max	34.53 (0.93)	34.73 (30.30–36.10)	34.43 (1.00)	34.58 (29.95–36.25)	
Neck_ave	33.56 (1.03)	33.60 (29.35–35.50)	33.57 (0.98)	33.68 (28.80–35.45)	
Upper back_min	30.53 (1.26)	30.58 (26.65–33.50)	30.39 (1.32)	30.28 (26.25–33.15)	
Upper back_max	33.75 (0.99)	33.85 (29.90–35.60)	33.63 (1.04)	33.75 (29.50–35.65)	
Upper back_ave	32.15 (1.09)	32.15 (28.20–34.45)	32.10 (1.18)	32.10 (27.95–34.35)	
Lower back_min	29.75 (1.44)	29.75 (26.60–32.45)	29.25 (1.42)*	29.20 (25.90–31.85)	0.16
Lower back_max	33.99 (1.08)	34.15 (30.55–35.85)	33.65 (1.13)*	33.78 (30.55–35.85)	0.15
Lower back_ave	31.68 (1.25)	31.55 (28.30–34.20)	31.21 (1.28)*	31.25 (28–33.85)	0.17
Chest_min	30.82 (1.20)	30.88 (27.30–33.90)	30.64 (1.22)	30.65 (27.75–33.35)	
Chest_max	34.28 (0.77)	34.25 (32.55–36.05)	34.09 (0.91)	34.15 (32.30–35.95)	
Chest_ave	32.34 (0.98)	32.28 (30.05–34.90)	32.19 (1.13)	32.15 (29.50–34.25)	
Knee_min	29.70 (1.12)	29.70 (27.35–32.25)	29.23 (1.03)*	29.28 (27.35–31.35)	0.19
Knee_max	31.86 (1.02)	31.83 (29.75–34.50)	31.49 (0.95)*	31.40 (29.70–34.65)	0.17
Knee_ave	30.80 (1.03)	30.75 (28.70–33.30)	30.21 (0.97)*	30.15 (28.35–32.25)	0.26
Elbow_min	30.28 (1.03)	30.45 (27.25–32.75)	30.05 (0.93)	30.15 (26.05–31.75)	
Elbow_max	32.79 (0.85)	32.80 (30.60–35.30)	32.61 (0.74)	32.70 (30.70–34.05)	
Elbow_ave	31.52 (0.89)	31.50 (29.15–33.95)	31.34 (0.80)	31.40 (29.70–33.15)	

FM: Fibromyalgia group; C: Control Group; min: Minimum temperature of the region; max: Maximum temperature of the region; ave: Average temperature of the region;

*: Statistical differences between groups ($p < 0.05$).

<https://doi.org/10.1371/journal.pone.0253281.t002>

Discussion

The present study was designed to identify possible differences in the skin temperature, at rest, between women with FM and healthy women in order to consider thermal skin imaging as a supplementary assessment of FM. Thermal assessment at rest is a simple and noninvasive tool that can provide a reliable non-fatiguing assessment in this population. Since previous studies evidenced inflammation of the small muscle fibers in people with FM [36–39] some authors predicted that infrared thermography could be useful to detect peripheral temperature changes derived from peripheral inflammation, as has been shown for other pathologies [40–47]. Further, we explored the thermal patterns of the body at rest because of the need to dispose of non-fatiguing and easy-to-use assessments. We took into consideration that people with FM refer fatigue when exertion is required [48], so we decided to use an assessment in a non-fatiguing environment. The results derived from our study revealed no statistical or clinical differences in the body surface temperature between groups in most of the analyzed areas. The temperature of the lower back and knee in FM women was higher than that obtained in healthy women, but this difference, although statistically significant, did not exceed 0.5° , either in the average or maximum temperature, as required to be considered a clinical finding [49]. Therefore, this study demonstrates that, in a resting condition, infrared thermography detects no changes in body heat in women with FM due to possible alterations in the surface blood flow or alterations in the microcirculation of ANS and therefore this tool should not be used to assess ANS disorders in this syndrome [9].

Our results are not consistent with those derived from two previous studies, conducted by the same research group, in which thermography was used as an assessment tool to

differentiate the thermal pattern in people with FM and their healthy controls [26,27]. Both studies analyzed whether there were hot spots in certain body areas in people with FM or myofascial pain, these hot spots being identified as differences in temperature of more than 0.5° with respect to the surrounding areas. Both studies reported that the FM group had a greater number of hot spots than the healthy group.

However, despite these positive results, thermography was considered as a specific but barely sensitive tool, although the exact sensitivity and specificity was not reported. However, such results are not exactly comparable with ours because, as noted, these studies used the fibromyalgia diagnosis criterion which involves 18 tender points [50], subsequently invalidated [1], unlike the approach used in our study, which explores the skin temperature pattern at rest. These studies measured the differences in the numbers of hot spots between people with and without FM. A hot spot is defined as a small area at least 5°C warmer than the surrounding area and therefore does not represent an increase in the temperature of larger areas included in the diagnostic criteria which are supposedly painful. We hypothesized that the skin temperature of these areas in FM population would be different than those of their healthy counterparts because of ANS impairment. Broader areas are expected to be affected due to the complex relationship of the different autonomic structures of the neuraxis [51]. Previous studies assessing people with arthritic diseases have indicated significant differences in heat distribution indices compared to healthy people as reported by the review of Jasti *et al.* [52]. Thus, our results are not completely comparable with those obtained from other previous studies.

In order to resolve the discrepancies between the results of previous studies on the usefulness of the tool, we designed a study ensuring a sufficient recruitment of participants to achieve a statistical power of 90% that would give greater reliability in the results than the previous studies, and therefore, the probability of committing a type II error or false negatives was low. Accordingly, the power of the study allows us to discard the hypothesis, since the thermal pattern between groups is similar.

Our results, therefore, suggest that there may be an equivalence of the local circulatory dynamics of the analyzed groups, while thermography fails to enable objectifying a reduction in the amount of blood in the areas evaluated in women with FM. This would be consistent with the results of the study by Dibai Filho's group, in which they also found no temperature differences between women with chronic neck pain and healthy women and which already suggested that the circulatory dynamics would be similar in both groups without a reduction of blood in subjects with pain [53].

Previous studies suggest that pain in FMS is caused by central and peripheral sensitization [54], altered pain mechanisms and the presence of inflammatory substances both in the spinal cord, brain and the peripheral tissues [55]. However, it is possible that this peripheral inflammation (neuropeptides, chemokines and cytokines), reported in previous studies [3,6,56], is not revealed at the depth of 5 mm where infra-red radiation is emitted by the skin [57]. In this regard, we did not determine any peripheral biomarker of inflammation, since it was beyond the scope of the manuscript. Likewise, there are other trends suggesting that central sensitivity could play a more important role in the perception of pain in FMS [58], due to a reduced central inhibition of pain, which perpetuates it and broadens its nociceptive signal [59]. Thus, people with FM can refer pain without any type of stimulus, something which could explain why this study fails to report differences in temperature between healthy women and women with fibromyalgia at rest.

We need to take into consideration that the assessments were conducted at rest, since our goal was to explore the usefulness of this method as an easy-to-use complementary clinical technique to help in the assessment of the FM. Further studies analyzing the thermal response

during sportive activities could provide interesting information regarding thermal patterns in this population.

Further, several study limitations need to be considered when evaluating our results. First, this is a cross-sectional study, so no cause-effect can be drawn from our results, since there were no follow-up of the participants. Second, people with FM conformed a heterogeneous population thus, even though the sample size calculation was well conducted and more participants than necessary were recruited, the heterogeneity may require even a larger sample size. Besides, a non-probabilistic sampling method of recruitment was used that may jeopardize the representativeness of the sample. Future studies using subgroups, exploring the effect of other possible confounder variables and performing other sportive or active protocols would provide more precise information.

Conclusions

There are no clinical differences in skin temperature at rest between women with FM and healthy women in the core regions of the body except for the low-back region in which the significant differences did not achieve a minimum of clinically detectable change. Further, women with FM showed a significant higher temperature in knees although the differences also did not reach the minimum of clinical detectable change. Infra-red thermography is not an effective supplementary assessment aid in women with fibromyalgia.

Supporting information

S1 Data.

(XLSX)

Acknowledgments

We thank the Spanish fibromyalgia associations for their willingness to contribute to the dissemination of the study.

Author Contributions

Conceptualization: Marta Inglés, Pilar Serra-Añó.

Data curation: Nuria Sempere-Rubio, Marta Aguilar-Rodríguez, Marta Inglés, Ruth Izquierdo-Alventosa, Pilar Serra-Añó.

Formal analysis: Marta Aguilar-Rodríguez, Marta Inglés, Pilar Serra-Añó.

Investigation: Nuria Sempere-Rubio.

Methodology: Nuria Sempere-Rubio.

Supervision: Pilar Serra-Añó.

Writing – original draft: Nuria Sempere-Rubio, Marta Inglés, Pilar Serra-Añó.

Writing – review & editing: Pilar Serra-Añó.

References

1. Wolfe F, Clauw DJ, Fitzcharles M-A., Goldenberg DL., Katz RS., Mease P. The American College of Rheumatology preliminary diagnostic criteria for fibromyalgia and measurement of symptom severity. *Arthritis Care Res.* 2010; 62:600–610. <https://doi.org/10.1002/acr.20140> PMID: 20461783

2. Thieme K, Turk DC, Flor H. Comorbid depression and anxiety in fibromyalgia syndrome: relationship to somatic and psychosocial variables. *Psychosom Med.* 2004; 66:837–844. <https://doi.org/10.1097/01.psy.0000146329.63158.40> PMID: 15564347
3. Groven N, Fors EA, Iversen VC, White LR, Reitan SK. Peripheral inflammation in fibromyalgia syndrome. *Eur Neuropsychopharmacol.* 2017; 27:S634.
4. Cordero MD, Díaz-Parrado E, Carrión AM, Alfonsi S, Sánchez-Alcazar JA, Bullón P, et al. Is inflammation a mitochondrial dysfunction-dependent event in fibromyalgia? *Antioxid Redox Signal* 2013; 18:800–807. <https://doi.org/10.1089/ars.2012.4892> PMID: 22938055
5. Bäckryd E, Tanum L, Lind A-L, Larsson A, Gordh T. Evidence of both systemic inflammation and neuroinflammation in fibromyalgia patients, as assessed by a multiplex protein panel applied to the cerebrospinal fluid and to plasma. *J Pain Res.* 2017; 10:515–525. <https://doi.org/10.2147/JPR.S128508> PMID: 28424559
6. Bazzichi L, Rossi A, Massimetti G, Giannaccini G, Giuliano T, De Feo F, et al. Cytokine patterns in fibromyalgia and their correlation with clinical manifestations. *Clin Exp Rheumatol.* 2007; 25(2):225. PMID: 17543146.
7. Kang JH, Kim JK, Hong SH, Lee CH, Choi BY. Heart Rate Variability for Quantification of Autonomic Dysfunction in Fibromyalgia. *Ann Rehabil Med.* 2016; 40(2):301–9. <https://doi.org/10.5535/arm.2016.40.2.301> PMID: 27152281
8. Rizzi M, Radovanovic D, Santus P, Airoldi A, Frassanito F, Vanni S, et al. Influence of autonomic nervous system dysfunction in the genesis of sleep disorders in fibromyalgia patients. *Clin Exp Rheumatol.* 2017; 35 Suppl 105(3):74–80. <https://doi.org/10.1136/annrheumdis-2017-eular.3365> PMID: 28681708
9. Clauw DJ. Fibromyalgia: A Clinical Review. *JAMA.* 2014; 311(15):1547–55. <https://doi.org/10.1001/jama.2014.3266> PMID: 24737367
10. Elmas O, Yildiz S, Bilgin S, Demirci S, Comlekci S, Koyuncuoglu HR, et al. Physiological parameters as a tool in the diagnosis of fibromyalgia syndrome in females: A preliminary study. *Life Sci.* 2016; 145:51–6. <https://doi.org/10.1016/j.lfs.2015.12.029> PMID: 26685758
11. Wolfe F, Clauw DJ, Fitzcharles M-A, Goldenberg DL, Häuser W, Katz RS, et al. Fibromyalgia criteria and severity scales for clinical and epidemiological studies: a modification of the ACR Preliminary Diagnostic Criteria for Fibromyalgia. *J Rheumatol.* 2011; 38(6):1113–1122. <https://doi.org/10.3899/jrheum.100594> PMID: 21285161
12. Borchers AT, Gershwin ME. Fibromyalgia: a critical and comprehensive review. *Clin Rev Allergy Immunol.* 2015; 49(2):100–151. <https://doi.org/10.1007/s12016-015-8509-4> PMID: 26445775
13. Wolfe F, Clauw DJ, Fitzcharles M-A, Goldenberg DL, Häuser W, Katz RL, et al. 2016 Revisions to the 2010/2011 fibromyalgia diagnostic criteria. *Semin Arthritis Rheum.* 2016; 46(3):319–329. <https://doi.org/10.1016/j.semarthrit.2016.08.012> PMID: 27916278
14. Rajapakse C, Grennan DM, Jones C, Wilkinson L, Jayson M. Thermography in the assessment of peripheral joint inflammation—a re-evaluation. *Rheumatology* 1981; 20:81–87. <https://doi.org/10.1093/rheumatology/20.2.81> PMID: 7280484
15. Denoble AE, Hall N, Pieper CF, Kraus VB. Patellar Skin Surface Temperature by Thermography Reflects Knee Osteoarthritis Severity. *Clin Med Insights Arthritis Musculoskelet Disord.* 2010; 3:69–75. <https://doi.org/10.4137/CMAMD.S5916> PMID: 21151853
16. Vecchio P, Adebajo A, Chard M, Thomas PP, Hazleman B. Thermography of frozen shoulder and rotator cuff tendinitis. *Clin Rheumatol.* 1992; 11:382–384. <https://doi.org/10.1007/BF02207197> PMID: 1458786
17. Miyakoshi N, Itoi E, Sato K, Suzuki K, Matsuura H. Skin temperature of the shoulder: Circadian rhythms in normal and pathologic shoulders. *J Shoulder Elbow Surg.* 1998; 7(6): 625–8. [https://doi.org/10.1016/s1058-2746\(98\)90012-x](https://doi.org/10.1016/s1058-2746(98)90012-x) PMID: 9883425
18. Vargas JVC, Brioschi ML, Dias FG, Parolin MB, Mulinari-Brenner FA, Ordonez JC, et al. Normalized methodology for medical infrared imaging. *Infrared Phys Technol.* 2009; 52(1):42–47. <https://doi.org/10.1016/j.infrared.2008.11.003>
19. Szentkuti A, Kavanagh HS, Grazio S. Infrared thermography and image analysis for biomedical use. *Period Biol.* 2011; 113(4):385–392.
20. Niehof SP, Huygen FJPM, Van der Weerd RWP, Westra M, Zijlstra FJ. Thermography imaging during static and controlled thermoregulation in complex regional pain syndrome type 1: diagnostic value and involvement of the central sympathetic system. *Biomed Eng Online.* 2006; 5:30. <https://doi.org/10.1186/1475-925X-5-30> PMID: 16689997
21. Iwase S, Cui J, Wallin BG, Kamiya A, Mano T. Effects of increased ambient temperature on skin sympathetic nerve activity and core temperature in humans. *Neurosci Lett.* 2002; 327(1):37–40. [https://doi.org/10.1016/s0304-3940\(02\)00374-9](https://doi.org/10.1016/s0304-3940(02)00374-9) PMID: 12098495

22. Larson AA, Pardo JV, Pasley JD. Review of Overlap Between Thermoregulation and Pain Modulation in Fibromyalgia: *Clin J Pain*. 2014; 30(6):544–555. <https://doi.org/10.1097/AJP.0b013e3182a0e383> PMID: 23887348
23. Moreira DG, Costello JT, Brito CJ, Adamczyk JG, Ammer K, Bach AJ, et al. Thermographic imaging in sports and exercise medicine: a Delphi study and consensus statement on the measurement of human skin temperature. *J Therm Biol*. 2017; 69:155–162. <https://doi.org/10.1016/j.jtherbio.2017.07.006> PMID: 29037377
24. Selvan AN, Childs C. Infrared Thermal Mapping, Analysis and Interpretation in Biomedicine. [Internet]. Springer Singapore; 2017: 377–394. http://link.springer.com/10.1007/978-981-10-3147-2_21.
25. Ludwig N, Formenti D, Gargano M, Alberti G. Skin temperature evaluation by infrared thermography: Comparison of image analysis methods. *Infrared Phys Technol*. 2014; 62:1–6. <https://doi.org/10.1016/j.infrared.2013.09.011>.
26. Ammer K, Schartermüller T, Melnizky P. Thermography in fibromyalgia. *Biomed Thermol*. 1995; 15:77–80.
27. Ammer K. Thermal imaging: a diagnostic aid for fibromyalgia? *Thermol Int*. 2008; 18:45–50.
28. Biasi G, Fioravanti A, Franci A, Marcolongo R. The role computerized telethermography in the diagnosis of fibromyalgia syndrome. *Minerva Med*. 1994; 85(9):451–4. PMID: 7936366.
29. Charkoudian N, Hart ECJ, Barnes JN, Joyner MJ. Autonomic control of body temperature and blood pressure: influences of female sex hormones. *Clin Auton Res*. 2017; 27(3):149–155. <https://doi.org/10.1007/s10286-017-0420-z> PMID: 28488202
30. Fernández-Cuevas I, Bouzas Marins JC, Arnáiz Lastras J, Gómez Carmona PM, Piñonosa Cano S, García-Concepción MA, et al. Classification of factors influencing the use of infrared thermography in humans: A review. *Infrared Phys Technol Elsevier*. 2015; 71:28–55. <http://dx.doi.org/10.1016/j.infrared.2015.02.007>.
31. Ammer K. The Glamorgan Protocol for recording and evaluation of thermal images of the human body. *Thermol Int*. 2008; 18(4):125–44.
32. Scolnik M, Vasta B, Hart DJ, Shipley JA, McHugh NJ, Pauling JD. Symptoms of Raynaud's phenomenon (RP) in fibromyalgia syndrome are similar to those reported in primary RP despite differences in objective assessment of digital microvascular function and morphology. *Rheumatol Int*. 2016; 36(10):1371–1377. <https://doi.org/10.1007/s00296-016-3483-6> PMID: 27136918
33. Neves EB, Vilaça-Alves J, Rosa C, Reis VM. Thermography in neurologic practice. *Open Neurol J*. 2015; 9:24–27. Neves EB, Vilaça-Alves J, Rosa C, Reis VM. Thermography in Neurologic Practice. *Open Neurol J*. 2015; 9:24–27. <https://doi.org/10.2174/1874205X01509010024> PMID: 26191090
34. Ring EFJ, Ammer K. Infrared thermal imaging in medicine. *Physiol Meas*. 1 de marzo de 2012; 33(3):33–46. <https://doi.org/10.1088/0967-3334/33/3/R33> PMID: 22370242
35. World Health Organization, editor. Obesity: preventing and managing the global epidemic: report of a WHO consultation. Geneva: World Health Organization; 2000. 253 p. (WHO technical report series). ISBN: 92 4 120894 5.
36. Ruggiero L, Manganelli F, Santoro L. Muscle pain syndromes and fibromyalgia: the role of muscle biopsy. *Curr Opin Support Palliat Care*. 2018; 12(3):382–387. <https://doi.org/10.1097/SPC.0000000000000355> PMID: 29912728
37. Olausson P, Ghafouri B, Gerdle B. Specific proteins of the trapezius muscle correlate with pain intensity and sensitivity—an explorative multivariate proteomic study of the trapezius muscle in women with chronic widespread pain. *J Pain Res*. 2016; 9:345–356. <https://doi.org/10.2147/JPR.S102275> eCollection 2016. PMID: 27330327
38. Üçeyler N., Zeller D., Kahn AK., Kewenig S., Kittel-Schneider S., Schmid A., et al. Small fibre pathology in patients with fibromyalgia syndrome. *Brain J Og Neurol*. 2013; 136:16857–1867. <https://doi.org/10.1093/brain/awt053> PMID: 23474848
39. Sánchez-Domínguez B, Bullón P, Román-Malo L, Marín-Aguilar F, Alcocer-Gómez E, Carrión AM, et al. Oxidative stress, mitochondrial dysfunction and, inflammation common events in skin of patients with Fibromyalgia. *Mitochondrion*. 2015; 21:69–75. <https://doi.org/10.1016/j.mito.2015.01.010> PMID: 25662535
40. Schürmann M, Zaspel J, Löhr P, Wizgall I, Tutic M, Manthey N, et al. Imaging in Early Posttraumatic Complex Regional Pain Syndrome: A Comparison of Diagnostic Methods. *Clin J Pain*. 2007; 23(5):449–457. <https://doi.org/10.1097/AJP.0b013e31805c9e66> PMID: 17515744
41. Nahm FS. Infrared Thermography in Pain Medicine. *Korean J Pain*. 2013; 26(3):219–222. <https://doi.org/10.3344/kjp.2013.26.3.219> PMID: 23861995

42. Choi E, Lee P-B, Nahm FS. Interexaminer reliability of infrared thermography for the diagnosis of complex regional pain syndrome. *Skin Res Technol*. 2013; 19(2):189–193. <https://doi.org/10.1111/srt.12032> PMID: 23331254
43. Uematsu S, Edwin DH, Jankel WR, Kozikowski J, Trattner M. Quantification of thermal asymmetry: part 1: normal values and reproducibility. *J Neurosurg*. 1988; 69(4):552–555. <https://doi.org/10.3171/jns.1988.69.4.0552> PMID: 3418388
44. Paeng SH, Jung YT, Pyo SY, Kim MS, Jeong YG. Is the use of digital infrared thermal imaging useful in whiplash injury? *Korean J Spine*. 2009; 6(4):274–279. <https://doi.org/10.3340/jkns.2015.57.4.283>
45. Brenner M. Thermal signature analysis as a novel method for evaluating inflammatory arthritis activity. *Ann Rheum Dis*. 2006; 65(3):306–311. <https://doi.org/10.1136/ard.2004.035246> PMID: 16150784
46. Haddad DS, Brioschi ML, Arita ES. Thermographic and clinical correlation of myofascial trigger points in the masticatory muscles. *Dentomaxillofacial Radiol*. 2012; 41(8):621–629. <https://doi.org/10.1259/dmfr/98504520> PMID: 23166359
47. Nahm FS, Lee PB, Park SY, Kim YC, Lee SC. Comparison of the Diagnostic Validity of Real and Absolute Skin Temperature Differences for Complex Regional Pain Syndrome. *Korean J Pain*. 2009; 22(2):146. <https://doi.org/10.3344/kjp.2009.22.2.146>.
48. Ericsson A, Mannerkorpi K. How to manage fatigue in fibromyalgia: nonpharmacological options. *Pain Manag*. 2016; 6(4):331–338. <https://doi.org/10.2217/pmt-2016-0015> PMID: 27297077
49. Selfe J, Whitaker J, Hardaker N. A narrative literature review identifying the minimum clinically important difference for skin temperature asymmetry at the knee. *Thermol Int*. 2008; 18(2):41–4.
50. Wolfe F, Smythe HA, Yunus MB, Bennett RM, Bombardier C, Goldenberg DL, et al. The American College of Rheumatology 1990 Criteria for the Classification of Fibromyalgia. Report of the Multicenter Criteria Committee. *Arthritis Rheum*. 1990; 33(2):160–172. PMID: 2306288
51. Al-Qudah ZA, Yacoub HA, Souayah N. Disorders of the autonomic nervous system after hemispheric cerebrovascular disorders: an update. *J Vasc Interv Neurol*. 2015; 8(4):43–52. PMID: 26576215.
52. Jasti N, Bista S, Bhargav H, Sinha S, Gupta S, Chaturvedi SK, et al. Medical applications of Infrared thermography: A narrative review. *J Stem Cells*. 2019; 14(1):35–53. ISSN: 1556-8539.
53. Dibai Filho AV, Packer AC, Costa AC, Berni-Schwarzenbeck KC, Rodrigues-Bigaton D. Assessment of the upper trapezius muscle temperature in women with and without neck pain. *J Manipulative Physiol Ther*. 2012; 413–417. <https://doi.org/10.1016/j.jmpt.2012.04.006> PMID: 22608286
54. Nielsen LA, Henriksson KG. Pathophysiological mechanisms in chronic musculoskeletal pain (fibromyalgia): the role of central and peripheral sensitization and pain disinhibition. *Best Pract Res Clin Rheumatol*. 2007; 21(3):465–480. <https://doi.org/10.1016/j.berh.2007.03.007> PMID: 17602994
55. Littlejohn G. Neurogenic neuroinflammation in fibromyalgia and complex regional pain syndrome. *Nat Rev Rheumatol*. 2015; 11(11):639–648. <https://doi.org/10.1038/nrrheum.2015.100> PMID: 26241184
56. Sluka KA, Clauw D. J. Neurobiology of fibromyalgia and chronic widespread pain. *Neuroscience*. 2016; 338:114–129. <https://doi.org/10.1016/j.neuroscience.2016.06.006> PMID: 27291641
57. Leando P. infrared thermal imaging (DITI) for referring health care practitioners. *Meditherm Inc*. 2003; 1–37. https://doi.org/10.1007/978-981-10-3147-2_21
58. Larsson A, Palstam A, Löfgren M, Ernberg M, Bjersing J, Bileviciute-Ljungar I, et al. Pain and fear avoidance partially mediate change in muscle strength during resistance exercise in women with fibromyalgia. *J Rehabil Med*. 2017; 49: 744–750. <https://doi.org/10.2340/16501977-2278> PMID: 29068036
59. Hermans L, Nijs J, Calders P. Influence of Morphine and Naloxone on Pain Modulation in Rheumatoid Arthritis, Chronic Fatigue Syndrome/Fibromyalgia, and Controls. A Double-Blind, Randomized, Placebo-Controlled, Cross-Over Study. *Pain Pract Off J World Inst Pain*. 2018; 18:418–430. <https://doi.org/10.1111/papr.12613> PMID: 28722815