

CORRECTION

Correction: Validation of the Surgical Fear Questionnaire in Adult Patients Waiting for Elective Surgery

Maurice Theunissen, Madelon L. Peters, Erik G. W. Schouten, Audrey A. A. Fiddlers, Mark G. A. Willemsen, Patricia R. Pinto, Hans-Fritz Gramke, Marco A. E. Marcus

There are three errors in the “Predictor” column of Table 7. For the last APSP, CPSP, and GSR “Outcome” rows, the correct predictor should be SFQ-1, not SGQ-1. Please view the corrected table below.



 OPEN ACCESS

Citation: Theunissen M, Peters ML, Schouten EGW, Fiddlers AAA, Willemsen MGA, Pinto PR, et al. (2016) Correction: Validation of the Surgical Fear Questionnaire in Adult Patients Waiting for Elective Surgery. PLoS ONE 11(9): e0162737. doi:10.1371/journal.pone.0162737

Published: September 9, 2016

Copyright: © 2016 Theunissen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Table 7. Predictive validity of the SFQ, SFQ-s and SFQ-I.

Outcome	Predictor	Study 1		Study 4	
		OR	(95 CI)	OR	(95 CI)
APSP	SFQ	2.73	(1.59–4.69)***	2.35	(1.81–3.04)***
APSP	SFQ-s	1.83	(1.08–3.10)	2.12	(1.64–2.73)***
APSP	SFQ-I	3.55	(1.99–6.32)***	2.62	(2.02–3.39)***
CPSP	SFQ	1.77	(1.25–2.51)**	2.28	(1.56–3.34)***
CPSP	SFQ-s	1.66	(1.16–2.37)**	1.66	(1.15–2.39)**
CPSP	SFQ-I	1.77	(1.24–2.51)**	3.05	(2.06–4.51)***
GSR	SFQ	0.44	(0.31–0.62)***	0.56	(0.41–0.77)***
GSR	SFQ-s	0.61	(0.43–0.87)**	0.77	(0.56–1.05)
GSR	SFQ-I	0.44	(0.31–0.63)***	0.40	(0.29–0.55)***

Bivariate logistic regression with median split SFQ, SFQ short term and SFQ long term as predictor. APSP: acute postsurgical pain on day 4. CPSP: chronic postsurgical pain after 6 months in Study 1, after one year in Study 4. Pain scores were dichotomized using a cut of value of 40 for the VAS/NRS. GSR: global surgical recovery on a scale of 0–100%, values of 80–100% were considered as good recovery; long term GSR after 6 months in Study 1, after one year in Study 4.

To adjust for multiple testing a Bonferroni correction was applied: a *p*-value <0.017 was considered statistically significant.

***p* <0.01,

****p* <0.001.

doi:10.1371/journal.pone.0162737.t001

Reference

1. Theunissen M, Peters ML, Schouten EGW, Fiddelaers AAA, Willemsen MGA, Pinto PR, et al. (2014) Validation of the Surgical Fear Questionnaire in Adult Patients Waiting for Elective Surgery. PLoS ONE 9(6): e100225. doi:10.1371/journal.pone.0100225 PMID: 24960025