


Effect of a ketogenic diet on pain and quality of life in patients with lipedema: The LIPODIET pilot study

Vilde Sørli¹ | Ann Kristin De Soysa² | Åsne Ask Hyldmo³ | Kjetil Retterstøl¹ |
Catia Martins^{3,4} | Siren Nymo^{3,4,5} 

¹Department of Nutrition, Institute of Basic Medical Sciences, Faculty of Medicine, University of Oslo, Oslo, Norway

²Department of Clinical Nutrition and Speech Pathology, Clinic of Clinical Services, St. Olav Hospital–Trondheim University Hospital, Trondheim, Norway

³Centre of Obesity Research and Innovation, Clinic of Surgery, St. Olav Hospital – Trondheim University Hospital, Trondheim, Norway

⁴Department of Clinical and Molecular Medicine, Centre for Obesity Research, Faculty of Medicine, Norwegian of Science and Technology, Trondheim, Norway

⁵Namsos Hospital, Clinic of Surgery, Nord-Trøndelag Hospital Trust, Namsos, Norway

Correspondence

Siren Nymo, Obesity Research Group, Department of Clinical and Molecular Medicine, Faculty of Medicine and Health Science, Norwegian University of Science and Technology, Forsyningscenteret, Prinsesse Kristinas gate 5, 7030 Trondheim, Norway.
Email: siren.nymo@ntnu.no

Funding information

The Norwegian Lymphoedema and Lipoedema association

Abstract

Background: Lipedema is an underdiagnosed condition in women, characterized by a symmetrical increase in subcutaneous adipose tissue (SAT) in the lower extremities, sparing the trunk. The lipedema SAT has been found to be resistant to diet, exercise and bariatric surgery, in regard to both weight loss (WL) and symptom relief. Current experience indicates that a low carbohydrate and high fat (LCHF-diet) might have a beneficial effect on weight and symptom management in lipedema.

Objective: To assess the impact of an eucaloric low carbohydrate, high fat (LCHF)-diet on pain and quality of life (QoL) in patients with lipedema.

Methods: Women diagnosed with lipedema, including all types and stages affecting the legs, (age 18–75 years, BMI 30–45 kg/m²) underwent 7 weeks of LCHF-diet and, thereafter 6 weeks of a diet following the Nordic nutrition recommendations. Pain (visual analog scale) and QoL (questionnaire for lymphedema of the limbs), weight and body composition were measured at baseline, week seven and 13.

Results: Nine women (BMI: 36.7 ± 4.5 kg/m² and age: 46.9 ± 7 years) were recruited. The LCHF diet induced a significant WL −4.6 ± 0.7 kg (−4.5 ± 2.4%), $p < 0.001$ for both, and reduction in pain (−2.3 ± 0.4 cm, $p = 0.020$). No correlation was found between WL and changes in pain at week seven ($r = 0.283$, $p = 0.460$). WL was maintained between week seven and 13 (0.3 ± 0.7 kg, $p = 0.430$), but pain returned to baseline levels at week 13 (4.2 ± 0.7 cm, $p = 0.690$). A significant increase in general QoL was found between baseline and week seven (1.0 (95% CI (2.0, 0.001)), $p = 0.050$) and 13 (1.0 95% CI (2.0, 0.001) $p = 0.050$), respectively.

Conclusion: A LCHF-diet is associated with reduction in perceived pain and improvement in QoL, in patients with lipedema. Larger randomized clinical trials are needed to confirm these findings.

KEYWORDS

bodyfat distribution, ketogenic diet, obesity, obesity phenotype, quality of life

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2021 The Authors. Obesity Science & Practice published by World Obesity and The Obesity Society and John Wiley & Sons Ltd.

1 | INTRODUCTION

Lipedema is an obesity related, underdiagnosed health condition affecting women exclusively, and is characterized by a symmetrical increase of subcutaneous adipose tissue (SAT) in the lower extremities, sparing the trunk.¹ As many as 50% of patients with lipedema may present with excess body weight, hence lipedema is often misdiagnosed as overweight or obesity.² Disease onset of lipedema is usually at puberty, during pregnancy or at menopause.^{2,3} Lipedema affects women almost exclusively, but it has also been observed in men with hormonal imbalances or liver disease.^{4,5} The diagnosis of lipedema is made clinically by family- and medical history, visual inspection and physical examination, as there are no blood or urine biomarkers for lipedema.³ Based on what body parts are involved, lipedema is differentiated into five types and it may progress through four stages.³ The main criteria to diagnose lipedema is bilateral and symmetrical fat deposits downward from the hips, while the feet are spared. Moreover, non-pitting edema, easy bruising, tenderness and pain in the affected regions are common in this patient group, even though they cannot be used as diagnostic criteria.⁴ However, clinical practice has shown that lipedema fat tissue can also be found in other areas of the body that not legs and arms, namely the abdomen.⁶ Most importantly, these patients have no or little response on loss of lipedema fat from traditional obesity treatments, such as lifestyle modification or bariatric surgery.^{1-3,7} Little is known about the etiology of lipedema, but the vascular and lymphatic systems seem to be implicated.⁸

Even though the etiology of pain in this patient group remains unknown, it might reflect an active state of inflammation within the SAT.^{8,9} The Lipedema UK and The British Lymphology Society (2012)¹⁰ reported that 87% of the 250 females interviewed agreed that lipedema had a negative effect on quality of life (QoL). Since it is unknown what is causing the pain, an optimal and effective treatment is unclear.^{11,12} Today's treatment options consist of conservative physical therapy, surgical liposuction and lifestyle modifications, including altering dietary intake and increasing physical activity.¹

Patients' experiences seems to indicate that a low-carbohydrate and high-fat diet (LCHF-diet), also called a ketogenic diet (KD), may have an effect on weight and symptom management in lipedema.¹³⁻¹⁵ KD are characterized by a reduction in carbohydrates, usually to less than 50 g/day,¹⁶ and a relative increase in the proportions of fat (70-75 E%), and protein (20 E%), inducing a ketogenic metabolic state.¹⁷ "The lipedema project", by Seo and Keith, promote a Ketogenic Way of Eating (Woe) for patients with lipedema, and has reported successful outcomes.^{14,18} Women on the ketogenic Woe report reduction or elimination of swelling and pain, and weight loss (WL).^{13,18} However, these reports came from patients' experiences, not clinical trials, therefore more research is needed to confirm these findings and to investigate if KD is a possible treatment option for lipedema. Recently, Renzo et al (2021) reported a reduction in fat mass (FM) and an improvement in QoL with a Mediterranean diet in patients with lipedema.¹⁹

The main objective of this study was to assess if an eucaloric LCHF-diet lead to change in pain and QoL in patients with lipedema. The second aim was to assess if a LCHF diet led to change in body composition in patients with lipedema. The hypothesis of the study were that a LCHF-diet would lead to a reduction in pain and increase in QoL, independently of WL, in lipedema patients.

2 | METHODS

2.1 | Selection and descriptions of participants

2.1.1 | Participants

Women (18-75 years old) with obesity (BMI: 30-45 kg/m²) and diagnosed with lipedema, including all types and stages affecting the legs, were included in the study. The participants had to be weight stable (<2 kg variation within the last 3 months) and not currently dieting to lose weight or on a KD. Exclusion criteria was pregnancy or breast feeding, history of infectious diseases, medication known to affect body weight and enrollment in any other obesity treatment. Moreover, those with a history of psychological disorders, not mastering a Scandinavian language, having a malign disease or any disease that leads to dietary advice that is not consistent with the intervention, or difficulties following the instructions, were not accepted into the study. Participants were recruited via announcement at obesity treatment facilities and Internet, (Lipedema-groups on Facebook and the intranet at the University Hospital in Trondheim), posters in the region, and through the Norwegian lymphedema- and lipedema association. All participants provided a written informed consent before commencement.

2.2 | Technical information

2.2.1 | Study design

This was a repeated measures study, where females with lipedema underwent 7 weeks of an eucaloric LCHF-diet, followed by 6 weeks of an isocaloric diet in line with the Nordic Nutrition Recommendations. The LIPODIET is a controlled, prospective clinical trial approved by the Regional Committee for Medical Research Ethics in Norway (2018/307) and registered in [ClinicalTrials.gov](https://www.clinicaltrials.gov/ct2/show/NCT03710798) (NCT03710798) (<https://www.clinicaltrials.gov/ct2/show/NCT03710798>).

The study design is illustrated in Figure 1.

2.2.2 | LCHF-phase

All participants followed a seven-week eucaloric LCHF-diet (fat: 70-75 E%, carbohydrate: 5-10 E%, protein: 20 E%). They received individual dietary plans at baseline, together with a recipe booklet. Dietary plans were based on individual calculated energy needs




FIGURE 1 Timeline of data collection throughout the LIPODIET-study

(resting energy expenditure (REE) \times physical activity levels (PAL)) aiming at maintaining baseline weight. REE was measured in the fasting state by indirect calorimetry (Vmax Encore 29N; CareFusion), at baseline, using a canopy system and following standard procedures.²⁰ PAL was measured with armbands (SenseWear, Body Media).

2.2.3 | Nordic nutrition recommendations-phase

During week 8, the participants were gradually introduced to a 6-week eucaloric diet (dietary plans and a recipe booklet as for the LCHF-phase) based on the Nordic nutrition recommendations (NNR). The control diet used in this study was based on the NNR that constitutes the scientific basis for national nutrient recommendations and dietary guidelines in Norway. The purpose is to help prevent chronic diet-related diseases in the population.²¹

2.2.4 | Compliance

The participants had 30-min weekly follow-ups with the dietitian aiming to ensure compliance with the diets. Moreover, patients were asked to fill out daily food records throughout the study period. Food records from baseline, weeks 6 and 11 were then analyzed for macro- and micronutrient composition using *Kostholdsplanleggeren* (Norwegian Food Safety Authority and Norwegian Directorate of Health, 2016).²² Acetoacetate (AcAc) was measured in the first urine of the day at the weekly follow-ups, using ketostix (Bayer Corp.), as a measure of compliance during the LCHF diet. Participants were asked not to change their activity throughout the study and armbands (SenseWear, Body Media) were used for 7 days, at pre-baseline, week six and week 12, to check for compliance. For data to be considered valid, participants had to wear the armbands for <7 days, including at least 1 weekend day, and more than 95% of data needed to be available over a 24 h period. PAL-value and number of steps/day were analyzed.²³

2.2.5 | Data collection

The following variables were measured at baseline, week seven and week 13.

2.2.6 | Pain

To measure the perceived pain, a visual analog scale (VAS) was used.^{24,25} Participants were asked to place a perpendicular line on the VAS-scale to represent their current pain-intensity at baseline, W7 and W13. The question used was: "In how much pain are you right now?" (0 = No pain at all 10 = As much pain you can possibly imagine).

2.2.7 | Quality of life

The Norwegian version of the QoL questionnaire for lymphedema of the legs, a validated health-related QoL questionnaire²⁶ was used. The questions cover four domains: symptoms, body image/appearance, function, and mood. Each item in each domain was scored after the Likert's scale, 1–4: not at all = 1, a little = 2, quite a bit = 3, a lot = 4.

2.2.8 | Weight and body composition

Height was measured following standard procedures,²⁷ using a measuring tape installed on the wall. Bioelectrical impedance analysis (BIA) was used (InBody 720 (BIA)) at baseline, week seven and week 13 to measure body weight (BW) and to analyze body composition.^{28–30} Waist, hip, thigh, and calf circumference were measured at baseline, week seven and week 13 using a measuring tape. The waist was defined as the midway between the lower rib and the iliac crest.³¹ The participants were asked to breathe normally, and the measurement was taken as they exhaled. Hip circumference was measured over the largest part of the buttocks.^{31–33} The thigh and calf circumferences were measured as the participants were lying on a bench and measured where the circumference of the thigh/calf apparently was largest. To ensure that the thigh and calf circumference was measured at approximately the same place each time, the length (cm) from the foot to the point, was measured and recorded for the next measurement. These circumference measures were taken only wearing underwear.

2.2.9 | Blood samples

Blood samples were collected in fasting at baseline, week seven and week 13, at the laboratory of the University Hospital in Trondheim,

to ensure that the participants tolerated the diet and that the diet did not lead to elevated blood lipids and/or deteriorated kidney function.^{17,34} The parameters measured in the blood were C-reactive protein (CRP), high-density lipoprotein (HDL) cholesterol, low-density protein (LDL) cholesterol, triglycerides, glucose, glycated hemoglobin (HbA1c), alanine transaminase (ALAT), gamma-glutamyl transferase (GT), alkaline phosphatase (ALP), natrium and potassium.

2.3 | Statistical analyses

Statistical analysis was carried out using IBM SPSS Statistics 26 (SPSS Inc.) and statistical significance assumed at 5%, unless otherwise stated, and data presented as mean \pm SEM, except for baseline anthropometric data, where mean \pm SD was used. Data was analyzed using linear mixed models (LLM) and Bonferroni adjustment were used for comparison between timepoints (baseline, week seven and week 13). Correlation analysis between WL at week seven and week 13 and change in perceived pain was performed using Spearman *r* correlation.

3 | RESULTS

Nine participants met study entry criteria and were included in the study with an average age of 46.9 ± 9.0 years and with a BMI of 36.7 ± 4.5 kg/m². One participant did not show up for testing at week 13, due to illness, but was included in the analysis. Mean attendance of weekly follow-up meetings throughout the study period was 85.7% (range: 71.4%–100.0%).

3.1 | Body weight and composition

Changes in body weight and composition overtime can be seen in Table 1. The LCHF induced a significant WL (-4.6 ± 0.7 kg ($-4.5 \pm 2.4\%$), $p < 0.001$), which was maintained at week 13 (-4.1 ± 0.7 kg ($-4.0 \pm 2.4\%$), $p < 0.001$). No significant change in BW was seen from week 7 to week 13 ($p = 0.430$).

There was a significant decrease in waist (98.3 ± 2.7 vs. 94.0 ± 2.7 cm, $p < 0.001$) and hip-circumference (125.2 ± 1.6 vs. 123.0 ± 1.6 cm, $p = 0.010$) from baseline to week 7, and this was maintained at week 13. There was a significant decrease in calf-circumference (48.0 ± 3.8 vs. 47.0 ± 3.8 cm, $p = 0.030$) from baseline to week seven, but no significant change in thigh-circumference (67.0 ± 3.0 vs. 65.0 ± 3.0 cm, $p = 0.200$) during the same period.

3.2 | Compliance

3.2.1 | Diet

All participants achieved ketosis, as positive AcAc (>0.5 mmol/L) was detected in urine during the LCHF-diet, according to objective

observations of color change on the ketostix. AcAc increased from baseline to week 7 (>0.05 – 8 mmol/L) and returned to baseline levels at week 13 (<0.03).

Changes in energy, macro- and micronutrients intake over time are shown in Table 2. There was no significant change in median total energy intake between any of the three time points. The total fat intake increased significantly from baseline to week 6 (88.0 ± 8.6 vs. 146.0 ± 8.6 g/day, $p < 0.001$), and decreased significantly from week 6 to week 11 (146.0 ± 8.6 vs. 73.0 ± 8.6 g/day, $p < 0.001$). Simultaneously the total intake of carbohydrates decreased significantly from baseline to week 6 (182.0 ± 21.6 vs. 28.0 ± 21.6 g/day, $p < 0.001$) and increased from week six to week 11 (28.0 ± 21.6 vs. 197 ± 21.6 g/day, $p < 0.001$). There were no significant differences between baseline and week 11 in any of the nutrients.

The daily intake of saturated fat, unsaturated fat, cholesterol, sugar, fiber, omega-3 and omega-6 are shown in Table 3. Between baseline and week six there was a significant increase in the total intake of saturated fat (32.0 ± 3.6 g/day vs. 58.8 ± 3.6 g/day, $p < 0.001$), unsaturated fat (46.5 ± 4.6 g/day vs. 67.7 ± 4.6 g/day, $p = 0.03$) cholesterol (325.4 ± 53.0 g/day vs. 594.8 ± 53.0 g/day, $p = 0.002$), omega-3 (2.4 ± 0.58 g/day vs. 6.5 ± 0.58 g/day, $p < 0.001$) and omega-6 fatty acids (10.2 ± 1.6 g/day vs. 16 ± 1.6 g/day, $p = 0.025$). There was also a significant decrease from week six to week 11 in total intake of saturated fat (58.8 ± 3.6 g/day vs. 27.5 ± 3.6 g/day, $p < 0.001$), unsaturated fat (67.7 ± 4.6 g/day vs. 37.5 ± 4.6 g/day, $p < 0.001$), cholesterol (594.8 ± 53.0 g/day vs. 271.6 ± 53.0 g/day, $p < 0.001$), omega-3 (6.5 ± 0.58 g/day vs. 2.7 ± 0.58 g/day, $p < 0.001$) and omega-6 fatty acids (15.6 ± 1.6 g/day vs. 9.7 ± 1.6 g/day, $p = 0.015$). There was a significant decrease in intake of sugar (33.0 ± 5.7 g/day vs. 3.1 ± 5.7 g/day, $p = 0.01$) from pre-baseline to week six, followed by a significant increase in intake of sugar (3.1 ± 5.7 g/day vs. 24.0 ± 5.7 g/day, $p < 0.05$) from week six to week 11.

There was a significant increase in the intake of main carbohydrate sources and a decreased intake of meat and eggs between weeks six and 11, when changing from the ketogenic to the NNR diet.

3.2.2 | Physical activity

There were no significant differences in PAL or number of steps per day over time. These data are presented in Table 4.

3.3 | Main outcome variables

3.3.1 | Pain

Changes in pain overtime can be seen in Table 5. The LCHF-diet induced a significant reduction in pain from baseline to week seven (4.6 ± 0.69 vs. 2.3 ± 0.69 cm, $p = 0.018$). Perceived pain returned to baseline levels at week 13 (4.2 ± 0.69 cm, $p = 0.69$).

TABLE 1 Body weight and composition and changes over time

	Baseline (n = 9)	W7 (n = 9)	W13 (n = 8)	Baseline to W7 Δ 95% CI p-value	Baseline to W13 Δ 95% CI p-value	W7 to W13 Δ 95% CI p-value
BW, kg	102.6 ± 4.1	98.0 ± 4.1	98.5 ± 4.1	Δ-4.6 (3.2, 6.2) <0.001	Δ-4.1 (2.7, 5.6) <0.001	Δ0.5 (-2.0, 0.9) 0.430
(%WL)				(-4.5% ± 2.4)	(-4.0% ± 2.2)	(-0.6% ± 0.9)
BMI, kg/m ²	36.7 ± 1.5	35.1 ± 1.5	35.3 ± 1.5	Δ-1.6 (1.2, 2.1) <0.001	Δ-1.4 (1.0, 1.9) <0.001	Δ0.2 (-0.7, 0.3) 0.350
Waist circumference, cm	98.3 ± 2.7	94.0 ± 2.7	96.0 ± 2.7	Δ-4.3 (2.8, 6.0) <0.001	Δ-2.3 (1.2, 4.4) 0.020	Δ2.0 (-3.2, 0.04) 0.060
Hip circumference, cm	125.2 ± 1.6	123.0 ± 1.6	123.0 ± 1.6	Δ-2.2 (1.3, 3.8) 0.010	Δ-2.2 (1.0, 3.6) 0.010	Δ0.0 (-1.5, 1.0) 0.700
Waist/hip ratio	0.8 ± 0.02	0.8 ± 0.02	0.8 ± 0.02	Δ-0.02 (0.004, 0.04) 0.017	Δ-0.01 (-0.01, 0.03) 0.300	Δ0.01 (-0.03, 0.004) 0.140
Thigh, cm	67.0 ± 3.0	65.0 ± 3.0	65.1 ± 3.0	Δ-2.0 (-0.6, 3.6) 0.200	Δ-1.9 (-0.3, 4.0) 0.080	Δ0.1 (-1.8, 2.5) 0.730
Calf, cm	48.0 ± 3.8	47.0 ± 3.8	47.5 ± 3.8	Δ-1.0 (0.1, 1.7) 0.030	Δ-0.5 (-0.4, 1.2) 0.330	Δ0.5 (-1.3, 0.3) 0.180
FM, kg	47.8 ± 3.1	46.4 ± 3.1	46.3 ± 3.1	Δ-1.4 (-0.5, 3.3) 0.140	Δ-1.5 (-0.5, 3.3) 0.140	Δ-0.1 (-1.9, 2.0) 0.970
FM, %	46.5 ± 1.4	46.8 ± 1.4	46.6 ± 1.4	Δ0.3 (-2.2, 1.6) 0.740	Δ0.1 (-2.0, 1.8) 0.920	Δ-0.2 (-1.7, 2.1) 0.810
FFM, kg	54.3 ± 1.7	51.9 ± 1.7	52.0 ± 1.7	Δ-2.4 (0.03, 4.7) 0.048	Δ-2.3 (-0.04, 4.7) 0.060	Δ0.1 (-2.4, 2.3) 0.950
SMM, kg	30.1 ± 1.0	28.7 ± 1.0	29.0 ± 1.0	Δ-1.4 (0.2, 2.7) 0.024	Δ-0.2 (-0.01, 2.5) 0.051	Δ0.3 (-1.5, 1.0) 0.700
Body water, l	41.0 ± 1.4	38.1 ± 1.4	38.6 ± 1.4	Δ-2.9 (-0.1, 5.0) 0.060	Δ-2.4 (-0.5, 4.6) 0.110	Δ0.5 (-3.0, 2.1) 0.750

Note: Data presented as estimated marginal means ± SEM. Bold is used to highlight the most significant p-values below 0.05.

Abbreviations: BMI, body mass index; BW, body weight; CI, confidence interval; E%, energy percent; FFM, fat free mass; FM, fat mass; SMM, skeletal muscle mass; W, week; Δ, Mean difference between the timepoints.

TABLE 2 Total energy and macronutrients intake and changes over time

	Baseline (n = 9)	W6 (n = 9)	W11 (n = 8)	Baseline to W6 Δ 95% CI p-value	Baseline to W11 Δ 95% CI p-value	W6 to W11 Δ 95% CI p-value
Energy, kcal/day	1927.0 ± 137.0	1863.0 ± 137.0	1878.0 ± 137.0	Δ-63.0 (-336.6, 463.4) 0.75	Δ-48.0 (-351.8, 448.2) 0.81	Δ15.0 (-415.3, 384.8) 0.94
Fat, g/day (E%)	88.0±8.6 (41)	146.0 ± 8.6 (71)	73.0±8.6 (35)	Δ57.0 (-80.7, -33.7) <0.001	Δ-15.0 (-8.2, 38.8) 0.19	Δ-72.0 (49.0, 96.0) <0.001
CHO, g/day (E%)	182.0±21.6 (38)	28.0 ± 21.6 (6)	197.0±21.6 (42)	Δ-154.0 (91.1, 217.1) <0.001	Δ15.0 (-78.4, 47.6) 0.6	Δ170.0 (232.5, 106.6) <0.001
Protein, g/day (E%)	85.0±5.8 (18)	99.0 ± 5.8 (21)	91.0±5.8 (19)	Δ14.0 (-31.3, 2.6) 0.09	Δ7.0 (-23.4, 10.5) 0.44	Δ-8.0 (-9.0, 25.0) 0.34

Note: Data presented as estimated marginal means ± SEM. Bold is used to highlight the most significant p-values below 0.05.

Abbreviations: CHO, carbohydrates; CI, confidence interval; E%, energy percent; W, week; Δ, Mean difference between the timepoints.

TABLE 3 Total daily intake and change of saturated fat, unsaturated fat, cholesterol, sugar, fiber, omega-3 and omega-6 over time

	Baseline (n = 9)	W6 (n = 9)	W11 (n = 8)	Baseline to W6 Δ 95% CI p-value	Baseline to W11 Δ 95% CI p-value	W6 to W11 Δ 95% CI p-value
Saturated fat, g (E%)	32.0 ± 3.6 (14.5)	58.8 ± 3.6 (28.5)	27.5 ± 3.6 (13.4)	Δ26.8 (37.2, 16.4) <0.001	Δ-4. (-5.8, 15.0) 0.370	Δ-31.3 (20.9, 41.7) <0.001
Unsaturated fat, g (E%)	46.5 ± 4.6 (22.0)	67.7 ± 4.6 (32.9)	37.5 ± 4.6 (18.2)	Δ21.2 (34.8, 7.8) 0.030	Δ-9.0 (-4.6, 22.5) 0.180	Δ-30.2 (16.7, 43.7) <0.001
Cholesterol, mg (E%)	325.4 ± 53.0 (0.15)	594.8 ± 53.0 (0.3)	271.6 ± 53.0 (0.13)	Δ269.4 (421.5, 117.3) 0.002	Δ-53.8 (-98.3, 205.9) 0.470	Δ-323.2 (171.1, 475.3) <0.001
Omega-3, g (E%)	2.4 ± 0.6 (0.9)	6.5 ± 0.6 (3.4)	2.7 ± 0.6 (1.4)	Δ4.1 (5.7, 2.5) Π < 0.001	Δ0.3 (-1.9, 1.3) 0.710	Δ-3.8 (2.2, 5.4) <0.001
Omega-6, g (E%)	10.2 ± 1.6 (4.7)	15.6 ± 1.6 (7.7)	9.7 ± 1.6 (4.8)	Δ5.4 (10.0, 0.7) 0.030	Δ-0.5 (-4.1, 5.2) 0.810	Δ-5.9 (1.3, 10.6) 0.020
Sugar, g (E%)	33.0 ± 5.7 (6.9)	3.1 ± 5.7 (0.7)	24.0 ± 5.7 (5.1)	Δ-30.0 (13.2, 46.7) 0.001	Δ-9.0 (-7.7, 25.8) 0.270	Δ20.9 (37.6, 4.1) 0.017
Fiber, g (E%)	25.8 ± 5.0 (5.4)	15.8 ± 5.0 (3.4)	26.4 ± 5.0 (5.5)	Δ-10.0 (-4.7, 24.6) 0.170	Δ0.6 (-15.4, 14.0) 0.930	Δ10.6 (-25.3, 4.1) 0.150

Note: Data presented as estimated marginal means ± SEM. Bold is used to highlight the most significant p-values below 0.05.

Abbreviations: CI, confidence interval; E%, energy percent; W, week; Δ, Mean difference between the timepoints.

TABLE 4 PAL and steps per day at the different timepoints and change over time

	Baseline (n = 9)	W7 (n = 9)	W13 (n = 8)	Baseline to W7 Δ 95% CI p-value	Baseline to W13 Δ 95% CI p-value	W7 to W13 Δ 95% CI p-value
PAL	1.4 ± 0.02	1.4 ± 0.02	1.4 ± 0.02	Δ-0.02 (-0.02, 0.06) 0.3	Δ-0.01 (-0.03, 0.05) 0.6	Δ0.01 (-0.05, 0.03) 0.6
Steps/day	5241.7 ± 465.8	5359.3 ± 465.8	5387.4 ± 465.8	Δ117.7 (-803.0, 567.7) 0.7	Δ145.8 (-831.1, 539.6) 0.7	Δ28.1 (-713.5, 657.3) 0.9

Note: Data presented as estimated marginal means ± SEM.

Abbreviations: CI, confidence interval; E%, energy percent; PAL, physical activity level; W, week; Δ, Mean difference between the timepoints.

TABLE 5 Pain at the different timepoints and changes over time

	Baseline (n = 9)	W7 (n = 9)	W13 (n = 8)	Baseline to W7 Δ 95% CI p-value	Baseline to W13 Δ 95% CI p-value	W7 to W13 Δ 95% CI p-value
VAS	4.6 ± 0.7	2.3 ± 0.7	4.2 ± 0.7	Δ-2.3 (0.4, 4.1) 0.018	Δ-0.4 (-1.5, 2.2) 0.690	Δ1.9 (3.7, 0.1) 0.041

Note: Data presented as estimated marginal means ± SEM. Bold is used to highlight the most significant p-values below 0.05.

Abbreviations: CI, confidence interval; VAS, visual analog scale; W, week; Δ, Mean difference between the timepoints.

3.3.2 | Quality of life

Changes in QoL can be seen in Table 6. There was a significant increase in *general life quality* from baseline to week seven (5.1 ± 0.61 vs. 6.1 ± 0.61 , $p = 0.05$) and week 13 (5.1 ± 0.61 vs. 6.1 ± 0.61 , $p = 0.05$). There was a significant decrease in median score in the QoL subcategory; *body image/appearance* from baseline to week seven (3.1 ± 0.18 vs. 2.7 ± 0.18 , $p = 0.03$). In the subcategory *symptoms*, there was a decrease in score from baseline to week 6 (2.8 ± 0.23 vs. 2.3 ± 0.23 , $p = 0.02$). There were no significant changes in any of the other subcategories of QoL (*function*, *body image*, *symptoms*, *feelings*, *total score*) between any of the time points.

3.3.3 | Markers of glucose, lipid and electrolyte metabolism in the blood

Changes in glucose, lipid and electrolyte markers in the blood can be seen in Table 7. The LCHF-diet induced a significant reduction in triglyceride levels (1.12 ± 0.12 vs. 0.73 ± 0.12 , $p = 0.003$), HbA1c (34.6 ± 1.1 vs. 31.8 ± 1.1 , $p = 0.01$) and ALP (73.4 ± 5.3 vs. 66.7 ± 5.36 , $p = 0.01$) measured in blood. Triglyceride levels then increase from week seven to week 13 (0.73 ± 0.12 vs. 1.1 ± 0.12 , $p = 0.01$), with no significant differences between week 13 and baseline. No significant changes were found in CRP, total cholesterol, HDL- and LDL-cholesterol, glucose, ALAT, GT, natrium or potassium at any timepoint.

3.3.4 | Correlations

No significant correlation was found between WL at week seven and change in perceived pain during the same time period ($r = 0.283$, $p = 0.46$), but there was a strong positive association between WL at week 13 and change in perceived pain ($r = 0.695$, $p = 0.04$).

4 | DISCUSSION

The aim of this pilot study was to investigate if an eucaloric LCHF-diet improved pain and QoL in females with lipedema, and if the diet caused any changes in weight and body composition. The main findings of this study were that a 7 week LCHF-diet induced a significant reduction in pain and WL. No correlation between WL and pain reduction was found, however during the NNR-phase the perceived pain increased, although WL was maintained. There was also an improvement in QoL for body image/appearance and symptoms in the LCHF-period.

The relief of pain with the 7 week LCHF-diet and maintenance of WL during the NNR-phase, with recurrence of pain supports our hypothesis that the pain reduction could be induced by the LCHF-diet itself, and not by the WL.

The mechanisms leading to pain reduction following KDs remain unknown. Whether a KD impacts on pain directly, through an increase in ketone bodies (KB), or indirectly, through changes in macronutrient and micronutrient intake remain to be established and further studies

TABLE 6 QoL scores at the different timepoints and changes over time

	Baseline (n = 9)	W7 (n = 9)	W13 (n = 8)	Baseline to W7 Δ 95% CI p-value	Baseline to W13 Δ 95% CI p-value	W7 to W13 Δ 95% CI p-value
QoL_GL	5.1 ± 0.6	6.1 ± 0.6	6.1 ± 0.6	Δ1.0 (2.0, 0.001) 0.050	Δ1.0 (2.0, 0.001) 0.050	Δ0.0 (-1.0, 1.0) 1.0
QoL_Fu	2.1 ± 14.4	2.0 ± 14.4	2.0 ± 14.4	Δ-0.2 (-0.1, 0.5) 0.230	Δ-0.1 (-0.3, 0.4) 0.790	Δ0.0 (-0.5, 0.2) 0.35
QoL_BI	3.1 ± 0.2	2.7 ± 0.2	2.9 ± 0.2	Δ-0.4 (0.2, 0.7) 0.030	Δ-0.2 (-0.08, 0.5) 0.100	Δ0.2 (-0.5, 0.03) 0.080
QoL_S	2.8 ± 0.2	2.3 ± 0.2	2.5 ± 0.2	Δ-0.5 (0.1, 1.0) 0.020	Δ-0.3 (-0.1, 0.8) 0.100	Δ0.2 (-0.7, 0.3) 0.400
QoL_Fe	1.9 ± 0.2	1.7 ± 0.2	1.7 ± 0.2	Δ-0.2 (-0.2, 0.6) 0.300	Δ-0.2 (-0.2, 0.6) 0.300	Δ0.01 (-0.4, 0.4) 1.000
QoL_T	11.0 ± 1.05	9.9 ± 1.05	10.0 ± 1.05	Δ-1.1 (-0.7, 2.9) 0.200	Δ-1.0 (-0.8, 2.8) 0.300	Δ0.1 (-1.9, 1.7) 0.900

Note: Data presented as estimated marginal means ± SEM. Bold is used to highlight the most significant p-values below 0.05.

Abbreviations: BI, body image; CI, confidence interval; E%, energy percent; Fe, Feelings; Fu, function; GL, general life quality; QoL, quality of life; S, symptoms; T, total; W, week; Δ, Mean difference between the timepoints.

are needed to elucidate this.¹⁸ The pain reduction may also be explained by the anti-inflammatory effects of an LCHF-diet, given the inflammation present in the lipedema fat.^{8,9} Nutrients affects our immune system, and various macronutrients can be either pro- or anti-inflammatory.³⁵ Omega-6, refined carbohydrates (white bread, pasta, and ultra-processed food) and saturated fats are nutrients with pro-inflammatory effects, while omega-3, polyphenols (fruits and vegetables) and whole-grain products have anti-inflammatory properties.³⁶ The intake of omega-3 fatty acids increased during the LCHF-period and decreased in the NNR-phase. Simultaneously the intake of sugar decreased in the LCHF-period. These findings may indicate that a decreased intake of refined carbohydrates, and increased intake of omega-3 fatty acids may be part of the explanation of the reduction in pain in these patients, due to their anti-inflammatory effects. Therefore, the Mediterranean diet could be more appropriate for lipedema patients, also allowing the consumption of more whole-grain products and some fruits.³⁷ This is in line with a recent study by Renzo et al, showing improvements in antioxidant capacity with a Mediterranean diet in patients with lipedema.¹⁹

This is one of the few studies investigating the effect of diet on pain in patients with lipedema, which is the dominating symptom in lipedema and the one with the strongest negative impact on QoL.

Liposuction has been shown to induce a reduction in pain (assessed with VAS), with a concomitant improvement in psychological stress³⁸ In an experimental study by Rapprich et al., 25 patients diagnosed with lipedema were examined before and after liposuction. These patients reported a significant reduction in pain from 7.2 to 2.1 cm on VAS-score, six months after liposuction.³⁸ Liposuction is a highly invasive treatment, not suitable to or affordable by all patients. The participants in our study reported a lower VAS-score initially, and therefore the reduced pain reported after 6 weeks on a LCHF-diet illustrates a great significant and clinical effect on pain.

The KD did induce a small, but significant reduction in BW (-4.6 kg CI (3.2,6.2 kg)). Loss of total body water (TBW), due to depletion of glycogen in the muscles and liver, is likely to account to some of the weight reduction seen (approximately 1.5 to 2 kg).^{39,40} Similarly, the small, although not significant, weight gain during the NNR-phase, is likely due to hydration of the lean tissue, which accompanies glycogen storage. Crescenzi R. et al. found that tissue sodium content is elevated in the skin and SAT in women with lipedema, and that the relative fat-to-water volume in the calf was elevated in lipedema, and skin sodium content was directly correlated with fat-to-water volume.¹ This indicates that the water volume in the legs of women with lipedema may be elevated.

TABLE 7 Markers of glucose, lipid and electrolytes metabolism in the blood at different timepoints and changes over time

	Baseline (n = 9)	W7 (n = 9)	W13 (n = 8)	Baseline to W7 Δ 95% CI p-value	Baseline to W13 Δ 95% CI p-value	W7 to W13 Δ 95% CI p-value
CRP (mg/L)	3.3 ± 0.9	4.0 ± 0.9	2.6 ± 0.9	Δ0.6 (-2.5, 1.2) 0.500	Δ-0.7 (-1.2, 2.5) 0.440	Δ-1.3 (-0.5, 3.1) 0.150
Total cholesterol (mmol/L)	5.0 ± 0.3	4.7 ± 0.3	4.8 ± 0.3	Δ-0.2 (-0.4, 0.8) 0.400	Δ-0.6 (-0.4, 0.8) 0.570	Δ0.08 (-0.7, 0.5) 0.790
HDL-cholesterol (mmol/L)	1.5 ± 0.07	1.5 ± 0.07	1.5 ± 0.07	Δ0.01 (-0.2, 0.1) 0.810	Δ0.003 (-0.2, 0.2) 0.960	Δ-0.01 (-0.1, 0.2) 0.840
LDL-cholesterol (mmol/L)	3.25 ± 0.3	2.9 ± 0.3	3.1 ± 0.3	Δ-0.2 (-0.3, 0.7) 0.320	Δ-0.1 (-0.4, 0.6) 0.660	Δ0.1 (-0.6, 0.4) 0.560
Triglyceride (mmol/L)	1.1 ± 0.1	0.7 ± 0.1	1.1 ± 0.1	Δ-0.4 (0.2, 0.6) 0.003	Δ-0.03 (-0.2, 0.3) 0.800	Δ0.4 (-0.6, -0.1) 0.010
Glucose (mmol/L)	3.2 ± 0.3	2.9 ± 0.3	3.1 ± 0.3	Δ-0.2 (-0.7, 0.3) 0.320	Δ-0.1 (-0.6, 0.4) 0.660	Δ0.1 (-0.4, 0.6) 0.580
HbA1c (mmol/mol)	34.6 ± 1.1	31.8 ± 1.1	33.9 ± 1.1	Δ-2.9 (-5.0, -0.7) 0.010	Δ-0.8 (-2.9, 1.4) 0.470	Δ2.1 (-0.03, 4.2) 0.050
ALAT (U/L)	22.2 ± 3.3	21.0 ± 3.23	17.8 ± 3.3	Δ-1.2 (-9.2, 6.8) 0.750	Δ-4.5 (-12.5, 3.6) 0.260	Δ-3.3 (-11.3, 0.8) 0.400
GT (U/L)	19.1 ± 4.02	18.0 ± 4.02	14.3 ± 4.02	Δ-1.1 (-7.7, 5.4) 0.720	Δ-4.9 (-11.4, 0.66) 0.130	Δ-3.7 (-10.3, 2.8) 0.250
ALP (U/L)	73.4 ± 5.3	66.7 ± 5.4	74.4 ± 5.4	Δ-6.7 (-11.9, -1.6) 0.010	Δ1.0 (-4.2, 6.2) 0.700	Δ7.7 (2.5, 12.9) 0.010
Sodium (mmol/L)	140.3 ± 0.5	140.3 ± 0.5	140.4 ± 0.5	Δ0.08 (-1.2, 1.3) 0.890	Δ0.1 (-1.1, 1.4) 0.830	Δ0.04 (-1.2, 1.3) 0.940
Potassium (mmol/L)	4.1 ± 0.07	4.0 ± 0.07	4.03 ± 0.07	Δ-0.11 (-0.3, 0.05) 0.170	Δ-0.04 (-0.12, 0.1) 0.620	Δ0.07 (-0.09, 0.2) 0.35

Note: Data presented as estimated marginal means ± SEM. Bold is used to highlight the most significant p-values below 0.05.

Abbreviations: ALAT, alanine transaminase; ALP, alkaline phosphatase; CI, confidence interval; CRP, C-reactive protein; GT, gamma-glutamyltransferase; HbA1c, glycated hemoglobin; HDL, high-density lipoprotein; LDL, low-density lipoprotein; W, week; Δ, Mean difference between the timepoints.

The reduction in waist- and hip-circumference being the most significantly compared to thigh- or calf-circumference during LCHF indicate that these were the areas mostly affected by the WL. This is in line with a review article by Herbst et al. (2012) that lipedema fat in the lower extremities of the body in lipedema patients is resistant to WL.³

Child et al. presented the hypothesis that adipocyte proliferation induce hypoxia due to a pressure on the capillaries, resulting in adipocyte necrosis followed by macrophage infiltration and inflammatory responses.² A LCHF-diet may reduce pain by reducing the amount of non-lipidemic fat in the legs, which then relieves the pressure on the blood vessels.⁴¹ The lack of change in thigh circumference in the present study is in contrast to the findings of another KD intervention study conducted in a mixed sample of patients with either lip- and/or lymphedema, which found a reduction in volume of the lower limbs.¹⁵ The diversity in findings could be due to differences in methods of assessment, participant characteristics (sex, age, disease type and stage), and sample size.

The early diagnosis of Lipedema followed by WL and dietary changes may reduce non-lipidemic fat and inflammation. This may also potentially slow down the progression of the condition. However, even with strict diet and exercise regimens, the disease may progress, and further treatment may be necessary.¹¹ People with obesity only need a modest WL of 5%–10% of initial weight to have a significant impact on health.⁹ Therefore, regardless of whether the WL impacts the lipedema symptoms or alter the body composition of the affected limbs, it will have a positive impact on health in general.

The presented findings are in line with the recently published review by Keith et al. The authors summarize that KDs can improve the clinical features of lipedema in three areas: reduction of weight and excessive adipose tissue deposition, pain reduction, and QoL improvement. These improvements are likely to be the result of changes in metabolism and hormonal function, reduction in edema or tissue water content, reduction in inflammation, and fibrosis prevention and/or reduction.¹⁸

This study has several strengths. First, compliance with the diet was excellent. Second, the measurements were performed at the same day and time each week in every participant. Third, the participants were relatively healthy with ongoing treatment, strengthening our findings of pain-reduction. Unfortunately, this study also has some limitations. Most importantly, the small sample size and lack of a control group. Without a control group, one cannot conclude on causality since it is possible that participation in the study by itself had a positive effect. The results should therefore be interpreted with caution. The awareness around lipedema is limited, and so is the number of women diagnosed with this disease. Therefore, the recruitment of participants was challenging, hence the small sample size. No power calculation was performed, and this should be seen as a pilot study. The measurement of circumference should have been performed several times and at several points on both left and right leg, as the affected areas vary. The body composition should preferably be measured with dual-energy x-ray absorptiometry (DEXA). Measurement of pain using the VAS could also have been done weekly, to better capture changes overtime. Finally, KB should

have been measured in blood and plasma, in addition to urine, as that is a more accurate measurement.

There might be an increase in number of women diagnosed with lipedema in the future, as the awareness among health care providers is increasing, and lipedema is underdiagnosed. To identify these patients, a physical examination is important. Therefore, the physical examination of all patients with overweight and obesity should be performed as a regular part of the clinical practice in obesity.

In summary, this study showed that there was a reduction in pain in patients with lipedema after 7 week on a KD. Even though the exact mechanisms mediating this effect remain unknown, several hypotheses have been proposed, namely ketosis per se,³⁵ the anti-inflammatory effects of a LCHF-diet or a potential effect of a reduction in TBW or WL.

5 | CONCLUSION

A LCHF-diet leads to a reduction in perceived pain and improvement in general QoL, independently of WL. Randomized clinical trials are needed to confirm these findings and explore the potential mechanisms involved.

ACKNOWLEDGEMENTS

Thanks to all participants for their time and commitment throughout the study. We have received funding from the Norwegian lymphedema- and lipedema association. And finally, thanks to Turid Follestad for helping with statistical analysis (at the Department of Clinical and Molecular Medicine, NTNU). The Norwegian patient association for lymphedema and lipedema supported the study with 10.000 NOK.

CONFLICT OF INTEREST

The authors declared no conflict of interest.

AUTHOR CONTRIBUTIONS

Siren Nymo and Vilde Sørli designed the study. Vilde Sørli conducted research, analyzed the data and wrote the paper. Åsne Ask Hyldmo was the medical doctor responsible for the requisition of the blood samples, and guided VS in the interpretation of the blood values. Ann Kristin De Soysa assisted with diet design. Catia Martins and Kjetil Retterstøl were co-supervisor of the project. All the authors contributed to the discussion of the results and writing up the manuscript. Siren Nymo had primary responsibility for final content.

ORCID

Siren Nymo  <https://orcid.org/0000-0002-7217-5913>

REFERENCES

1. Crescenzi R, Marton A, Donahue PMC, et al. Tissue sodium content is elevated in the skin and subcutaneous adipose tissue in women with lipedema. *Obes.* 2018;26(2):310-317.
2. Child AH, Gordon KD, Sharpe P, et al. Lipedema: an inherited condition. *Am J Med Genet.* 2010;152A(4):970-976.

3. Herbst KL. Rare adipose disorders (RADs) masquerading as obesity. *Acta Pharmacol Sin.* 2012;33(2):155-172.
4. Wold LE, Hines EA, Jr., Allen EV. Lipedema of the legs; a syndrome characterized by fat legs and edema. *Ann Intern Med.* 1951;34(5):1243-1250.
5. Földi EFM. *Földi's Textbook of Lymphology.* 3rd ed. Elsevier GmbH; 2012:364-369.
6. Herbst K, Mirkovskaya L, Bharhagava A, Chava Y, Te C. Lipedema fat and signs and symptoms of illness, increase with advancing stage. *Archives Med.* 2015;7.
7. Bast JH, Ahmed L, Engdahl R. Lipedema in patients after bariatric surgery. *Surg Obes Relat Dis.* 2016;12(5):1131-1132.
8. Lontok ET. *Lipedema: A Giving Smarter Guide.* Milken Institute; 2017.
9. Suga H, Araki J, Aoi N, Kato H, Higashino T, Yoshimura K. Adipose tissue remodeling in lipedema: adipocyte death and concurrent regeneration. *J Cutan Pathol.* 2009;36(12):1293-1298.
10. Evans S. Lipoedema: the first UK patient survey. *Br J Community Nurs.* 2013;18(4 Suppl):S26-S27.
11. Peled AW, Kappos EA. Lipedema: diagnostic and management challenges. *Int J women's health.* 2016;8:389-395.
12. Langendoen SI, Habbema L, Nijsten TE, Neumann HA. Lipoedema: from clinical presentation to therapy. A review of the literature. *Br J Dermatol.* 2009;161(5):980-986.
13. Seo C. *Lipedema and Keto 2017.* Available from. <https://keto.lipedema-simplified.org>
14. Mark L, Smith CS. *The Lipedema Project 2016.* Available from. <https://lipedemaproject.org/treatment-for-lipedema>
15. Keith L, Rowsemitt C, Richards LG. Lifestyle modification group for lymphedema and obesity results in significant health outcomes. *Am J Lifestyle Med.* 2017;14(4):420-428. <https://doi.org/10.1177/1559827617742108>
16. Kelly T, Unwin D, Finucane F. Low-carbohydrate diets in the management of obesity and type 2 Diabetes: a review from clinicians using the approach in practice. *Int J Environ Res Publ Health.* 2020;17(7).
17. Paoli A, Rubini A, Volek JS, Grimaldi KA. Beyond weight loss: a review of the therapeutic uses of very-low-carbohydrate (ketogenic) diets. *Eur J Clin Nutr.* 2013;67(8):789-796.
18. Keith L, Seo CA, Rowsemitt C, et al. Ketogenic diet as a potential intervention for lipedema. *Med Hypotheses.* 2021;146:110435.
19. Di Renzo L, Cinelli G, Romano L, et al. Potential effects of a modified mediterranean diet on body composition in lipoedema. *Nutrients.* 2021;13(2).
20. Compher C, Frankenfield D, Keim N, Roth-Yousey L, Evidence Analysis Working Group. Best practice methods to apply to measurement of resting metabolic rate in adults: a systematic review. *J Am Diet Assoc.* 2006;106(6):881-903.
21. Helsedirektoratet. *Kostråd for Å Fremme Folkehelsen Og Forebygge Kroniske Sykdommer.* Metodologi Og Vitenskapelig Kunnskapsgrunnlag. Nasjonalt råd for ernæring 2011; 2011.
22. *Kostholdsplanleggeren Oslo Mattilsynet.* Helsedirektoratet; 2014 Available from: <https://www.kostholdsplanleggeren.no>
23. Jakicic JM, Marcus M, Gallagher KI, et al. Evaluation of the Sense-Wear Pro Armband to assess energy expenditure during exercise. *Med Sci Sports Exerc.* 2004;36(5):897-904.
24. Hawker GA, Mian S, Kendzerska T, French M. Measures of adult pain. *American College of Rheumatology.* 2011;63(11):240-252.
25. Ferreira-Valente MA, Pais-Ribeiro JL, Jensen MP. Validity of four pain intensity rating scales. *Pain.* 2011;152(10):2399-2404.
26. Keeley V, Crooks S, Locke J, Veigas D, Riches K, Hilliam R, et al. The development of a condition-specific quality of life measure for lymphoedema (LYMQOL). *Eur J Lymphol.* 2004;12(41):36.
27. Helsedirektoratet. *Kosthåndboken – Veileder I Ernæringsarbeid I Helse-Og Omsorgstjenesten;* 2012:288.
28. Das SK. Body composition measurement in severe obesity. *Curr Opin Clin Nutr Metab Care.* 2005;8(6):602-606.
29. Strain GW, Wang J, Gagner M, Pomp A, Inabnet WB, Heymsfield SB. Bioimpedance for severe obesity: comparing research methods for total body water and resting energy expenditure. *Obes.* 2008;16(8):1953-1956.
30. Volgyi E, Tylavsky FA, Lyytikainen A, Suominen H, Alen M, Cheng S. Assessing body composition with DXA and bioimpedance: effects of obesity, physical activity, and age. *Obes.* 2008;16(3):700-705.
31. Nishida C, Ko GT, Kumanyika S. Body fat distribution and non-communicable diseases in populations: overview of the 2008 WHO expert consultation on waist circumference and waist-hip ratio. *Eur J Clin Nutr.* 2010;64(1):2-5.
32. Molarius A, Seidell JC, Sans S, Tuomilehto J, Kuulasmaa K. Waist and hip circumferences, and waist-hip ratio in 19 populations of the WHO MONICA Project. *Int J Obes Relat Metab Disord.* 1999;23(2):116-125.
33. Bernritter JA, Johnson JL, Woodard SL. Validation of a novel method for measuring waist circumference. *Plast Surg Nurs.* 2011;31(1):9-15.
34. Rosenbaum M, Hall KD, Guo J, et al. Glucose and lipid homeostasis and inflammation in humans following an isocaloric ketogenic diet. *Obes.* 2019;27(6):971-981.
35. Strath LJ, Jones CD, Philip George A, et al. The effect of low-carbohydrate and low-fat diets on pain in individuals with knee osteoarthritis. *Pain Med.* 2020;21(1):150-160.
36. Sears B. Anti-inflammatory diets. *J Am Coll Nutr.* 2015;34(Suppl 1):14-21.
37. Estruch R, Ros E, Salas-Salvado J, et al. Primary prevention of cardiovascular disease with a mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med.* 2018;378(25):e34.
38. Rapprich S, Dingler A, Podda M. Liposuction is an effective treatment for lipedema-results of a study with 25 patients. *J Dtsch Dermatol Ges.* 2011;9(1):33-40.
39. Kreitzman SN, Coxon AY, Szaz KF. Glycogen storage: illusions of easy weight loss, excessive weight regain, and distortions in estimates of body composition. *Am J Clin Nutr.* 1992;56(1 Suppl):292s-3s.
40. Yang MU, Van Itallie TB. Composition of weight lost during short-term weight reduction. Metabolic responses of obese subjects to starvation and low-calorie ketogenic and nonketogenic diets. *J Clin Invest.* 1976;58(3):722-730.
41. Szel E, Kemeny L, Groma G, Szolnoky G. Pathophysiological dilemmas of lipedema. *Med Hypotheses.* 2014;83(5):599-606.

How to cite this article: Sørlie V, De Soysa AK, Hyldmo ÅÅ, Retterstøl K, Martins C, Nymo S. Effect of a ketogenic diet on pain and quality of life in patients with lipedema: the LIPODIET pilot study. *Obes Sci Pract.* 2022;8(4):483-493. <https://doi.org/10.1002/osp4.580>