

Diethyl 4-methoxyoxalyl-3,5-diphenyl-pyrrolidine-2,2-dicarboxylate

Yan-Qin He* and Jin-Quan Chen

Chemical Synthesis and Pollution Control Key Laboratory of Sichuan Province, China
West Normal University, Nanchong 637002, People's Republic of China
Correspondence e-mail: heyanqin123@yahoo.com.cn


Received 28 March 2012; accepted 30 March 2012

Key indicators: single-crystal X-ray study; $T = 290\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$;
 R factor = 0.051; wR factor = 0.133; data-to-parameter ratio = 16.0.

In the title compound, $\text{C}_{25}\text{H}_{27}\text{NO}_7$, the pyrrolidine ring exhibits an envelope conformation and the benzene rings form a dihedral angle of $33.47(11)^\circ$. In the crystal, pairs of $\text{N}-\text{H}\cdots\text{O}$ hydrogen bonds link the molecules into centrosymmetric dimers. Weak $\text{C}-\text{H}\cdots\text{O}$ interactions link the dimers into layers parallel to the bc plane.

Related literature

For applications of pyrrolidine derivatives, see: Shih *et al.* (1995); Enyedy *et al.* (2001); Kravchenko *et al.* (2005); Lack *et al.* (2011).



Experimental

Crystal data

$\text{C}_{25}\text{H}_{27}\text{NO}_7$	$V = 2396.12(14)\text{ \AA}^3$
$M_r = 453.48$	$Z = 4$
Monoclinic, $P2_1/c$	Mo $K\alpha$ radiation
$a = 8.8779(3)\text{ \AA}$	$\mu = 0.09\text{ mm}^{-1}$
$b = 20.3438(6)\text{ \AA}$	$T = 290\text{ K}$
$c = 13.5740(5)\text{ \AA}$	$0.38 \times 0.35 \times 0.30\text{ mm}$
$\beta = 102.213(3)^\circ$	

Data collection

Oxford Diffraction Gemini S Ultra diffractometer	14990 measured reflections
Absorption correction: multi-scan (<i>CrysAlis PRO</i> ; Oxford Diffraction, 2010)	4887 independent reflections
$T_{\min} = 0.966$, $T_{\max} = 0.973$	3129 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.033$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.051$	H atoms treated by a mixture of independent and constrained refinement
$wR(F^2) = 0.133$	$\Delta\rho_{\text{max}} = 0.28\text{ e \AA}^{-3}$
$S = 1.02$	$\Delta\rho_{\text{min}} = -0.26\text{ e \AA}^{-3}$
4887 reflections	8 restraints
305 parameters	

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N \cdots O1 ⁱ	0.88 (2)	2.27 (2)	3.061 (2)	150 (2)
C23—H23 \cdots O3 ⁱⁱ	0.93	2.54	3.461 (3)	170
C25—H25 \cdots O7 ⁱⁱⁱ	0.93	2.60	3.526 (3)	172

Symmetry codes: (i) $-x + 2, -y, -z + 2$; (ii) $-x + 2, y - \frac{1}{2}, -z + \frac{3}{2}$; (iii) $-x + 2, -y, -z + 1$.

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97*.

The authors thank the Centre for Testing and Analysis, Cheng Du Branch Chinese Academy of Sciences, for analytic support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5272).

References

- Enyedy, I. J., Zaman, W. A., Sakamuri, S., Kozikowski, A. P., Johnson, K. M. & Wang, S. (2001). *Bioorg. Med. Chem. Lett.* **11**, 1113–1118.
- Kravchenko, D. V., Kysil, V. M., Tkachenko, S. E., Maliarchouk, S., Okun, I. M. & Ivachtchenko, A. V. (2005). *Eur. J. Med. Chem.* **40**, 1377–1383.
- Lack, N. A., Axerio-Cilies, P., Tavassoli, P., Han, F. Q., Chan, K. H., Feau, C., LeBlanc, E., Guns, E. T., Guy, R. K., Rennie, P. S. & Cherkasov, A. (2011). *J. Med. Chem.* **54**, 8563–8573.
- Oxford Diffraction (2010). *CrysAlis PRO*. Oxford Diffraction Ltd, Yarnton, England.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Shih, N.-Y., Lupo, A. T., Aslanian, R., Orlando, S., Piwinski, J. J., Green, M. J., Ganguly, A. K., Clark, M. A. & Tozzi, S. (1995). *J. Med. Chem.* **38**, 1593–1599.

supplementary materials

Acta Cryst. (2012). E68, o1325 [doi:10.1107/S160053681201392X]

Diethyl 4-methoxyoxalyl-3,5-diphenylpyrrolidine-2,2-dicarboxylate

Yan-Qin He and Jin-Quan Chen

Comment

Pyrrolidine derivatives have been found to be the core of numerous natural products and pharmaceutical candidates, see: Shih *et al.* (1995); Enyedy *et al.* (2001); Kravchenko *et al.* (2005); Lack *et al.* (2011). Herein, we report the crystal structure of the title compound (I).

The bond lengths and angles in (I) are normal. The pyrrolidine ring exhibits an envelope conformation, with the C2 atom occupying the flap position (Fig. 1). Two benzene rings - C12—C17 and C21—C26, respectively - form a dihedral angle of 33.47 (11)°. In the crystal structure, intermolecular N—H···O hydrogen bonds (Table 1) link the molecules into centrosymmetric dimers, and weak C—H···O interactions (Table 1) link further these dimers into layers parallel to *bc* plane.

Experimental


To a flame-dried test tube was added *N,N*-dimethyl-1-[2-(diphenylphosphino) ferrocenyl]ethylamine (0.04 mmol), copper(II) trifluoromethanesulfonate (0.04 mmol), THF (0.5 ml), benzaldehyde (0.24 mmol) and diethyl *A*-amino-malonate (0.24 mmol) under an argon atmosphere. After the mixture was stirred at room temperature for 2 h, a solution of 2-Oxo-4-phenyl-but-3-enoic acid methyl ester (0.2 mmol) in THF (1.0 ml) was added through a syringe in one portion and the resulting mixture was stirred at room temperature until TLC showed no starting material left. Then, solvent was removed and the residue was purified directly by silica gel column chromatography (eluent: petroleum ether / ethyl acetate = 20/1 to 5/1) to yield product. The colourless single crystals suitable for X-ray diffraction were obtained in methanol solvent by slow evaporation.

Refinement

N-bound H atoms were located in a difference Fourier map and refined isotropically [N—H = 0.878 (16) Å]. The C-bound H atoms were placed in calculated positions, with C—H = 0.93 - 0.98 Å, and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2$ or $1.5U_{\text{eq}}(\text{C})$.

Computing details

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO* (Oxford Diffraction, 2010); data reduction: *CrysAlis PRO* (Oxford Diffraction, 2010); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *XP* in *SHELXTL* (Sheldrick, 2008); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

**Figure 1**

The molecular structure of (I) showing 30% probability displacement ellipsoids and the atomic numbering.

Diethyl 4-methoxyxaryl-3,5-diphenylpyrrolidine-2,2-dicarboxylate

Crystal data

$C_{25}H_{27}NO_7$
 $M_r = 453.48$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 8.8779 (3)$ Å
 $b = 20.3438 (6)$ Å
 $c = 13.5740 (5)$ Å
 $\beta = 102.213 (3)^\circ$
 $V = 2396.12 (14)$ Å³
 $Z = 4$

$F(000) = 960$
 $D_x = 1.257 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 4380 reflections
 $\theta = 3.2\text{--}29.1^\circ$
 $\mu = 0.09 \text{ mm}^{-1}$
 $T = 290 \text{ K}$
Block, colorless
 $0.38 \times 0.35 \times 0.30 \text{ mm}$

Data collection

Oxford Diffraction Gemini S Ultra
diffractometer
Radiation source: Enhance (Mo) X-ray Source
Graphite monochromator
Detector resolution: 15.9149 pixels mm⁻¹
 ω scans
Absorption correction: multi-scan
(CrysAlis PRO; Oxford Diffraction, 2010)
 $T_{\min} = 0.966$, $T_{\max} = 0.973$

14990 measured reflections
4887 independent reflections
3129 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.033$
 $\theta_{\text{max}} = 26.4^\circ$, $\theta_{\text{min}} = 3.2^\circ$
 $h = -11 \rightarrow 11$
 $k = -25 \rightarrow 25$
 $l = -16 \rightarrow 14$

*Refinement*Refinement on F^2

Least-squares matrix: full

$$R[F^2 > 2\sigma(F^2)] = 0.051$$

$$wR(F^2) = 0.133$$

$$S = 1.02$$

4887 reflections

305 parameters

8 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0554P)^2 + 0.3662P]$$

where $P = (F_o^2 + 2F_c^2)/3$

$$(\Delta/\sigma)_{\max} < 0.001$$

$$\Delta\rho_{\max} = 0.28 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
N1	0.9482 (2)	0.04749 (8)	0.86894 (15)	0.0445 (4)
O1	0.75874 (18)	0.01366 (7)	1.00477 (12)	0.0587 (4)
O2	0.68505 (17)	0.11814 (7)	1.01209 (11)	0.0537 (4)
O3	0.90888 (18)	0.20277 (8)	0.85551 (13)	0.0627 (5)
O4	0.99367 (18)	0.16006 (7)	1.00924 (12)	0.0606 (5)
O5	0.66663 (2)	0.05129 (8)	0.58463 (13)	0.0646 (5)
O6	0.5961 (2)	-0.10174 (9)	0.66768 (15)	0.0833 (6)
O7	0.6680 (2)	-0.07505 (8)	0.52526 (13)	0.0672 (5)
C2	0.8329 (2)	0.09267 (9)	0.89093 (15)	0.0390 (5)
C3	0.7135 (2)	0.09231 (9)	0.78941 (15)	0.0393 (5)
H3	0.7590	0.1168	0.7408	0.047*
C4	0.7112 (2)	0.01965 (9)	0.75916 (16)	0.0416 (5)
H4	0.6324	-0.0027	0.7877	0.050*
C5	0.8726 (2)	-0.00897 (10)	0.81100 (16)	0.0427 (5)
H5	0.8575	-0.0436	0.8582	0.051*
C6	0.7574 (2)	0.06980 (10)	0.97687 (16)	0.0418 (5)
C7	0.5949 (3)	0.10022 (14)	1.0861 (2)	0.0740 (8)
H7A	0.5254	0.0644	1.0608	0.089*
H7B	0.6625	0.0862	1.1483	0.089*
C8	0.5078 (5)	0.15783 (19)	1.1041 (3)	0.1267 (13)
H8A	0.4510	0.1746	1.0409	0.190*
H8B	0.5773	0.1910	1.1372	0.190*
H8C	0.4374	0.1460	1.1460	0.190*
C9	0.9123 (2)	0.15908 (11)	0.91495 (18)	0.0453 (5)
C10	1.0665 (3)	0.22220 (14)	1.0448 (2)	0.0809 (9)

H10A	1.1018	0.2439	0.9902	0.097*
H10B	1.1555	0.2140	1.0986	0.097*
C11	0.9557 (4)	0.26554 (15)	1.0823 (3)	0.1000 (11)
H11A	0.8684	0.2741	1.0287	0.150*
H11B	1.0053	0.3063	1.1056	0.150*
H11C	0.9219	0.2442	1.1369	0.150*
C12	0.5595 (2)	0.12441 (10)	0.79130 (15)	0.0412 (5)
C13	0.4339 (2)	0.09043 (12)	0.81017 (19)	0.0570 (6)
H13	0.4396	0.0450	0.8180	0.068*
C14	0.3000 (3)	0.12289 (14)	0.8176 (2)	0.0704 (7)
H14	0.2169	0.0991	0.8307	0.084*
C15	0.2887 (3)	0.18935 (14)	0.8060 (2)	0.0683 (7)
H15	0.1988	0.2110	0.8118	0.082*
C16	0.4104 (3)	0.22371 (13)	0.7859 (2)	0.0671 (7)
H16	0.4033	0.2691	0.7776	0.080*
C17	0.5440 (3)	0.19164 (10)	0.77789 (18)	0.0545 (6)
H17	0.6255	0.2157	0.7631	0.065*
C18	0.6739 (2)	0.00889 (10)	0.64679 (17)	0.0459 (5)
C19	0.6410 (3)	-0.06290 (12)	0.61527 (19)	0.0533 (6)
C20	0.6394 (4)	-0.14192 (14)	0.4894 (2)	0.0933 (10)
H20A	0.6611	-0.1457	0.4232	0.140*
H20B	0.5335	-0.1530	0.4866	0.140*
H20C	0.7048	-0.1714	0.5345	0.140*
C21	0.9705 (2)	-0.03601 (10)	0.74167 (17)	0.0466 (5)
C22	0.9944 (3)	-0.10277 (12)	0.7365 (2)	0.0651 (7)
H22	0.9494	-0.1311	0.7757	0.078*
C23	1.0847 (4)	-0.12831 (15)	0.6737 (2)	0.0820 (9)
H23	1.0998	-0.1734	0.6709	0.098*
C24	1.1512 (4)	-0.08695 (18)	0.6163 (2)	0.0861 (9)
H24	1.2127	-0.1040	0.5748	0.103*
C25	1.1283 (3)	-0.02117 (16)	0.6190 (2)	0.0783 (8)
H25	1.1730	0.0067	0.5790	0.094*
C26	1.0380 (3)	0.00470 (12)	0.68185 (19)	0.0611 (6)
H26	1.0228	0.0499	0.6836	0.073*
H1N	1.016 (2)	0.0357 (10)	0.9232 (14)	0.054 (7)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
N1	0.0426 (10)	0.0438 (10)	0.0464 (12)	0.0056 (8)	0.0077 (9)	-0.0022 (9)
O1	0.0719 (11)	0.0459 (9)	0.0627 (11)	0.0056 (7)	0.0240 (8)	0.0149 (8)
O2	0.0648 (10)	0.0496 (9)	0.0533 (10)	0.0054 (7)	0.0277 (8)	0.0016 (7)
O3	0.0745 (11)	0.0482 (9)	0.0640 (11)	-0.0176 (8)	0.0113 (9)	0.0053 (9)
O4	0.0645 (10)	0.0603 (10)	0.0520 (10)	-0.0165 (8)	0.0009 (8)	-0.0057 (8)
O5	0.0861 (12)	0.0569 (10)	0.0495 (10)	0.0043 (8)	0.0110 (9)	0.0064 (8)
O6	0.1233 (17)	0.0567 (11)	0.0763 (14)	-0.0250 (10)	0.0357 (12)	-0.0118 (10)
O7	0.0863 (12)	0.0633 (11)	0.0497 (11)	0.0081 (9)	0.0093 (9)	-0.0136 (9)
C2	0.0406 (11)	0.0356 (10)	0.0417 (12)	-0.0003 (8)	0.0104 (9)	0.0006 (9)
C3	0.0429 (11)	0.0374 (11)	0.0384 (12)	-0.0001 (8)	0.0106 (9)	0.0025 (9)
C4	0.0456 (12)	0.0371 (11)	0.0438 (13)	-0.0015 (8)	0.0131 (9)	0.0003 (9)

C5	0.0505 (12)	0.0372 (11)	0.0418 (12)	0.0025 (9)	0.0131 (10)	0.0036 (9)
C6	0.0425 (11)	0.0418 (12)	0.0402 (12)	0.0015 (9)	0.0066 (9)	0.0012 (10)
C7	0.0804 (18)	0.0894 (19)	0.0637 (18)	0.0056 (14)	0.0409 (15)	0.0081 (15)
C8	0.142 (2)	0.132 (2)	0.129 (2)	0.0343 (16)	0.0808 (17)	-0.0039 (16)
C9	0.0423 (12)	0.0439 (12)	0.0519 (15)	-0.0020 (9)	0.0152 (10)	-0.0022 (11)
C10	0.087 (2)	0.0787 (19)	0.072 (2)	-0.0357 (16)	0.0045 (16)	-0.0169 (16)
C11	0.141 (3)	0.072 (2)	0.085 (2)	-0.012 (2)	0.019 (2)	-0.0180 (18)
C12	0.0449 (12)	0.0424 (11)	0.0368 (12)	0.0007 (9)	0.0094 (9)	0.0013 (9)
C13	0.0479 (13)	0.0519 (13)	0.0731 (18)	-0.0032 (10)	0.0171 (12)	-0.0031 (12)
C14	0.0472 (14)	0.0823 (19)	0.085 (2)	-0.0060 (13)	0.0218 (13)	-0.0096 (16)
C15	0.0543 (15)	0.085 (2)	0.0671 (18)	0.0218 (13)	0.0154 (13)	-0.0022 (15)
C16	0.0718 (17)	0.0548 (15)	0.0766 (19)	0.0199 (13)	0.0201 (14)	0.0073 (13)
C17	0.0546 (14)	0.0469 (13)	0.0642 (16)	0.0049 (10)	0.0171 (12)	0.0077 (11)
C18	0.0439 (12)	0.0464 (12)	0.0471 (13)	0.0020 (9)	0.0087 (10)	-0.0010 (11)
C19	0.0553 (14)	0.0530 (14)	0.0491 (15)	0.0012 (11)	0.0056 (11)	-0.0052 (12)
C20	0.131 (3)	0.0688 (18)	0.072 (2)	0.0200 (17)	0.0035 (19)	-0.0275 (16)
C21	0.0476 (12)	0.0470 (13)	0.0442 (13)	0.0067 (9)	0.0072 (10)	-0.0009 (10)
C22	0.0812 (18)	0.0505 (14)	0.0649 (17)	0.0158 (12)	0.0183 (14)	-0.0015 (12)
C23	0.102 (2)	0.0648 (18)	0.080 (2)	0.0265 (16)	0.0218 (18)	-0.0160 (16)
C24	0.083 (2)	0.103 (2)	0.077 (2)	0.0192 (18)	0.0290 (17)	-0.025 (2)
C25	0.0768 (19)	0.100 (2)	0.0669 (19)	-0.0079 (16)	0.0358 (15)	-0.0105 (17)
C26	0.0670 (16)	0.0576 (15)	0.0631 (17)	-0.0018 (12)	0.0236 (13)	-0.0036 (13)

Geometric parameters (\AA , $^\circ$)

N1—C2	1.453 (2)	C10—H10A	0.9700
N1—C5	1.472 (3)	C10—H10B	0.9700
N1—H1N	0.878 (16)	C11—H11A	0.9600
O1—C6	1.203 (2)	C11—H11B	0.9600
O2—C6	1.318 (2)	C11—H11C	0.9600
O2—C7	1.457 (3)	C12—C13	1.381 (3)
O3—C9	1.196 (3)	C12—C17	1.383 (3)
O4—C9	1.330 (3)	C13—C14	1.382 (3)
O4—C10	1.455 (3)	C13—H13	0.9300
O5—C18	1.198 (3)	C14—C15	1.362 (4)
O6—C19	1.187 (3)	C14—H14	0.9300
O7—C19	1.317 (3)	C15—C16	1.362 (4)
O7—C20	1.449 (3)	C15—H15	0.9300
C2—C9	1.527 (3)	C16—C17	1.378 (3)
C2—C6	1.536 (3)	C16—H16	0.9300
C2—C3	1.551 (3)	C17—H17	0.9300
C3—C12	1.520 (3)	C18—C19	1.533 (3)
C3—C4	1.533 (3)	C20—H20A	0.9600
C3—H3	0.9800	C20—H20B	0.9600
C4—C18	1.507 (3)	C20—H20C	0.9600
C4—C5	1.569 (3)	C21—C22	1.379 (3)
C4—H4	0.9800	C21—C26	1.382 (3)
C5—C21	1.513 (3)	C22—C23	1.388 (4)
C5—H5	0.9800	C22—H22	0.9300
C7—C8	1.453 (4)	C23—C24	1.364 (4)

C7—H7A	0.9700	C23—H23	0.9300
C7—H7B	0.9700	C24—C25	1.355 (4)
C8—H8A	0.9600	C24—H24	0.9300
C8—H8B	0.9600	C25—C26	1.392 (4)
C8—H8C	0.9600	C25—H25	0.9300
C10—C11	1.489 (4)	C26—H26	0.9300
C2—N1—C5	110.00 (16)	C10—C11—H11A	109.5
C2—N1—H1N	112.7 (15)	C10—C11—H11B	109.5
C5—N1—H1N	112.8 (14)	H11A—C11—H11B	109.5
C6—O2—C7	116.37 (18)	C10—C11—H11C	109.5
C9—O4—C10	116.42 (19)	H11A—C11—H11C	109.5
C19—O7—C20	115.8 (2)	H11B—C11—H11C	109.5
N1—C2—C9	106.88 (16)	C13—C12—C17	117.3 (2)
N1—C2—C6	113.85 (16)	C13—C12—C3	123.52 (18)
C9—C2—C6	111.21 (17)	C17—C12—C3	119.14 (18)
N1—C2—C3	101.17 (16)	C12—C13—C14	121.0 (2)
C9—C2—C3	113.06 (16)	C12—C13—H13	119.5
C6—C2—C3	110.31 (15)	C14—C13—H13	119.5
C12—C3—C4	117.24 (16)	C15—C14—C13	120.6 (2)
C12—C3—C2	115.56 (17)	C15—C14—H14	119.7
C4—C3—C2	102.13 (15)	C13—C14—H14	119.7
C12—C3—H3	107.1	C16—C15—C14	119.4 (2)
C4—C3—H3	107.1	C16—C15—H15	120.3
C2—C3—H3	107.1	C14—C15—H15	120.3
C18—C4—C3	113.50 (17)	C15—C16—C17	120.4 (2)
C18—C4—C5	112.60 (17)	C15—C16—H16	119.8
C3—C4—C5	106.19 (15)	C17—C16—H16	119.8
C18—C4—H4	108.1	C16—C17—C12	121.4 (2)
C3—C4—H4	108.1	C16—C17—H17	119.3
C5—C4—H4	108.1	C12—C17—H17	119.3
N1—C5—C21	111.04 (17)	O5—C18—C4	125.16 (19)
N1—C5—C4	102.72 (15)	O5—C18—C19	120.7 (2)
C21—C5—C4	116.51 (17)	C4—C18—C19	114.14 (19)
N1—C5—H5	108.8	O6—C19—O7	125.5 (2)
C21—C5—H5	108.8	O6—C19—C18	122.4 (2)
C4—C5—H5	108.8	O7—C19—C18	112.1 (2)
O1—C6—O2	124.6 (2)	O7—C20—H20A	109.5
O1—C6—C2	123.36 (19)	O7—C20—H20B	109.5
O2—C6—C2	111.96 (17)	H20A—C20—H20B	109.5
C8—C7—O2	107.7 (2)	O7—C20—H20C	109.5
C8—C7—H7A	110.2	H20A—C20—H20C	109.5
O2—C7—H7A	110.2	H20B—C20—H20C	109.5
C8—C7—H7B	110.2	C22—C21—C26	118.0 (2)
O2—C7—H7B	110.2	C22—C21—C5	120.3 (2)
H7A—C7—H7B	108.5	C26—C21—C5	121.66 (19)
C7—C8—H8A	109.5	C21—C22—C23	121.0 (3)
C7—C8—H8B	109.5	C21—C22—H22	119.5
H8A—C8—H8B	109.5	C23—C22—H22	119.5

C7—C8—H8C	109.5	C24—C23—C22	119.7 (3)
H8A—C8—H8C	109.5	C24—C23—H23	120.1
H8B—C8—H8C	109.5	C22—C23—H23	120.1
O3—C9—O4	124.67 (19)	C25—C24—C23	120.5 (3)
O3—C9—C2	124.7 (2)	C25—C24—H24	119.7
O4—C9—C2	110.54 (18)	C23—C24—H24	119.7
O4—C10—C11	110.4 (2)	C24—C25—C26	120.0 (3)
O4—C10—H10A	109.6	C24—C25—H25	120.0
C11—C10—H10A	109.6	C26—C25—H25	120.0
O4—C10—H10B	109.6	C21—C26—C25	120.7 (2)
C11—C10—H10B	109.6	C21—C26—H26	119.6
H10A—C10—H10B	108.1	C25—C26—H26	119.6
C5—N1—C2—C9	-159.71 (17)	C9—O4—C10—C11	84.5 (3)
C5—N1—C2—C6	77.1 (2)	C4—C3—C12—C13	28.5 (3)
C5—N1—C2—C3	-41.2 (2)	C2—C3—C12—C13	-92.0 (2)
N1—C2—C3—C12	168.35 (16)	C4—C3—C12—C17	-154.38 (19)
C9—C2—C3—C12	-77.7 (2)	C2—C3—C12—C17	85.1 (2)
C6—C2—C3—C12	47.5 (2)	C17—C12—C13—C14	-1.5 (4)
N1—C2—C3—C4	39.90 (18)	C3—C12—C13—C14	175.6 (2)
C9—C2—C3—C4	153.85 (16)	C12—C13—C14—C15	0.3 (4)
C6—C2—C3—C4	-80.93 (19)	C13—C14—C15—C16	0.6 (4)
C12—C3—C4—C18	82.3 (2)	C14—C15—C16—C17	-0.3 (4)
C2—C3—C4—C18	-150.29 (17)	C15—C16—C17—C12	-1.0 (4)
C12—C3—C4—C5	-153.43 (17)	C13—C12—C17—C16	1.9 (4)
C2—C3—C4—C5	-26.1 (2)	C3—C12—C17—C16	-175.4 (2)
C2—N1—C5—C21	149.70 (17)	C3—C4—C18—O5	9.5 (3)
C2—N1—C5—C4	24.5 (2)	C5—C4—C18—O5	-111.1 (2)
C18—C4—C5—N1	127.43 (18)	C3—C4—C18—C19	-169.00 (17)
C3—C4—C5—N1	2.6 (2)	C5—C4—C18—C19	70.3 (2)
C18—C4—C5—C21	5.8 (2)	C20—O7—C19—O6	-0.1 (4)
C3—C4—C5—C21	-118.94 (19)	C20—O7—C19—C18	-179.9 (2)
C7—O2—C6—O1	-3.3 (3)	O5—C18—C19—O6	-153.7 (2)
C7—O2—C6—C2	173.36 (18)	C4—C18—C19—O6	24.9 (3)
N1—C2—C6—O1	-19.7 (3)	O5—C18—C19—O7	26.1 (3)
C9—C2—C6—O1	-140.5 (2)	C4—C18—C19—O7	-155.34 (19)
C3—C2—C6—O1	93.2 (2)	N1—C5—C21—C22	133.7 (2)
N1—C2—C6—O2	163.56 (16)	C4—C5—C21—C22	-109.2 (2)
C9—C2—C6—O2	42.8 (2)	N1—C5—C21—C26	-46.4 (3)
C3—C2—C6—O2	-83.5 (2)	C4—C5—C21—C26	70.7 (3)
C6—O2—C7—C8	-171.9 (3)	C26—C21—C22—C23	0.5 (4)
C10—O4—C9—O3	8.3 (3)	C5—C21—C22—C23	-179.6 (2)
C10—O4—C9—C2	-175.66 (19)	C21—C22—C23—C24	0.1 (4)
N1—C2—C9—O3	96.2 (2)	C22—C23—C24—C25	-0.8 (5)
C6—C2—C9—O3	-139.0 (2)	C23—C24—C25—C26	0.8 (5)
C3—C2—C9—O3	-14.3 (3)	C22—C21—C26—C25	-0.5 (4)
N1—C2—C9—O4	-79.9 (2)	C5—C21—C26—C25	179.6 (2)
C6—C2—C9—O4	45.0 (2)	C24—C25—C26—C21	-0.1 (4)
C3—C2—C9—O4	169.70 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1—H1N \cdots O1 ⁱ	0.88 (2)	2.27 (2)	3.061 (2)	150 (2)
C23—H23 \cdots O3 ⁱⁱ	0.93	2.54	3.461 (3)	170
C25—H25 \cdots O7 ⁱⁱⁱ	0.93	2.60	3.526 (3)	172

Symmetry codes: (i) $-x+2, -y, -z+2$; (ii) $-x+2, y-1/2, -z+3/2$; (iii) $-x+2, -y, -z+1$.