

2,4-Bis(2-bromophenyl)-7-*tert*-pentyl-3-azabicyclo[3.3.1]nonan-9-one

Dong Ho Park,^a V. Ramkumar^b and P. Parthiban^{a*}

^aDepartment of Biomedicinal Chemistry, Inje University, Gimhae, Gyeongnam 621 749, Republic of Korea, and ^bDepartment of Chemistry, IIT Madras, Chennai 600 036, TamilNadu, India

Correspondence e-mail: parthisivam@yahoo.co.in


Received 9 September 2012; accepted 13 September 2012

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.008\text{ \AA}$; R factor = 0.062; wR factor = 0.212; data-to-parameter ratio = 23.2.

The title compound, $\text{C}_{25}\text{H}_{29}\text{Br}_2\text{NO}$, is a *tert*-pentyl analog of 2,4-bis(2-bromophenyl)-3-azabicyclo[3.3.1]nonan-9-one [Parthiban *et al.* (2008)]. Similar to its analog, the title compound exists in a twin-chair conformation with an equatorial orientation of the 2-bromophenyl groups. The benzene rings are inclined to each other at a dihedral angle of $29.6(3)^\circ$. The *tert*-pentyl group on the cyclohexanone ring also adopts an exocyclic equatorial disposition.

Related literature

For the synthesis, stereochemistry and biological activity of 3-azabicyclo[3.3.1]nonan-9-ones, see: Park *et al.* (2011, 2012a). For the crystal structure of closely related compound, see: Parthiban *et al.* (2008). For examples of azabicycles with different conformations, see: Parthiban *et al.* (2010); Park *et al.* (2012b); Padegimas & Kovacic (1972).



Experimental

Crystal data

$\text{C}_{25}\text{H}_{29}\text{Br}_2\text{NO}$	$\gamma = 97.112(4)^\circ$
$M_r = 519.31$	$V = 1151.08(18)\text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 7.7342(7)\text{ \AA}$	Mo $K\alpha$ radiation
$b = 10.6409(10)\text{ \AA}$	$\mu = 3.54\text{ mm}^{-1}$
$c = 15.0924(12)\text{ \AA}$	$T = 298\text{ K}$
$\alpha = 105.856(4)^\circ$	$0.18 \times 0.15 \times 0.10\text{ mm}$
$\beta = 101.242(4)^\circ$	

Data collection

Bruker APEXII CCD area-detector diffractometer	16189 measured reflections
Absorption correction: multi-scan (<i>SADABS</i> ; Bruker, 2004)	6006 independent reflections
$T_{\min} = 0.569$, $T_{\max} = 0.719$	3081 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.062$	1 restraint
$wR(F^2) = 0.212$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 1.24\text{ e \AA}^{-3}$
6006 reflections	$\Delta\rho_{\text{min}} = -1.26\text{ e \AA}^{-3}$
259 parameters	

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97*.

This research was supported by the Brain Korea 21 (BK 21) Research Program (a human resource development program of the Korean government). The authors acknowledge the Department of Chemistry, IIT Madras, for the X-ray data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: CV5341).

References

- Bruker (2004). *APEX2, XPREP, SAINT-Plus* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Padegimas, S. J. & Kovacic, P. (1972). *J. Org. Chem.* **37**, 2672–2676.
- Park, D. H., Jeong, Y. T. & Parthiban, P. (2011). *J. Mol. Struct.* **1005**, 31–44.
- Park, D. H., Ramkumar, V. & Parthiban, P. (2012b). *Acta Cryst. E68*, o1481.
- Park, D. H., Venkatesan, J., Kim, S. K. & Parthiban, P. (2012a). *Bioorg. Med. Chem. Lett.* **22**, 6004–6009.
- Parthiban, P., Ramkumar, V. & Jeong, Y. T. (2010). *Acta Cryst. E66*, o194–o195.
- Parthiban, P., Ramkumar, V., Kim, M. S., Son, S. M. & Jeong, Y. T. (2008). *Acta Cryst. E64*, o2385.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o2946 [doi:10.1107/S1600536812039128]

2,4-Bis(2-bromophenyl)-7-*tert*-pentyl-3-azabicyclo[3.3.1]nonan-9-one

Dong Ho Park, V. Ramkumar and P. Parthiban

Comment

The bicyclic compounds posses three major conformations, *viz.*, chair-chair (Parthiban *et al.*, 2010), chair-boat (Park *et al.*, 2012*b*) and boat-boat (Padegimas & Kovacic, 1972), depending upon the nature and position of the substituents on the bicycle. The aim of the present study was to explore the stereochemistry as well as the impact of *tert*-pentyl on the twin-chair conformation of the 2,4-bis(2-bromophenyl)-3-azabicyclo[3.3.1]nonan-9-one (Parthiban *et al.*, 2008).

Examination of the asymmery parameters and torsion angles of the title compound (Fig. 1) reveals that the values are very similar to those in its analog. In detail, the torsion angles of the title compound C2—C8—C6—C7, C1—C2—C8—C6, C5—C6—C8—C2 and C3—C2—C8—C6 are -61.2 (5), 61.9 (5), 63.4 (5) and -63.8 (5) $^{\circ}$, respectively. These clearly assign slightly distorted chair conformation to both six-membered rings, and the cyclohexanone ring is comparatively flattened. Furthermore, the orientation of the bromophenyl groups on both sides of the secondary amino group is identified by their torsion angles. The torsion angle of C9—C1—C2—C8 and C8—C6—C7—C15 are 178.0 (4) and -178.6 (4) $^{\circ}$, respectively. This clearly conform their euatorial orientations, and it is similar to its non-*tert*-pentyl analog [C9—C1—C2—C8 and C8—C6—C7—C15 are 177.8 (4) and -179.4 (6) $^{\circ}$, respectively]. Also the orientation of *tert*-pentyl group on the cyclohexanone ring is identified by its torsion angles [C21—C4—C5—C6 and C2—C3—C4—C21 are 171.4 (4) and -172.2 (4) $^{\circ}$, respectively]. In addition to the above similarities, the title compound and its analog's benzene rings orientations are very similar. In titile compound, the benzene rings are inclined to echh other with an angle of 29.6 (3) $^{\circ}$; it is 29.1 $^{\circ}$ for its analog. Hence, the title compound, C₂₅H₂₉Br₂NO, exists in a twin-chair conformation with an euatorial orientation of the *ortho*-bromophenyl groups as its non-*tert*-pentyl analog. The *tert*-pentyl group on the cyclohexnone also adopts an exocyclic equitorial disposition.

Experimental

The 2,4-bis(2-bromophenyl)-7-(*tert*-pentyl) -3-azabicyclo[3.3.1]nonan-9-one was synthesized by a modified and an optimized Mannich condensation in one-pot, using 2-bromobenzaldehyde (0.1 mol, 18.50 g/11.58 ml), 4-*tert*-pentylcyclohexanone (0.05 mol, 8.41 g/9.15 ml) and ammonium acetate (0.075 mol, 5.78 g) in a 50 ml of absolute ethanol (Park *et al.*, 2011). The mixture was gently warmed on a hot plate at 303–308 K (30–35 $^{\circ}$ C) with moderate stirring till the complete consumption of the starting materials, which was monitored by TLC. At the end, the crude azabicyclic ketone was separated by filtration and gently washed with 1:5 cold ethanol-ether mixture. X-ray diffraction quality crystals of the title compound were obtained by slow evaporation from ethanol.

Refinement

All hydrogen atoms were fixed geometrically and allowed to ride on the parent carbon atoms with aromatic C—H = 0.93 Å, aliphatic C—H = 0.98 Å, methylene C—H = 0.97 Å. The displacement parameters were set for phenyl, methylene and aliphatic H atoms at $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$, methyl H atoms at $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C})$ and the hydrogen atoms were fixed

geometrically and allowed to ride on the parent nitrogen atom with N—H = 0.86 Å and the displacement parameter was set at $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{N})$.

Computing details

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT-Plus* (Bruker, 2004); data reduction: *SAINT-Plus* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).




Figure 1

The molecular structure of the title compound showing the atomic numbering and displacement ellipsoids drawn at the 30% probability level.

2,4-Bis(2-bromophenyl)-7-tert-pentyl-3-azabicyclo[3.3.1]nonan-9-one

Crystal data

$\text{C}_{25}\text{H}_{29}\text{Br}_2\text{NO}$
 $M_r = 519.31$
Triclinic, $P\bar{1}$
Hall symbol: -P 1
 $a = 7.7342 (7)$ Å
 $b = 10.6409 (10)$ Å
 $c = 15.0924 (12)$ Å
 $\alpha = 105.856 (4)^\circ$
 $\beta = 101.242 (4)^\circ$
 $\gamma = 97.112 (4)^\circ$
 $V = 1151.08 (18)$ Å³

$Z = 2$
 $F(000) = 528$
 $D_x = 1.498 \text{ Mg m}^{-3}$
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 4160 reflections
 $\theta = 2.7\text{--}22.8^\circ$
 $\mu = 3.54 \text{ mm}^{-1}$
 $T = 298 \text{ K}$
Prism, colourless
 $0.18 \times 0.15 \times 0.10 \text{ mm}$

Data collection

Bruker APEXII CCD area-detector
diffractometer
Radiation source: fine-focus sealed tube
Graphite monochromator
phi and ω scans
Absorption correction: multi-scan
(*SADABS*; Bruker, 2004)
 $T_{\min} = 0.569$, $T_{\max} = 0.719$

16189 measured reflections
6006 independent reflections
3081 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$
 $\theta_{\max} = 29.5^\circ$, $\theta_{\min} = 1.4^\circ$
 $h = -10 \rightarrow 10$
 $k = -14 \rightarrow 14$
 $l = -20 \rightarrow 19$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.062$
 $wR(F^2) = 0.212$
 $S = 1.02$
6006 reflections
259 parameters
1 restraint
Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map
Hydrogen site location: inferred from
neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.117P)^2]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 1.24 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -1.26 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.86819 (8)	0.90205 (5)	0.32088 (5)	0.0717 (3)
Br2	0.85323 (8)	0.28410 (7)	0.51276 (4)	0.0732 (3)
C1	0.6565 (6)	0.6164 (4)	0.3068 (3)	0.0411 (10)
H1	0.7614	0.6679	0.3573	0.049*
C2	0.7228 (6)	0.5383 (4)	0.2223 (3)	0.0440 (11)
H2	0.8022	0.6010	0.2042	0.053*
C3	0.5766 (7)	0.4547 (4)	0.1333 (3)	0.0486 (12)
H3A	0.4990	0.5124	0.1143	0.058*
H3B	0.6342	0.4219	0.0822	0.058*
C4	0.4601 (7)	0.3360 (4)	0.1455 (3)	0.0456 (11)
H4	0.3820	0.3730	0.1857	0.055*
C5	0.5749 (7)	0.2611 (4)	0.1995 (3)	0.0474 (12)
H5A	0.6321	0.2059	0.1558	0.057*

H5B	0.4959	0.2021	0.2195	0.057*
C6	0.7214 (6)	0.3469 (4)	0.2870 (3)	0.0441 (11)
H6	0.7977	0.2894	0.3093	0.053*
C7	0.6514 (6)	0.4244 (4)	0.3698 (3)	0.0408 (10)
H7	0.7552	0.4716	0.4223	0.049*
C8	0.8320 (7)	0.4438 (4)	0.2566 (3)	0.0462 (11)
C9	0.5400 (6)	0.7115 (4)	0.2822 (3)	0.0405 (10)
C10	0.6147 (7)	0.8408 (4)	0.2862 (3)	0.0461 (11)
C11	0.5088 (10)	0.9304 (5)	0.2667 (4)	0.0670 (16)
H11	0.5618	1.0158	0.2708	0.080*
C12	0.3248 (9)	0.8927 (6)	0.2414 (5)	0.0755 (18)
H12	0.2527	0.9530	0.2298	0.091*
C13	0.2472 (8)	0.7627 (6)	0.2332 (4)	0.0676 (16)
H13	0.1235	0.7341	0.2130	0.081*
C14	0.3574 (7)	0.6776 (5)	0.2555 (4)	0.0531 (13)
H14	0.3042	0.5925	0.2521	0.064*
C15	0.5335 (6)	0.3345 (4)	0.4043 (3)	0.0384 (10)
C16	0.3459 (6)	0.3148 (5)	0.3759 (3)	0.0483 (11)
H16	0.2941	0.3628	0.3382	0.058*
C17	0.2358 (8)	0.2263 (6)	0.4020 (4)	0.0596 (14)
H17	0.1119	0.2140	0.3809	0.072*
C18	0.3103 (8)	0.1554 (5)	0.4598 (4)	0.0586 (14)
H18	0.2364	0.0954	0.4773	0.070*
C19	0.4926 (8)	0.1744 (4)	0.4908 (3)	0.0509 (12)
H19	0.5431	0.1276	0.5298	0.061*
C20	0.6015 (6)	0.2631 (4)	0.4641 (3)	0.0415 (10)
N1	0.5541 (5)	0.5240 (3)	0.3436 (3)	0.0413 (9)
H1A	0.4463	0.5282	0.3493	0.050*
O1	0.9910 (5)	0.4487 (4)	0.2582 (3)	0.0695 (10)
C21	0.3326 (9)	0.2448 (5)	0.0486 (4)	0.0732 (18)
C24	0.2293 (15)	0.1222 (10)	0.0634 (7)	0.155 (4)
H24A	0.3131	0.0784	0.0950	0.186*
H24B	0.1656	0.0598	0.0023	0.186*
C22	0.1993 (13)	0.3238 (9)	0.0052 (6)	0.151 (5)
H22A	0.1380	0.2718	-0.0585	0.226*
H22B	0.2649	0.4062	0.0046	0.226*
H22C	0.1133	0.3417	0.0429	0.226*
C25	0.0936 (15)	0.1639 (10)	0.1242 (7)	0.155 (4)
H25A	0.1470	0.2459	0.1736	0.232*
H25B	0.0624	0.0959	0.1521	0.232*
H25C	-0.0126	0.1758	0.0847	0.232*
C23	0.4459 (12)	0.1841 (8)	-0.0222 (5)	0.113 (3)
H23A	0.5272	0.1372	0.0065	0.169*
H23B	0.5130	0.2542	-0.0373	0.169*
H23C	0.3671	0.1237	-0.0792	0.169*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.0725 (4)	0.0449 (3)	0.0923 (5)	-0.0143 (3)	0.0195 (3)	0.0238 (3)

Br2	0.0604 (4)	0.0956 (5)	0.0726 (5)	0.0205 (3)	0.0054 (3)	0.0452 (4)
C1	0.045 (3)	0.026 (2)	0.051 (3)	-0.0006 (18)	0.012 (2)	0.0140 (18)
C2	0.047 (3)	0.035 (2)	0.056 (3)	0.0005 (19)	0.024 (2)	0.021 (2)
C3	0.067 (3)	0.035 (2)	0.049 (3)	0.005 (2)	0.021 (2)	0.019 (2)
C4	0.060 (3)	0.034 (2)	0.045 (3)	0.002 (2)	0.015 (2)	0.018 (2)
C5	0.070 (3)	0.028 (2)	0.050 (3)	0.007 (2)	0.021 (2)	0.0159 (19)
C6	0.049 (3)	0.039 (2)	0.054 (3)	0.013 (2)	0.017 (2)	0.023 (2)
C7	0.050 (3)	0.034 (2)	0.040 (3)	0.0032 (19)	0.011 (2)	0.0167 (19)
C8	0.047 (3)	0.041 (3)	0.053 (3)	0.009 (2)	0.020 (2)	0.014 (2)
C9	0.054 (3)	0.028 (2)	0.044 (3)	0.0075 (19)	0.020 (2)	0.0130 (18)
C10	0.068 (3)	0.031 (2)	0.045 (3)	0.008 (2)	0.024 (2)	0.0141 (19)
C11	0.111 (5)	0.039 (3)	0.066 (4)	0.019 (3)	0.037 (3)	0.027 (3)
C12	0.080 (4)	0.077 (4)	0.103 (5)	0.043 (4)	0.042 (4)	0.055 (4)
C13	0.061 (4)	0.079 (4)	0.080 (4)	0.027 (3)	0.028 (3)	0.040 (3)
C14	0.053 (3)	0.041 (3)	0.072 (4)	0.007 (2)	0.021 (3)	0.025 (2)
C15	0.045 (3)	0.031 (2)	0.040 (3)	0.0051 (18)	0.013 (2)	0.0095 (18)
C16	0.043 (3)	0.044 (3)	0.063 (3)	0.007 (2)	0.015 (2)	0.024 (2)
C17	0.050 (3)	0.064 (3)	0.064 (3)	-0.003 (3)	0.014 (3)	0.025 (3)
C18	0.075 (4)	0.046 (3)	0.055 (3)	-0.007 (3)	0.025 (3)	0.018 (2)
C19	0.076 (4)	0.039 (3)	0.043 (3)	0.008 (2)	0.018 (2)	0.019 (2)
C20	0.053 (3)	0.036 (2)	0.041 (3)	0.014 (2)	0.016 (2)	0.0135 (19)
N1	0.046 (2)	0.0310 (18)	0.057 (2)	0.0084 (16)	0.0239 (18)	0.0205 (17)
O1	0.050 (2)	0.078 (3)	0.096 (3)	0.019 (2)	0.033 (2)	0.039 (2)
C21	0.095 (5)	0.049 (3)	0.062 (4)	-0.017 (3)	0.001 (3)	0.021 (3)
C24	0.197 (9)	0.131 (6)	0.100 (5)	-0.054 (6)	0.005 (4)	0.034 (5)
C22	0.179 (9)	0.106 (6)	0.120 (7)	-0.064 (6)	-0.080 (6)	0.078 (5)
C25	0.197 (9)	0.131 (6)	0.100 (5)	-0.054 (6)	0.005 (4)	0.034 (5)
C23	0.160 (8)	0.107 (6)	0.041 (4)	-0.016 (5)	0.017 (4)	-0.004 (3)

Geometric parameters (\AA , $\text{^{\circ}}$)

Br1—C10	1.908 (5)	C12—H12	0.9300
Br2—C20	1.904 (5)	C13—C14	1.380 (7)
C1—N1	1.476 (5)	C13—H13	0.9300
C1—C9	1.507 (6)	C14—H14	0.9300
C1—C2	1.537 (7)	C15—C20	1.400 (6)
C1—H1	0.9800	C15—C16	1.402 (6)
C2—C8	1.521 (6)	C16—C17	1.379 (7)
C2—C3	1.542 (7)	C16—H16	0.9300
C2—H2	0.9800	C17—C18	1.390 (8)
C3—C4	1.534 (6)	C17—H17	0.9300
C3—H3A	0.9700	C18—C19	1.367 (7)
C3—H3B	0.9700	C18—H18	0.9300
C4—C5	1.528 (6)	C19—C20	1.381 (6)
C4—C21	1.574 (7)	C19—H19	0.9300
C4—H4	0.9800	N1—H1A	0.8600
C5—C6	1.538 (6)	C21—C24	1.537 (10)
C5—H5A	0.9700	C21—C22	1.565 (11)
C5—H5B	0.9700	C21—C23	1.567 (10)
C6—C8	1.483 (6)	C24—C25	1.553 (12)

C6—C7	1.531 (6)	C24—H24A	0.9700
C6—H6	0.9800	C24—H24B	0.9700
C7—N1	1.468 (6)	C22—H22A	0.9600
C7—C15	1.503 (6)	C22—H22B	0.9600
C7—H7	0.9800	C22—H22C	0.9600
C8—O1	1.220 (6)	C25—H25A	0.9600
C9—C14	1.364 (7)	C25—H25B	0.9600
C9—C10	1.406 (6)	C25—H25C	0.9600
C10—C11	1.385 (7)	C23—H23A	0.9600
C11—C12	1.376 (8)	C23—H23B	0.9600
C11—H11	0.9300	C23—H23C	0.9600
C12—C13	1.401 (8)		
N1—C1—C9	108.6 (4)	C14—C13—C12	119.0 (6)
N1—C1—C2	110.2 (3)	C14—C13—H13	120.5
C9—C1—C2	113.0 (4)	C12—C13—H13	120.5
N1—C1—H1	108.3	C9—C14—C13	123.5 (5)
C9—C1—H1	108.3	C9—C14—H14	118.2
C2—C1—H1	108.3	C13—C14—H14	118.2
C8—C2—C1	107.1 (4)	C20—C15—C16	115.7 (4)
C8—C2—C3	107.6 (4)	C20—C15—C7	123.0 (4)
C1—C2—C3	116.3 (4)	C16—C15—C7	121.3 (4)
C8—C2—H2	108.5	C17—C16—C15	122.0 (4)
C1—C2—H2	108.5	C17—C16—H16	119.0
C3—C2—H2	108.5	C15—C16—H16	119.0
C4—C3—C2	115.2 (4)	C16—C17—C18	120.0 (5)
C4—C3—H3A	108.5	C16—C17—H17	120.0
C2—C3—H3A	108.5	C18—C17—H17	120.0
C4—C3—H3B	108.5	C19—C18—C17	119.7 (5)
C2—C3—H3B	108.5	C19—C18—H18	120.1
H3A—C3—H3B	107.5	C17—C18—H18	120.1
C5—C4—C3	111.0 (4)	C18—C19—C20	119.8 (5)
C5—C4—C21	114.0 (4)	C18—C19—H19	120.1
C3—C4—C21	112.0 (4)	C20—C19—H19	120.1
C5—C4—H4	106.4	C19—C20—C15	122.7 (5)
C3—C4—H4	106.4	C19—C20—Br2	116.5 (3)
C21—C4—H4	106.4	C15—C20—Br2	120.8 (3)
C4—C5—C6	116.3 (4)	C7—N1—C1	114.5 (4)
C4—C5—H5A	108.2	C7—N1—H1A	122.7
C6—C5—H5A	108.2	C1—N1—H1A	122.7
C4—C5—H5B	108.2	C24—C21—C22	110.5 (7)
C6—C5—H5B	108.2	C24—C21—C23	103.6 (6)
H5A—C5—H5B	107.4	C22—C21—C23	111.4 (7)
C8—C6—C7	108.2 (4)	C24—C21—C4	109.9 (5)
C8—C6—C5	107.5 (4)	C22—C21—C4	110.9 (5)
C7—C6—C5	114.9 (4)	C23—C21—C4	110.3 (5)
C8—C6—H6	108.7	C21—C24—C25	110.3 (8)
C7—C6—H6	108.7	C21—C24—H24A	109.6
C5—C6—H6	108.7	C25—C24—H24A	109.6

N1—C7—C15	109.9 (4)	C21—C24—H24B	109.6
N1—C7—C6	111.0 (3)	C25—C24—H24B	109.6
C15—C7—C6	112.2 (3)	H24A—C24—H24B	108.1
N1—C7—H7	107.9	C21—C22—H22A	109.5
C15—C7—H7	107.9	C21—C22—H22B	109.5
C6—C7—H7	107.9	H22A—C22—H22B	109.5
O1—C8—C6	125.0 (4)	C21—C22—H22C	109.5
O1—C8—C2	123.3 (4)	H22A—C22—H22C	109.5
C6—C8—C2	111.7 (4)	H22B—C22—H22C	109.5
C14—C9—C10	116.3 (4)	C24—C25—H25A	109.5
C14—C9—C1	122.2 (4)	C24—C25—H25B	109.5
C10—C9—C1	121.5 (4)	H25A—C25—H25B	109.5
C11—C10—C9	122.0 (5)	C24—C25—H25C	109.5
C11—C10—Br1	116.8 (4)	H25A—C25—H25C	109.5
C9—C10—Br1	121.2 (4)	H25B—C25—H25C	109.5
C12—C11—C10	119.8 (5)	C21—C23—H23A	109.5
C12—C11—H11	120.1	C21—C23—H23B	109.5
C10—C11—H11	120.1	H23A—C23—H23B	109.5
C11—C12—C13	119.4 (5)	C21—C23—H23C	109.5
C11—C12—H12	120.3	H23A—C23—H23C	109.5
C13—C12—H12	120.3	H23B—C23—H23C	109.5
N1—C1—C2—C8	-56.2 (5)	C10—C11—C12—C13	-1.7 (9)
C9—C1—C2—C8	-178.0 (4)	C11—C12—C13—C14	3.2 (9)
N1—C1—C2—C3	64.1 (5)	C10—C9—C14—C13	-0.1 (8)
C9—C1—C2—C3	-57.7 (5)	C1—C9—C14—C13	179.5 (5)
C8—C2—C3—C4	52.9 (5)	C12—C13—C14—C9	-2.4 (9)
C1—C2—C3—C4	-67.2 (5)	N1—C7—C15—C20	-156.6 (4)
C2—C3—C4—C5	-43.5 (5)	C6—C7—C15—C20	79.4 (5)
C2—C3—C4—C21	-172.2 (4)	N1—C7—C15—C16	25.0 (6)
C3—C4—C5—C6	43.8 (5)	C6—C7—C15—C16	-99.0 (5)
C21—C4—C5—C6	171.4 (4)	C20—C15—C16—C17	-2.7 (7)
C4—C5—C6—C8	-53.5 (5)	C7—C15—C16—C17	175.8 (5)
C4—C5—C6—C7	67.0 (5)	C15—C16—C17—C18	1.3 (8)
C8—C6—C7—N1	55.2 (5)	C16—C17—C18—C19	0.3 (8)
C5—C6—C7—N1	-64.8 (5)	C17—C18—C19—C20	-0.4 (8)
C8—C6—C7—C15	178.6 (4)	C18—C19—C20—C15	-1.1 (7)
C5—C6—C7—C15	58.6 (5)	C18—C19—C20—Br2	179.1 (4)
C7—C6—C8—O1	118.8 (5)	C16—C15—C20—C19	2.6 (6)
C5—C6—C8—O1	-116.6 (5)	C7—C15—C20—C19	-175.9 (4)
C7—C6—C8—C2	-61.2 (5)	C16—C15—C20—Br2	-177.6 (3)
C5—C6—C8—C2	63.4 (5)	C7—C15—C20—Br2	3.9 (6)
C1—C2—C8—O1	-118.1 (5)	C15—C7—N1—C1	-178.9 (3)
C3—C2—C8—O1	116.2 (5)	C6—C7—N1—C1	-54.2 (5)
C1—C2—C8—C6	61.9 (5)	C9—C1—N1—C7	179.5 (3)
C3—C2—C8—C6	-63.8 (5)	C2—C1—N1—C7	55.1 (5)
N1—C1—C9—C14	-28.3 (6)	C5—C4—C21—C24	48.8 (8)
C2—C1—C9—C14	94.4 (5)	C3—C4—C21—C24	175.8 (6)
N1—C1—C9—C10	151.3 (4)	C5—C4—C21—C22	171.4 (6)

supplementary materials

C2—C1—C9—C10	−86.0 (5)	C3—C4—C21—C22	−61.6 (7)
C14—C9—C10—C11	1.8 (7)	C5—C4—C21—C23	−64.8 (6)
C1—C9—C10—C11	−177.9 (4)	C3—C4—C21—C23	62.3 (6)
C14—C9—C10—Br1	−179.2 (4)	C22—C21—C24—C25	−54.6 (10)
C1—C9—C10—Br1	1.1 (6)	C23—C21—C24—C25	−174.0 (8)
C9—C10—C11—C12	−0.9 (8)	C4—C21—C24—C25	68.2 (10)
Br1—C10—C11—C12	−179.9 (5)		
