

EPV0685

Social media as a psychoeducative and preventive tool in mental health

R. Molina Ruiz^{1*}, N. Nuñez Morales² and A. González Vazquez³

¹Hospital clínico san carlos, Psychiatry, Madrid, Spain; ²ITA SALUD MENTAL, Psychiatry, Zaragoza, Spain and ³Hospital Universitario de A Coruña, Psychiatry, A Coruña, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1475

Introduction: The use of social networks is an integral part of our daily life as a means of communication. Scientific dissemination through these kind of platforms has expanded enormously in recent years, for example in networks such as Instagram, a free photo, text messaging and video sharing social media application. Instagram has been used extensively in different fields of medicine. Initially it was limited to visually rich fields but it has been extended to others such as mental health. This tool has enormous potential as a means of more effective communication and prevention.

Objectives: To analyze the role of Instagram as a tool of psychoeducation and prevention in mental health

Methods: An analysis of different mental health profiles was carried based on different items: publications, interactions, likes, commentaries, shares, accounts reached, accounts engaged and hashtags.

Results: Mental health has become one of the sanitary fields generating more traffic on social media reaching a great number of publications per day. Topics that generated more interaction were: depression, anxiety, trauma, sleep and emotion regulation difficulties among others. However most mental health information available in social media had not been provided by professionals of the mental health but by many others.

Conclusions: Instagram is presented as a valuable mental health prevention tool and professionals should engage and adapt to new scenarios of communication. In the era where information is easy to get but knowledge is difficult to find, experts of mental health should more involved.

Disclosure: No significant relationships.

Keywords: mental health; social media; prevention; Instagram

EPV0684

What can we say about the impact of teleconsultations on preventing psychiatric inpatient treatment during COVID-19 pandemic?

I. Figueiredo*, I. Cargaleiro and M. Nascimento

Centro Hospitalar Psiquiátrico de Lisboa, Clinic 3, Lisboa, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1476

Introduction: Telepsychiatry was proved effective and satisfactory in settings like the emergency department and mental health services, but its use is historically restricted. Although there are several studies about telepsychiatry pros and cons, more is needed to know about its effect on patient's follow-up and its influence on inpatient treatment rates, specifically during COVID-19 pandemic.

Objectives: The aim is to look for an eventual change on psychiatric inpatient admissions, during COVID-19 pandemic, when psychiatric patients are in follow-up through teleconsultation.

Methods: We compared the number of hospitalizations for 3 different 6 months periods of time: the 2nd semester of 2019 with no teleconsultations, March-August 2020 only with teleconsultations (except some few 1st consultations) and the 1st semester of 2021 with face-to-face and teleconsultations. A one-way repeated measures ANOVA was conducted on a 1050 patients sample.

Results: The statistics showed that the type of approach in consultations didn't lead to statistically significant differences in hospitalizations (F test-statistic = 0.33086, p = 0.718345).

Conclusions: There is a plethora of advantages about telepsychiatry and it was already shown to be as effective as in-person contact. Some articles show an association of telepsychiatry with a decrease in hospitalization rates, but mostly display similar clinical outcomes. In this study, the authors found that the results follow the latter tendency, although we must consider the COVID-19 pandemic as a possible decompensation and worsening clinical factor. More studies on this matter are important to better understand the potential benefits (and risks) of this treatment setting.

Disclosure: No significant relationships.

EPV0685

YOU'VE BEEN CATFISHED: An exploration of social deception on online platforms

S. Jesus^{1*}, A. Costa², G. Simões¹, G. Dias Dos Santos¹, M. Almeida¹ and P. Garrido¹

¹Baixo Vouga Hospital Centre - EPE, Psychiatry And Mental Health Department, Aveiro, Portugal and ²Centro Hospitalar do Baixo Vouga, Psiquiatria E Saúde Mental, Aveiro, Portugal

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1477

Introduction: Life is a stage in which we are all actors and online we can choose who we want to be. Catfishing is a modern phenomenon in which individuals present themselves online as someone they are not as a means of engaging with others through an idealized avatar. This term has gained prominence since its portrayal in documentary and television series. With the emergence of catfishing, an expectation of betrayal in online relationships is anticipated with increasing caution being exercised by those that engage in online forums.

Objectives: The authors aim to explore this phenomenon and explore what personality traits might be associated with those who engage in catfishing others and in those that fall for the dupe.

Methods: A review of the recent literature on the topic with focus on that which is most relevant to the theme was included.

Results: The literature demonstrates that catfishing is an increasing trend as our online social interaction also increases. Catfishing appears to exist on a scale, where approximately 80% of the online population engage in some form, by means of amplifying their social status. Those with low self-esteem, poor self-worth with and a need to connect and to be validated were most susceptible.

Conclusions: The internet permits anonymity where trading "real world" skins with digital ones creating avatars in order to seek what is desired. Catfishing raises questions about the nature of the human self and the role it plays in deception. Understanding how patients use the internet may provide insight into how personality acts on a stage of total anonymity.

Disclosure: No significant relationships.

Keywords: social interaction; personality; e-mental health; catfish

EPV0686

Griefing reaction and social networks

M.D.C. Molina Liétor^{1*}, M. Martín², I. Cuevas Iñiguez¹,
I. Romero Gerechter¹ and A. Sanz Giancola²

¹Hospital Universitario Príncipe de Asturias, Psiquiatría, Alcalá de Henares, Spain and ²Hospital Universitario Príncipe de Asturias, Psychiatry, Alcalá de Henares, Spain

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1478

Introduction: Grief is a normal and not necessarily pathological psychological process that occurs after the loss of a family member or loved one with its psycho-affective consequences, external manifestations and rituals. Although mourning processes can be associated with losses of different types (employment, housing, baseline situation, housing), we will refer to mourning for the loss of a loved one. For some people, social networks facilitate the expression of feelings and experiences of grief, connecting with the emotional support of other friends and loved ones. However, the presence of accounts belonging to these deceased persons, the persistence of photos and memories that periodically appear on the screen without the person being able to choose them, can make it difficult to process the mourning.

Objectives: The aim of this paper is to consider the beneficial and detrimental factors of social media during a grieving reaction after the loss of a loved one.

Methods: For the preparation of this work, a bibliographic review on the subject has been carried out. Likewise, the clinical information provided by patients during our evaluations has provided critical views on what has been published in this regard.

Results: Support through social networks can help to feel more affectionate, but there are other harmful factors that must be taken into account: permanence of photos, appearance of memories and reminder of the deceased person's birthday.

Conclusions: Social networks can have favorable but also detrimental factors in the elaboration of a grief and should be considered in the psychiatric exploration and intervention.

Disclosure: No significant relationships.

Keywords: Instagram; Griefing reaction; Facebook

EPV0688

Prediction and classification of anxiety-related psychological scale and VR sickness based on autonomic physiological responses during VR treatment in patients with social anxiety disorder

J.Y. Chun¹, H.-J. Kim², S. Lee², M. Kim², T. Cheong¹ and C.-H. Cho^{2*}

¹Korea University, School Of Industrial And Management Engineering, Seoul, Korea, Republic of and ²Chungnam National University Sejong Hospital, Psychiatry, Sejong, Korea, Republic of

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1479

Introduction: Social anxiety disorder (SAD) can accompany emotional symptoms as well as physical reactions. The assessment and real-time measurement of SAD is difficult in real-world.

Objectives: This study aims to predict the severity of specific anxiety states and virtual reality (VR) sickness in SAD patients by a machine learning model based on only quantitative measuring of autonomic physiologic signals during VR therapy sessions.

Methods: In total, 32 individuals with SAD symptoms were enrolled in VR participatory sessions. We assessed patients' specific anxiety symptoms through Internalized Shame Scale (ISS) and Post-Event Rumination Scale (PERS), and VR sickness through Simulator Sickness Questionnaire (SSQ). Specific anxiety symptoms and VR sickness were divided into severe and non-severe states based on the total score of each scale by K-means clustering. Logistic regression, Random Forest, Naïve Bayes classifier, and Support Vector Machine were used based on the physiological signal data to predict the severity group in subdomains of ISS, PERS, and SSQ.

Results: Prediction performance (F1 score) for the severity of the ISS mistake anxiety subdomain was higher than other scales with 0.8421. For VR sickness, prediction performance for the severity of the physical subdomain was higher than the non-physical subdomain with 0.7692.

Conclusions: The study findings present that mistake anxiety and physical sickness could be predicted more accurately by only autonomic physiological signals, suggesting these features are probably associated with autonomic responses. Based on the present study results, we could provide the evidence for predicting the severity of specific anxiety or VR adverse effects only based on in-situ physiological signals.

Disclosure: No significant relationships.

Keywords: physiological response; social anxiety disorder; virtual reality; Anxiety

Eating Disorders

EPV0689

Emotion dysregulation, dissociation and body dissatisfaction mediate the relationship between trauma exposure and ED symptoms

L. Lev-Ari^{1*}, A. Zohar² and R. Bachner Melman²

¹Ruppim Academic Center, The Lior Tsfaty Center For Suicide And Mental Pain Studies, Emek Hefer, Israel and ²Ruppim Academic Center, Clinical Psychology, Emek Hefer, Israel

*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1480

Introduction: The current study tests the relationship between eating disorder (ED) symptoms and trauma exposure. The mechanisms via which trauma is related to ED symptoms have not been sufficiently examined. This study examines the complex role of dissociation and emotional dysregulation in the context of trauma, BMI, ED symptoms and body dissatisfaction (BD).

Objectives: We hypothesized that dissociation and emotional dysregulation would mediate the relationship between trauma exposure and ED symptoms / BD. We further hypothesized that BMI would play a moderating role in this association.