

Evidence-Based Medicine of Traumatic Thoracolumbar Burst Fractures: A Systematic Review of Operative Management across 20 Years

Justin K. Scheer¹ Joshua Bakhsheshian¹ Shayan Fakurnejad¹ Taemin Oh¹ Nader S. Dahdaleh¹
Zachary A. Smith¹

¹Department of Neurological Surgery, Northwestern University, Feinberg School of Medicine, Chicago, Illinois, United States

Address for correspondence Justin K. Scheer, BS, Department of Neurological Surgery, Northwestern University, Feinberg School of Medicine, 676 N. St. Clair Street, Suite 2210, Chicago, IL 60611, United States (e-mail: jscheer@ucsd.edu).

Global Spine J 2015;5:73–82.

Abstract

Study Design Systematic literature review.

Objective The management of traumatic thoracolumbar burst fractures (TLBF) remains challenging, and analyzing the levels of evidence (LOEs) for treatment practices can reform the decision-making process. However, no review has yet evaluated the operative management of traumatic thoracolumbar burst fractures with particular attention placed on LOE from an established methodology. The objective of the present study was to characterize the literature evidence for TLBF, specifically for operative management.

Methods A comprehensive search of the English literature over the past 20 years was conducted using PubMed (MEDLINE). The inclusion criteria consisted of (1) traumatic burst fractures (2) in the thoracic or lumbar spine. Exclusion criteria included (1) osteoporotic burst fractures, (2) pathologic burst fractures, (3) cervical fractures, (4) biomechanical studies or those involving cadavers, and (5) computer-based studies. Studies were assigned an LOE and those meeting level 1 or 2 were included.

Results From 1,138 abstracts, 272 studies met the criteria. Twenty-three studies (8.5%) met level 1 ($n = 4$, 1.5%) or 2 ($n = 19$, 7.0%) criteria. All 23 studies were reported.

Conclusions The literature contains a high LOE to support the operative management of traumatic thoracolumbar burst fractures. For patients who are neurologically intact, a high LOE demonstrated similar functional outcomes, lower complication rates, and less costs with conservative management when compared with surgical management. There is a high LOE for short- or long-segment pedicle instrumentation without fusion and less invasive (percutaneous and paraspinal) approaches. Furthermore, the posterior approaches are associated with lower complications as opposed to the anterior or combined approaches.

Keywords

- ▶ burst fracture
- ▶ thoracolumbar
- ▶ spine
- ▶ trauma
- ▶ level of evidence

received
May 6, 2014
accepted after revision
October 12, 2014
published online
November 24, 2014

© 2015 Georg Thieme Verlag KG
Stuttgart · New York

DOI <http://dx.doi.org/10.1055/s-0034-1396047>.
ISSN 2192-5682.

Introduction

Traumatic spinal fractures can cause spinal instability with neurologic deficits. The majority of such fractures occur at the thoracolumbar junction (T11–L2), largely due to regional biomechanics in which the rigid, kyphotic thoracic spine is juxtaposed with the more mobile, lordotic lumbar spine.¹ Burst fractures account for 30 to 64% of the thoracolumbar spinal fractures.^{1,2} Historically, burst fractures have been defined as involving the anterior and middle spinal columns based on the Denis three-column spinal model.³ However, the more current classification of burst fractures utilizes the load-sharing classification.⁴ Under the Magerl fracture classification system, burst fractures are designated as type A3 compression fractures.² The etiology is typically related to powerful axial compression forces, such as in falls from heights (34 to 54%),^{1,5} motor vehicle accidents (51 to 65%),^{1,5} or high-intensity sports (9%).^{5–9}

Despite an overwhelming amount of descriptive and clinical studies addressing traumatic thoracolumbar burst fractures, a significant controversy surrounds the optimum management strategies. This is reflected by the lack of uniformity in the treatment modalities. Typically, the management can be either operative or conservative: the former involves a combination of spinal fixation and fusion that may include interbody devices, with or without decompression,^{1,10–17} and the latter incorporates the use of braces/casts, bed rest, analgesics, or observation with no bed rest or casts.^{12,13,17–24} By extension, the algorithms to guide management decisions are limited, with one example being the Thoracolumbar Injury Classification and Severity Score.²⁵

With ambiguity regarding the best management, compiling and analyzing the levels of evidence (LOEs) for treatment practices can reform the decision-making process. The practice of evidence-based health care has gradually become more prevalent in modern medicine, as the pay-for-performance interventions with high LOE lead to greater reimbursements.^{26–31} Systematic reviews (SRs) and meta-analyses provide the means to rapidly analyze a study's quality, ultimately facilitating the introduction of recommendation grades and providing a foundation for establishing practice guidelines.³² To the author's best knowledge, there are currently seven SRs involving various aspects of traumatic thoracolumbar burst fractures.^{13,16,33–37} However, no review has yet evaluated the operative management of traumatic thoracolumbar burst fractures with particular attention placed on the LOE from an established methodology. The current LOE for the operative management of traumatic thoracolumbar burst fractures remains unknown. Therefore, the objective of the present study was to characterize the literature evidence for traumatic thoracolumbar burst fractures, specifically by detailing the studies involving the operative management, with an LOE of 2 or greater.

Methods

A systematic literature review was conducted according to the Cochrane guidelines to capture all studies pertaining to

traumatic thoracolumbar burst fractures over the past 20 years.³⁸ This time frame was arbitrarily predetermined, as the surgical management of burst fractures has greatly evolved during this period. An electronic search of PubMed (MEDLINE) was done on April 17, 2014 with permutations of the following terms: (“spine AND burst”) or “burst fracture” or (“burst and fracture”) or (A3 and fracture and spine). The inclusion criteria included: (1) burst fractures resulting from traumatic mechanism and (2) fractures of the thoracic or lumbar spine. The exclusion criteria were: (1) osteoporotic burst fractures, (2) pathologic burst fractures, (3) cervical fractures, (4) biomechanical studies or those involving cadavers, and (5) computer-based studies such as finite element analyses. Search results were blinded such that only the titles and abstracts were reviewed. When inclusion or exclusion was unclear, full texts were retrieved; if doubt remained, a decision was made through discussion and addition of a third reviewer if needed. Following the search, only those involving operative management were selected to be reviewed in the present study. They were categorized as operative versus nonoperative or as treatment/technique studies.

LOEs were assigned using criteria from the Journal of Clinical Orthopaedics and Related Research,³⁰ which is adapted from the Oxford Centre of Evidence Based Medicine LOEs.^{39,40} The LOE scale ranges from 1 being the highest to 5 the lowest. Studies with level 1 or 2 LOE were classified as high LOE. Ambiguities regarding LOE were resolved through discussion and addition of a third reviewer if needed. Once all the studies were assigned a LOE, only those rated as level 1 and 2 were included. All data were stored in Microsoft Excel 2013 (Microsoft Corp., Redmond, Washington, United States).

Results

Overall Composition of the Literature

The search yielded a total of 1,138 abstracts, of which 272 met the criteria, involved operative management, as well as were published between January 1, 1992, and April 17, 2014. Twenty-three of the studies (8.5%) met level 1 ($n = 4$, 1.5%) or 2 ($n = 19$, 7.0%) criteria (– **Table 1**).

Operative versus Nonoperative Management

Fifteen of the studies (5.5%) investigated the differences between operative versus nonoperative treatment. Of these,

Table 1 The number of studies found for each level of evidence and the corresponding percentage

Level of evidence	<i>n</i>	Percentage
Total studies	274	100.0
Level 1	4	1.5
Level 2	19	7.0
Level 3	38	14.0
Level 4	210	77.2
Level 5	1	0.4

seven qualified as high LOE (two level 1, five level 2; ► **Table 2**). All others were classified as level 3 ($n = 6$) and level 4 ($n = 2$).

Two studies—one SR and one randomized controlled trial (RCT)—met level 1 criteria. Abudou et al reported an SR through the Cochrane collaboration in which they searched for RCTs comparing operative and nonoperative treatment in patients without neurologic deficit.⁴¹ Due to strict inclusion and exclusion criteria, only two studies with contrasting results were identified (Siebenga et al and Wood et al, discussed below),^{14,42} leading the authors to conclude that there is insufficient evidence to conclude whether surgical or conservative treatment yields superior functional outcomes.⁴¹ The second study was an RCT reported by Siebenga et al that compared the radiographic and functional outcomes in neurologically intact patients with burst fractures. The operative cohort underwent short-segment posterior pedicle screw fixation above and below the fracture level with an autologous fusion; the nonoperative treatment involved rehabilitation and bed rest for ≥ 5 days following the injury. All patients wore a Jewett hyperextension orthosis for 3 months. Although both the patient groups had similar baseline local and regional sagittal angles and visual analog scale scores, the surgery resulted in significantly lower angles and less pain at the final follow-up. Further, the authors reported that the conservative treatment led to higher Roland Morris Disability Questionnaire-24 (RMDQ-24) scores at the final follow-up, with fewer patients capable of returning to work.⁴²

Five studies met the level 2 criteria. In 1992, Dr. Gertzbein and the Scoliosis Research Society prospectively analyzed spinal fractures in 1,019 patients across 12 countries. Operative intervention was administered to 545 patients and 96 received conservative therapy. The results revealed a statistically significant, albeit weak, relationship between the initial canal compromise and the neurologic deficit in burst fractures. However, there were no differences in the rates of the neurologic improvement between the treatment groups. Of the operative patients, the neurologic outcomes were equivalent for anterior versus posterior surgery. Among patients with preoperative neurologic deterioration, surgical management, particularly anterior surgery, led to greater improvement. At 2-year follow-up, a higher proportion of patients managed conservatively complained of “severe/moderate” pain compared with “none/mild” pain.¹ Hitchon et al performed a prospective study to outline the indications, benefits, and costs of bed rest compared with surgical stabilization with pedicle screws, rods, and plates; Luque or Harrington rods with sublaminar wiring; or an “anterior spinal device.” All patients were immobilized in thoracolumbar orthoses for 3 to 5 months. Although both cohorts improved neurologically, a progressive angular deformity was present, with rates twice as high in the surgery group.¹¹

Shen et al conducted an RCT to examine nonoperative treatment (a hyperextension brace for 3 months without fracture reduction) versus short-segment posterior fixation. Although the surgical group had better pain and clinical outcomes in the early postoperative period (3 to 6 months), these differences were not observed at final follow-up (2

years). Regarding kyphosis correction, the operative group had an initial correction of 17 degrees, which was lost over time, and the nonoperative group had a loss of 4 degrees. Also, hospital charges were 4 times greater with surgery.¹² Wood et al reported another RCT in which nonoperative patients underwent a fracture reduction supplemented by a body cast or a thoracolumbar orthosis for 3 months. The operative patients underwent either a posterolateral spinal arthrodesis with pedicle screw and hook instrumentation and autologous bone grafting or an anterior two-level fibular and rib-strut construct arthrodesis with local autologous bone grafting and instrumentation. Both groups were similar at final follow-up with respect to the radiographic and clinical outcomes.¹⁴ A level 2 meta-analysis reported by Gnanenthiran et al pooled the results from the four aforementioned trials. No differences in the clinical outcomes of pain and function were seen. The surgical group, however, did show better correction radiographically at follow-up, albeit this came with higher complication rates and costs.¹³

Treatment and Technique

Two hundred fifty-seven studies (9.5%) examined the operative treatment and technique of thoracolumbar burst fractures. There were 16 high LOE studies (two level 1, 14 level 2; ► **Table 2**), and the rest were level 3 ($n = 32$), level 4 ($n = 208$), and level 5 studies ($n = 1$).

Both level 1 studies were prospective RCTs. Dai et al investigated posterolateral fusion in combination with short-segment posterior pedicle screw instrumentation into the levels cephalad and caudad to the fractured vertebra. A total of 37 patients underwent fusion and 36 did not, and no differences in the radiographic or clinical outcomes were seen.⁴³ In the second study, two muscle-sparing surgical approaches—the percutaneous and paraspinal approaches for pedicle screw and rod fixation—were compared by Jiang et al for efficacy and safety. The percutaneous approach led to lower intraoperative blood loss, shorter surgical time, and shorter hospitalization. In the short-term postoperative period, this approach also yielded less pain and greater improvements in the functional scores. In contrast, the paraspinal approach resulted in better correction of the kyphosis and restoration of the vertebral height. The long-term clinical outcomes were equivalent.⁴⁴

The first level 2 study was published by Gertzbein and colleagues, in which 25 patients were treated with the AO internal fixator before or after the first 4 days postinjury. Earlier intervention led to greater improvement in canal encroachment.⁴⁵ Vornanen et al compared three methods of internal fixation (Harrington rods, AO internal fixators, posterior segmental fixators) with respect to improvement in canal encroachment. All methods improved encroachment, with Harrington rods providing the best reduction.⁴⁶ Next, Alanay et al reported an RCT comparing short-segment pedicle instrumentation with or without transpedicular grafting. No differences in the radiographic, operative, pain, or functional outcomes were seen.⁴⁷

Stancić et al compared anterior and posterior surgical approaches for overall clinical outcomes and operative

Table 2 Studies ($n = 23$) with high level of evidence

Author	Study topic	Follow-up	Population/etiology	Assessment measures	Results
Xu et al 2013 ³⁷	Technique	NA	4 RCT and 3 controlled clinical trials	Study design, patient demographics, inclusion and exclusion criteria, interventions, outcomes, follow-up duration, and rate of lost to follow-up	The posterior approach may be more effective than the anterior. More high-level evidence is required.
Schmid et al 2012 ⁵⁶	Technique	20.3 ^a	21 TLIF and 14 combined posteroanterior fusion	Surgical complications, VAS, ODI, MPQ, RMDQ, FbHR, and radiographic	There were no significant differences between the two treatment groups regarding clinical and radiologic outcome.
Jindal et al 2012 ⁵⁵	Technique	23.9 (18–30)	Short-segment pedicle screw fixation: 23 with fusion and 24 without fusion	Surgical complications, VAS, Greenough Low Back Outcome Scale, neurologic status (AIS), radiographic	Adjunctive fusion is unnecessary when managing patients suffering from burst fractures of the thoracolumbar spine with short-segment pedicle screw fixation.
Jiang et al 2012 ⁴⁴	Technique	58.6 ^a (37–84)	31 percutaneous and 30 paraspinous	Surgical complications, instrumentation accuracy, VAS, ODI, and radiographic	The percutaneous approach is recommended in cases with successful postural reduction, whereas the paraspinous approach is associated with better surgical correction.
Gnanenthiran et al 2012 ¹³	Operative vs. nonoperative	NA	4 RCT studies	Study design, methodological quality, clinical outcomes, costs	Operative management for patients without neurologic deficit may improve residual kyphosis but does not appear to improve pain or function and is associated with higher complication rates and costs.
Wei et al 2010 ⁵⁴	Technique	27.8 (19–52)	47 MSPI and 38 SSPI	ODI, LBOS, LSC, radiographic	MSPI yields decreased operative time and blood loss, and offers better clinical results.
Tezeren et al 2009 ⁵³	Technique	34.6 (24–60)	Long-segment posterior instrumentation: 21 with fusion and 21 without fusion	LBOS, radiographic	Spinal fusion is not necessary in long-segment posterior instrumentation.
Pang et al 2009 ⁵²	Technique	18.6 (12–26)	34 paravertebral and 28 traditional	Operative data, VAS	The paravertebral approach is recommended because it is much less invasive, can reduce blood loss, and accelerates rehabilitation.
Dai et al 2009 ⁴³	Technique	72 (60–84)	Posterior short-segment fixation: 37 with fusion and 36 without fusion	Operative data, neurologic status (Frankel, ASIA motor score), load-sharing score, SF-36, VAS, radiographic	Fusion is not necessary with short-segment pedicle screw fixation for Denis type-B burst fractures with load-sharing scores of ≤ 6 .
Dai et al 2009 ⁵¹	Technique	64.8 ^a (48–84)	32 iliac graft and 33 titanium mesh cage	Operative data, neurologic status (Frankel, ASIA motor score), SF-36, VAS, radiographic	No significant difference was found between the treatment groups, and both approaches are effective surgical treatments.
Abudou et al 2013 ⁴¹	Operative vs. nonoperative	NA	2 RCT studies	Study design, methodological quality, clinical outcomes, duration of hospitalization	One trial found better pain improvement and functional outcomes and the other trial found the opposite. Two contradictory studies was not enough to conclude whether operative or nonoperative management was better.

Table 2 (Continued)

Author	Study topic	Follow-up	Population/etiology	Assessment measures	Results
Korovessis et al 2006 ⁵⁰	Technique	47 ^a (27–74)	Short-segment transpedicular fixation: 20 with anterior and posterior stabilization and 20 with solely posterior	Operative data, surgical complications, neurologic status (Frankel), VAS, SF-36, radiographic	SSTF is not recommended for operative stabilization of such fractures.
Siebenga et al 2006 ⁴²	Operative vs. nonoperative	51.6 (24–79.2)	17 operative (posterior stabilization with transpedicular grafting or posterolateral fusion) and 15 nonoperative (orthosis)	Treatment complications, fracture classification, VAS, RMDQ, LSC, duration of hospitalization, radiographic	Patients with a type A3 fractures without neurologic deficit should be treated by short-segment posterior stabilization.
Wood et al 2005 ⁵⁷	Technique	43.5 (24–108)	20 anterior and 18 posterior fusion	Operative data, surgical complications, mechanism of injury, VAS, RMDQ, ODI, SF-36, radiographic	Anterior fusion and instrumentation for thoracolumbar burst fractures may present fewer complications or additional surgeries.
Tezeren and Kuru 2005 ⁴⁹	Technique	29.6 (23–40)	9 short-segment pedicle fixation and 9 long-segment pedicle fixation	Operative data, LBOS, radiographic	LS fixation is a more effective management of thoracolumbar burst fractures. Nevertheless, clinical outcome was the same between the two groups.
Wood et al 2003 ¹⁴	Operative vs. nonoperative	44 (24–118)	24 operative (posterior or anterior arthrodesis and instrumentation) and 23 nonoperative (body cast or orthosis)	Treatment complications, VAS, RMDQ, ODI, SF-36, radiographic, cost	Operative treatment of patients with stable fractures without neurologic deficits provided no significant long-term advantage compared with nonoperative treatment.
Stancić et al 2001 ⁴⁸	Technique	12	13 anterior neurodecompression and fixation and 12 posterior reposition and semirigid fixation by hook-rod pedicle screw fixation	Operative data, cosmetic result, Prolo functional rating scale, neurologic status (ASIA), radiographic, cost	No significant differences were found between the techniques in terms of neurologic improvement and functional outcome. The posterior approach is recommended in emergency neurodecompression and fixation of unstable thoracolumbar fractures because of the shorter operation time and smaller blood loss.
Shen et al 2001 ¹²	Operative vs. nonoperative	24	33 operative (fixation with instrumentation) and 47 nonoperative (fitted brace)	Employment status (Denis), patient satisfaction, VAS, LBOS, radiographic, cost	Surgical intervention provides partial kyphosis correction and earlier pain relief, but functional outcomes for both interventions were similar at 2 y.
Alanay et al 2001 ⁴⁷	Technique	32 (24–72)	Short-segment posterior instrumentation: 10 with transpedicular grafting and 10 without grafting	Operative data, fracture classification (Denis), LSC, radiographic	Grafting did not significantly decrease the high rate of failure in short-segment instrumentation.
Hitchon et al 1998 ¹¹	Operative vs. nonoperative	15.4 ^a	36 operative (various techniques) and 32 nonoperative (orthosis)	Neurologic status (Frankel), pain, employment, radiographic, cost	Selection of operative or nonoperative intervention should be based on severity of clinical and radiologic findings.

(Continued)

Table 2 (Continued)

Author	Study topic	Follow-up	Population/etiology	Assessment measures	Results
Vornanen et al 1995 ⁴⁶	Technique	NA	22 fixed with Harrington rods, 21 with AO internal, and 24 with posterior segmental	Fracture classification, canal encroachment, radiographic	There were no significant differences between the three instrumentation devices.
Gertzbein et al 1992 ⁴⁵	Technique	NA	AO internal fixator: 16 patients treated within 4 d of injury and 9 treated 5 d or more after injury	Fracture classification, canal encroachment, radiographic	The AO internal fixator is most effective when used within the first 4 d after injury in patients with an initial canal compromise of 34–66%.
Gertzbein 1992 ¹	Operative vs. nonoperative	24	820 operative and 199 nonoperative	Neurologic status (Frankel, Motor Index Score), pain, radiographic	Many conclusions were drawn from this large prospective study, pertaining to prognosis, diagnosis, and treatment of burst fractures.

Abbreviations: AIS, American Spinal Injury Association Impairment Scale; ASIA, American Spinal Injury Association; FFbH-R, Hannover Functional Questionnaire; LBOS, Low Back Outcome Score; LS, long-segment; LSC, Load-Sharing Classification; MPQ, McGill Pain Questionnaire; MSPi, monosegmental pedicle instrumentation; NA, not applicable; ODI, Oswestry Disability Index; RCT, randomized controlled trial; RMDQ, Roland-Morris Disability Questionnaire; SF-36, Short Form-36; SSPi, short-segment pedicle instrumentation; TILF, transforaminal lumbar interbody fusion; VAS, visual analog scale.
^aWeighted mean.

efficacy. Of the 25 patients enrolled, 13 received anterior neurodecompression and fixation and 12 underwent posterior fixation by hook and rod and pedicle screw fixation. Neither the neurologic outcomes nor economic/functional outcomes proved to be different. However, the posterior group suffered less blood loss and had shorter operative times, better aesthetic outcomes, and lower costs, complications, and donor site pain.⁴⁸

Tezeren and Kuru performed a prospective study in which patients were treated with short- or long-segment pedicle fixation.⁴⁹ The short-segment fixation involved fixation one level above and below the fracture; long-segment instrumentation involved hook fixation two or three levels above plus pedicle fixation two levels below the fracture. At final follow-up, patients in the long-segment group showed significantly better correction of the sagittal index and anterior body compression.⁴⁹ Wood and colleagues compared anterior reconstruction, fusion, and instrumentation to posterior fusion with instrumentation.⁵⁷ Although the posterior approach had significantly less blood loss, no differences in radiographic, clinical, and functional outcomes were observed. In a prospective randomized trial by Korovessis et al, patients received a combined anterior and posterior stabilization technique or standard posterior short-segment transpedicular fixation. Both resulted in radiographic improvement but the combined group faced greater blood loss, operative time, complications, and duration of hospital stay. Furthermore, visual analog scale and SF-36 scores did not correlate with the anatomical correction in either treatment group.⁵⁰

Dai et al prospectively evaluated the outcomes following anterior-only stabilization with either structural autografts or titanium mesh cages. Apart from the differing instrumentation, the surgical approaches were similar between the groups. In the autograft group, the operative times were longer but no other differences were observed. Both the treatment groups improved postoperatively and there were no significant differences.⁵¹

In 2009, Pang et al prospectively compared the outcomes and surgical efficiency/safety between an approach via the paravertebral muscle or the traditional posterior approach. The paravertebral muscle approach led to significantly better outcomes in terms of blood loss, drainage, duration of recumbency, and clinical outcome scores.⁵² Tezeren et al prospectively investigated long-segment instrumentation with or without fusion. The average operation time and blood loss were significantly lower in the nonfusion group, but the two groups were otherwise similar in terms of the radiologic and clinical assessment measures that were analyzed.⁵³

Wei and colleagues published a clinical trial in 2010 comparing the clinical and radiologic outcomes between monosegmental transpedicular fixation and short-segment pedicle instrumentation. Patients were analyzed based on the operative data, Low Back Outcome Score (LBOS), Oswestry Disability Index, Load-Sharing Classification, sagittal index, and percentage of anterior body height compression. Monosegmental fixation resulted in

lower operative times and blood loss but did lead to higher LBOS scores.⁵⁴ The groups were also similar at final follow-up for the sagittal index, local kyphosis, and average correction loss. Following this, in 2012, Jindal et al assessed the benefits of short-segment pedicle fixation with or without fusion in an RCT. Ultimately, the fusion did not improve the radiographic or functional outcomes but did increase the operative time and necessity for blood transfusion.⁵⁵ Schmid et al prospectively compared thoracoscopy or transforaminal lumbar interbody fusion (TLIF) in the anterior reconstruction of burst fractures. The TLIF group received a monocortical strut graft and the thoracoscopy included tricortical strut grafts or titanium vertebral body-replacing implants. Satisfactory outcomes were achieved in both groups, and no differences were seen in the radiographic or clinical outcomes. Surgical decompression, however, was higher with TLIF.⁵⁶

In 2013, Xu et al performed a meta-analysis of RCTs, quasi-RCTs, or controlled trials investigating anterior versus posterior approaches in attempt to identify the optimal approach.³⁷ The authors analyzed seven studies comparing 179 and 152 patients in the anterior and posterior groups, respectively. No significant differences were found between the approaches for neurologic recovery, return to work, complications, and deformity correction. However, the anterior group was associated with longer operative time, greater blood loss, and a higher cost than the posterior approach.

Discussion

Traumatic thoracolumbar burst fractures remain a challenge to treat, and optimal operative and nonoperative techniques have not yet been established. The goals of treatment are to obtain early patient mobilization and a painless, balanced, stable vertebral column with maximum spine mobility and optimal neurologic function. To achieve these goals, numerous studies have investigated different diagnostic, prognostic, and management approaches, with a greater emphasis in the latter. The present study attempted to characterize the high LOE studies pertaining to the operative management of traumatic thoracolumbar burst fractures and found that the overall literature contained 23 studies (8.5%) in PubMed that met an LOE of 1 or 2.

When treating neurologically intact patients with thoracolumbar burst fractures, there is no difference in the clinical outcomes when comparing a conservative or open operative approach. This is supported by five of the seven high-level studies investigating this cohort. Although better correction of the spinal deformity can be achieved surgically, three level 2 clinical studies demonstrated higher rates of complications and costs with this approach. Two high-level SRs confirmed no difference in the functional outcomes when the approaches were compared. These studies utilized different operative and conservative approaches, follow-up times, and inclusion/exclusion criteria. It is worth noting that there are limited numbers of patients in these studies and thus their results are inherently limited.

There is a high LOE for short- or long-segment pedicle instrumentation without fusion. Tezeren and Kuru compared short- versus long-segment pedicle fixation in a level 2 study and concluded that better correction can be achieved with the long construct, and LBOS outcomes were similar.⁴⁹ Longer constructs provide the necessary rigidity to decrease failure rates, but they also decrease patient mobility due to the more rigid fixation, which may impact quality-of-life parameters that were not assessed. Four high-level studies questioned the necessity of additional fusion to posterior short-segment pedicle instrumentation. Dai et al (level 1), Alanay et al (level 2), and Jindal et al (level 2) demonstrated that bone grafting was not necessary in short-segment pedicle instrumentation in RCTs.^{43,47,55} Tezeren et al (level 2) also demonstrated that fusion did not change the radiologic or clinical outcomes in long-segment instrumentation.⁵³ The operative time would decrease by forgoing fusion, and because bone graft is not needed, this would collectively decrease the risk of infections. As mentioned previously with studies involving neurologically intact patients, the discussed RCTs also have small sample sizes and the conclusions from these studies should be considered within the context of that limitation.

Open approaches to spine surgery are associated with higher morbidity, and there has been a recent trend toward preserving thoracolumbar muscle attachments. The percutaneous and paraspinal approaches to thoracolumbar burst fractures are supported by level 1 and 2 studies. Pang et al (level 2) demonstrated that the paravertebral approach results in better clinical outcome scores than the traditional posterior approach.⁵² Jiang et al (level 1) demonstrated that shorter hospitalization time and equal 3-year clinical outcomes can be achieved with the percutaneous versus paraspinal approach in pedicle screw and rod fixation.⁴⁴ However, better deformity correction can be achieved with the paraspinal approach.

When investigating an anterior, posterior, or combined approach in instrumentation, four level 2 studies reported similar radiographic, clinical, and functional outcomes.^{37,48,50,57} Furthermore, when compared with the anterior or combined approach, the posterior approach correlated with fewer complications and operative risks. When considering anterior reconstruction, Dai et al demonstrated no significant differences in the use of titanium or autologous grafting for fusion.⁵¹ Furthermore, Schmid et al demonstrated that transforaminal lumbar interbody fusion may provide further surgical decompression than thoracoscopy.⁵⁶ And finally, Xu et al performed a meta-analysis and found similar results with the exception that the anterior approach also had a higher cost associated with it.³⁷ The choice of the operative approach can be decided by the preference of the surgeon and patient, while acknowledging that greater risks and possible higher costs are associated with the anterior approach.

Additional considerations in the operative management of traumatic thoracolumbar burst fractures include both the correction of a kyphotic deformity and the potential loss of a kyphotic correction over time. The results regarding whether

operative management maintained the correction more successfully than nonoperative management are mixed.^{12,14,42} All studies show a significant improvement postoperatively; however, in the study by Siebenga et al (level 1),⁴² the operative group maintained the correction and the kyphosis was significantly less than in the nonoperative group at final follow-up (a mean of 4.3 years). On the other hand, Shen et al (level 2) and Wood et al (level 2) did not show any differences in the deformity correction between operative and nonoperative management at the final follow-up (2 years).^{12,14} The clinical implication of kyphotic correction is questionable, as Wood et al (level 2) and Korovessis et al (level 2) both did not find any correlation between the degree of kyphosis at the final follow-up and the degree of pain or disability.^{14,50} Operative management does improve pain and stability.^{12–14,16,42} Most studies had similar radiographic results between the various operative techniques.^{37,43,47,51,53–57} Specifically, fusion does not appear to affect the maintenance of radiographic correction as demonstrated by four high LOE studies.^{43,47,53,55} However, Jiang and colleagues (level 1) found better long-term local kyphosis correction with a paraspinous approach versus a percutaneous.⁴⁴ Similarly, Tezzer and Kuruet (level 2) found better kyphotic correction at final follow-up for patients who underwent long-segment pedicle fixation compared with short-segment fixation.⁴⁹

The heterogeneity nature of burst fractures due to the different mechanisms of injury results in a variety of clinical presentations, and makes it difficult to formulate a universal approach. Other SRs were limited to select studies by the strict inclusion/exclusion criteria and therefore did not include most of the literature covered in this report. This report was conducted in accordance with the Cochrane Collaboration Handbook on SRs; it highlights the current evidence and provides further insight into future investigations needed. This review is limited by studies found only in the PubMed database (MEDLINE). However, the MEDLINE database is well accepted as the comprehensive source of relevant literature.

Conclusion

The literature contains a high LOE to support the operative management of traumatic thoracolumbar burst fractures. The highest degree of controversy involves the decision whether to operate on neurologically intact patients. For patients who are neurologically intact, a high LOE demonstrated similar functional outcomes and lower complication rates and costs with conservative management when compared with surgical management. However, the evidence is unclear when there is a neurologic deficit. If a patient is determined to require surgery, there is no general consensus on the appropriate technique to be used. There is a high LOE for short- or long-segment pedicle instrumentation without fusion and less invasive (percutaneous and paraspinous) approaches. Furthermore, posterior approaches are associated with fewer complications as opposed to anterior or combined approaches.

Funding

No funding was used for this study.

Disclosures

Justin K. Scheer, none
 Joshua Bakhsheshian, none
 Shayan Fakurnejad, none
 Taemin Oh, none
 Nader S. Dahdaleh, none
 Zachary A. Smith, none

References

- Gertzbein SD. Scoliosis Research Society. Multicenter spine fracture study. *Spine (Phila Pa 1976)* 1992;17(5):528–540
- Magerl F, Aebi M, Gertzbein SD, Harms J, Nazarian S. A comprehensive classification of thoracic and lumbar injuries. *Eur Spine J* 1994;3(4):184–201
- Denis F. The three column spine and its significance in the classification of acute thoracolumbar spinal injuries. *Spine (Phila Pa 1976)* 1983;8(8):817–831
- McCormack T, Karaikovic E, Gaines RW. The load sharing classification of spine fractures. *Spine (Phila Pa 1976)* 1994;19(15):1741–1744
- Bensch FV, Koivikko MP, Kiuru MJ, Koskinen SK. The incidence and distribution of burst fractures. *Emerg Radiol* 2006;12(3):124–129
- Donald S, Chalmers D, Theis JC. Are snowboarders more likely to damage their spines than skiers? Lessons learned from a study of spinal injuries from the Otago ski fields in New Zealand. *N Z Med J* 2005;118(1217):U1530
- Shorter NA, Mooney DP, Harmon BJ. Snowboarding injuries in children and adolescents. *Am J Emerg Med* 1999;17(3):261–263
- Koo DW, Fish WW. Spinal cord injury and snowboarding—the British Columbia experience. *J Spinal Cord Med* 1999;22(4):246–251
- Gertzbein SD, Khoury D, Bullington A, St John TA, Larson AI. Thoracic and lumbar fractures associated with skiing and snowboarding injuries according to the AO Comprehensive Classification. *Am J Sports Med* 2012;40(8):1750–1754
- Butt MF, Farooq M, Mir B, Dhar AS, Hussain A, Mumtaz M. Management of unstable thoracolumbar spinal injuries by posterior short segment spinal fixation. *Int Orthop* 2007;31(2):259–264
- Hitchon PW, Torner JC, Haddad SF, Follett KA. Management options in thoracolumbar burst fractures. *Surg Neurol* 1998;49(6):619–626, discussion 626–627
- Shen WJ, Liu TJ, Shen YS. Nonoperative treatment versus posterior fixation for thoracolumbar junction burst fractures without neurologic deficit. *Spine (Phila Pa 1976)* 2001;26(9):1038–1045
- Gnanenthiran SR, Adie S, Harris IA. Nonoperative versus operative treatment for thoracolumbar burst fractures without neurologic deficit: a meta-analysis. *Clin Orthop Relat Res* 2012;470(2):567–577
- Wood K, Buttermann G, Mehdob A, Garvey T, Jhanjee R, Sechriest V. Operative compared with nonoperative treatment of a thoracolumbar burst fracture without neurological deficit. A prospective, randomized study. *J Bone Joint Surg Am* 2003;85-A(5):773–781
- Khattak MJ, Syed S, Lakdawala RH. Operative management of unstable thoracolumbar burst fractures. *J Coll Physicians Surg Pak* 2010;20(5):347–349
- Yi L, Jingping B, Gele J, Baolier X, Taixiang W. Operative versus nonoperative treatment for thoracolumbar burst fractures without neurological deficit. *Cochrane Database Syst Rev* 2006;(4):CD005079

- 17 Bailey CS, Urquhart JC, Dvorak MF, et al. Orthosis versus no orthosis for the treatment of thoracolumbar burst fractures without neurologic injury: a multicenter prospective randomized equivalence trial. *Spine J* 2014;14(11):2557–2564
- 18 Cantor JB, Lebowitz NH, Garvey T, Eismont FJ. Nonoperative management of stable thoracolumbar burst fractures with early ambulation and bracing. *Spine (Phila Pa 1976)* 1993;18(8):971–976
- 19 Chan DP, Seng NK, Kaan KT. Nonoperative treatment in burst fractures of the lumbar spine (L2–L5) without neurologic deficits. *Spine (Phila Pa 1976)* 1993;18(3):320–325
- 20 Ağuş H, Kayali C, Arslantaş M. Nonoperative treatment of burst-type thoracolumbar vertebra fractures: clinical and radiological results of 29 patients. *Eur Spine J* 2005;14(6):536–540
- 21 Stadhouders A, Buskens E, Vergroesen DA, Fidler MW, de Nies F, Oner FC. Nonoperative treatment of thoracic and lumbar spine fractures: a prospective randomized study of different treatment options. *J Orthop Trauma* 2009;23(8):588–594
- 22 Hartman MB, Chrin AM, Rehtine GR. Non-operative treatment of thoracolumbar fractures. *Paraplegia* 1995;33(2):73–76
- 23 Moller A, Hasserijs R, Redlund-Johnell I, Ohlin A, Karlsson MK. Nonoperatively treated burst fractures of the thoracic and lumbar spine in adults: a 23- to 41-year follow-up. *Spine J* 2007;7(6):701–707
- 24 Shen WJ, Shen YS. Nonsurgical treatment of three-column thoracolumbar junction burst fractures without neurologic deficit. *Spine (Phila Pa 1976)* 1999;24(4):412–415
- 25 Vaccaro AR, Lehman RA Jr, Hurlbert RJ, et al. A new classification of thoracolumbar injuries: the importance of injury morphology, the integrity of the posterior ligamentous complex, and neurologic status. *Spine (Phila Pa 1976)* 2005;30(20):2325–2333
- 26 Rosenthal MB, Frank RG, Li Z, Epstein AM. Early experience with pay-for-performance: from concept to practice. *JAMA* 2005;294(14):1788–1793
- 27 Doran T, Fullwood C, Gravelle H, et al. Pay-for-performance programs in family practices in the United Kingdom. *N Engl J Med* 2006;355(4):375–384
- 28 Wright JG, Swiontkowski MF, Heckman JD. Introducing levels of evidence to the journal. *J Bone Joint Surg Am* 2003;85-A(1):1–3
- 29 Sullivan D, Chung KC, Eaves FF III, Rohrich RJ. The level of evidence pyramid: indicating levels of evidence in Plastic and Reconstructive Surgery articles. *Plast Reconstr Surg* 2011;128(1):311–314
- 30 [Anonymous]. Levels of Evidence. In: *Clinical Orthopaedics and Related Research*; 2013. Available at: http://www.clinorthop.org/author_and_reviewer_information.html
- 31 Ebell MH, Siwek J, Weiss BD, et al. Simplifying the language of evidence to improve patient care: strength of recommendation taxonomy (SORT): a patient-centered approach to grading evidence in medical literature. *J Fam Pract* 2004;53(2):111–120
- 32 Gopalakrishnan S, Ganeshkumar P. Systematic reviews and meta-analysis: understanding the best evidence in primary healthcare. *J Family Med Prim Care* 2013;2(1):9–14
- 33 Thomas KC, Bailey CS, Dvorak MF, Kwon B, Fisher C. Comparison of operative and nonoperative treatment for thoracolumbar burst fractures in patients without neurological deficit: a systematic review. *J Neurosurg Spine* 2006;4(5):351–358
- 34 Oner FC, Wood KB, Smith JS, Shaffrey CI. Therapeutic decision making in thoracolumbar spine trauma. *Spine (Phila Pa 1976)* 2010;35(21, Suppl):S235–S244
- 35 Boerger TO, Limb D, Dickson RA. Does “canal clearance” affect neurological outcome after thoracolumbar burst fractures? *J Bone Joint Surg Br* 2000;82(5):629–635
- 36 Alpantaki K, Bano A, Pasku D, et al. Thoracolumbar burst fractures: a systematic review of management. *Orthopedics* 2010;33(6):422–429
- 37 Xu GJ, Li ZJ, Ma JX, Zhang T, Fu X, Ma XL. Anterior versus posterior approach for treatment of thoracolumbar burst fractures: a meta-analysis. *Eur Spine J* 2013;22(10):2176–2183
- 38 Higgins JPT, Green S, eds. *Cochrane Handbook for Systematic Reviews of Interventions*, Version 5.1.0 [updated March 2011]. The Cochrane Collaboration, 2011. Available at: www.cochrane-handbook.org
- 39 Howick J, Chalmers I, Glasziou P, et al. The 2011 Oxford CEBM Levels of Evidence (Introductory Document). Available at: <http://www.cebm.net/index.aspx?o=5653>
- 40 Howick J, Chalmers I, Glasziou P, et al. Explanation of the 2011 Oxford Centre for Evidence-Based Medicine (OCEBM) Levels of Evidence (Background Document). Available at: <http://www.cebm.net/index.aspx?o=5653>
- 41 Abudou M, Chen X, Kong X, Wu T. Surgical versus non-surgical treatment for thoracolumbar burst fractures without neurological deficit. *Cochrane Database Syst Rev* 2013;6:CD005079
- 42 Siebenga J, Leferink VJ, Segers MJ, et al. Treatment of traumatic thoracolumbar spine fractures: a multicenter prospective randomized study of operative versus nonsurgical treatment. *Spine (Phila Pa 1976)* 2006;31(25):2881–2890
- 43 Dai LY, Jiang LS, Jiang SD. Posterior short-segment fixation with or without fusion for thoracolumbar burst fractures: a five to seven-year prospective randomized study. *J Bone Joint Surg Am* 2009;91(5):1033–1041
- 44 Jiang XZ, Tian W, Liu B, et al. Comparison of a paraspinous approach with a percutaneous approach in the treatment of thoracolumbar burst fractures with posterior ligamentous complex injury: a prospective randomized controlled trial. *J Int Med Res* 2012;40(4):1343–1356
- 45 Gertzbein SD, Crowe PJ, Fazl M, Schwartz M, Rowed D. Canal clearance in burst fractures using the AO internal fixator. *Spine (Phila Pa 1976)* 1992;17(5):558–560
- 46 Vornanen MJ, Böstman OM, Myllynen PJ. Reduction of bone retracted into the spinal canal in thoracolumbar vertebral body compression burst fractures. A prospective randomized comparative study between Harrington rods and two transpedicular devices. *Spine (Phila Pa 1976)* 1995;20(15):1699–1703
- 47 Alanay A, Acaroglu E, Yazici M, Ozgur A, Surat A. Short-segment pedicle instrumentation of thoracolumbar burst fractures: does transpedicular intracorporeal grafting prevent early failure? *Spine (Phila Pa 1976)* 2001;26(2):213–217
- 48 Stancić MF, Gregorović E, Nozica E, Penezić L. Anterior decompression and fixation versus posterior reposition and semirigid fixation in the treatment of unstable burst thoracolumbar fracture: prospective clinical trial. *Croat Med J* 2001;42(1):49–53
- 49 Tezeren G, Kuru I. Posterior fixation of thoracolumbar burst fracture: short-segment pedicle fixation versus long-segment instrumentation. *J Spinal Disord Tech* 2005;18(6):485–488
- 50 Korovessis P, Baikousis A, Zacharatos S, Petsinis G, Koureas G, Iliopoulos P. Combined anterior plus posterior stabilization versus posterior short-segment instrumentation and fusion for mid-lumbar (L2–L4) burst fractures. *Spine (Phila Pa 1976)* 2006;31(8):859–868
- 51 Dai LY, Jiang LS, Jiang SD. Anterior-only stabilization using plating with bone structural autograft versus titanium mesh cages for two- or three-column thoracolumbar burst fractures: a prospective randomized study. *Spine (Phila Pa 1976)* 2009;34(14):1429–1435
- 52 Pang W, Zhang GL, Tian W, et al. Surgical treatment of thoracolumbar fracture through an approach via the paravertebral muscle. *Orthop Surg* 2009;1(3):184–188
- 53 Tezeren G, Bulut O, Tukenmez M, Ozturk H, Ozturk Z, Ozturk A. Long segment instrumentation of thoracolumbar burst fracture: fusion versus nonfusion. *J Back Musculoskeletal Rehabil* 2009;22(2):107–112
- 54 Wei FX, Liu SY, Liang CX, et al. Transpedicular fixation in management of thoracolumbar burst fractures: monosegmental fixation versus short-segment instrumentation. *Spine (Phila Pa 1976)* 2010;35(15):E714–E720

- 55 Jindal N, Sankhala SS, Bachhal V. The role of fusion in the management of burst fractures of the thoracolumbar spine treated by short segment pedicle screw fixation: a prospective randomised trial. *J Bone Joint Surg Br* 2012;94(8):1101–1106
- 56 Schmid R, Lindtner RA, Lill M, Blauth M, Krappinger D, Kammerlander C. Combined posteroanterior fusion versus transforaminal lumbar interbody fusion (TLIF) in thoracolumbar burst fractures. *Injury* 2012;43(4):475–479
- 57 Wood KB, Bohn D, Mehbod A. Anterior versus posterior treatment of stable thoracolumbar burst fractures without neurologic deficit: a prospective, randomized study. *J Spinal Disord Tech* 2005;18 (Suppl):S15–S23