



Long-term associations of PM₁ versus PM_{2.5} and PM₁₀ with asthma and asthma-related respiratory symptoms in the middle-aged and elderly population

Xue-yan Zheng^{1,9}, Shu-jun Guo^{2,9}, Jian-xiong Hu^{3,9}, Rui-lin Meng¹, Yan-jun Xu¹, Yun-hong Lv¹, Ye Wang¹, Ni Xiao¹, Chuan Li¹, Xiao-jun Xu¹, De-jian Zhao¹, Hong-ye Zhou⁴, Jia-hui He², Xiao-min Tan⁴, Jing Wei⁵, Li-feng Lin^{1,6,10} and Wei-jie Guan^{2,7,8,10}

¹Guangdong Provincial Center for Disease Control and Prevention, Guangzhou, China. ²State Key Laboratory of Respiratory Disease, National Clinical Research Center for Respiratory Disease, Department of Respiratory and Critical Care Medicine, Guangzhou Institute of Respiratory Health, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China. ³Department of Public Health and Preventive Medicine, School of Medicine, Jinan University, Guangzhou, China. ⁴Department of Epidemiology and Biostatistics, School of Public Health, Guangdong Pharmaceutical University, Guangzhou, China. ⁵Department of Atmospheric and Oceanic Science, Earth System Science Interdisciplinary Center, University of Maryland, College Park, MD, USA. ⁶School of Public Health, Southern Medical University, Guangzhou, China. ⁷Department of Thoracic Surgery, Guangzhou Institute of Respiratory Health, The First Affiliated Hospital of Guangzhou Medical University, Guangzhou, China. ⁸Guangzhou National Laboratory, Guangzhou, China. ⁹Xue-yan Zheng, Shu-jun Guo and Jian-xiong Hu contributed equally to this article as joint first authors. ¹⁰Li-feng Lin and Wei-jie Guan contributed equally to this article as lead authors and supervised the work.

Corresponding author: Wei-jie Guan (battery203@163.com)



Shareable abstract (@ERSpublications)

Long-term exposure to particulate matter increases the risks of asthma and respiratory symptoms in the middle-aged and elderly population <https://bit.ly/3wE05CB>

Cite this article as: Zheng X-y, Guo S-j, Hu J-x, *et al.* Long-term associations of PM₁ versus PM_{2.5} and PM₁₀ with asthma and asthma-related respiratory symptoms in the middle-aged and elderly population. *ERJ Open Res* 2024; 10: 00972-2023 [DOI: 10.1183/23120541.00972-2023].

Copyright ©The authors 2024

This version is distributed under the terms of the Creative Commons Attribution Non-Commercial Licence 4.0. For commercial reproduction rights and permissions contact permissions@ersnet.org

Received: 4 Dec 2023
Accepted: 25 Feb 2024

Abstract

Background Few studies have compared the associations between long-term exposures to particulate matters (aerodynamic diameter ≤ 1 , ≤ 2.5 and ≤ 10 μm : PM₁, PM_{2.5} and PM₁₀, respectively) and asthma and asthma-related respiratory symptoms. The objective of the present study was to compare the strength of the aforementioned associations in middle-aged and elderly adults.

Methods We calculated the mean 722-day personal exposure estimates of PM₁, PM_{2.5} and PM₁₀ at 1 km \times 1 km spatial resolution between 2013 and 2019 at individual levels from China High Air Pollutants (CHAP) datasets. Using logistic regression models, we presented the associations as odds ratios and 95% confidence intervals, for each interquartile range (IQR) increase in PM₁/PM_{2.5}/PM₁₀ concentration. Asthma denoted a self-reported history of physician-diagnosed asthma or wheezing in the preceding 12 months.

Results We included 7371 participants in COPD surveillance from Guangdong, China. Each IQR increase in PM₁, PM_{2.5} and PM₁₀ was associated with a greater odds (OR (95% CI)) of asthma (PM₁: 1.22 (1.02–1.45); PM_{2.5}: 1.24 (1.04–1.48); PM₁₀: 1.30 (1.07–1.57)), wheeze (PM₁: 1.27 (1.11–1.44); PM_{2.5}: 1.30 (1.14–1.48); PM₁₀: 1.34 (1.17–1.55)), persistent cough (PM₁: 1.33 (1.06–1.66); PM_{2.5}: 1.36 (1.09–1.71); PM₁₀: 1.31 (1.02–1.68)) and dyspnoea (PM₁: 2.10 (1.84–2.41); PM_{2.5}: 2.17 (1.90–2.48); PM₁₀: 2.29 (1.96–2.66)). Sensitivity analysis results were robust after excluding individuals with a family history of allergy. Associations of PM₁, PM_{2.5} and PM₁₀ with asthma and asthma-related respiratory symptoms were slightly stronger in males.

Conclusion Long-term exposure to PM is associated with increased risks of asthma and asthma-related respiratory symptoms.

Introduction

Air pollution, an important environmental component, has been associated with blunted therapeutic responses to inhaled corticosteroids, more frequent use of healthcare resources and a poorer control of chronic respiratory diseases, including asthma and asthma-respiratory symptoms [1–8].



Accumulating evidence has demonstrated that exposure to ambient particulate matters (PMs) with an aerodynamic diameter ≤ 2.5 and ≤ 10 μm ($\text{PM}_{2.5}$ and PM_{10} , respectively) was adversely associated with respiratory health, including asthma and asthma-related symptoms [8–12], despite inconsistent conclusions. This might be, at least partially, interpreted by the regional differences in the compositions and concentrations of PMs [10–12]. Furthermore, previous studies were mainly conducted in Europe and America where there were comparatively low exposure levels. Studies regarding the associations between air pollution and asthma and asthma-related respiratory symptoms in higher exposure settings (such as mainland China) have been scarce [13], where there was rapid population growth and industrial development and urbanisation. Guangdong, one of the most developed provinces of China, is characterised by lower concentrations of PMs with more toxic components [14, 15]. Therefore, it would be worthwhile to explore the associations between air pollutants and asthma and asthma-related respiratory symptoms in Guangdong province.

Ambient PMs, including PM with an aerodynamic diameter ≤ 1 μm (PM_1) [16], $\text{PM}_{2.5}$ [17] and PM_{10} , are major drivers of urban air pollution. PM_1 is smaller than $\text{PM}_{2.5}$ and PM_{10} , and is characterised by a higher surface area to volume ratio, which might elicit greater adverse impact on the respiratory tract and other health outcomes [18, 19]. PM_1 conferred greater cellular toxicity and elicited more notable inflammation compared with $\text{PM}_{2.5}$ and PM_{10} [20]. The association between asthma and asthma-related respiratory symptoms and PM_1 (the smaller and potentially more toxic PM) in China has not been well characterised, except for a single study focusing on children and adolescents in Northeast China [13].

We conducted a population-based study of 7371 middle-aged and elderly residents in Guangdong province. We compared the strength of associations of long-term ambient exposure to PM_1 versus $\text{PM}_{2.5}$ and PM_{10} on asthma and asthma-related respiratory symptoms, and identified the more vulnerable subpopulations (e.g. males).

Methods

Study design and participants

We employed a multistage, probability-based sampling strategy in six surveillance points: Panyu district (Guangzhou city), Doumen district (Zhuhai city), Yingde (Qingyuan city), Taishan (Jiangmen city), Luoding (Yunfu city) and Wengyuan county (Shaoguan city). We included adults who were aged ≥ 40 years, and who had participated in the 2014–2015 and 2019–2020 COPD surveillance in Guangdong [21, 22]. We combined the data from the two surveys as a larger cross-sectional study. Briefly, we included residents who had been living in their current residence for >6 months within the year of the survey. We excluded residents with paraplegia, malignancy, cognitive defects, linguistic or mental disorders and pregnant or breastfeeding women. Data were collected during an interview in a healthcare facility by trained staff from the local health stations or community clinics.

The study protocol was approved by the Ethics Review Committee of the National Center for Chronic and Noncommunicable Disease Control and Prevention, China Center for Disease Control and Prevention (201410 and 201901). All participants provided written informed consent.

Ambient and personal air pollution assessment

Daily concentrations of ambient PM_1 , $\text{PM}_{2.5}$ and PM_{10} at 1 km \times 1 km (km 2) spatial resolution between 2013 and 2019 were obtained from the China High Air Pollutants (CHAP) datasets. Home addresses of all participants were geocoded using the Baidu Maps API (<https://lbsyun.baidu.com>) for matching the geocodes of each pollutant from the CHAP datasets. The first participant was recruited on 23 December 2014, while the concentration of air pollutants could be obtained on the earliest date (1 January 2013) by matching the geocode in the CHAP datasets.

In light of the fact that long-term exposure has been conventionally defined among participants who have been exposed to air pollutants for >1 year [23–25], the total exposure window was extended to 722 days (the longest duration within the CHAP database), which allowed us to investigate the associations between the 722-day averaged exposure levels of PMs and asthma and asthma-related respiratory symptoms. Details of PM_1 , $\text{PM}_{2.5}$ and PM_{10} estimates have been described previously [26–28]. Briefly, a space-time extremely randomised tree model was developed to generate the daily time-series dataset of PM_1 , $\text{PM}_{2.5}$ and PM_{10} concentrations at 1 km 2 grids from 2013 to 2019 in China. The data (model input) mainly included satellite remote sensing (MAIAC AOD), Multiresolution Emission Inventory for China (MEIC) pollution emissions, meteorological characteristics, land use and urban form (land use, road, population), and spatiotemporal terms. Hence, a time-series and full-coverage dataset of PM_1 , $\text{PM}_{2.5}$ and PM_{10} at 1 km 2 cell grids from 2013 to 2019 was generated. The spatiotemporal terms were included to capture the

spatiotemporal autocorrelation of PM_{10} , $PM_{2.5}$ and PM_{10} concentrations. Based on 10-fold cross-validation [26–28], greater consistency was observed between the estimated PM_{10} (coefficient of determination (R^2) = 0.83, root-mean-square error (RMSE) = $6.2 \mu\text{g}\cdot\text{m}^{-3}$), $PM_{2.5}$ (R^2 = 0.92, RMSE = $10.8 \mu\text{g}\cdot\text{m}^{-3}$) and PM_{10} (R^2 = 0.90, RMSE = $21.13 \mu\text{g}\cdot\text{m}^{-3}$) and ground-level measurements at a daily scale. Geocoding was performed at each residential location of study participants to estimate the exposure concentration of PMs, which allowed for the calculation of the 722-day averaged concentration prior to the survey for each participant.

For other criteria pollutants, the daily concentrations of ambient sulfur dioxide (SO_2), nitrogen dioxide (NO_2), carbon monoxide (CO) and ozone (O_3) at 10 km^2 grids between 2013 and 2019 were also obtained from the CHAP datasets. As reported previously [26–31], the consistency between the estimated SO_2 , NO_2 , CO and O_3 data and ground-based measurements was high.

Clinical assessments

Respiratory symptoms (wheeze, persistent cough, dyspnoea and persistent phlegm) were captured from the Epidemiologic Standardization Project Questionnaire of the American Thoracic Society (ATS-DLD-78-A) [32], which has been extensively validated.

Participants were considered as having wheezing if they had a positive response to “Does your chest ever sound wheezy or whistling, including the time when you had a cold?” [13].

Persistent cough denoted coughing on most days (≥ 4 days per week) for ≥ 3 months annually in the absence or presence of cold (positive responses to several items on cough) [13].

Persistent phlegm denoted phlegm, sputum or mucus brought up from the chest in the morning on most days (≥ 4 days per week) for ≥ 3 months of the previous year in the absence or presence of cold (positive responses to several items on phlegm) [13].

The questionnaire inquired about shortness of breath during daily activity and its severity based on the modified Medical Research Council (mMRC) dyspnoea scale. Only 8.2% of participants had a documented mMRC dyspnoea scale because it was only applied to the participants with COPD. To account for the heterogeneity of participants and study-specific assessment items, the harmonised dyspnoea symptom variables were defined as “shortness of breath during daily life activity” [33].

Asthma was defined on the basis of a self-reported history of asthma diagnosis by physicians or by the presence of wheezing in the preceding 12 months, which has also been adopted in the China Pulmonary Health (CPH) Study [34]. The European Community Respiratory Health Survey (ECRHS) asthma questionnaire served as the source of reference for this definition [35].

A family history of allergic rhinitis or asthma was defined as any biological parent who had ever been diagnosed as having allergic rhinitis or asthma. If participants answered “Yes” to a family history of allergic rhinitis or asthma, then she/he would be defined as having a family history of allergy (supplementary table E1) [36].

Covariates

A standardised questionnaire was employed to examine the potential confounding and mediating effects. We captured the following variables from the questionnaire: sex (male, female), age, region (urban, rural), educational level (primary school or lower, middle school, university or higher), marital status (unmarried, married, widowed/divorced), biomass fuel exposure (yes, no), household ventilation (yes, no), smoking status (yes, no), passive smoking (yes, no), occupational exposure (yes, no), family history of allergy (yes, no) and body mass index (BMI) category (underweight, normal weight, overweight, obesity). The definition of the covariates is summarised in supplementary table E2 [22, 37].

Statistical analysis

The odds ratios and 95% confidence intervals were calculated to determine the association between each interquartile range (IQR) change in ambient $PM_{10}/PM_{2.5}/PM_{10}$ concentration and asthma and asthma-related respiratory symptoms by using logistic regression models [6]. We compared the Akaike Information Criterion value of three models to avoid over-fitting. The magnitude of collinearity was assessed based on the variance inflation factor (VIF). $VIF \geq 5$ indicated collinearity among the variables. Variables with a significant collinearity were excluded. Finally, we included age, sex, education, marital status, region, occupational exposure, household ventilation, passive smoking, biomass fuel exposure, smoking status,

family history of allergy and BMI categories to adjust for confounding. The statistical significance for the difference between the groups was examined by using a two-sample z-test, whose 95% confidence interval was calculated as: $(\beta_1 - \beta_2) \pm 1.96 \sqrt{(SE_1^2 + SE_2^2)}$, where β_1 and β_2 are the coefficient estimates from the model for each stratum and SE_1 and SE_2 are the standard errors.

Spearman's rank correlation test was used to determine the association of air pollutants. Strong, moderate and weak correlations were defined as coefficients (r_s) >0.60 , $0.30-0.60$ and <0.30 , respectively. In light of the moderate-to-high correlation between $PM_1/PM_{2.5}/PM_{10}$ and other pollutant models (except for O_3), we applied a two-pollutant model of $PM_1/PM_{2.5}/PM_{10}$ and O_3 to exclude the inclusion of parameters with a significant collinearity.

For subgroup analysis, we further explored the effect modification according to sex. To evaluate the robustness of our estimates, we conducted the sensitivity analyses by excluding participants with a family history of allergy.

Statistical analyses were performed with R version 4.2.2 (<https://cran.r-project.org>). The threshold of statistical significance for main effects and interactions was set at $p < 0.05$.

Results

Participant characteristics

Of 7510 participants screened, 139 had ineligible data (three did not answer the symptom-related questions and 136 did not report height). Therefore, 7371 participants were finally included, whose characteristics are presented in table 1. The mean age was 57.26 years and 3764 participants (51.06%) were males. The overall prevalence of wheeze (9.96% versus 7.37%), persistent cough (4.33% versus 1.19%) and persistent phlegm (7.62% versus 2.27%) was significantly higher in males than in females (all $p < 0.05$). The overall prevalence of asthma (4.57% versus 3.91%) and dyspnoea (8.55% versus 7.98%) did not differ significantly between males and females (both $p > 0.05$).

Air pollutant concentrations

The mean daily ambient concentration of PM_1 , $PM_{2.5}$, PM_{10} from each surveillance point is presented in table 2. The mean daily ambient concentration of PM_1 , $PM_{2.5}$ and PM_{10} was 20.85, 34.92 and $54.35 \mu\text{g}\cdot\text{m}^{-3}$, respectively. The IQR of ambient PM_1 , $PM_{2.5}$ and PM_{10} concentrations was 6.72, 10.81 and $16.39 \mu\text{g}\cdot\text{m}^{-3}$, respectively. Overall, the concentration of PM_1 , $PM_{2.5}$ and PM_{10} yielded a medium-to-high correlation with that of the other pollutants, with r_s ranging from 0.40 to 0.99, except for the weak correlation between O_3 and PM_1 , $PM_{2.5}$ and PM_{10} (figure 1). The $PM_1:PM_{2.5}$ and $PM_1:PM_{10}$ ratios at the residential address were 59.69% and 38.36%, respectively. There was a strong positive correlation between PM_1 and $PM_{2.5}$ (Spearman's γ correlation coefficient (γ_s)=0.99) and PM_{10} (γ_s =0.96).

Associations between PMs and asthma and asthma-related respiratory symptoms

Based on the single-pollutant model, each IQR increase in the 722-day mean exposure of PM_1 , $PM_{2.5}$ and PM_{10} was associated with a significantly higher odds (OR (95% CI)) of having asthma (PM_1 : 1.22 (1.02–1.45); $PM_{2.5}$: 1.24 (1.04–1.48); PM_{10} : 1.30 (1.07–1.57)), wheeze (PM_1 : 1.27 (1.11–1.44); $PM_{2.5}$: 1.30 (1.14–1.48); PM_{10} : 1.34 (1.17–1.55)), persistent cough (PM_1 : 1.33 (1.06–1.66); $PM_{2.5}$: 1.36 (1.09–1.71); PM_{10} : 1.31 (1.02–1.68)) and dyspnoea (PM_1 : 2.10 (1.84–2.41); $PM_{2.5}$: 2.17 (1.90–2.48); PM_{10} : 2.29 (1.96–2.66)) in the overall population. We did not identify significant positive associations between $PM_1/PM_{2.5}/PM_{10}$ exposures and persistent phlegm (table 3). The strengths of associations between the 722-day means of PMs concentration and asthma and asthma-related respiratory symptoms in single-pollutant models and two-pollutant models were similar with minor differences. Therefore, we present the results of single-pollutant models only hereafter. Overall, the associations of PM_1 , $PM_{2.5}$ and PM_{10} with asthma and asthma-related respiratory symptoms were comparable. When stratified by sex, the associations were slightly stronger in males (table 4). The plots of the exposure–response relationships of PM_1 , $PM_{2.5}$ and PM_{10} exposure with asthma and asthma-related respiratory symptoms are shown in supplementary figure E1.

Sensitivity analyses

Overall, we found similar results when excluding participants with a family history of allergy in sensitivity analyses (supplementary tables E3 and E4, and figure 2). There were significant associations between $PM_1/PM_{2.5}/PM_{10}$ exposures and asthma, wheeze and dyspnoea in the participants without a family history of allergy. Stratified analyses among individuals with no family history of allergy showed that the associations of asthma and dyspnoea were slightly stronger in males.

TABLE 1 Baseline characteristics of the study participants

	All participants (n=7371)	Males (n=3764)	Females (n=3607)
Age (years)[#]	57.26±9.26	57.90±9.35	56.58±9.12
Educational level[#]			
Primary school or lower	4231 (57.40)	1756 (46.65)	2475 (68.62)
Middle school	2989 (40.55)	1901 (50.50)	1088 (30.16)
University or higher	151 (2.05)	107 (2.84)	44 (1.22)
Marriage status[#]			
Unmarried	94 (1.28)	68 (1.81)	26 (0.72)
Married	6720 (91.17)	3536 (93.94)	3184 (88.27)
Widowed/divorced	557 (7.56)	160 (4.25)	397 (11.01)
Region of residence[#]			
Urban	2745 (37.24)	1291 (34.30)	1454 (40.31)
Rural	4626 (62.76)	2473 (65.70)	2153 (59.69)
Occupational exposure[#]			
Yes	3676 (49.87)	2009 (53.37)	1667 (46.22)
No	3695 (50.13)	1755 (46.63)	1940 (53.78)
Household ventilation[#]			
Yes	5835 (79.16)	2893 (76.86)	2942 (81.56)
No	1536 (20.84)	871 (23.14)	665 (18.44)
Passive smoking[#]			
Yes	4200 (56.98)	2256 (59.94)	1944 (53.90)
No	3171 (43.02)	1508 (40.06)	1663 (46.10)
Biomass fuel exposure[#]			
Yes	2738 (37.15)	1331 (35.36)	1407 (39.01)
No	4633 (62.85)	2433 (64.64)	2200 (60.99)
Smoking status[#]			
Yes	3183 (43.18)	3107 (82.55)	76 (2.11)
No	4188 (56.82)	657 (17.45)	3531 (97.89)
BMI category[#]			
Underweight	295 (4.00)	165 (4.38)	130 (3.60)
Normal	3684 (49.98)	2035 (54.06)	1649 (45.72)
Overweight	2547 (34.55)	1212 (32.20)	1335 (37.01)
Obesity	845 (11.46)	352 (9.35)	493 (13.67)
Family history of allergy[#]			
Yes	502 (6.81)	300 (7.97)	202 (5.60)
No	6869 (93.19)	3464 (92.03)	3405 (94.40)
Asthma			
Yes	313 (4.25)	172 (4.57)	141 (3.91)
No	7058 (95.75)	3592 (95.43)	3466 (96.09)
Wheeze[#]			
Yes	641 (8.70)	375 (9.96)	266 (7.37)
No	6730 (91.30)	3389 (90.04)	3341 (92.63)
Persistent cough[#]			
Yes	206 (2.79)	163 (4.33)	43 (1.19)
No	7165 (97.21)	3601 (95.67)	3564 (98.81)
Persistent phlegm[#]			
Yes	369 (5.01)	287 (7.62)	82 (2.27)
No	7002 (94.99)	3477 (92.38)	3525 (97.73)
Dyspnoea			
Yes	610 (8.28)	322 (8.55)	288 (7.98)
No	6761 (91.72)	3442 (91.45)	3319 (92.02)

Data are presented as mean±SD or n (%). BMI: body mass index. #: significant difference between males and females (p<0.05), tested with the Chi-squared test for categorical variables and the t-test for continuous variables.

Discussion

This is the first study which has compared the associations of long-term exposure of PM₁ versus PM_{2.5} and PM₁₀ with asthma and asthma-related respiratory symptoms in the middle-aged and elderly population in China. Our study has added to the existing evidence by demonstrating robust associations between

TABLE 2 Descriptive statistics of particulate matter concentrations stratified by surveillance points

	Mean±sd	Median	Minimum	Maximum	IQR
Panyu, Guangzhou (n=1230)					
PM ₁ (μg·m ⁻³)	23.49±5.43	18.97	17.44	31.54	10.88
PM _{2.5} (μg·m ⁻³)	40.09±7.94	32.74	31.87	48.74	15.95
PM ₁₀ (μg·m ⁻³)	61.59±9.42	52.92	51.03	73.05	18.41
Doumen, Zhuhai (n=1184)					
PM ₁ (μg·m ⁻³)	18.86±2.92	17.50	15.36	22.99	5.34
PM _{2.5} (μg·m ⁻³)	32.09±4.51	28.07	27.19	37.07	9.00
PM ₁₀ (μg·m ⁻³)	52.58±10.72	44.18	39.50	66.42	22.85
Yingde, Qingyuan (n=1231)					
PM ₁ (μg·m ⁻³)	22.37±4.72	18.15	17.36	28.24	9.38
PM _{2.5} (μg·m ⁻³)	37.74±7.89	30.58	29.62	47.34	15.72
PM ₁₀ (μg·m ⁻³)	56.88±11.07	47.13	45.18	70.88	21.98
Taishan, Jiangmen (n=1353)					
PM ₁ (μg·m ⁻³)	19.84±3.23	17.98	14.96	24.81	6.42
PM _{2.5} (μg·m ⁻³)	32.58±5.03	29.22	24.13	41.04	9.90
PM ₁₀ (μg·m ⁻³)	51.61±7.28	47.67	39.83	63.70	13.83
Luoding, Yunfu (n=1210)					
PM ₁ (μg·m ⁻³)	20.11±3.01	22.59	15.89	23.74	5.78
PM _{2.5} (μg·m ⁻³)	33.34±4.91	37.31	26.67	39.23	9.21
PM ₁₀ (μg·m ⁻³)	51.91±6.80	57.21	41.97	59.78	13.70
Wengyuan, Shaoguan (n=1163)					
PM ₁ (μg·m ⁻³)	20.39±4.34	17.05	15.31	25.50	8.59
PM _{2.5} (μg·m ⁻³)	33.75±6.71	28.47	26.13	42.09	13.12
PM ₁₀ (μg·m ⁻³)	51.56±9.04	44.30	41.08	63.04	16.69

PM₁, PM_{2.5} and PM₁₀: particulate matter with an aerodynamic diameter ≤1, ≤2.5 and ≤10 μm, respectively; IQR: interquartile range.

long-term exposure to ambient PM₁, PM_{2.5} and PM₁₀ and asthma and asthma-related respiratory symptoms in Guangdong province. The associations were slightly stronger in males than females.

Our findings have further complemented the studies which demonstrated the detrimental associations of PM_{2.5} exposure with respiratory health [8–12]. Per-IQR-increase analyses have been essential pertaining to




FIGURE 1 Pairwise correlation (r_s) of different ambient air pollutants. PM₁, PM_{2.5} and PM₁₀: particulate matter with an aerodynamic diameter ≤1, ≤2.5 and ≤10 μm, respectively; SO₂: sulfur dioxide; NO₂: nitrogen dioxide; O₃: ozone; CO: carbon monoxide.

TABLE 3 Associations between per interquartile range increase in particulate matter with an aerodynamic diameter ≤ 1 , ≤ 2.5 and $\leq 10 \mu\text{m}$ (PM_{10} , $\text{PM}_{2.5}$ and PM_{10} , respectively) and asthma and asthma-related respiratory symptoms for the whole study population (n=7371 for all analyses)

	Single-pollutant model			Two-pollutant model ($\text{PM}+\text{O}_3$)		
	OR (95% CI)	AIC	Adjusted p-value	OR (95% CI)	AIC	Adjusted p-value
Asthma						
PM_1	1.22 (1.02–1.45)	2523.99	0.03*	1.20 (1.00–1.44)	2525.70	0.05*
$\text{PM}_{2.5}$	1.24 (1.04–1.48)	2523.00	0.01*	1.23 (1.03–1.47)	2524.68	0.02*
PM_{10}	1.30 (1.07–1.57)	2521.71	0.008**	1.28 (1.06–1.56)	2523.33	0.01*
Wheeze						
PM_1	1.27 (1.11–1.44)	4156.12	<0.001***	1.25 (1.10–1.43)	4157.73	<0.001***
$\text{PM}_{2.5}$	1.30 (1.14–1.48)	4153.10	<0.001***	1.29 (1.13–1.47)	4154.68	<0.001***
PM_{10}	1.34 (1.17–1.55)	4151.66	<0.001***	1.33 (1.15–1.53)	4153.05	<0.001***
Persistent cough						
PM_1	1.33 (1.06–1.66)	1770.09	0.01*	1.29 (1.02–1.63)	1771.36	0.03*
$\text{PM}_{2.5}$	1.36 (1.09–1.71)	1768.79	0.007**	1.33 (1.06–1.68)	1770.03	0.01*
PM_{10}	1.31 (1.02–1.68)	1771.40	0.03*	1.28 (0.99–1.64)	1772.27	0.06
Persistent phlegm						
PM_1	1.07 (0.91–1.27)	2780.95	0.42	1.06 (0.89–1.26)	2782.53	0.55
$\text{PM}_{2.5}$	1.09 (0.93–1.29)	2780.51	0.29	1.08 (0.91–1.28)	2782.12	0.38
PM_{10}	1.10 (0.92–1.31)	2780.60	0.32	1.08 (0.90–1.30)	2782.15	0.39
Dyspnoea						
PM_1	2.10 (1.84–2.41)	3933.66	<0.001***	1.98 (1.73–2.27)	3921.06	<0.001***
$\text{PM}_{2.5}$	2.17 (1.90–2.48)	3923.48	<0.001***	2.05 (1.79–2.35)	3909.25	<0.001***
PM_{10}	2.29 (1.96–2.66)	3935.27	<0.001***	2.15 (1.84–2.51)	3917.53	<0.001***

AIC: Akaike Information Criterion; O_3 : ozone. The model was adjusted by age, educational level, marital status, body mass index category, region, passive smoking, occupational exposure, household ventilation, biomass fuel exposure, smoking status, sex and family history of allergy. *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$.

TABLE 4 Associations between per interquartile range increase in particulate matter with an aerodynamic diameter ≤ 1 , ≤ 2.5 and $\leq 10 \mu\text{m}$ (PM_{10} , $\text{PM}_{2.5}$ and PM_{10} , respectively) and asthma and asthma-related respiratory symptoms among males and females

	Males (n=3764)			Females (n=3607)		
	OR (95% CI)	AIC	Adjusted p-value	OR (95% CI)	AIC	Adjusted p-value
Asthma						
PM_1	1.32 (1.04–1.68)	1358.67	0.02*	1.12 (0.86–1.45)	1184.86	0.41
$\text{PM}_{2.5}$	1.31 (1.03–1.66)	1359.11	0.03*	1.18 (0.91–1.53)	1184.03	0.22
PM_{10}	1.40 (1.07–1.81)	1357.60	0.01*	1.20 (0.91–1.60)	1183.92	0.20
Wheeze						
PM_1	1.28 (1.08–1.52)	2343.80	0.004**	1.25 (1.03–1.52)	1827.40	0.02*
$\text{PM}_{2.5}$	1.30 (1.10–1.54)	2342.97	0.003**	1.31 (1.08–1.59)	1825.21	0.007**
PM_{10}	1.37 (1.14–1.65)	2340.57	<0.001***	1.31 (1.06–1.62)	1826.21	0.01*
Persistent cough						
PM_1	1.35 (1.05–1.75)	1313.56	0.02*	1.22 (0.75–1.98)	466.35	0.43
$\text{PM}_{2.5}$	1.38 (1.07–1.78)	1312.78	0.01*	1.30 (0.80–2.10)	465.87	0.29
PM_{10}	1.34 (1.01–1.77)	1314.70	0.04*	1.21 (0.72–2.06)	466.45	0.47
Persistent phlegm						
PM_1	1.03 (0.84–1.25)	1993.21	0.78	1.26 (0.91–1.75)	794.51	0.17
$\text{PM}_{2.5}$	1.04 (0.86–1.26)	1993.12	0.68	1.32 (0.95–1.84)	793.71	0.10
PM_{10}	1.04 (0.84–1.28)	1993.17	0.73	1.37 (0.95–1.97)	793.58	0.10
Dyspnoea						
PM_1	2.23 (1.85–2.70)	2054.39	<0.001***	2.01 (1.66–2.43)	1890.82	<0.001***
$\text{PM}_{2.5}$	2.23 (1.85–2.69)	2053.06	<0.001***	2.13 (1.76–2.58)	1882.12	<0.001***
PM_{10}	2.44 (1.97–3.03)	2054.38	<0.001***	2.16 (1.74–2.69)	1892.03	<0.001***

AIC: Akaike Information Criterion. The model was adjusted by age, educational level, marital status, body mass index category, region, passive smoking, occupational exposure, household ventilation, biomass fuel exposure, smoking status and family history of allergy. *: $p < 0.05$; **: $p < 0.01$; ***: $p < 0.001$.




FIGURE 2 Sensitivity analysis on the associations between particulate matter with an aerodynamic diameter ≤ 1 , ≤ 2.5 and $\leq 10 \mu\text{m}$ (PM₁, PM_{2.5} and PM₁₀, respectively) and asthma and asthma-related respiratory symptoms among participants without a family history of allergy: a) asthma, b) wheeze, c) persistent cough, d) persistent phlegm and e) dyspnoea. The associations were adjusted for age, sex, educational level, marriage status, body mass index, region, passive smoking, occupational exposure, ventilation, biomass fuel exposure and smoking status. Adjusted odds ratios and 95% confidence intervals were estimated per interquartile range increase in PM₁, PM_{2.5} and PM₁₀ for asthma and asthma-related respiratory symptoms.

air pollutants such as PMs in the published epidemiological studies. The rationale is that PM₁ is the essential component of both PM_{2.5} and PM₁₀, and that it would be impossible under any real-life conditions for an individual to be exposed to an equal concentration of all three PMs. We have analysed the association per IQR increase in concentration of the respective pollutant, showing that the associations of PM₁, PM_{2.5} and PM₁₀ were comparable. The association of the respective pollutant per $10 \mu\text{g}\cdot\text{m}^{-3}$ increase also supported a similar conclusion (supplementary tables E5–E7). This finding was consistent with the results among children and adolescents residing in 33 communities in cities of Northeast China, in which the adjusted OR (95% CI) of asthma and asthma-related respiratory symptoms per $10 \mu\text{g}\cdot\text{m}^{-3}$ increase in PM₁ (asthma: 1.56 (1.46–1.66)) was similar to PM_{2.5} (asthma: 1.50 (1.41–1.59)) [13]. However, the drivers of the associations for PM₁, PM_{2.5} and PM₁₀ were notably different between the two studies. Consistent with most previous studies in China [38–40], PM₁ accounted for >85% and >50% for the ambient PM_{2.5} and PM₁₀ in seven cities in Northeast China [13] and Beijing [39]. Therefore, PM₁ was an essential driver of the PM pollution, accounting for the main components and concentrations of PM_{2.5} and PM₁₀ in most areas (more than 26 cities) across mainland China [38–40]. However, the PM₁:PM_{2.5} (59.69%) and PM₁:PM₁₀ (38.36%) ratios in our study were similar to those reported in other European countries (PM₁:PM_{2.5} ratio: 48–69%; PM₁:PM_{2.5} ratio: 40%) [41, 42], but remained much lower compared with other areas in China. We posited that the adverse health effects of PM_{2.5} and PM₁₀ in Guangdong province could be partially attributed to PM₁ [38]. Additional studies comparing the associations of PM₁, PM_{2.5} and PM₁₀ (particularly their specific components) are needed to shed light on how the different PM sizes and compositions could have affected asthma and asthma-related respiratory symptoms in humans.

We have identified slightly stronger associations between PM concentrations and asthma and asthma-related respiratory symptoms in males than in females. Consistent with the study focusing on children in Northeast China, stronger associations between PM concentration and asthma and asthma-related respiratory symptoms were observed among boys compared with girls [13]. A study of 12 Southern California communities has evaluated the associations between different ambient air pollutant concentrations and asthma [43]. The study showed that asthma-related symptoms among 3676 children correlated positively with PM levels in boys, but not in girls. In three studies reporting a greater impact of

air pollution among males, the higher levels of work-related exposures and the more time spent outdoors might help collectively explain the higher likelihood of developing asthma and asthma-related respiratory symptoms [13, 44, 45]. Other sex-related traits, such as hormonal status and body size, could have influenced the biological transportation of environmental chemicals, potentially leading to different associations between males and females [9, 46].

Notably, PM exposures did not significantly increase the risk of asthma among participants with a family history of allergy. The discordant findings between participants with *versus* without a family history of allergy could be partly interpreted by the non-equivalence between the history of documented diseases and the currently symptomatic diseases. Participants with a family history of allergy might be more likely to avoid contact with risk factors of allergic diseases and more proactively seek prescription refills to maintain asthma or allergic symptom control. Identification of the underlying causes, however, would benefit from more in-depth analysis *via* longitudinal assessments within our study setting.

Although our study did not aim to identify the underlying mechanisms of the noxious effects of PMs on asthma and asthma-related respiratory symptoms, the systemic inflammation, oxidative stress responses, immune injury, disease-associated gene polymorphisms and DNA methylation, regulation of the gene expression, and altered signalling pathways associated with oxidative stress responses might collectively help explain our findings [47–52]. First, compared with PM_{2.5} and PM₁₀, the smaller diameter of PM₁ could elicit inflammatory responses *via* inducing oxidative stress after internalisation by airway epithelial cells [53, 54]. PM₁ could penetrate deeper to the more distal airways and be better adsorbed to the cellular surface, resulting in similar levels of toxicity despite the lower concentration. Second, the effects of airborne PMs on lung function might depend primarily on the pro-inflammatory responses [55, 56]. Third, PMs may also absorb aeroallergens released by pollen grains, partially contributing to the enhanced IgE antibody-mediated immune response with the prolonged retention of the allergens [57]. Moreover, both B-cells and T-cells could sense intercellular signals and utilise cellular and humoral immunity to induce potent inflammatory responses. Studies have shown that PMs can cause an imbalance in Th1/Th2 T-helper cell ratios and functional dysregulation by skewing towards the Th2 inflammatory responses and inhibiting Th1 inflammatory responses [18, 58].

Some limitations should be considered. First, the causality between long-term exposure to ambient PMs and asthma and asthma-related respiratory symptoms cannot be inferred because of the cross-sectional study design. Second, the definitions of self-reported asthma and asthma-related respiratory symptoms could only be regarded as a proxy for the typical clinical diagnosis; however, these definitions were consistent with those adopted in major large-scale epidemiological studies [8, 13, 32, 34, 35, 59, 60]. Third, information on multiple food intake, physical activity and vaccination was lacking. Fourth, PMs might interact with multiple indoor air pollutants (*e.g.* mould, household fuels, allergens, furniture, paints and cleaning agents) [61], which cannot be readily disentangled. Fifth, our study has addressed the association between PMs and asthma and asthma-related respiratory symptoms in Guangdong only, and hence a wider distribution of geographic locations should be taken into account in future studies. We will also conduct long-term analysis and follow-up studies to further verify the current findings. In addition, our study has highlighted the need for ongoing research on PM₁, especially its constituents and toxicity, which might help broaden the scope of the study. Finally, exposure measurement errors have not been thoroughly considered. We have enrolled participants who had been living at the current location for >6 months within the year of the survey. The 722-day averaged exposure levels of the air pollutants might have been biased for those living at the location for <722 days. However, high-quality PM estimates at finer spatiotemporal scales were lacking at the global and regional levels, which has hampered more comparative analyses using various exposure datasets.

Despite these limitations, our study has included a relatively large sample of the middle-aged and elderly population, allowing for a robust analysis to detect the associations for PMs with asthma and asthma-related respiratory symptoms based on a comprehensive panel of covariates for adjustment. Second, we have estimated the residential concentrations of PMs at a spatial resolution of 1 km², a reasonably fine precision of exposure modelling which has been validated in previous studies [26–28] to safeguard the accuracy of the exposure assessment. Additionally, despite excluding the participants with a known family history of allergy, the overall results were not altered materially. Results of the single- and two-pollutant models remained robust.

Our study has ascertained the associations between the respiratory health status of middle-aged and elderly residents in Guangdong province and the multiple airborne PMs, thus providing a more solid basis for policymakers to formulate updated policies for continuous air quality improvement. Our findings call

for continued monitoring of the dynamics between atmospheric PMs pollution and the burden of chronic respiratory diseases at the population level.

Conclusions

Long-term ambient PM₁ air pollution is associated with asthma, wheeze, persistent cough and dyspnoea in the middle-aged and elderly population. The strength of associations of PM₁, PM_{2.5} and PM₁₀ is comparable. Males appear more susceptible to the adverse associations of PMs than females. Future longitudinal studies incorporating individual-level monitoring for outdoor and indoor air pollutants as well as comprehensive measures of pollen are needed to confirm the findings.

Provenance: Submitted article, peer reviewed.

Acknowledgements: We thank Li-wen Fang and He-ling Bao (National Center for Chronic and Noncommunicable Disease Control and Prevention, Chinese Center for Disease Control and Prevention, Beijing, China) for their organisation and instruction of the programme. We thank the working staff from the local Centres for Disease Control and Prevention, health stations and community clinics for the programme organisation and implementation, epidemiological survey, and lung function testing.

Author contributions: Xue-yan Zheng, Shu-jun Guo, Jian-xiong Hu, Rui-lin Meng, Li-feng Lin and Wei-jie Guan made substantial contributions to the conception and design of the study. Xue-yan Zheng, Rui-lin Meng, Yan-jun Xu, Yun-hong Lv, Ye Wang, Ni Xiao, Chuan Li, Xiao-jun Xu, De-jian Zhao, Jing Wei, Li-feng Lin and Wei-jie Guan all contributed significantly to the acquisition of data. Xue-yan Zheng, Shu-jun Guo, Jian-xiong Hu, Rui-lin Meng, Hong-ye Zhou, Jia-hui He, Xiao-min Tan, Li-feng Lin and Wei-jie Guan contributed to the analysis and interpretation of the data. All authors read and approved the final manuscript.

Conflict of interest: The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper. Wei-jie Guan is an Associate Editor of *ERJ Open Research*.

Support statement: This study was supported by the National Natural Science Foundation Outstanding Youth Fund (82222001), Natural Science Foundation of Guangdong Province (2023A1515012328), and Guangdong Provincial Medical Science and Technology Research Funding (C2021083). Funding information for this article has been deposited with the Crossref Funder Registry.

Ethics statement: The study protocol was approved by the Ethics Review Committee of the National Center for Chronic and Noncommunicable Disease Control and Prevention, China Center for Disease Control and Prevention (201410 and 201901). All participants provided written informed consent.

References

- 1 Hansbro PM, Kim RY, Starkey MR, *et al.* Mechanisms and treatments for severe, steroid-resistant allergic airway disease and asthma. *Immunol Rev* 2017; 278: 41–62.
- 2 Andersen ZJ, Hvidberg M, Jensen SS, *et al.* Chronic obstructive pulmonary disease and long-term exposure to traffic-related air pollution: a cohort study. *Am J Respir Crit Care Med* 2011; 183: 455–461.
- 3 Liu C, Chen R, Sera F, *et al.* Ambient particulate air pollution and daily mortality in 652 cities. *N Engl J Med* 2019; 381: 705–715.
- 4 Beelen R, Raaschou-Nielsen O, Stafoggia M, *et al.* Effects of long-term exposure to air pollution on natural-cause mortality: an analysis of 22 European cohorts within the multicentre ESCAPE project. *Lancet* 2014; 383: 785–795.
- 5 Hansel NN, McCormack MC, Kim V. The effects of air pollution and temperature on COPD. *COPD* 2016; 13: 372–379.
- 6 Young MT, Sandler DP, DeRoo LA, *et al.* Ambient air pollution exposure and incident adult asthma in a nationwide cohort of U.S. women. *Am J Respir Crit Care Med* 2014; 190: 914–921.
- 7 Bowatte G, Erbas B, Lodge CJ, *et al.* Traffic-related air pollution exposure over a 5-year period is associated with increased risk of asthma and poor lung function in middle age. *Eur Respir J* 2017; 50: 1602357.
- 8 GBD 2019 Risk Factors Collaborators. Global burden of 87 risk factors in 204 countries and territories, 1990–2019: a systematic analysis for the Global Burden of Disease Study 2019. *Lancet* 2020; 396: 1223–1249.
- 9 Khreis H, Kelly C, Tate J, *et al.* Exposure to traffic-related air pollution and risk of development of childhood asthma: a systematic review and meta-analysis. *Environ Int* 2017; 100: 1–31.
- 10 Fuertes E, Heinrich J. The influence of childhood traffic-related air pollution exposure on asthma, allergy and sensitization. *Allergy* 2015; 70: 1350–1351.

- 11 Bowatte G, Lodge C, Lowe AJ, *et al.* The influence of childhood traffic-related air pollution exposure on asthma, allergy and sensitization: a systematic review and a meta-analysis of birth cohort studies. *Allergy* 2015; 70: 245–256.
- 12 Jacquemin B, Siroux V, Sanchez M, *et al.* Ambient air pollution and adult asthma incidence in six European cohorts (ESCAPE). *Environ Health Perspect* 2015; 123: 613–621.
- 13 Yang M, Chu C, Bloom MS, *et al.* Is smaller worse? New insights about associations of PM₁ and respiratory health in children and adolescents. *Environ Int* 2018; 120: 516–524.
- 14 Li J, Liu X, Yu LL, *et al.* Comparing polybrominated diphenyl ethers (PBDEs) in airborne particles in Guangzhou and Hong Kong: sources, seasonal variations and inland outflow. *J Environ Monit* 2009; 11: 1185–1191.
- 15 Wang C, Li W, Chen J, *et al.* Summer atmospheric polybrominated diphenyl ethers in urban and rural areas of northern China. *Environ Pollut* 2012; 171: 234–240.
- 16 Li N, Chen G, Liu F, *et al.* Associations of long-term exposure to ambient PM₁ with hypertension and blood pressure in rural Chinese population: the Henan rural cohort study. *Environ Int* 2019; 128: 95–102.
- 17 Kirrane EF, Luben TJ, Benson A, *et al.* A systematic review of cardiovascular responses associated with ambient black carbon and fine particulate matter. *Environ Int* 2019; 127: 305–316.
- 18 Mei M, Song H, Chen L, *et al.* Early-life exposure to three size-fractionated ultrafine and fine atmospheric particulates in Beijing exacerbates asthma development in mature mice. *Part Fibre Toxicol* 2018; 15: 13.
- 19 Bornehag CG, Lundgren B, Weschler CJ, *et al.* Phthalates in indoor dust and their association with building characteristics. *Environ Health Perspect* 2005; 113: 1399–1404.
- 20 Jalava PI, Wang Q, Kuusalo K, *et al.* Day and night variation in chemical composition and toxicological responses of size segregated urban air PM samples in a high air pollution situation. *Atmos Environ* 2015; 120: 427–437.
- 21 Fang L, Gao P, Bao H, *et al.* Chronic obstructive pulmonary disease in China: a nationwide prevalence study. *Lancet Respir Med* 2018; 6: 421–430.
- 22 Zheng XY, Li ZL, Li C, *et al.* Effects of cigarette smoking and biomass fuel on lung function and respiratory symptoms in middle-aged adults and the elderly in Guangdong province, China: a cross-sectional study. *Indoor Air* 2020; 30: 860–871.
- 23 Zhang S, Wolf K, Breitner S, *et al.* Long-term effects of air pollution on ankle-brachial index. *Environ Int* 2018; 118: 17–25.
- 24 Zhang JS, Gui ZH, Zou ZY, *et al.* Long-term exposure to ambient air pollution and metabolic syndrome in children and adolescents: a national cross-sectional study in China. *Environ Int* 2021; 148: 106383.
- 25 Tran PM, Warren JL, Leifheit EC, *et al.* Associations between long-term air pollutant exposure and 30-day all-cause hospital readmissions in US patients with stroke. *Stroke* 2023; 54: e126–e129.
- 26 Wei J, Li Z, Guo J, *et al.* Satellite-derived 1-km-resolution PM₁ concentrations from 2014 to 2018 across China. *Environ Sci Technol* 2019; 53: 13265–13274.
- 27 Wei J, Li Z, Lyapustin A, *et al.* Reconstructing 1-km-resolution high-quality PM_{2.5} data records from 2000 to 2018 in China: spatiotemporal variations and policy implications. *Remote Sens Environ* 2021; 252: 112136.
- 28 Wei J, Li Z, Xue W, *et al.* The ChinaHighPM₁₀ dataset: generation, validation, and spatiotemporal variations from 2015 to 2019 across China. *Environ Int* 2021; 146: 106290.
- 29 Wei J, Li Z, Li K, *et al.* Full-coverage mapping and spatiotemporal variations of ground-level ozone (O₃) pollution from 2013 to 2020 across China. *Remote Sens Environ* 2022; 270: 112775.
- 30 Wei J, Liu S, Li Z, *et al.* Ground-level NO₂ surveillance from space across China for high resolution using interpretable spatiotemporally weighted artificial intelligence. *Environ Sci Technol* 2022; 56: 9988–9998.
- 31 Wei J, Li Z, Wang J, *et al.* Ground-level gaseous pollutants (NO₂, SO₂, and CO) in China: daily seamless mapping and spatiotemporal variations. *Atmos Chem Phys* 2023; 23: 1511–1532.
- 32 Ferris BG. Epidemiology standardization project (American Thoracic Society). *Am Rev Respir Dis* 1978; 118: 1–120.
- 33 Kochovska S, Ekström M, Hansen-Flaschen J, *et al.* Hiding in plain sight: the evolving definition of chronic breathlessness and new ICD-11 wording. *Eur Respir J* 2023; 61: 2300252.
- 34 Huang K, Yang T, Xu J, *et al.* Prevalence, risk factors, and management of asthma in China: a national cross-sectional study. *Lancet* 2019; 394: 407–418.
- 35 de Marco R, Zanolin ME, Accordini S, *et al.* A new questionnaire for the repeat of the first stage of the European Community Respiratory Health Survey: a pilot study. *Eur Respir J* 1999; 14: 1044–1048.
- 36 Dong GH, Wang J, Liu MM, *et al.* Allergic predisposition modifies the effects of pet exposure on respiratory disease in boys and girls: the seven northeast cities of China (SNECC) study. *Environ Health* 2012; 11: 50.
- 37 Chen H, Zhang Y, Zhang L, *et al.* Indoor air pollution from coal combustion and tobacco smoke during the periconceptional period and risk for neural tube defects in offspring in five rural counties of Shanxi Province, China, 2010–2016. *Environ Int* 2023; 171: 107728.
- 38 Chen G, Li S, Zhang Y, *et al.* Effects of ambient PM₁ air pollution on daily emergency hospital visits in China: an epidemiological study. *Lancet Planet Health* 2017; 1: e221–e229.

- 39 Li H, Duan F, He K, *et al.* Size-dependent characterization of atmospheric particles during winter in Beijing. *Atmosphere* 2016; 7: 36.
- 40 Wang YQ, Zhang XY, Sun JY, *et al.* Spatial and temporal variations of the concentrations of PM₁₀, PM_{2.5} and PM₁ in China. *Atmos Chem Phys* 2015; 15: 13585–13598.
- 41 Koulouri E, Grivas G, Gerasopoulos E, *et al.* Study of size-segregated particle (PM₁, PM_{2.5}, PM₁₀) concentrations over Greece. *Global NEST J* 2008; 10: 132–139.
- 42 Vecchi R, Marcazzan G, Valli G, *et al.* The role of atmospheric dispersion in the seasonal variation of PM₁ and PM_{2.5} concentration and composition in the urban area of Milan (Italy). *Atmos Environ* 2004; 38: 4437–4446.
- 43 Peters JM, Avol E, Navidi W, *et al.* A study of twelve Southern California communities with differing levels and types of air pollution. I. Prevalence of respiratory morbidity. *Am J Respir Crit Care Med* 1999; 159: 760–767.
- 44 Abbey DE, Burchette RJ, Knutsen SF, *et al.* Long-term particulate and other air pollutants and lung function in nonsmokers. *Am J Respir Crit Care Med* 1998; 158: 289–298.
- 45 Galizia A, Kinney PL. Long-term residence in areas of high ozone: associations with respiratory health in a nationwide sample of nonsmoking young adults. *Environ Health Perspect* 1999; 107: 675–679.
- 46 Clougherty JE. A growing role for gender analysis in air pollution epidemiology. *Cien Saude Colet* 2011; 16: 2221–2238.
- 47 Haberzettl P, O'Toole TE, Bhatnagar A, *et al.* Exposure to fine particulate air pollution causes vascular insulin resistance by inducing pulmonary oxidative stress. *Environ Health Perspect* 2016; 124: 1830–1839.
- 48 Van Den Heuvel R, Den Hond E, Govarts E, *et al.* Identification of PM₁₀ characteristics involved in cellular responses in human bronchial epithelial cells (Beas-2B). *Environ Res* 2016; 149: 48–56.
- 49 Chen M, Qin X, Qiu L, *et al.* Concentrated ambient PM_{2.5}-induced inflammation and endothelial dysfunction in a murine model of neural IKK2 deficiency. *Environ Health Perspect* 2018; 126: 027003.
- 50 Wang Y, Wu T, Zou L, *et al.* Genome-wide identification and functional analysis of long non-coding RNAs in human endothelial cell line after incubation with PM_{2.5}. *Chemosphere* 2019; 216: 396–403.
- 51 Zhu XM, Wang Q, Xing WW, *et al.* PM_{2.5} induces autophagy-mediated cell death via NOS2 signaling in human bronchial epithelium cells. *Int J Biol Sci* 2018; 14: 557–564.
- 52 Gu LZ, Sun H, Chen JH. Histone deacetylases 3 deletion restrains PM_{2.5}-induced mice lung injury by regulating NF-κB and TGF-β/Smad2/3 signaling pathways. *Biomed Pharmacother* 2017; 85: 756–762.
- 53 Gong J, Zhu T, Kipen H, *et al.* Comparisons of ultrafine and fine particles in their associations with biomarkers reflecting physiological pathways. *Environ Sci Technol* 2014; 48: 5264–5273.
- 54 Cigankova H, Mikuska P, Hegrova J, *et al.* Comparison of oxidative potential of PM₁ and PM_{2.5} urban aerosol and bioaccessibility of associated elements in three simulated lung fluids. *Sci Total Environ* 2021; 800: 149502.
- 55 Watanabe M, Noma H, Kurai J, *et al.* A panel study of airborne particulate matter composition versus concentration: potential for inflammatory response and impaired pulmonary function in children. *Allergol Int* 2017; 66: 52–58.
- 56 Feng S, Gao D, Liao F, *et al.* The health effects of ambient PM_{2.5} and potential mechanisms. *Ecotoxicol Environ Saf* 2016; 128: 67–74.
- 57 Melén E, Standl M, Gehring U, *et al.* Air pollution and IgE sensitization in 4 European birth cohorts – the MeDALL project. *J Allergy Clin Immunol* 2021; 147: 713–722.
- 58 Huang KL, Liu SY, Chou CC, *et al.* The effect of size-segregated ambient particulate matter on Th1/Th2-like immune responses in mice. *PLoS One* 2017; 12: e0173158.
- 59 Porsbjerg C, Melén E, Lehtimäki L, *et al.* Asthma. *Lancet* 2023; 401: 858–873.
- 60 Berhane K, Chang CC, McConnell R, *et al.* Association of changes in air quality with bronchitic symptoms in children in California, 1993–2012. *JAMA* 2016; 315: 1491–1501.
- 61 Devien L, Giovannelli J, Cuny D, *et al.* Sources of household air pollution: the association with lung function and respiratory symptoms in middle-aged adult. *Environ Res* 2018; 164: 140–148.