



Correction

Correction: Schwarzkopf et al. Weaning Holstein Calves at 17 Weeks of Age Enables Smooth Transition from Liquid to Solid Feed. *Animals* 2019, 9, 1132

Sarah Schwarzkopf ¹, Asako Kinoshita ¹, Jeannette Kluess ², Susanne Kersten ², Ulrich Meyer ², Korinna Huber ¹, Sven Dänicke ² and Jana Frahm ²,*

- Institute of Animal Science, Faculty of Agricultural Sciences, University of Hohenheim, 70599 Stuttgart, Germany; sarah.schwarzkopf@uni-hohenheim.de (S.S.); asakokinoshita@googlemail.com (A.K.); korinna.huber@uni-hohenheim.de (K.H.)
- ² Institute of Animal Nutrition, Friedrich-Loeffler-Institute, 38116 Braunschweig, Germany; jeannette.kluess@fli.de (J.K.); Susanne.kersten@fli.de (S.K.); ulrich.meyer@fli.de (U.M.); sven.daenicke@fli.de (S.D.)
- * Correspondence: jana.frahm@fli.de

The authors wish to make the following correction to their paper [1]. In the Materials and Method Section, Table 3 was incorrect due to some numerical errors. The correct version of the table appears below. The authors apologize for any inconvenience; the change does not affect the scientific results on the animal level.

Table 3. Dry matter content and chemical composition of milk replacer, concentrate feed and roughage (hay) fed prior to weaning and total mixed ration (TMR) fed after weaning of the calves.

Feed	DM %	CA g/kg DM	CP g/kg DM	EE g/kg DM	CF g/kg DM	NDF g/kg DM	ADF g/kg DM	Starch g/kg DM	Sugar g/kg DM
Prior to weaning									
milk replacer	97	79	225	184					
concentrate feed	87	63	230	51	63	188	77	371	47
roughage (hay)	88	75	97	18	322	666	379		
After weaning									
TMR	44	73	134	36	212	421	242	182	12

All ingredients were assessed by Weender analysis. Dry matter (DM), crude ash (CA), crude protein (CP) and ether extract (EE) were analyzed in all feedstuff. Crude fiber (CF), neutral detergent fiber (NDF) and acid detergent fiber (ADF) were analyzed in the solid feed. In concentrate feed and TMR starch was analyzed and in concentrate feed additionally sugar was analyzed.

Reference

 Schwarzkopf, S.; Kinoshita, A.; Kluess, J.; Kersten, S.; Meyer, U.; Huber, K.; Dänicke, S.; Frahm, J. Weaning Holstein Calves at 17 Weeks of Age Enables Smooth Transition from Liquid to Solid Feed. *Animals* 2019, 9, 1132. [CrossRef] [PubMed]



Citation: Schwarzkopf, S.; Kinoshita, A.; Kluess, J.; Kersten, S.; Meyer, U.; Huber, K.; Dänicke, S.; Frahm, J. Correction: Schwarzkopf et al. Weaning Holstein Calves at 17 Weeks of Age Enables Smooth Transition from Liquid to Solid Feed. *Animals* 2019, 9, 1132. *Animals* 2021, 11, 3044. https://doi.org/10.3390/ani11113044

Received: 16 April 2021 Accepted: 9 July 2021 Published: 25 October 2021

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https://creativecommons.org/licenses/by/4.0/).