

## Reduced port laparoscopic surgery for colon cancer is safe and feasible in terms of short-term outcomes: comparative study with conventional multiport laparoscopic surgery

Ju Myung Song, Ji Hoon Kim, Yoon Suk Lee, Ho Young Kim, In Kyu Lee, Seung Teak Oh, Jun Gi Kim

Department of Surgery, College of Medicine, The Catholic University of Korea, Seoul, Korea

**Purpose:** Laparoscopic surgery was previously accepted as an alternative surgical option in treatment for colorectal cancer. Nowadays, single-port laparoscopic surgery (SPLS) is introduced as a method to maximize advantages of minimally invasive surgery. However, SPLS has several limitations compared to conventional multiport laparoscopic surgery (CMLS). To overcome those limitations of SPLS, reduced port laparoscopic surgery (RPLS) was introduced. This study aimed at evaluating the short-term outcomes of RPLS.

**Methods:** Patients who underwent CMLS and RPLS of colon cancer between August 2011 and December 2013 were included in this study. Short-term clinical and pathological outcome were compared between the 2 groups.

**Results:** Thirty-two patients underwent RPLS and 217 patients underwent CMLS. Shorter operation time, less blood loss, and faster bowel movement were shown in RPLS group in this study. In terms of postoperative pain, numeric rating scale (NRS) of RPLS was lower than that of CMLS. Significant differences were shown in terms of tumor size, harvested lymph node, perineural invasion, and pathological stage. No significant differences were confirmed in terms of other surgical outcomes.

**Conclusion:** In this study, RPLS was technically feasible and safe. Especially in terms of postoperative pain, RPLS was comparable to CMLS. RPLS may be a feasible alternative option in selected patients with colon cancer.

[Ann Surg Treat Res 2016;91(4):195-201]

**Key Words:** Colon cancer, Laparoscopic surgery, Minimally invasive surgery

### INTRODUCTION

Several prospective randomized studies of laparoscopic surgery for colorectal cancer had shown both short-term benefits such as less pain, better cosmetic effect, faster recovery, and long-term oncological safety [1-5]. So nowadays, laparoscopic surgery for colorectal cancer has been accepted as an alternative to conventional open surgery. So many colorectal surgeons have considered laparoscopic surgery as a useful surgical tool for treatment of colorectal cancer. However, conventional multiport

laparoscopic surgery (CMLS) for colorectal cancer requires 4 or 5 abdominal incisions for trocars and each incision could be associated with wound complications and pain. Single port laparoscopic surgery (SPLS) permits surgeons to reduce such wound related complications and achieve better cosmetic effects compared to CMLS. The potential advantages of SPLS are less pain and earlier recovery than CMLS. Actually, several studies had already reported acceptable short-term outcomes of SPLS compared to CMLS [6-9]. However, SPLS is a highly demanding procedure that also has several technical challenges such as

Received April 15, 2016, Revised June 1, 2016, Accepted June 21, 2016

Corresponding Author: Ji Hoon Kim

Department of Surgery, Incheon St. Mary's Hospital, The Catholic University of Korea, 56 Dongsu-ro, Bupyeong-gu, Incheon 21431, Korea

Tel: +82-32-280-5609, Fax: +82-32-280-5988

E-mail: samryong@catholic.ac.kr

Copyright © 2016, the Korean Surgical Society

© Annals of Surgical Treatment and Research is an Open Access Journal. All articles are distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

handling conventional laparoscopic instruments through small incisions, which could decrease the range of motion; and, there could be conflict between instruments and camera. As a result, there are also potential disadvantages such as longer operation time, increased surgeon fatigue and steep learning curve.

Reduced port laparoscopic surgery (RPLS), which is single port plus one additional port, may overcome the limitations of SPLS while maintaining the advantages of SPLS.

The aim of this study is to evaluate the short-term outcomes of RPLS and assess the safety and feasibility of RPLS in colon cancer.

## METHODS


This study is a retrospective review of prospectively collected data of patient who underwent CMLS and RPLS for colon cancer between August 2011 and December 2013 at Incheon St. Mary's Hospital, The Catholic University of Korea by a single surgeon. All patients underwent preoperative work-up including colonoscopy with biopsy, CT, laboratory test, and chest radiography. All patients with colorectal cancer admitted to Incheon St. Mary's Hospital were considered for laparoscopic surgery, both CMLS and RPLS. Because of the nature of the retrospective study, the indication for RPLS for colon cancer was not so strict and changed case by case. In general, patients with ascending or sigmoid colon cancer with small sized tumor and relatively early stage were considered for RPLS. In our institute, rectal cancer and transverse colon cancer were also considered for laparoscopic surgery. However, in this study, rectal cancer and transverse colon cancer were excluded; because, in our institute, rectal cancer and transverse colon cancer were only considered for CMLS but not for RPLS or SILS. Informed consent was obtained from all patients. In our institute, critical pathway was used for all patients who

underwent colorectal cancer surgery. In the present study, we compared short-term clinical outcomes including pain score and pathologic outcomes between CMLS and RPLS. Postoperative pain was categorized into 0–3; 0 (no pain), 1 (mild pain), 2 (moderate pain), 3 (severe pain). And pain level was checked at 2:00 PM on postoperative day 1, 3, and 5. In our institution, we routinely use patient-controlled analgesia (PCA), with a regimen containing fentanyl. But in case where patients refuse PCA, we use continuous infusion combined with bolus injection of pethidine hydrochloride. We analyzed the total amount of infused analgesics as converted into fentanyl (fentanyl 1,000  $\mu$ g was considered equal to 750 mg of pethidine).

This study was approved by the Institutional Review Board of Incheon St. Mary's Hospital (OC14RISI0087).

## Operative procedure

For both CMLS and RPLS, patients were placed in the lithotomy position. For CMLS, a 10-mm trocar was inserted at umbilicus for the camera, and 4 trocars were inserted at RLQ (right lower quadrant), RUQ (right upper quadrant), LLQ (left lower quadrant), and LUQ (left upper quadrant) sites. For RPLS, a single port was inserted using transumbilical incision and an additional port insert in the dominant-hand side of the operator (Fig. 1). All RPLS were performed using an OCTO port (Dalim SurgNET, Seoul, Korea), single port with a 4-channel system. For right colon cancer, we performed complete mesocolonic excision (CME) and central vessel ligation (CVL) independent to tumor stages in both RPLS and CMLS. For sigmoid colon cancer, high ligation of inferior mesenteric artery was routinely performed in both RPLS and CMLS. In RPLS of anterior resection, proximal rectal transection was performed using a linear stapler, which was introduced via additional RLQ port, and intracorporeal stapled anastomosis was performed. For right hemicolectomy, extracorporeal resection and anastomosis was performed. In



**Fig. 1.** Trocars placement of reduced port laparoscopic surgery for colon cancer. (A) Trocars placement in right-side colon cancer. (B) Trocars placement in left-side colon cancer.

RPLS, all the specimens were retrieved through the single port and in CMLS, all specimens were extracted through extension of the umbilical incision. After anastomosis, irrigation was done in the abdominal cavity and a drain was inserted through the port incision.

### Statistics

Continuous variables were compared using t-test. Categorical variables were compared using chi-square and Fisher exact test. Postoperative pain score was compared using t-test. A P-value

less than 0.05 was considered statistically significant.

## RESULTS

During the study period, 32 cases were enrolled in RPLS group and 217 cases were enrolled in CMLS group. In the present study, no significant differences were shown in terms of demographics and clinical characteristics of patients between the 2 groups (Table 1). In terms of operative outcomes, longer operation time and large amount of blood loss were shown

**Table 1.** Demographics and clinical characteristics of reduced port and conventional multiport laparoscopic surgery for colon cancer

Variable	RPLS (n = 32)	CMLS (n = 217)	P-value
Sex			0.967
Male	19 (59.4)	128 (59.0)	
Female	13 (40.6)	89 (41.0)	
Age (yr)	65.2 ± 10.8 (44–87)	64.6 ± 12.7 (32–87)	0.804
Body mass index (kg/m <sup>2</sup> )	23.3 ± 2.8 (17.2–29.5)	23.9 ± 3.9 (14.9–42.6)	0.381
ASA PS classification			0.960
I	1 (3.1)	8 (3.7)	
II	29 (90.6)	193 (88.9)	
III	2 (6.3)	16 (7.4)	0.909
Tumor location			
Right-sided colon cancer	10 (31.2)	70 (32.3)	
Left-sided colon cancer	22 (68.8)	147 (67.7)	
Operation title			0.712
Right colectomy	10 (31.2)	70 (32.3)	
Left colectomy	2 (6.3)	19 (8.7)	
Anterior resection	13 (40.6)	97 (44.7)	
Low anterior resection	7 (21.9)	31 (14.3)	

Values are presented as number (%) or mean ± standard deviation (range).

RPLS, reduced port laparoscopic surgery; CMLS, conventional multiport laparoscopic surgery; ASA PS, American Society of Anesthesiologists physical status.

**Table 2.** Operative and pathologic findings of reduced port and conventional multiport laparoscopic surgery

Variable	RPLS (n = 32)	CMLS (n = 217)	P-value
Operation time (min)	104.1 ± 33.8 (70–200)	128.4 ± 37.3 (65–290)	<0.001
Blood loss (mL)	23.4 ± 17.3 (0–80)	56.1 ± 82.5 (0–600)	0.008
Tumor size (cm)	3.6 ± 3.0 (0–13.5)	5.6 ± 3.0 (0–20)	<0.001
PRM (cm)	13.2 ± 5.9 (4–30)	14.0 ± 7.3 (5–60)	0.542
DRM (cm)	11.6 ± 5.8 (5–28)	11.8 ± 5.8 (2–37)	0.825
Harvested lymph nodes	18.7 ± 9.5 (3–40)	24.1 ± 12.8 (1–76)	0.022
Lymphovascular invasion	10 (31.2)	92 (42.4)	0.399
Perineural invasion	5 (15.6)	74 (34.1)	0.036
Stage			0.009
I	12 (37.5)	31 (14.3)	
II	7 (21.9)	71 (32.7)	
III	7 (21.9)	79 (36.4)	
IV	6 (18.7)	36 (16.6)	

Values are presented as mean ± standard deviation (range) or number (%).

RPLS, reduced port laparoscopic surgery; CMLS, conventional multiport laparoscopic surgery; PRM, proximal resection margin; DRM, distal resection margin.

in CMLS group (Table 2). In terms of pathological outcomes, tumor size and harvested number of lymph nodes were larger in CMLS group compared to RPLS group. When we compared the pathologic tumor stage, stage I was more frequent in RPLS group and stages II and III were more frequent in CMLS group. The rate of stage IV was similar between the 2 groups.

In terms of postoperative clinical outcomes, time to pass flatus was shorter in RPLS group compared to CMLS ( $2.1 \pm 0.9$  vs.  $2.4 \pm 1.3$ ,  $P = 0.011$ ) (Table 3). Surgical morbidity was 9 cases (28.1%) in RPLS group and 44 cases (20.3%) in CMLS group. Anastomosis leak was encountered in 0 cases in RPLS group and 5 cases in CMLS group but showed no difference. And in

terms of conversion, there was no significant difference. There was no surgical mortality in this study.

In the present study, we estimated postoperative pain between the 2 groups using numeric rating scale (NRS). NRS was significantly different between RPLS and CMLS. NRS of RPLS at postoperative day (POD) 1, 3, and 5 were significantly lower than those of CMLS (Table 4). There was no significant difference in terms of total amount of infused analgesics.

In subgroup analysis of stage I colon cancer, pathologic outcomes such as PRM, DRM, or harvested lymph nodes did not differ between the 2 groups (Table 5). A shorter operative time was observed in the RPLS group. However, in terms of postoperative pain, only the NRS score of POD 3 was significantly lower in the RPLS group than the CMLS group.

**Table 3.** Postoperative outcomes of reduced port and conventional multiport laparoscopic surgery

Variable	RPLS (n = 32)	CMLS (n = 217)	P-value
Time to pass flatus (day)	$2.1 \pm 0.9$ (0–4)	$2.4 \pm 1.3$ (0–8)	0.011
Diet start (day)	$2.3 \pm 0.7$ (2–5)	$2.4 \pm 0.6$ (2–6)	0.777
Hospital stay (day)	$7.9 \pm 4.9$ (6–32)	$7.7 \pm 3.6$ (5–32)	0.765
Morbidity	9 (28.1)	44 (20.3)	0.311
Urinary retension	5 (15.6)	13 (6.0)	
Anastomosis leak	0 (0)	5 (2.3)	
SSI	0 (0)	5 (2.3)	
Ileus	1 (3.1)	13 (6.0)	
Others	3 (9.4)	8 (3.7)	
Conversion	0 (0)	10 (4.6)	0.215

Values are presented as mean  $\pm$  standard deviation (range) or number (%).

RPLS, reduced port laparoscopic surgery; CMLS, conventional multiport laparoscopic surgery; SSI, surgical site infection.

**Table 4.** Numeric rating scale scores of reduced port and conventional multiport laparoscopic surgery

Numeric rating scale	RPLS (n = 32)	CMLS (n = 217)	P-value
NRS of POD 1	$1.58 \pm 0.83$	$1.92 \pm 0.62$	0.027
NRS of POD 3	$1.58 \pm 0.61$	$1.82 \pm 0.54$	0.039
NRS of POD 5	$1.50 \pm 0.51$	$1.72 \pm 0.54$	0.029
Total dose of analgesics (converted into fentanyl, $\mu\text{g}$ )	$1595.8 \pm 792.0$	$1629.2 \pm 507.7$	0.815

Values are presented as mean  $\pm$  standard deviation.

RPLS, reduced port laparoscopic surgery; CMLS, conventional multiport laparoscopic surgery; NRS, numeric rating scale; POD, postoperative day.

**Table 5.** Subgroup analysis for stage I colon cancer of reduced port and conventional multiport laparoscopic surgery

Variable	RPLS (n = 32)	CMLS (n = 217)	P-value
Sex			0.075
Male:female	11:1	20:11	
Age (yr)	$66.1 \pm 7.5$ (59–87)	$65.5 \pm 9.6$ (45–85)	0.855
Body mass index ( $\text{kg}/\text{m}^2$ )	$24.1 \pm 2.7$ (20.0–27.2)	$24.9 \pm 3.1$ (19.2–31.1)	0.399
Operation time (min)	$100.2 \pm 33.0$ (70–195)	$126.6 \pm 30.8$ (80–200)	0.017
PRM (cm)	$14.0 \pm 7.7$ (4.0–30.0)	$13.4 \pm 5.2$ (6.0–27.0)	0.758
DRM (cm)	$12.4 \pm 3.9$ (5.0–17.0)	$9.8 \pm 4.7$ (2.0–19.0)	0.108
Harvested lymph nodes	$15.9 \pm 7.5$ (8–32)	$15.9 \pm 10.1$ (1–36)	0.997
Time to pass flatus (day)	$2.0 \pm 1.0$ (1–4)	$2.8 \pm 1.1$ (0–6)	0.033
Diet start (day)	$2.3 \pm 0.9$ (2–5)	$2.4 \pm 0.6$ (2–4)	0.706
Hospital stay (day)	$8.8 \pm 7.4$ (6–32)	$8.8 \pm 4.9$ (5–26)	0.990
Morbidity	5 (41.7)	8 (26.0)	0.240
NRS of POD 1	$1.83 \pm 0.83$	$1.94 \pm 0.57$	0.702
NRS of POD 3	$1.58 \pm 0.51$	$1.94 \pm 0.36$	0.046
NRS of POD 5	$1.72 \pm 0.47$	$1.77 \pm 0.56$	0.805

Values are presented as mean  $\pm$  standard deviation (range) or number (%).

RPLS, reduced port laparoscopic surgery; CMLS, conventional multiport laparoscopic surgery; PRM, proximal resection margin; DRM, distal resection margin; NRS, numeric rating scale; POD, postoperative day.

## DISCUSSION

During the past decades, several large prospective randomized trials have demonstrated better short-term clinical outcomes of laparoscopic surgery for colorectal cancer, and also established the long-term oncological safety of laparoscopic surgery for colorectal cancer [10-14]. So nowadays, many surgeons consider laparoscopic surgery as a better surgical option to treat patients suffering from colorectal cancer than conventional open surgery.

However, conventional laparoscopic surgery requires 3 or more ports and also need an additional incision for specimen removal. Additionally, each trocar site incision has potential risk for morbidities such as postoperative wound pain and negative effects on cosmetic outcomes. Accumulation of experience in advanced laparoscopic surgery combined with development of laparoscopic surgical instruments has induced to use less trocars or single trocar. Many surgeons had applied single incision laparoscopic surgery to the era of colorectal cancer expecting potential advantages including better cosmetic outcomes, less postoperative pain, and faster recovery, and several studies have reported such short-term outcomes [15-17].

However, single incision laparoscopic surgery is technically difficulty and results in increasing surgeon fatigue and operation time compared to CMLS [18-20].

RPLS, a single port with one additional port, may decrease the technical difficulties of SILS while maintaining its cosmetic outcomes and can be consider as a bridge to SILS. However, there are few studies evaluating the safety or technical feasibility of RPLS.

In the present study, 7 cases of low anterior resection for distal sigmoid colon cancer were enrolled in RPLS group. Although the safety and technical feasibility of SILS low anterior resection has been reported [21,22], until now those factors had not yet been established. Actually, applying laparoscopic linear stapler through a single port for rectal transection does not permit sufficient distal resection margin and is sometimes physically and technically impossible in a narrow pelvis. Inserting an additional port at RLQ permits the surgeon to apply a laparoscopic linear stapler and perform rectal transection more easily.

One of the disadvantages of SILS for colon cancer is longer operation time compared to conventional laparoscopic multiport surgery [23,24]. There are several possible explanations for the longer operation time of SILS. The most important reason is limitation of motion and reduced ability of triangular dissection. Because a camera and 2 laparoscopic instruments are introduced into the peritoneal cavity through a small incision or the umbilicus, collision and conflict between the camera and laparoscopic instruments are inevitable. Also, the steep learning curve of SILS is another possible explanation.

Inserting one additional port at the surgeon's dominant-hand side might resolve those drawbacks of SILS, including collision and difficulty of triangular dissection. Several studies of RPLS reported that operation time of RPLS was comparable to CMLS [25,26]

In the present study, the operation time was significantly shorter in RPLS group compared to CMLS group. We considered it as selection bias; because tumor size was also significantly smaller and tumor stage I was significantly more frequent in RPLS group. However, in the subgroup analysis for stage I cancer, the operation time of RPLS was significantly shorter than CMLS group. Another reason for shorter operative time of RPLS might be caused by the operator's advanced techniques. CMLS was generally performed earlier during the study period than RPLS, therefore the surgeon's technique might have improved with the passage of time. In our institute, the extent of surgery was the same, independent of tumor stage: CME and CVL for right colon cancer and high ligation of inferior mesenteric artery for sigmoid colon cancer. So, considering difference of tumor stage, we thought operation time of RPLS for colon cancer was comparable to CMLS for colon cancer.

When introducing a new surgical technique, intraoperative morbidity and postoperative morbidity are very important issues. And one of the parameters of intraoperative morbidity is the amount of intraoperative blood loss. In the present study, blood loss during surgery was about 24 mL in RPLS group and 56 mL in CMLS group, which showed statistical difference. And in terms of postoperative morbidities, there were no significant difference between the 2 groups. Anastomosis leakage is one of the most dreadful complications of colorectal surgery and could result in severe surgical morbidity or mortality. Although it is still controversial, reducing the number of linear stapler firings was necessary to avoid anastomosis leakage after laparoscopic colorectal surgery [27]. During SILS anterior or low anterior resection, applying a linear stapler and proximal rectum transection is a very challenging procedure. Sometimes more linear staplers are needed, because it is difficult to achieve the right angle between the bowel and linear stapler. Using one additional port for the linear stapler permitted the surgeon to transect the proximal rectum easier. In the present study, authors used an average of 1.3 linear staplers in anterior resection and 2.0 in low anterior resection, and there was no anastomosis leakage in RPLS group. Authors thought that insertion of an additional port could reduce the number of linear staplers for anterior or low anterior resection, and also risk of anastomosis leakage.

In terms of postoperative recovery, time to pass flatus of this study was shorter in RPLS group compared to CMLS group. The authors used the same critical pathway for all patients who underwent laparoscopic colorectal cancer surgery in both RPLS and CMLS. We thought less postoperative pain contributed

to faster bowel movement because if the patients had less postoperative pain, earlier ambulation could be possible resulting in faster bowel movement.

When single incision laparoscopic surgery was introduced, many surgeons expected potential advantages, especially less postoperative pain compared to CMLS. However, that potential advantage is still controversial and several studies have reported that postoperative pain of SPLS was similar to or more severe than CMLS [28-30]. The authors thought that although SPLS has the advantage of a single incision, this single incision site might be strongly lengthened and stretched by inserting a single port during the operation since SPLS has several technical challenges such as handling conventional laparoscopic instruments. So, that could be the major cause of postoperative pain of SILS.

The advantages of RPLS are a relative decrease of technical difficulties of SILS while maintaining cosmetic outcomes of SILS, and comparable operation time compared to CMLS.

Therefore, the single incision site of RPLS is less lengthened and stretched than that of SPLS during operation and the lengthening and stretching time of the single incision site of RPLS is shorter than that of SPLS. These factors can explain why patients who underwent RLPS could experience less pain. However, there are few studies evaluating the postoperative pain of RPLS and reported favorable result. In the present study,

we evaluated postoperative pain of RPLS group and CMLS group using NRS. Although authors used the same critical pathway for all patients, NRS of RPLS at POD 1, 3, and 5 were significantly lower than those of CMLS. We assumed that relative decrease of technical difficulties and shorter operation time of RPLS contributed to less postoperative pain.

There were several limitations in this study. First, this study is retrospective and has selection bias. We performed RPLS in the relatively early stage of colon cancer compared with CMLS. Second, this study did not compare the cosmetic outcomes or patient satisfaction of cosmetic effects. Large-scale prospective randomized studies are needed to establish the benefits of RPLS in colon cancer.

In conclusion, the present study showed that RPLS for colon cancer is safe and technically feasible. And in terms of postoperative pain, RPLS was more favorable than CMLS. We thought RPLS for colon cancer could be a good surgical option for selected patients suffering from colon cancer.

## CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

## REFERENCES

- Lorenzon L, La Torre M, Ziparo V, Montebelli F, Mercantini P, Balducci G, et al. Evidence based medicine and surgical approaches for colon cancer: evidences, benefits and limitations of the laparoscopic vs open resection. *World J Gastroenterol* 2014;20:3680-92.
- Scarpa M, Erroi F, Ruffolo C, Mollica E, Polese L, Pozza G, et al. Minimally invasive surgery for colorectal cancer: quality of life, body image, cosmesis, and functional results. *Surg Endosc* 2009;23:577-82.
- Baek JH, Lee GJ, Lee WS. Comparison of long-term oncologic outcomes of stage III colorectal cancer following laparoscopic versus open surgery. *Ann Surg Treat Res* 2015;88:8-14.
- Sammour T, Jones IT, Gibbs P, Chandra R, Steel MC, Shedda SM, et al. Comparing oncological outcomes of laparoscopic versus open surgery for colon cancer: analysis of a large prospective clinical database. *J Surg Oncol* 2015;111:891-8.
- Cianchi F, Trallori G, Mallardi B, Macri G, Biagini MR, Lami G, et al. Survival after laparoscopic and open surgery for colon cancer: a comparative, single-institution study. *BMC Surg* 2015;15:33.
- Kim SJ, Ryu GO, Choi BJ, Kim JG, Lee KJ, Lee SC, et al. The short-term outcomes of conventional and single-port laparoscopic surgery for colorectal cancer. *Ann Surg* 2011;254:933-40.
- Egi H, Hattori M, Hinoi T, Takakura Y, Kawaguchi Y, Shimomura M, et al. Single-port laparoscopic colectomy versus conventional laparoscopic colectomy for colon cancer: a comparison of surgical results. *World J Surg Oncol* 2012;10:61.
- Papaconstantinou HT, Thomas JS. Single-incision laparoscopic colectomy for cancer: assessment of oncologic resection and short-term outcomes in a case-matched comparison with standard laparoscopy. *Surgery* 2011;150:820-7.
- Poon JT, Cheung CW, Fan JK, Lo OS, Law WL. Single-incision versus conventional laparoscopic colectomy for colonic neoplasm: a randomized, controlled trial. *Surg Endosc* 2012;26:2729-34.
- Kim IY, Kim BR, Kim YW. The short-term and oncologic outcomes of laparoscopic versus open surgery for T4 colon cancer. *Surg Endosc* 2016;30:1508-18.
- Allaix ME, Giraud G, Mistrangelo M, Arezzo A, Morino M. Laparoscopic versus open resection for colon cancer: 10-year outcomes of a prospective clinical trial. *Surg Endosc* 2015;29:916-24.

12. Zeng WG, Liu MJ, Zhou ZX, Hou HR, Liang JW, Wang Z, et al. Outcome of laparoscopic versus open resection for transverse colon cancer. *J Gastrointest Surg* 2015;19:1869-74.
13. Wang CL, Qu G, Xu HW. The short- and long-term outcomes of laparoscopic versus open surgery for colorectal cancer: a meta-analysis. *Int J Colorectal Dis* 2014;29:309-20.
14. Desiderio J, Trastulli S, Ricci F, Penzo J, Cirocchi R, Farinacci F, et al. Laparoscopic versus open left colectomy in patients with sigmoid colon cancer: prospective cohort study with long-term follow-up. *Int J Surg* 2014;12:745-50.
15. Katsuno G, Fukunaga M, Nagakari K, Yoshikawa S, Azuma D, Kohama S. Short-term and long-term outcomes of single-incision versus multi-incision laparoscopic resection for colorectal cancer: a propensity-score-matched analysis of 214 cases. *Surg Endosc* 2016;30:1317-25.
16. Tei M, Wakasugi M, Akamatsu H. Comparison of perioperative and short-term oncological outcomes after single- or multiport surgery for colorectal cancer. *Colorectal Dis* 2015;17:O141-7.
17. Markar SR, Wiggins T, Penna M, Paraskeva P. Single-incision versus conventional multiport laparoscopic colorectal surgery: systematic review and pooled analysis. *J Gastrointest Surg* 2014;18:2214-27.
18. Levic K, Bulut O. The short-term outcomes of conventional and single-port laparoscopic surgery for rectal cancer: a comparative non-randomized study. *Minim Invasive Ther Allied Technol* 2014; 23:214-22.
19. Champagne BJ, Lee EC, Leblanc F, Stein SL, Delaney CP. Single-incision vs straight laparoscopic segmental colectomy: a case-controlled study. *Dis Colon Rectum* 2011; 54:183-6.
20. Yun JA, Kim HC, Park JS, Cho YB, Yun SH, Lee WY. Perioperative and oncologic outcomes of single-incision laparoscopy compared with conventional laparoscopy for colon cancer: an observational propensity score-matched study. *Am Surg* 2015; 81:316-23.
21. Gash K, Bicsak M, Dixon A. Single-incision laparoscopic surgery for rectal cancer: early results and medium-term oncological outcome. *Colorectal Dis* 2015; 17:1071-8.
22. Uematsu D, Akiyama G, Narita M, Magishi A. Single-access laparoscopic low anterior resection with vertical suspension of the rectum. *Dis Colon Rectum* 2011;54:632-7.
23. Kwag SJ, Kim JG, Oh ST, Kang WK. Single incision vs conventional laparoscopic anterior resection for sigmoid colon cancer: a case-matched study. *Am J Surg* 2013; 206:320-5.
24. Lim SW, Kim HR, Kim YJ. Single incision laparoscopic colectomy for colorectal cancer: comparison with conventional laparoscopic colectomy. *Ann Surg Treat Res* 2014;87:131-8.
25. Lim SW, Kim HJ, Kim CH, Huh JW, Kim YJ, Kim HR. Umbilical incision laparoscopic colectomy with one additional port for colorectal cancer. *Tech Coloproctol* 2013;17:193-9.
26. Kawamata F, Homma S, Minagawa N, Kawamura H, Takahashi N, Taketomi A. Comparison of single-incision plus one additional port laparoscopy-assisted anterior resection with conventional laparoscopy-assisted anterior resection for rectal cancer. *World J Surg* 2014;38:2716-23.
27. Kim JS, Cho SY, Min BS, Kim NK. Risk factors for anastomotic leakage after laparoscopic intracorporeal colorectal anastomosis with a double stapling technique. *J Am Coll Surg* 2009;209:694-701.
28. Wolthuis AM, Penninckx F, Fieuws S, D'Hoore A. Outcomes for case-matched single-port colectomy are comparable with conventional laparoscopic colectomy. *Colorectal Dis* 2012;14:634-41.
29. Kim SJ, Choi BJ, Lee SC. Successful total shift from multiport to single-port laparoscopic surgery in low anterior resection of colorectal cancer. *Surg Endosc* 2014;28: 2920-30.
30. Park SJ, Lee KY, Kang BM, Choi SI, Lee SH. Initial experience of single-port laparoscopic surgery for sigmoid colon cancer. *World J Surg* 2013;37:652-6.