



Genetic Factors in Nonsyndromic Orofacial Clefts

Mahamad Irfanulla Khan¹ Prashanth C.S.² Narasimha Murthy Srinath³

¹ Department of Orthodontics & Dentofacial Orthopedics, The Oxford Dental College, Bangalore, Karnataka, India

² Department of Orthodontics & Dentofacial Orthopedics, DAPM R.V. Dental College, Bangalore, Karnataka, India

³ Department of Oral & Maxillofacial Surgery, Krishnadevaraya College of Dental Sciences, Bangalore, Karnataka, India

Address for correspondence Mahamad Irfanulla Khan, BDS, MDS, Department of Orthodontics & Dentofacial Orthopedics, The Oxford Dental College, Bangalore, Karnataka 560068, India (e-mail: drirfankhanmds@gmail.com).

Global Med Genet 2020;7:101–108.

Abstract

Orofacial clefts (OFCs) are the most common congenital birth defects in humans and immediately recognized at birth. The etiology remains complex and poorly understood and seems to result from multiple genetic and environmental factors along with gene–environment interactions. It can be classified into syndromic (30%) and nonsyndromic (70%) clefts. Nonsyndromic OFCs include clefts without any additional physical or cognitive deficits. Recently, various genetic approaches, such as genome-wide association studies (GWAS), candidate gene association studies, and linkage analysis, have identified multiple genes involved in the etiology of OFCs.

This article provides an insight into the multiple genes involved in the etiology of OFCs. Identification of specific genetic causes of clefts helps in a better understanding of the molecular pathogenesis of OFC. In the near future, it helps to provide a more accurate diagnosis, genetic counseling, personalized medicine for better clinical care, and prevention of OFCs.

Keywords

- ▶ orofacial clefts
- ▶ nonsyndromic
- ▶ genetics
- ▶ gene mutation
- ▶ genome-wide association study
- ▶ linkage analysis

Introduction

Orofacial clefts (OFC) are the most common congenital malformations in the orofacial region causing a significant personal and social encumbrance.^{1–3} The etiology remains complex involving multiple genetic and environmental factors along with gene–environment interactions.^{4–7} Children born with clefts may have difficulties in sucking, speech, hearing, and unaesthetic facial features due to the anatomical deformities.^{8,9} OFCs generally require surgical reconstruction of the lip and palate at different stages from birth to adulthood,¹⁰ and their rehabilitation involves a multidisciplinary approach such as, pediatric care, speech and hearing therapy, dental and orthodontic treatment, genetic counseling, and other mental health therapy.^{11,12}

Epidemiology

The prevalence of OFCs ranges from 1 in 700 to 1,000 newborns worldwide¹³ and include cleft lip only (CLO), cleft palate only (CPO), and cleft lip and palate (CLP).¹⁴ They may occur as unilateral or bilateral, complete, or incomplete, and may involve the lip only, the palate only, or both.¹⁵

The prevalence of OFC varies according to geographical location, ethnicity, race, gender and socioeconomic status,^{16–19} Asians have the highest prevalence rate (1:500), the intermediate prevalence in Europeans (1:1000), and lowest in Africans (1:2500).^{20–22} In India, the incidence of clefts is around 1:800 to 1:1000, and three infants are born with some type of cleft every hour.²³ These differences appear to persist even after migration, suggesting that they are mediated by genetic, rather than environmental

published online
February 12, 2021

DOI <https://doi.org/10.1055/s-0041-1722951>.
ISSN 2699-9404.

© 2021. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

factors.²⁴ Overall, 70% of the OFCs are nonsyndromic (NS) and occur as isolated cases without any additional physical or cognitive deficits. In contrast, 30% of clefts are syndromic and are associated with a few other developmental anomalies.²⁵

The frequency of occurrence of OFC differs with regard to gender and side of clefting. Cleft lip is more common in males at a 2:1 male to female ratio, whereas a cleft palate is more common in females.²⁶ Approximately 90% of OFCs are unilateral with primarily left-sided involvement.²⁷

Development of Cleft Lip and Palate

The development of lip and palate begins during the fourth week of gestation where migrating neural crest cells combine with mesodermal cells to establish facial primordia which consist of five different processes (medial nasal, lateral nasal, frontonasal, maxillary, and mandibular processes) derived from the first pharyngeal arch.²⁸ Once facial prominences are formed, the nasal placodes invaginate to form the medial nasal process (MNP) and lateral nasal process (LNP). The maxillary processes initially grow medially, pushing the LNP toward the upper side. During the sixth and seventh weeks of gestation fusion of maxillary processes with each other and also with lateral and MNPs takes place to form upper lip and primary palate.²⁹ Defect in fusion or failure in the growth of these processes results in clefts involving the upper lip, alveolus, and/ or primary palate.

The secondary palate begins to develop in the seventh week of embryogenesis; the maxillary processes initially outgrow as palatal shelves which move toward each other vertically. After proper growth they settle at a horizontal position above the tongue which entails much extracellular remodeling.^{30,31} The palatal shelves then fuse in the midline both anteriorly and posteriorly like a zipper that forms a midline epithelial seam (MES). The disintegration of MES is required to maintain palatal confluency, which may involve apoptosis, epithelial-mesenchymal transition (EMT), and cell migration.³² Effective fusion of the secondary palate results in complete separation of the oral and nasal cavities. Cleft palate can result from failure at any of the steps, including palatal shelves elevation, cell migration, or fusion.

In summary, a variety of cellular mechanisms such as cell proliferation, cell migration, cell growth, cell fusion, apoptosis, EMT, and extracellular remodeling are involved in a coordinated manner during the development of lip and palate. Therefore, disruption in the gene/s involved in these processes during lip and palate development may lead to an OFC.

Glimpse into the History of Genetic Etiology of Orofacial Clefts

Although the familiarity of OFC has long been noted, Fogh-Andersen was the first to provide the evidence for genetic factors contributing to the etiology of CLP from family-based studies where it was observed that the siblings of CLP patients had an increased frequency of cleft lip with or without cleft palate.³³ This observation was further confirmed by studies on the familial distribution of congenital clefts of the lip and

palate,³⁴ and Dr. Clark Fraser published a review paper highlighting the conclusions of a workshop on CLP sponsored by the National Institutes of Health of the United States, and he mentioned the etiology is indeed multifactorial.³⁵ Later, more evidence in favor of contribution of genetic factors to the etiology of CLP accrued from segregation analysis³⁶ and twin studies^{37,38} where the monozygotic twins showed high prevalence rate (40%) than the dizygotic twins (4%).

Role of Genetic Factors in the Etiology of Orofacial Clefts

The etiology of OFCs involves genetic factors, environmental influences, and gene–environment interactions, all contributing to its susceptibility. Scientific literature evidence suggests that environmental factors such as maternal tobacco smoking and alcohol consumption, antiepileptic medications, maternal folate deficiency, infections, consanguinity, and geographical location are risk factors for NS cleft lip and palate (NSCLP).^{39–42}

Advances in genetics and molecular biology techniques have discovered multiple genes and loci associated with CLP. This article provides an overview of the genes implicated in the etiology of NSCLP. Identification of specific genetic variation contributing to NSCLP has led to an increase in our understanding of the molecular pathogenesis of OFC.

Genes Involved in the Etiology of Nonsyndromic Orofacial Clefts

Special AT-Rich Sequence-Binding Protein

Special AT-rich sequence-binding protein (SATB2) is a DNA binding protein which binds with nuclear matrix attachment regions. It is involved in transcription regulation and chromatin remodeling process. In an animal study, mouse SATB2 is strongly expressed in the developing cleft palate and is similar to the human SATB2 protein.⁴³ The identification of SATB2 gene responsible for the craniofacial dysmorphologies associated with deletions and translocations at 2q32–q33, only one region of the genome has been significantly associated with the development of isolated cleft palate.⁴⁴ Glass syndrome characterized by cleft palate, gum hyperplasia, slight micrognathia, generalized osteoporosis, and mental retardation reported from Thai patient,⁴⁵ was caused by SATB2 gene mutation. Recently, using salivary miRNAs showed that the SATB2 genes are involved in the development of the cleft palate and lip development.⁴⁶ **Table 1** provides the list of genes involved in the etiology of NS OFC in humans.

B-Cell Leukemia/Lymphoma 3

B-cell leukemia/lymphoma 3 (BCL3) is a proto-oncogene, acts as a transcriptional co-activator through NF-kappa-B target genes and located on 19q13.2. BCL3 gene has also shown a strong association with NS orofacial clefts (NSOCs).⁴⁷ A case-parent trio study showed BCL3 influence risk of CL/P through a parent-of-origin effect with the excess maternal transmission.⁴⁸ Several studies in different

Table 1 Genes involved in the etiology of nonsyndromic orofacial clefts in humans

Gene	Gene symbol	Loci	OMIM	Evidence	References
Special AT-rich sequence-binding protein	<i>SATB2</i>	2q33.1	608148	M	43,46
B-cell leukemia/lymphoma 3	<i>BCL3</i>	19q13.32	109560	LD, L	47,49
Distal-less homeobox 4	<i>DLX4</i>	17q21.33	601911	M	53,55,56
Paired box gene 9	<i>PAX9</i>	14q13.3	167416	L, GWAS	6,60,62,64
Netrin 1	<i>NTN1</i>	17p13.1	601614	GWAS	70,71
T-box transcription factor 22	<i>TBX22</i>	Xq21.1	300307	M, LD	75,77,79
Poliovirus receptor like-1	<i>PVRL1</i>	11q23.3	600644	M, LD,GWAS	81,83,85
Cleft lip and palate associated transmembrane protein 1	<i>CLPTM1</i>	19q13.32	604783	M, L, LD	89,90
MAF bZIP transcription factor B	<i>MAFB</i>	20q12	608968	GWAS	98,101
Fibroblast growth factor receptor 1	<i>FGFR1</i>	8p11.23	136350	M, D	102,103
Transcription factor AP2- α	<i>TFAP2A</i>	6p24.3	107580	D, L	104
S-glutathione transferase T1	<i>GSTT1</i>	22q11.2	600436	LD	108,109
Receptor-like tyrosine kinase	<i>RYK</i>	3q22.2	600524	M	110,111
Gamma-aminobutyric acid receptor, Beta-3	<i>GABRB3</i>	15q12	137192	GWAS	112
ATP-binding cassette, subfamily A, member 4	<i>ABCA4</i>	1p22.1	601691	GWAS	115,117,118

Abbreviations: D, deletions; GWAS, genome-wide association studies; L, linkage; LD, linkage disequilibrium; M, mutations; OMIM, Online Mendelian Inheritance in Man.

populations have implicated the role of *BCL3* in development of NSCL/P. It was found that *BCL3* contributes to the regulation of cell proliferation, and cell cycle regulation can cause a disturbance in facial formation.⁴⁹ A study also reported *BCL3* contribution in angiogenesis-related genes in the etiology of CLP.⁵⁰ However, no association of *BCL3* gene with NSCLP was found in multigenerational families of Indian population.⁵¹

Distal-Less Homeobox 4

Distal-Less Homeobox 4 (*DLX4*) belongs to *DLX* gene family containing a homeobox transcription factor which plays an important role in craniofacial development and palatogenesis. It is located on chromosome 17q21.33 and causes orofacial cleft 15 (*OFC15*) and cleft lip/palate. In an animal study, the *DLX* genes caused cleft palate showing the importance of these genes in craniofacial morphogenesis.⁵² Whole-exome sequencing study in a Hispanic mother and son with bilateral CLP confirmed the *DLX4* as a potential cause of oral clefts.⁵³⁻⁵⁵ Recently, a study showed that none of the distal-less 4 (*DLX4*) gene SNPs were associated with NSOCs, so it should be interpreted with a caution in the etiology of nonsyndromic orofacial clefts.⁵⁶

Paired Box Gene 9

Paired Box Gene 9 (*PAX9*) is a member of the paired box (*PAX*) family of transcription factors and contains a paired box domain, an octapeptide, and a paired-type homeodomain. It plays a critical role during neural crest and fetal development. *PAX9* is located on chromosome 14q13.3, consists of five exons, and associated with the formation of the teeth and palate.^{57,58} Tooth agenesis and the formation of a cleft palate in *PAX9*-deficient mice have been reported.⁵⁹

Schuffenhauer et al, first reported the role of *PAX9* in a patient presented with bilateral CLP.⁶⁰ A linkage analysis of *PAX9* gene in two large families and four unrelated families showed that the hypodontia primarily involving molars, suggested that the mutant *PAX9* protein acquires functional defects in DNA binding, as well as the loss of function of *PAX9* resulting in haploinsufficiency during the morphogenesis of the dentition and the subsequent tooth agenesis.^{61,62} Several studies identified mutations at *PAX9* may increase the risk of NS cleft lip with or without palate.⁶³⁻⁶⁶

A combined genome-wide association study (GWAS) used unmixed NS CPO and NS CLO subtypes suggested that *PAX9* is a strong genetic factor for NS CPO in the Chinese population. Mutation analysis of *PAX9* SNPs rs12885612 and rs12881248 revealed that *PAX9* is a promising susceptible gene for NS CLO in Western Han Chinese population.^{5,6}

Netrin 1 (NTN1)

The netrin 1 (OMIM: 601614) located on chromosome 17p13.1 is a family of laminin-related secreted proteins. Its functions include axon guidance and cell migration in the central nervous system, angiogenesis, and semicircular canal formation.

NTN1 is expressed at high levels in cells that will come together to form a fusion plate, a prerequisite for the formation of semicircular canals. In netrin 1 mutant mice, fusion plate formation is severely affected, and it stimulates proliferation of the periotic mesenchymal cells which then push the epithelial cell walls together to form the fusion plate.^{67,68} It affects the development of the craniofacial region and has been shown to play a vital role in regulating cell migration during embryogenesis, and it is also expressed in the medial edges and oral sides of the palatal shelves.⁶⁹

A GWAS included 1,409 case-parent trios by several research groups with samples of Asian or European ancestry from Europe, the United States, China, and the Philippines found NTN1 role in the etiology of NSCLP.⁷⁰ A case-control study of NTN1, identified SNP (rs9788972) as a risk locus for NSOCs susceptibility in a northern Chinese population.⁷¹

T-Box Transcription Factor 22

The *T-box 22* (TBX22) gene encodes transcription factors involved in the regulation of developmental processes and plays a major role in human palatogenesis. It contains eight coding exons. TBX22 cause X-linked cleft palate and is characterized by isolated cleft palate and ankyloglossia.⁷² It is expressed in the developing palatal shelves and at the base of the tongue prior to elevation to a horizontal position above the tongue.⁷³

Genome-wide linkage analysis showed the association of TBX22 role in the development of NSCLP.⁷⁴ In addition, mutations in TBX22 were found in individuals with isolated CPO.⁷⁵⁻⁷⁷ It plays a significant role in tooth and upper lip development and causes hypodontia and cleft lip.⁷⁸ DNA methylation study suggests that cleft palate-susceptible gene *Tbx22* is associated with gene expression and might be responsible for the developmental failure of palatal fusion, eventually resulting in the formation of cleft palate.⁷⁹

Poliovirus Receptor-Like 1

Poliovirus receptor-like 1 (PVRL1), also known as NECTIN1 (nectin cell adhesion molecule 1) belongs to the nectin subfamily of immunoglobulin-like adhesion molecules that involve cell-cell adhesion. It plays a vital role in the organization of adherens junctions and tight junctions in epithelial and endothelial cells.⁸⁰ During the developmental process, the palatal shelves and palatal epithelium come in close contact and fuse together. PVRL1 plays a major role in these developments during palatogenesis and genetic variations reported to have a significant relationship with CLP. Diseases associated with PVRL1 include cleft lip/palate-ectodermal dysplasia syndrome (CLPED) and herpes simplex.

In animal experiments, PVRL1 expressed at the medial edge epithelium of the palatal shelves and the skin surface epithelium locations that corresponded to the clinical phenotypes of CLPED and in humans, mutations of the *PVRL1* gene resulting CLPED in families from Israel and Brazil.⁸¹ Interestingly, heterozygous mutation of PVRL1 (W185X) associated with NSCLP in northern Venezuela^{82,83} and two novel variants of the *PVRL1* gene were identified in Turkish NSCLP patients.⁸⁴ In an experimental animal studies, PVRL1 expressed at the medial edge epithelium of the palatal shelves and the skin surface epithelium locations that corresponded to the clinical phenotypes of CLPED and in humans, mutations of the *PVRL1* genes caused CLPED in Israel and Brazilian population.⁸⁵⁻⁸⁸

Cleft Lip and Palate-Associated Transmembrane Protein 1

Cleft lip and palate-associated transmembrane protein 1 (CLPTM1) is a multipass transmembrane protein that regulates

GABA-A receptors (e.g., GABRA1) and modulates inhibitory synaptic strength. It is located on chromosome 19q13.3 and plays a role in T-cell development. Mutation of *CLPTM1* genes suggested that a regulatory element in this gene region get affected and which is responsible for the development of orofacial clefts.^{89,90} However, some studies contradict the role of CLPTM1 in the etiology of NSCLP/P as no evidence of an association with oral clefts was found among the SNPs of CLPTM1 selected for testing in Japanese and Irish population.^{91,92}

MAF bZIP Transcription Factor B

The MAF bZIP Transcription Factor B (*MAFB*) gene encodes a basic leucine zipper (bZIP) transcription factor that plays an important role in the regulation of lineage-specific hematopoiesis. It is located on chromosome 20q11.2, consists of a single exon and spans approximately 3 kb.⁹³ In a mouse study, *MAFB* was expressed in the palatal shelves and the medial edge epithelia (or MEE) during palatal fusion.⁹⁴ Mutations in the *MAFB* gene reported causing multicentric carpotarsal osteolysis syndrome and Duane retraction syndrome 3 with or without deafness.^{95,96}

Several GWAS in European and Asian populations identified the role of *MAFB* in NSCLP.^{97,98} Different systematic review and meta-analyses confirmed that the *MAFB* gene SNP (rs13041247) is associated with NSCLP/P risk in different population; however, this association is not significant in East Asian or Caucasian populations,^{99,100} whereas, the SNPs rs17820943 and rs6072081 of *MAFB* found to be associated with NSCLP in an East Asian population.¹⁰¹

Other Candidate Genes

A variety of genetic approaches have identified several genes located on different chromosomes, contributing to etiology of NSCLP. Advances in genetics and molecular biology techniques have led the way to the discovery of genetic variation involved in NSCLP. Genes such as *FGFR1*, *TFAP2A*, *GSTT1*, receptor-like tyrosine kinase (*RYK*), and *ABCA4* have also been found to be associated with NSCLP.

Riley et al assessed the genes involved in the fibroblast growth factor (FGF) signaling pathway and identified the functional impairment in the *FGFR1* gene in NSCLP families and suggested that the FGF signaling pathway may contribute to as much as 3 to 5% of NS cleft lip or palate.¹⁰² Xu et al reported mutations of the *FGFR1* gene in Chinese Kallmann syndrome males with cleft lip/palate.¹⁰³ *TFAP2A* located on 6p24 region has been associated with orofacial clefting. Davies et al reported a patient with cleft palate, microretrognathia, frontal bossing, hypertelorism, flat, broad nasal bridge, low set ears, and developmental delay.¹⁰⁴ However, it was found that there was no significant association between *TFAP2A* and NSCLP in this northern Chinese¹⁰⁵ and Indian population.¹⁰⁶ S-glutathione transferase T1 (*GSTT1*) play an important role in the detoxification and secretion of smoking byproducts and deficiency of this enzyme may cause a greater risk of NSCLP if the individuals were exposed to smoking byproducts during pregnancy.^{107,108} Hozyasz et al suggested that homozygous deletion of *GSTT1* in mother genome might increase the risk of having a child with NSCLP.¹⁰⁹

Further, mutations in the *RYK* gene were also found to be associated with orofacial clefting.^{110,111} Significant linkage disequilibrium between Gamma-aminobutyric acid receptor, Beta-3 (*GABRB3*), and *CLP* was reported by Scapoli et al and this finding in humans is in agreement with previously reported data obtained with the murine model.¹¹² In addition, Baroni et al and Carter et al reported the association of *GABRB3* with oral clefts in different populations.^{113,114} *ATP-binding cassette, subfamily A, member 4 (ABCA4)* encodes an ATP-binding cassette transporter. Several linkages and GWAS showed the role of *ABCA4* in NSCLP with stronger evidence among Asian samples. *ABCA4* is known to cause the autosomal-recessive retinal degenerative disease Stargardt's disease. A GWAS and a case-parent trio approach by Beaty et al identified *ABCA4* gene association with NSCLP^{115,116} and several other studies also reported a potential role of *ABCA4* in the etiology of *CL/P* in Brazilian and northern Chinese Han population.^{117,118}

Conclusion

Genetic studies provide insight into the multiple genes involved in the etiology of OFCs. Identification of specific genetic causes of clefts helps for better understanding of the molecular pathogenesis of OFC. In the near future, it helps to provide a more accurate diagnosis, genetic counseling, personalized medicine for better clinical care and prevention of OFCs.

Conflict of Interest

None declared.

References

- Wong FK, Hagg U. An update on the aetiology of orofacial clefts. *Hong Kong Med J* 2004;10(05):331–336
- Mossey P. Epidemiology underpinning research in the aetiology of orofacial clefts. *Orthod Craniofac Res* 2007;10(03):114–120
- Rahimov F, Jugessur A, Murray JC. Genetics of nonsyndromic orofacial clefts. *Cleft Palate Craniofac J* 2012;49(01):73–91
- Worley ML, Patel KG, Kilpatrick LA. Cleft lip and palate. *Clin Perinatol* 2018;45(04):661–678
- Huang L, Jia Z, Shi Y, et al. Genetic factors define CPO and CLO subtypes of nonsyndromic orofacial cleft. *PLoS Genet* 2019;15(10):e1008357
- Yang CW, Shi JY, Yin B, Shi B, Jia ZL. Mutation at paired box gene 9 is associated with non-syndromic cleft lip only from Western Han Chinese population. *Arch Oral Biol* 2020;117:104829
- Martinelli M, Palmieri A, Carinci F, Scapoli L. Non-syndromic cleft palate: an overview on human genetic and environmental risk factors. *Front Cell Dev Biol* 2020;8:592271
- Jugessur A, Farlie PG, Kilpatrick N. The genetics of isolated orofacial clefts: from genotypes to subphenotypes. *Oral Dis* 2009;15(07):437–453
- do Rego Borges A, Sá J, Hoshi R, et al. Genetic risk factors for nonsyndromic cleft lip with or without cleft palate in a Brazilian population with high African ancestry. *Am J Med Genet A* 2015;167A(10):2344–2349
- Saad AN, Parina RP, Tokin C, Chang DC, Gosman A. Incidence of oral clefts among different ethnicities in the state of California. *Ann Plast Surg* 2014;72(Suppl 1):S81–S83
- Ghazali N, Rahman NA, Kannan TP, Jaafar S. Screening of transforming growth factor Beta 3 and *Jagged2* genes in the Malay population with nonsyndromic cleft lip with or without cleft palate. *Cleft Palate Craniofac J* 2015;52(04):e88–e94
- Hoffman-Andrews L, Tarnowski JM, Lee S, et al. Characteristics of orofacial clefting in Hawai'i. *Hawaii J Health Soc Welf* 2019;78(08):258–261
- Dixon MJ, Marazita ML, Beaty TH, Murray JC. Cleft lip and palate: understanding genetic and environmental influences. *Nat Rev Genet* 2011;12(03):167–178
- Leslie EJ, Marazita ML. Genetics of cleft lip and cleft palate. *Am J Med Genet C Semin Med Genet* 2013;163C(04):246–258
- Neela PK, Gosla SR, Husain A, Mohan V. *CRISPLD2* gene polymorphisms with nonsyndromic cleft lip palate in Indian population. *Glob Med Genet* 2020;7(01):22–25
- Hagberg C, Larson O, Milerad J. Incidence of cleft lip and palate and risks of additional malformations. *Cleft Palate Craniofac J* 1998;35(01):40–45
- Ogle OE. Incidence of cleft lip and palate in a newborn Zairian sample. *Cleft Palate Craniofac J* 1993;30(02):250–251
- Wyszynski DF, Wu T. Use of US birth certificate data to estimate the risk of maternal cigarette smoking for oral clefting. *Cleft Palate Craniofac J* 2002;39(02):188–192
- Yang J, Carmichael SL, Canfield M, Song J, Shaw GM. National Birth Defects Prevention Study. Socioeconomic status in relation to selected birth defects in a large multicentered US case-control study. *Am J Epidemiol* 2008;167(02):145–154
- Croen LA, Shaw GM, Wasserman CR, Tolarová MM. Racial and ethnic variations in the prevalence of orofacial clefts in California, 1983–1992. *Am J Med Genet* 1998;79(01):42–47
- Mossey PA, Little J, Munger RG, Dixon MJ, Shaw WC. Cleft lip and palate. *Lancet* 2009;374(9703):1773–1785
- Parker SE, Mai CT, Canfield MA, et al; National Birth Defects Prevention Network. Updated National Birth Prevalence estimates for selected birth defects in the United States, 2004–2006. *Birth Defects Res A Clin Mol Teratol* 2010;88(12):1008–1016
- Neela PK, Gosla SR, Husain A, Mohan V, Thumuju S, Rajeshwari BV. Analysis of single nucleotide polymorphisms on locus 13q33.1–34 in multigenerational families of cleft lip palate using mass array. *Indones Biomed J*. 2020; In press
- Bender PL. Genetics of cleft lip and palate. *J Pediatr Nurs* 2000;15(04):242–249
- Neela PK, Gosla SR, Husain A, Mohan V, Thumuju S, Bv R. Association of *MAPK4* and *SOX1-OT* gene polymorphisms with cleft lip palate in multiplex families: a genetic study. *J Dent Res Dent Clin Dent Prospect* 2020;14(02):93–96
- Wyszynski DF, Wu T. Prenatal and perinatal factors associated with isolated oral clefting. *Cleft Palate Craniofac J* 2002;39(03):370–375
- Cobourne MT. Cleft Lip and Palate. *Epidemiology, Aetiology and Treatment*. *Front Oral Biol London*: Karger; 2012:60–70
- Singh I, Pal GP. *Human Embryology*. 7th ed. Delhi: Macmillan India Ltd; 2001:130–131
- Jiang R, Bush JO, Lidral AC. Development of the upper lip: morphogenetic and molecular mechanisms. *Dev Dyn* 2006;235(05):1152–1166
- Stanier P, Moore GE. Genetics of cleft lip and palate: syndromic genes contribute to the incidence of non-syndromic clefts. *Hum Mol Genet* 2004;13(Spec No 1):R73–R81
- Gritli-Linde A. Molecular control of secondary palate development. *Dev Biol* 2007;301(02):309–326
- Bush JO, Jiang R. Palatogenesis: morphogenetic and molecular mechanisms of secondary palate development. *Development* 2012;139(02):231–243
- Fogh Andersen P. *Inheritance of harelip and cleft palate*. Copenhagen: Busck; 1942
- Fraser FC, Baxter H. The familial distribution of congenital clefts of the lip and palate; a preliminary report. *Am J Surg* 1954;87(05):656–659

- 35 Fraser FC. The genetics of cleft lip and cleft palate. *Am J Hum Genet* 1970;22(03):336–352
- 36 Marazita ML, Spence MA, Melnick M. Genetic analysis of cleft lip with or without cleft palate in Danish kindreds. *Am J Med Genet* 1984;19(01):9–18
- 37 Mitchell LE. Mode of inheritance of oral clefts. In: Wyszyski DF, ed. *Cleft Lip and Palate: From Origin to Treatment*. New York, NY: Oxford University Press; 2002:234–239
- 38 Grosen D, Bille C, Pedersen JK, Skytthe A, Murray JC, Christensen K. Recurrence risk for offspring of twins discordant for oral cleft: a population-based cohort study of the Danish 1936–2004 cleft twin cohort. *Am J Med Genet A* 2010;152A(10):2468–2474
- 39 Chung KC, Kowalski CP, Kim HM, Buchman SR. Maternal cigarette smoking during pregnancy and the risk of having a child with cleft lip/palate. *Plast Reconstr Surg* 2000;105(02):485–491
- 40 Bhaskar LV, Murthy J, Venkatesh Babu G. Polymorphisms in genes involved in folate metabolism and orofacial clefts. *Arch Oral Biol* 2011;56(08):723–737
- 41 Sabbagh HJ, Innes NP, Sallout BI, et al. Birth prevalence of nonsyndromic orofacial clefts in Saudi Arabia and the effects of parental consanguinity. *Saudi Med J* 2015;36(09):1076–1083
- 42 Neela PK, Reddy SG, Husain A, Mohan V. Association of cleft lip and/or palate in people born to consanguineous parents: a 13-year retrospective study from a very high volume cleft center. *J Cleft Lip Palate Craniofacial Anomalies* 2019;6:33–37
- 43 FitzPatrick DR, Carr IM, McLaren L, et al. Identification of SATB2 as the cleft palate gene on 2q32–q33. *Hum Mol Genet* 2003;12(19):2491–2501
- 44 Britanova O, Depew MJ, Schwark M, et al. SATB2 haploinsufficiency phenocopies 2q32–q33 deletions, whereas loss suggests a fundamental role in the coordination of jaw development. *Am J Hum Genet* 2006;79(04):668–678
- 45 Leoyklang P, Suphapeetiporn K, Siriwan P, et al. Heterozygous nonsense mutation SATB2 associated with cleft palate, osteoporosis, and cognitive defects. *Hum Mutat* 2007;28(07):732–738
- 46 Grassia V, Lombardi A, Kawasaki H, et al. Salivary microRNAs as new molecular markers in cleft lip and palate: a new frontier in molecular medicine. *Oncotarget* 2018;9(27):18929–18938
- 47 Martinelli M, Scapoli L, Pezzetti F, et al. Suggestive linkage between markers on chromosome 19q13.2 and nonsyndromic orofacial cleft malformation. *Genomics* 1998;51(02):177–181
- 48 Park BY, Sull JW, Park JY, Jee SH, Beatty TH. Differential parental transmission of markers in BCL3 among Korean cleft case-parent trios. *J Prev Med Public Health* 2009;42(01):1–4
- 49 Lace B, Kempa I, Klovinis J, et al. BCL3 gene role in facial morphology. *Birth Defects Res A Clin Mol Teratol* 2012;94(11):918–924
- 50 François-Fiquet C, Poli-Merol ML, Nguyen P, Landais E, Gaillard D, Doco-Fenzy M. Role of angiogenesis-related genes in cleft lip/palate: review of the literature. *Int J Pediatr Otorhinolaryngol* 2014;78(10):1579–1585
- 51 Neela PK, Gosla SR, Husain A, Mohan V, Thumoju S, Rajeshwari BV. Association of nucleotide variants of GRHL3, IRF6, NAT2, SDC2, BCL3 and PVRL1 genes with non-syndromic cleft lip with/without cleft palate in multigenerational families: a retrospective study. *Contemp Clin Dent* 2020; In press
- 52 Talbot JC, Johnson SL, Kimmel CB. hand2 and Dlx genes specify dorsal, intermediate and ventral domains within zebrafish pharyngeal arches. *Development* 2010;137(15):2507–2517
- 53 Murthi P, Said JM, Doherty VL, et al. Homeobox gene DLX4 expression is increased in idiopathic human fetal growth restriction. *Mol Hum Reprod* 2006;12(12):763–769
- 54 Zhang L, Yang M, Gan L, et al. DLX4 upregulates TWIST and enhances tumor migration, invasion and metastasis. *Int J Biol Sci* 2012;8(08):1178–1187
- 55 Wu D, Mandal S, Choi A, et al. DLX4 is associated with orofacial clefting and abnormal jaw development. *Hum Mol Genet* 2015;24(15):4340–4352
- 56 He M, Bian Z. Association between DLX4 polymorphisms and nonsyndromic orofacial clefts in a Chinese Han population. *Cleft Palate Craniofac J* 2019;56(03):357–362
- 57 Bonczek O, Balcar VJ, Šerý O. PAX9 gene mutations and tooth agenesis: a review. *Clin Genet* 2017;92(05):467–476
- 58 Li R, Chen Z, Yu Q, Weng M, Chen Z. The function and regulatory network of PAX9 gene in palate development. *J Dent Res* 2019;98(03):277–287
- 59 Peters H, Neubüser A, Kratochwil K, Balling R. PAX9-deficient mice lack pharyngeal pouch derivatives and teeth and exhibit craniofacial and limb abnormalities. *Genes Dev* 1998;12(17):2735–2747
- 60 Schuffenhauer S, Leifheit HJ, Lichtner P, Peters H, Murken J, Emmerich P. De novo deletion (14)(q11.2q13) including PAX9: clinical and molecular findings. *J Med Genet* 1999;36(03):233–236
- 61 Das P, Hai M, Elcock C, et al. Novel missense mutations and a 288-bp exonic insertion in PAX9 in families with autosomal dominant hypodontia. *Am J Med Genet A* 2003;118A(01):35–42
- 62 Zhao JL, Chen YX, Bao L, Xia QJ, Wu TJ, Zhou L. Novel mutations of PAX9 gene in Chinese patients with oligodontia. *Zhonghua Kou Qiang Yi Xue Za Zhi* 2005;40(04):266–270
- 63 Sull JW, Liang KY, Hetmanski JB, et al. Maternal transmission effects of the PAX genes among cleft case-parent trios from four populations. *Eur J Hum Genet* 2009;17(06):831–839
- 64 Nakatomi M, Wang XP, Key D, et al. Genetic interactions between PAX9 and Msx1 regulate lip development and several stages of tooth morphogenesis. *Dev Biol* 2010;340(02):438–449
- 65 Song T, Wu D, Wang Y, Li H, Yin N, Zhao Z. SNPs and interaction analyses of IRF6, MSX1 and PAX9 genes in patients with nonsyndromic cleft lip with or without palate. *Mol Med Rep* 2013;8(04):1228–1234
- 66 de Araujo TK, Secolin R, Félix TM, et al. A multicentric association study between 39 genes and nonsyndromic cleft lip and palate in a Brazilian population. *J Craniomaxillofac Surg* 2016;44(01):16–20
- 67 Serafini T, Colamarino SA, Leonardo ED, et al. Netrin-1 is required for commissural axon guidance in the developing vertebrate nervous system. *Cell* 1996;87(06):1001–1014
- 68 Salminen M, Meyer BI, Bober E, Gruss P. Netrin 1 is required for semicircular canal formation in the mouse inner ear. *Development* 2000;127(01):13–22
- 69 Srinivasan K, Strickland P, Valdes A, Shin GC, Hinck L. Netrin-1/neogenin interaction stabilizes multipotent progenitor cap cells during mammary gland morphogenesis. *Dev Cell* 2003;4(03):371–382
- 70 Leslie EJ, Taub MA, Liu H, et al. Identification of functional variants for cleft lip with or without cleft palate in or near PAX7, FGFR2, and NOG by targeted sequencing of GWAS loci. *Am J Hum Genet* 2015;96(03):397–411
- 71 Guo Q, Li D, Meng X, et al. Association between PAX7 and NTN1 gene polymorphisms and nonsyndromic orofacial clefts in a northern Chinese population. *Medicine (Baltimore)* 2017;96:19 (e6724)
- 72 Braybrook C, Doudney K, Marçano AC, et al. The T-box transcription factor gene TBX22 is mutated in X-linked cleft palate and ankyloglossia. *Nat Genet* 2001;29(02):179–183
- 73 Braybrook C, Liso S, Doudney K, et al. Craniofacial expression of human and murine TBX22 correlates with the cleft palate and ankyloglossia phenotype observed in CPX patients. *Hum Mol Genet* 2002;11(22):2793–2804
- 74 Prescott NJ, Lees MM, Winter RM, Malcolm S. Identification of susceptibility loci for nonsyndromic cleft lip with or without cleft palate in a two stage genome scan of affected sib-pairs. *Hum Genet* 2000;106(03):345–350
- 75 Marçano AC, Doudney K, Braybrook C, et al. TBX22 mutations are a frequent cause of cleft palate. *J Med Genet* 2004;41(01):68–74
- 76 Suphapeetiporn K, Tongkobpetch S, Siriwan P, Shotelersuk V. TBX22 mutations are a frequent cause of non-syndromic cleft palate in the Thai population. *Clin Genet* 2007;72(05):478–483

- 77 Dai J, Xu C, Wang G, et al. Novel *TBX22* mutations in Chinese nonsyndromic cleft lip/palate families. *J Genet* 2018;97(02):411–417
- 78 Kantaputra PN, Paramee M, Kaewkhampa A, et al. Cleft lip with cleft palate, ankyloglossia, and hypodontia are associated with *TBX22* mutations. *J Dent Res* 2011;90(04):450–455
- 79 Shu X, Dong Z, Cheng L, Shu S. DNA hypermethylation of *Fgf16* and *Tbx22* associated with cleft palate during palatal fusion. *J Appl Oral Sci* 2019;27:e20180649
- 80 Takahashi K, Nakanishi H, Miyahara M, et al. Nectin/PRR: an immunoglobulin-like cell adhesion molecule recruited to cadherin-based adherens junctions through interaction with Afaadin, a PDZ domain-containing protein. *J Cell Biol* 1999;145(03):539–549
- 81 Suzuki K, Hu D, Bustos T, et al. Mutations of *PVRL1*, encoding a cell-cell adhesion molecule/herpesvirus receptor, in cleft lip/palate-ectodermal dysplasia. *Nat Genet* 2000;25(04):427–430
- 82 Sözen MA, Suzuki K, Tolarova MM, Bustos T, Fernández Iglesias JE, Spritz RA. Mutation of *PVRL1* is associated with sporadic, non-syndromic cleft lip/palate in northern Venezuela. *Nat Genet* 2001;29(02):141–142
- 83 Scapoli L, Palmieri A, Martinelli M, et al. Study of the *PVRL1* gene in Italian nonsyndromic cleft lip patients with or without cleft palate. *Ann Hum Genet* 2006;70(Pt 3):410–413
- 84 Oner DA, Tastan H. Identification of novel variants in the *PVRL1* gene in patients with nonsyndromic cleft lip with or without cleft palate. *Genet Test Mol Biomarkers* 2016;20(05):269–272
- 85 Avila JR, Jezewski PA, Vieira AR, et al. *PVRL1* variants contribute to non-syndromic cleft lip and palate in multiple populations. *Am J Med Genet A* 2006;140(23):2562–2570
- 86 Sözen MA, Hecht JT, Spritz RA. Mutation analysis of the *PVRL1* gene in Caucasians with nonsyndromic cleft lip/palate. *Genet Test Mol Biomarkers* 2009;13(05):617–621
- 87 Cheng HQ, Huang EM, Xu MY, Shu SY, Tang SJ. *PVRL1* as a candidate gene for nonsyndromic cleft lip with or without cleft palate: no evidence for the involvement of common or rare variants in southern Han Chinese patients. *DNA Cell Biol* 2012;31(07):1321–1327
- 88 Shu SY, Zhang MJ, Cheng HQ, et al. Mutation analysis of *PVRL1* in patients with non-syndromic cleft of the lip and/or palate in Guangdong. *Genet Mol Res* 2015;14(02):3400–3408
- 89 Yoshiura K, Machida J, Daack-Hirsch S, et al. Characterization of a novel gene disrupted by a balanced chromosomal translocation t(2;19)(q11.2;q13.3) in a family with cleft lip and palate. *Genomics* 1998;54(02):231–240
- 90 Warrington A, Vieira AR, Christensen K, et al. Genetic evidence for the role of loci at 19q13 in cleft lip and palate. *J Med Genet* 2006;43(06):e26
- 91 Ichikawa E, Watanabe A, Nakano Y, et al. *PAX9* and *TGFB3* are linked to susceptibility to nonsyndromic cleft lip with or without cleft palate in the Japanese: population-based and family-based candidate gene analyses. *J Hum Genet* 2006;51(01):38–46
- 92 Carter TC, Molloy AM, Pangilinan F, et al. Testing reported associations of genetic risk factors for oral clefts in a large Irish study population. *Birth Defects Res A Clin Mol Teratol* 2010;88(02):84–93
- 93 Wang PW, Eisenbart JD, Cordes SP, Barsh GS, Stoffel M, Le Beau MM. Human *KRML* (*MAFB*): cDNA cloning, genomic structure, and evaluation as a candidate tumor suppressor gene in myeloid leukemias. *Genomics* 1999;59(03):275–281
- 94 Beaty TH, Murray JC, Marazita ML, et al. A genome-wide association study of cleft lip with and without cleft palate identifies risk variants near *MAFB* and *ABCA4*. *Nat Genet* 2010;42(06):525–529
- 95 Zankl A, Duncan EL, Leo PJ, et al. Multicentric carpotarsal osteolysis is caused by mutations clustering in the amino-terminal transcriptional activation domain of *MAFB*. *Am J Hum Genet* 2012;90(03):494–501
- 96 Park JG, Tischfield MA, Nugent AA, et al. Loss of *MAFB* function in humans and mice causes Duane syndrome, aberrant extraocular muscle innervation, and inner-ear defects. *Am J Hum Genet* 2016;98(06):1220–1227
- 97 Mi N, Hao Y, Jiao X, et al. Association study of single nucleotide polymorphisms of *MAFB* with non-syndromic cleft lip with or without cleft palate in a population in Heilongjiang Province, northern China. *Br J Oral Maxillofac Surg* 2014;52(08):746–750
- 98 He Y, Huang L, Zheng Y, Chen JH, Tang S. Association of single nucleotide polymorphisms at 20q12 with nonsyndromic cleft lip with or without cleft palate in a Southern Chinese Han cohort. *Mol Genet Genomic Med* 2020;8(01):e1028
- 99 Imani MM, Lopez-Jornet P, Pons-Fuster López E, Sadeghi M. Polymorphic variants of *V-Maf* musculoaponeurotic fibrosarcoma oncogene homolog B (rs13041247 and rs11696257) and risk of non-syndromic cleft lip/palate: systematic review and meta-analysis. *Int J Environ Res Public Health* 2019;16(15):2792
- 100 Huang L, Liang X, Ou Y, Tang S, He Y. Association between 20q12 rs13041247 polymorphism and risk of nonsyndromic cleft lip with or without cleft palate: a meta-analysis. *BMC Oral Health* 2020;20(01):39
- 101 Liang X, Huang L, Ou Y, He Y, Tang S. Association between *MAFB* rs17820943 and rs6072081 polymorphism and risk of nonsyndromic cleft lip with or without cleft palate: a meta-analysis. *Br J Oral Maxillofac Surg* 2020;58(09):1065–1072
- 102 Riley BM, Mansilla MA, Ma J, et al. Impaired FGF signaling contributes to cleft lip and palate. *Proc Natl Acad Sci U S A* 2007;104(11):4512–4517
- 103 Xu H, Niu Y, Wang T, et al. Novel *FGFR1* and *KISS1R* mutations in Chinese Kallmann syndrome males with cleft lip/palate. *BioMed Res Int* 2015;2015:649698
- 104 Davies AF, Imaizumi K, Mirza G, et al. Further evidence for the involvement of human chromosome 6p24 in the aetiology of orofacial clefting. *J Med Genet* 1998;35(10):857–861
- 105 Shi J, Song T, Jiao X, Qin C, Zhou J. Single-nucleotide polymorphisms (SNPs) of the *IRF6* and *TFAP2A* in non-syndromic cleft lip with or without cleft palate (NSCLP) in a northern Chinese population. *Biochem Biophys Res Commun* 2011;410(04):732–736
- 106 Babu Gurrakonda V, Syed AH, Murthy J, V K S Lakkakula B. Association of *TFAP2A* gene polymorphism with susceptibility to non-syndromic cleft lip with or without palate risk in south Indian population. *Meta Gene* 2016;9:181–184
- 107 Cobourne MT. The complex genetics of cleft lip and palate. *Eur J Orthod* 2004;26(01):7–16
- 108 Shi M, Christensen K, Weinberg CR, et al. Orofacial cleft risk is increased with maternal smoking and specific detoxification-gene variants. *Am J Hum Genet* 2007;80(01):76–90
- 109 Hozyasz KK, Mostowska A, Surowiec Z, Jagodziński PP. [Genetic polymorphisms of *GSTM1* and *GSTT1* in mothers of children with isolated cleft lip with or without cleft palate]. *Przegl Lek* 2005;62(10):1019–1022
- 110 Watanabe A, Akita S, Tin NT, et al. A mutation in *RYK* is a genetic factor for nonsyndromic cleft lip and palate. *Cleft Palate Craniofac J* 2006;43(03):310–316
- 111 Smane L, Pilmann M. *IRF6*, *RYK*, and *PAX9* expression in facial tissue of children with cleft palate. *Int J Morphol* 2015;33:647–652
- 112 Scapoli L, Martinelli M, Pezzetti F, et al. Linkage disequilibrium between *GABRB3* gene and nonsyndromic familial cleft lip with or without cleft palate. *Hum Genet* 2002;110(01):15–20
- 113 Baroni T, Bellucci C, Lilli C, et al. Retinoic acid, GABA-ergic, and TGF-beta signaling systems are involved in human cleft palate fibroblast phenotype. *Mol Med* 2006;12(9-10):237–245
- 114 Vieira AR, Howe A, Murray JC. Studies of γ -aminobutyric acid type A receptor $\beta 3$ (*GABRB3*) and glutamic acid decarboxylase 67

- (GAD67) with oral clefts. *Am J Med Genet A* 2008;146A(21):2828–2830
- 115 Fontoura C, Silva RM, Granjeiro JM, Letra A. Further evidence of association of the *ABCA4* gene with cleft lip/palate. *Eur J Oral Sci* 2012;120(06):553–557
- 116 Beaty TH, Taub MA, Scott AF, et al. Confirming genes influencing risk to cleft lip with/without cleft palate in a case-parent trio study. *Hum Genet* 2013;132(07):771–781
- 117 Mi N, Hao Y, Jiao X, Zheng X, Shi J, Chen Y. A polymorphic marker associated with non-syndromic cleft lip with or without cleft palate in a population in Heilongjiang Province, northern China. *Arch Oral Biol* 2015;60(02):357–361
- 118 Peng HH, Chang NC, Chen KT, et al. Nonsynonymous variants in *MYH9* and *ABCA4* are the most frequent risk loci associated with nonsyndromic orofacial cleft in Taiwanese population. *BMC Med Genet* 2016;17(01):59

Web References

- National Center for Biotechnology Information (NCBI). Available at: <http://www.ncbi.nih.gov/>. Accessed August 20, 2020
- Online Mendelian Inheritance in Man (OMIM). Available at: <http://www.ncbi.nlm.nih.gov/omim/>. Accessed August 20, 2020