

# Polyadenylation of a functional mRNA controls gene expression in *Escherichia coli*

Géraldine Joanny<sup>1</sup>, Jacques Le Derout<sup>1</sup>, Dominique Bréchemier-Baey<sup>1</sup>, Valérie Labas<sup>2</sup>, Joelle Vinh<sup>2</sup>, Philippe Régnier<sup>1</sup> and Eliane Hajnsdorf<sup>1,\*</sup>

<sup>1</sup>Régulation de l'Expression Génétique chez les Microorganismes, UPR CNRS n° 9073, conventionnée avec l'Université Paris 7—Denis Diderot, Paris, France and <sup>2</sup>Neurobiologie et Diversité Cellulaire, UMR CNRS n° 7637, Paris, France

Received December 14, 2006; Revised and Accepted February 12, 2007

## ABSTRACT

Although usually implicated in the stabilization of mRNAs in eukaryotes, polyadenylation was initially shown to destabilize RNA in bacteria. All the data are consistent with polyadenylation being part of a quality control process targeting folded RNA fragments and non-functional RNA molecules to degradation. We report here an example in *Escherichia coli*, where polyadenylation directly controls the level of expression of a gene by modulating the stability of a functional transcript. Inactivation of poly(A)polymerase I causes over-expression of glucosamine-6-phosphate synthase (GlmS) and both the accumulation and stabilization of the *glmS* transcript. Moreover, we show that the *glmS* mRNA results from the processing of the *glmU-glmS* cotranscript by RNase E. Interestingly, the *glmU-glmS* cotranscript and the mRNA fragment encoding GlmU only slightly accumulated in the absence of poly(A)polymerase, suggesting that the endonucleolytically generated *glmS* mRNA harbouring a 5' monophosphate and a 3' stable hairpin is highly susceptible to poly(A)-dependent degradation.

## INTRODUCTION

It is well established now that mRNA decay has an integral role in the control of gene expression. mRNAs can be degraded by a number of mechanisms that act independently, in parallel or sequentially. Several cis- and trans-acting factors including higher order RNA structures (1,2), RNA-binding proteins (3–5) and translating ribosomes (6,7) control the sensitivity of an mRNA

molecule to degradation. Hairpin structures located at the 3' end of operons or in intercistronic regions protect mRNA molecules from digestion by 3'–5' exoribonucleases. To overcome these barriers, bacteria initiate RNA degradation by endoribonucleolytic cleavages mostly carried out by RNase E in *E.coli*. RNase III and RNase G have also been found to initiate RNA decay in certain cases (8,9). RNA fragments resulting from endonucleolytic cleavages can be either further degraded endonucleolytically or by the 3'–5' exonucleases, polynucleotide phosphorylase (PNPase), RNase II and RNase R [reviewed in (10)]. Oligo(A) tails can be added to RNA fragments resulting from endo- or exonucleolytic cleavages and promote their exonucleolytic degradation (11).

As opposed to their predominant role in eukaryotes, poly(A) tails act as RNA-destabilizing elements in bacteria (12). A destabilizing function for poly(A) tails has also been subsequently reported in mitochondria of higher plants (13), chloroplasts (14), and the nucleus of yeast and humans [reviewed in (15,16)]. The enzyme responsible for the majority of polyadenylation in *E.coli*, poly(A)polymerase (PAP I), is encoded by the *pcnB* gene (17). The consensus idea emerging from numerous studies of *pcnB* mutations is that the main function of polyadenylation is to facilitate degradation of short folded RNAs such as RNA fragments produced during the processing of longer RNA molecules (17–19) and of a non-functional mutated tRNA (20). This latter example suggested that poly(A)-dependent decay is involved in quality surveillance of bacterial RNA, as previously reported in yeast (21). A special function of polyadenylation is in the control of copy number of ColEI plasmids. The decay of the regulatory RNA I is initiated by RNase E cleavage five nucleotides from the 5' end, which activates very fast decay via the PAP I-PNPase pathway (22). There are also data indicating that PAP I may control the stability of primary transcripts (23,24).

\*To whom correspondence should be addressed. Tel: +33 1 58 41 51 26; Fax: +33 1 58 41 50 20; Email: Eliane.Hajnsdorf@ibpc.fr  
Present address:

Géraldine Joanny, Laboratory of Diabetes and Experimental Endocrinology Lab.4136 Clinical Science Department, IDIBELL — University of Barcelona Hospital Universitari de Bellvitge, Spain

For instance, mutations that inactivate RNase E-dependent decay render mRNA more sensitive to poly(A)-dependent degradation (25–27). In addition, Aiso *et al.* showed that the *pcnB* mutation affected degradation of *rmf* mRNA in stationary phase, but this effect could be indirect as it required *de novo* mRNA synthesis (24). However, there has been no evidence so far that polyadenylation directly controls protein levels by modulating mRNA stability.

In an attempt to identify genes whose expression is affected by mRNA polyadenylation, we compared proteins of a wild-type and PAP I-deficient strains on 1D and 2D PAGE gels (2D-gel). There were several differences and, in particular, one polypeptide; glucosamine-6-phosphate synthase (GlmS) was significantly more abundant in the mutant strain. In this work, we demonstrate that the overexpression of GlmS is correlated with the accumulation and stabilization of the monocistronic *glmS* mRNA resulting from the processing of the *glmU-glmS* cotranscript.

## MATERIALS AND METHODS

### Bacterial strains

The name and origin of the strains used here are listed in Table 1. New strains were constructed by transduction of the  $\Delta pcnB$  (Kan<sup>R</sup>) and *rng::Cm<sup>R</sup>* alleles of strains SK7988 and CA244 *cafA*, respectively, with bacteriophage P1. Cells were grown in a MOPS medium supplemented with 0.2% glucose (w/v) and all amino acids and vitamins (28,29), or in an LB medium at the temperature specified in the text.

### Functional lifetime of bulk mRNA

Cells were grown in MOPS medium supplemented with glucose and amino acids lacking methionine and cysteine and then treated with rifampicin (500 µg/ml; zero time). At timed intervals, 1 ml aliquots were labelled for 1 min with 25 µCi [<sup>35</sup>S] methionine (>1000 Ci/mmol) at 37°C. Cells were then treated with excess cold methionine and sodium azide for 5 min and pelleted. Frozen cells were resuspended in 0.1 M Tris-HCl, pH 8, 0.4 M NaCl, 0.1 mM EDTA, 1 mM 2-β-mercaptoethanol, lysed by sonication on ice, and total proteins were separated on 10% SDS-PAGE.

**Table 1.** *E. coli* strains used in this work

Strains	Relevant markers	Source/origin
N3433	HfrH, <i>lacZ43, λ, relA1, spoT1, thi1</i>	(52)
N3431	N3433 <i>rne3071<sup>ts</sup></i>	(52)
IBPC 903	N3433 $\Delta pcnB$ (Kan <sup>R</sup> )	This work
IBPC 904	N3431 $\Delta pcnB$ (Kan <sup>R</sup> )	This work
CA244	<i>lacZ, trp, relA, spoT</i>	(53)
CA244-PAP	CA244 $\Delta pcnB$ (Kan <sup>R</sup> )	(53)
IBPC 694	MG1693 <i>zad::Tn10</i> (Tet <sup>R</sup> )	(25)
IBPC 690	MG1693 <i>pcnB80, zad::Tn10</i> (Tet <sup>R</sup> )	(25)
IBPC 633	N3433 <i>rnc105</i>	(54)
IBPC 935	N3433 <i>rng::Cm<sup>R</sup></i>	(55)
IBPC 934	N3431 <i>rng::Cm<sup>R</sup></i>	This work
CA244- <i>cafA</i>	CA244 <i>rng::Cm<sup>R</sup></i>	(56)
SK7988	$\Delta pcnB$ (Kan <sup>R</sup> )	(26)

Gels were stained with Coomassie blue and dried, and the radioactive pattern was analyzed with a PhosphorImager.

### Immunoblotting

Total proteins were separated on 10% SDS-PAGE, the proteins were transferred to Hybond C-super (Amersham) by electroblotting. The membrane was incubated with polyclonal anti-GFAT (GFAT=human glucosamine-6P synthase enzyme) antibodies, which cross-react against *E. coli* GlmS. The bound antibodies were detected by reaction with [<sup>125</sup>I]-labelled protein A.

### RNA extraction and analysis

Templates for the synthesis of the different RNA probes were obtained by PCR amplification using the primers described in Table 2. The upstream primer includes the T7 promoter. RNA were synthesized by T7 RNA polymerase yielding uniformly labelled RNA with [ $\alpha$ -<sup>32</sup>P] UTP (30). 5S rRNA was probed with the 5'-labelled ACTACCATCGGCGCTACGGC oligonucleotide.

RNA preparation, reverse transcription (using the 5'-CGGTATTCCAGACGACGTAA primer complementary to the *glmS* mRNA) and Northern blotting on 10% agarose formaldehyde gel were previously described (31,32).

### 2D gel electrophoresis

Three independent cultures of N3433 and IBPC 903 were grown in LB at 37°C. 50 ml were withdrawn at OD<sub>650</sub>=0.5. Soluble protein extracts were prepared from cells sonicated in water, 20 mg/ml DNase and RNase were added after centrifugation at 10000 g for 30 min. First-dimension electrophoresis was carried out on a non-linear immobilized pH 3–11 gradient IPG strip (18 cm BioRad), and the second dimension by 12% SDS-PAGE. The six samples were analyzed in parallel. Densitometric quantification of the six blue-colloidal-stained 2D gels was performed using the PDQuest<sup>TM</sup> 2D Analysis software. The analysis of these gels revealed that expression of two polypeptides was greatly increased in the mutant. These protein spots were excised from gels for trypsin in-gel digestion and MALDI-TOF analysis.


## RESULTS

### PAP I deficiency modifies gene expression

The presence of 3' terminal hairpins has been shown to protect mRNAs against exonucleolytic degradation.

**Table 2.** PCR primers. Italics correspond to RNA polymerase sequence promoter of the T7 phage

Primer name	Primer sequence
Reverse GlmS	5'-ATGTGTGGAATTGTTGGCGCG
Forward GlmS	5'- <i>TAATACGACTCACTATAGGGCAGAGTCCC</i> GCCTTGTTTCA
Reverse GlmU	5'-ATGAGCGTAGTGATCCTTGCC
Forward GlmU	5'- <i>TAATACGACTCACTATAGGGCATCTTTG</i> TGCTCAACAAT



**Figure 1.** Effect of the *pcnB* mutation on functional mRNA stability and protein synthesis. (A) Cultures of N3433 and IBPC 903 (*pcnB*) were grown in MOPS medium containing glucose and amino acids lacking methionine and cysteine until mid-log phase. Cells were treated with rifampicin (zero time), and their protein-synthesizing capacity was monitored after 1, 3, 5, 7 and 9 min by pulse-labelling for 1 min with radioactive methionine. Soluble proteins were analyzed on SDS-PAGE. Left panel: Coomassie blue staining; right panel: autoradiograph. The positions of polypeptides overproduced in the *pcnB* strain are indicated by an arrow for the 66-kDa protein and by asterisks. (B) Comparison of the protein profile of N3433 (lane 1), CA244 (lane 3) and IBPC 694 (lane 5) wild-type strains to their *pcnB::kan<sup>R</sup>* (IBPC 903 and CA244-PAP, respectively) (lanes 2 and 4) and *pcnB80* (IBPC 690) (lane 6) derivatives. Cells were grown in LB medium until mid-log phase and soluble proteins analyzed by SDS-PAGE electrophoresis and revealed by Coomassie Blue staining.

The fact that the addition of poly(A) tails facilitates exonucleolytic degradation of tightly folded RNA prompted us to investigate whether polyadenylation affects stability of normal cellular mRNAs that possess stable 3' terminal hairpins. To do this, we examined the functional half-life of mRNAs by comparing the incorporation of radiolabelled methionine into proteins at fixed times after inhibiting RNA synthesis with rifampicin. Newly synthesized proteins were pulse-labelled with [<sup>35</sup>S] methionine at different times after the addition of rifampicin to exponentially growing cultures of the PAP<sup>+</sup> N3433 strain and of its *pcnB::kan<sup>R</sup>* derivative IBPC 903. Total proteins were then separated on SDS-polyacrylamide gels that were autoradiographed. Strikingly, the PAP I-deficient strain synthesized a

66-kDa polypeptide for at least 3 min after inhibition of transcription initiation, the protein was hardly visible in PAP I-proficient cells (Figure 1A, right panel). Consistent with this observation, Coomassie blue staining of the gel shows that a 66-kDa polypeptide dramatically accumulates in the absence of PAP I (Figure 1A, left panel). These data suggest that the mRNA coding this 66-kDa polypeptide is either more efficiently translated or more likely that it is more stable and hence more abundant in the absence of polyadenylation. Failure to detect the radioactive polypeptide in the mutant 5 min after transcription inhibition indicates that the corresponding mRNA is not exceptionally stable; its functional half-life is shorter than 2 min even in the absence of polyadenylation.


There were a few other differences in the pulse-labelled pattern of wt and *pcnB* strains. Two polypeptides whose expression is enhanced in the *pcnB* strain are indicated by asterisks (Figure 1A). It has been reported that overexpression of PAP I induces functional stabilization of *rne* and *pnp* mRNAs (33). However co-migration of the labelled proteins with a degradosome preparation did not detect any changes in expression of proteins corresponding to the sizes of these degradosome components.

In this work, we concentrate on the 66-kDa protein, which shows the strongest effect of *pcnB*-induced overproduction. In order to rule out possible strain or allele-specific effects of PAP I inactivation, we checked whether the 66-kDa polypeptide also accumulates in a strain carrying another *pcnB* allele and in different genetic backgrounds (Table 1 and Figure 1B). Indeed, the 66-kDa polypeptide accumulates in strains harbouring the *pcnB::kan<sup>R</sup>* (IBPC 903 and CA244-PAP) and the *pcnB80* point mutation (IBPC 694), but is not detected in their *pcnB<sup>+</sup>* parents. Moreover, this polypeptide is overproduced in the IBPC 903 mutant strain grown either in MOPS medium containing glucose and amino acids (Figure 1A) and in LB medium (Figure 1B). Strain N3433 and its *pcnB::kan<sup>R</sup>* derivative grown in LB medium were used in the experiments below.

### GlmS is upregulated in PAP I mutant

In order to identify the 66-kDa polypeptide overproduced in the mutant, soluble proteins extracted from wild-type and *pcnB* strains were analyzed by 2D gel electrophoresis. Figure 2A shows that two polypeptides of about 66 kDa, migrating to adjacent positions on the gel, were much more abundant in the mutant than in the wild-type strain. Both spots were excised, and mass spectrometry identified these proteins as glucosamine-6-phosphate synthase (GlmS) (molecular weight 66.9 kDa), one of which may correspond to a modified GlmS form, although no modification of this protein was previously reported in *E.coli*. Densitometric quantification of the blue-colloidal-stained 2D gels revealed that the two spots of GlmS are 10- and 30-fold more abundant in the *pcnB* mutant compared to the wild-type strains. Finally, Western blotting with anti-GlmS antibodies confirmed that a 66-kDa polypeptide is produced much more efficiently in the PAP I mutant than in the wild-type strain (Figure 2B).






**Figure 2.** The *pcnB* mutation increases the amount of GlmS. (A) Protein profile of N3433 (wt) and IBPC 903 (*pcnB*) strains grown in LB medium were analyzed by 2D-PAGE. Representative gels from experiments performed in triplicate were presented. Soluble proteins with isoelectric point (pI) values in the range 3.0–11.0 and molecular mass in the range of 26–170 kDa were resolved in 12% SDS-PAGE gels and colloidal blue stained for comparative analysis. Corresponding sections of representative gels show the increased synthesis of two spots in the *pcnB* strain compared with the wt. The two spots identified as GlmS by MALDI-TOF are indicated by arrows. Protein level were determined by densitometric quantification (PDQuest<sup>TM</sup> 2D analysis software, Biorad). (B) Western blot showing the GlmS polypeptide synthesized in wt and its *pcnB* derivative. Total proteins were loaded on 10% SDS-PAGE. Position of the GlmS protein used as a control is indicated on the left.

GlmS encodes the enzyme that synthesizes glucosamine-6-phosphate (GlcN6P), an essential precursor of lipopolysaccharides and peptidoglycan. The preceding gene on the chromosome, *glmU*, encodes the bifunctional enzyme glucosamine-1-phosphate acetyl transferase: UDP-GlcNAc synthetase (34,35), which synthesizes UDP-GlcNAc, the first dedicated component for the cell wall components. Previous work has shown that expression of both *glmS* and *glmU* is subject to regulation by the availability of amino sugars (35,36). Both genes are believed to be transcribed from two promoters located upstream of the first gene of the operon, *glmU*, and no promoter has been identified in the intergenic region.


#### The *glmU*-*glmS* primary transcript is processed in the intercistronic region

To investigate whether the overproduction of GlmS in cells lacking PAP I results from an accumulation of its mRNA, we compared the abundance of the *glmS* transcript in wild-type and PAP I-deficient strains. Total RNA was extracted from cells growing exponentially at 37°C, and amounts of this transcript were monitored on Northern blot. Probing for *glmS* detected a band of 1.9 kb that was about 16-fold more abundant in the *pcnB* strain than in the wild-type cells (Figure 3). A 1.9 kb transcript is sufficient to code for a protein of 66 kDa, but not for the *glmU*-*glmS* (*glmUS*) cotranscript. The very low amount of the 3.5-kb mRNA visible in Figure 3 led us to postulate that the expected *glmUS* cotranscript is very rapidly cleaved into the shorter 1.9-kb *glmS* mRNA detected in Figure 3.




**Figure 3.** The *pcnB* mutation increases amounts of the *glmS* mRNA. Total RNA (12 µg) from N3433 (wt) and IBPC 903 (*pcnB*) strains grown in LB medium at 37°C were analyzed on Northern blot and probed for the *glmS* mRNA (upper part). Reprobing of the same membrane with the 5S rRNA probe is shown on the lower part. Positions of rRNA markers are indicated on the left.

We therefore investigated whether the endonucleases involved in mRNA maturation and degradation of *E.coli* mRNAs, namely RNase III (*rnc*), RNase E (*rne*) and RNase G (*rng*), cleave the *glmUS* cotranscript. Isogenic strains deficient for each of these RNases were used as well as a *rne*-*rng* double mutant. The effect of RNase E inactivation was examined by using the thermosensitive *rne3071* allele. For comparison, RNAs from all strains were prepared from bacteria grown at 30°C (time 0) and 15 and 30 min after a temperature shift to 44°C required to inactivate thermosensitive RNase E. Total RNAs were then analyzed on Northern blots probed for *glmS* (Figure 4B). Neither the *rnc105*, nor the *rng::Cm<sup>R</sup>* mutations allowed detection of the *glmUS* primary transcript or an increased level of *glmS* mRNA. In contrast, RNase E inactivation caused the accumulation of a long transcript with the size of the expected *glmUS* mRNA (3.5 kb). This result demonstrated that the *glmUS* cotranscript is cleaved by RNase E. Moreover, inactivation of RNase G in the absence of RNase E increases the level of the dicistronic transcript, suggesting that in the absence of RNase E there is some processing of the long primary transcript by RNase G but that normally rapid degradation by RNase E eliminates any effect of *rng* mutation. Probing the Northern blot for the *glmU* mRNA confirms that the full-length 3.5-kb transcript was only



**Figure 4.** Fate of the transcripts of the *glmU-glmS* cotranscript. (A) Structure of the *glmU-glmS* transcript and of the processed mRNAs. The map of the *glmU-glmS* genes of *E. coli* shows the location of the 2 *glmU* promoters (arrows), of the transcription terminator (t) and the major RNase E processing site (black triangle). The structure of the primary transcript is indicated beneath the map, together with the mRNA species that were detected by Northern blot. The Northern blots do not distinguish between mRNAs starting at the two promoters. The small black box locates the oligonucleotide used in the primer extension experiment. Locations of the RNA used to probe for *glmU* and *glmS* are indicated by rectangles. (B) Northern blot analysis of transcripts of the *glmU-glmS* operon. Strains indicated at the top of the autoradiograph were grown at 30°C in LB medium and shifted to 44°C for the times indicated in minutes at the top of the autoradiograph. Total RNA (12 µg) were analyzed on Northern blot, which were probed for *glmS*, *glmU* mRNAs and 5S rRNA. 9S rRNA precursors accumulate after RNase E inactivation. As reported previously, *rne*<sup>+</sup> strains contain higher amounts of 5S rRNA after the shift to 44°C (51). (C) Overexposure of a Northern blot probed for *glmU* transcripts. (D) Primer extension analysis of the *glmU-glmS* intercistronic region. Total RNA (12 µg) from N3433 (wt) and IBPC 903 (*pcnB*) strains were used as templates for AMV reverse transcriptase with the 5' labelled *glmS* oligonucleotide (1 pmol). The extension products were analyzed on a 6% denaturing polyacrylamide gel. Molecular sizing markers are shown on the right side of the autoradiograph. The *glmS* primer was used with the Sequenase kit (Amersham) on a PCR amplified fragment to precisely locate the 5' extremities of the reverse transcriptase reactions (data not shown).



**Figure 5.** PAP I destabilizes the *glmS* transcript. (A) Autoradiograph comparing the decay kinetics of *glmS* mRNA in strains N3433 (wt) and IBPC 903 (*pcnB*). Times after inhibition of transcription at which *glmS* (upper part) and 5S (lower part) RNA contents were determined are indicated above each lane. (B) RNA levels were quantified by phosphorimager and plotted as a function of time. Quantification of the results.

present in the *rne* and *rne-rng* strains, and that it was more abundant in the double mutant (Figure 4B).

Surprisingly, thermal inactivation of RNase E leads not only to the accumulation of the uncleaved *glmUS* mRNA but also to an increased level of the downstream *glmS* monocistronic mRNA. The detection of the *glmS* monocistronic transcript implies that residual cleavage still occurs in the intergenic region of the *glmUS* transcript after the temperature shift that inactivates RNase E. RNase G is not responsible for this cleavage since the 1.9-kb *glmS* transcript was still detected in the double *rne-rng* strain. We previously observed residual processing of another RNase E site after thermal inactivation, implying that thermal inactivation of RNase E carrying the *rne3071* mutation is not complete (31). Alternatively, the *glmS* transcript could be due to another unidentified endonuclease, or to the presence of a *glmS*-specific promoter located near the intergenic region. We note that the monocistronic *glmS* transcript is somewhat diffuse in the *rne* and *rne-rng* strains, which might be consistent with it being due to something other than cleavage at the major RNase E site. The observation that the *glmS* transcript is less abundant in *rnc* cells than in wt while no *glmUS* transcript is detected (Figure 4B lanes 1–3 and 13–15, respectively) suggests that RNase III may indirectly participate to *glmS* stability.

To localize the processing site that generates the 1.9-kb *glmS* mRNA, primer extension was performed with an oligonucleotide hybridizing to the beginning of the *glmS* structural gene. Three 5' extremities were located, all mapping in the *glmU-glmS* intergenic region. The most abundant (band a on Figure 4D) is located just downstream of the stop codon of *glmU* mRNA. The AU-rich sequence surrounding this extremity

(AAAAAGUGA↓UUCU) suggests that it might be generated by RNase E. Two minor 5' ends located in the intercistronic region were also observed when PAP I was inactive. One (band b) maps 20 nt downstream of the major cleavage (band a) in the sequence CACAU↓GGG, and the other (band c) maps in a GC-rich sequence that does not conform to the usual RNase E cleavage site sequence (Figure 4D).

### Polyadenylation promotes degradation of the *glmS* transcript

We next investigated whether the abundance of the *glmS* mRNA in the *pcnB* strain reflected the stabilization of this molecule. For this purpose, we measured the half-life of the *glmS* transcript in wild-type and *pcnB* strains at 37°C. The very low level of the bicistronic transcript detected under these conditions prevents bias due to the production of new *glmS* molecules during the experiment. The intracellular concentration of the monocistronic *glmS* mRNA was followed versus time after rifampicin addition (Figure 5A and B). The same Northern blot was probed for 5S rRNA to normalize the data. Figure 5A and B show that the *glmS* mRNA was more stable in the *pcnB* strain where its half-life was about  $2.95 \pm 0.17$  min. The very low amount of *glmS* mRNA in the wild-type strain disappeared very rapidly; it was no longer detected 1.5 min after rifampicin addition. Its half-life was estimated to be  $0.72 \pm 0.17$  min. The relatively short half-life of the *glmS* transcript in the *pcnB* strain agrees with the functional half-life measurement showing that GlmS synthesis was already considerably reduced 3 min after inhibition of transcription (Figure 1A). These data clearly demonstrate that PAP I inactivation stabilizes the *glmS* monocistronic mRNA. However, the ratio of half-lives is only a factor of about four, which is much less than the ratio of the abundance of the *glmS* transcript levels observed in the *pcnB* and the wild-type strains (about 16-fold) (Figures 3 and 5A) or of the GlmS protein on 2D gels (Figure 2A). It is clear that the *pcnB* mutation has some additional effect on the level of the *glmS* transcript. Whether this involves the RNase E processing of the bicistronic transcript or activation of additional transcription from the *glmU* promoters or a *glmS* specific promoter has not been clarified.

### Poly(A)polymerase specifically affects the stability of the *glmS* mRNA

A typical Rho-independent terminator is located after the *glmS* gene, and one interpretation of the data described above is that polyadenylation facilitates exonucleolytic degradation of the 3'-terminal G-C rich hairpin of the transcription terminator, which otherwise protects the *glmS* transcript. Surprisingly, however, the *glmUS* cotranscript, which should have the same 3' terminal-protecting motif, is much less abundant than the *glmS* mRNA in the absence of PAP I. RNAs extracted from wild-type cells and isogenic bacteria deficient for RNase E or PAP I or both RNase E and PAP I were analyzed on Northern blots and probed for *glmS* and *glmU* mRNAs (Figure 4B and C). The fact that the amount of the dicistronic mRNA detected in the *rne-pcnB* strain was



slightly more abundant than in *rne* cells suggested that the long primary transcript was, to some extent, sensitive to poly(A)-assisted exonucleolytic decay. In agreement with this hypothesis, a smear of *glmS* mRNAs, ranging from 1.9 to 3.5 kb, which presumably results from exonucleolytic degradation of the cotranscript, was detected when PAP I was active but not when it was inactive. However, the monocistronic *glmS* mRNA accumulated much more strongly than the dicistronic transcript under identical conditions (i.e. after thermal inactivation of RNase E in the *pcnB* strain). These data reinforce the above observation that thermal inactivation of *rne3071* is not complete or that some other mechanism exists to generate the 5' end of the *glmS* mRNA. Irrespective of its origin, these data demonstrated that the monocistronic *glmS* mRNA is much more sensitive to poly(A)-dependent degradation than the *glmUS* cotranscript. Moreover, accumulation of both the *glmS* monocistronic and the *glmUS* dicistronic transcripts in a *pcnB* strain after RNase E inactivation confirms (see above) that they are both degraded by RNase E. Another unexpected observation was that the *glmS* mRNA, which accumulates in the *pcnB-rne*<sup>+</sup> strain growing at 30°C (Figure 4B, lane 16) and 37°C (Figure 3), disappeared after the temperature shift to 44°C (Figure 4B, lanes 17–18), suggesting that the RNase E-mediated degradation of *glmS* mRNA is more efficient at 44°C than at 37°C.

#### Degradation of the *glmU* transcript

Cleavage of the *glmUS* cotranscript would be expected to generate a *glmU*-specific transcript that is not detected in the wt strain (Figure 4B, lane 13). However in the *rne* strain after temperature shift, an additional transcript of about 1.5 kb is observed with the *glmU* probe (Figure 4B, lanes 11–12 and 4C lanes 2–3). Again, this transcript must be either due to residual RNase E activity in the thermal inactivated cultures, or the result of another endonuclease. As the transcript is only slightly more abundant in the *rne-pcnB* strain than the *rne* strain (Figure 4C, lanes 5–6 and 2–3, respectively), it is not dependent on polyadenylation for its degradation, unlike the *glmS* monocistronic transcript.

#### DISCUSSION

We present here evidence that poly(A)polymerase can negatively affect gene expression. Previously, it was thought that the role of the poly(A)-dependent pathway was to increase the turnovers of defective tRNAs (37) and non-processed RNA precursors (38), and to clear the cell of endonucleolytically produced, tightly folded mRNA fragments resistant to further degradation by exonucleases (39,40). Here, we show that overproduction of glucosamine-6P synthase, an essential enzyme in *E.coli* in the absence of exogenous aminosugars, occurs upon inactivation of PAP I, and that this accumulation is correlated with the accumulation of the *glmS* mRNA and the reduction in its decay-rate. This led us to conclude that poly(A)-dependent degradation of the functional *glmS*

mRNA determines the yield of GlmS. In agreement with our observation, *glmS* ORF was recently shown, using macroarrays, to be highly polyadenylated when PAP I is overproduced (41). One cannot exclude, however, that polyadenylation acts indirectly on GlmS expression, e.g. by modulating the stability of a small regulatory RNA or another trans-acting factor.

We also demonstrate that the *glmS* monocistronic mRNA, stabilized in PAP I-deficient cells, likely results from the endonucleolytic maturation of the *glmUS* dicistronic primary transcript. Although the dicistronic transcript was only detected in cells lacking RNase E, the fact that the monocistronic *glmS* transcript is still produced in these bacteria suggests that *glmUS* could be processed by another enzyme. The 5' mRNA extremity mapping just downstream of the UGA stop codon suggests that ribosome stalling could be implicated in the mechanism of *glmUS* RNA processing (42–45).

Interestingly, in spite of the fact that the *glmS* and *glmUS* mRNAs have identical 3' terminal structures, we found that the primary transcript is much less sensitive to the poly(A)-assisted exonucleolytic route of decay than the monocistronic processed molecule. An important question arising is how the poly(A)-assisted machinery of decay recognizes its targets and, in the case described above, how it distinguishes the *glmS* monocistronic mRNA from *glmUS* primary transcript and other primary transcripts. The fact that the *glmS* and *glmUS* mRNA correspond to the 3' part of the dicistronic transcript implies that it is the nature of the 5' part of the RNA that determines its sensitivity to this pathway. One possibility is that, as described earlier for RNA I, the 5' monophosphorylated terminus generated by RNase E facilitates polyadenylation of the *glmS* mRNA and thus promotes its degradation (22). Such a model could explain why the RNase E-processed mRNA is more sensitive to the poly(A)-dependent pathway than the primary transcript harbouring a 5' triphosphate extremity. At the moment, the mechanism by which the 5' end of the long 1.9-kb *glmS* mRNA could affect a reaction occurring at its 3' end is not known. A greater accessibility of the 3' end has been proposed to explain why the precursor of a mutated tRNA<sup>Trp</sup> is degraded by a poly(A)-dependent pathway (37). One can imagine that refolding of the *glmS* mRNA consecutive to the processing of the primary transcript improves accessibility of PAP I and/or exoribonucleases at its 3' end. It is possible, for example, that translation of *glmS* is coupled to that of the upstream gene *glmU* and that a reduction of ribosome loading of the *glmS* message consecutive to the removal of the upstream *glmU* cistron modifies the folding of the molecule.

It is worth mentioning here that the 5' monophosphate extremity that may trigger the poly(A)-dependent decay of the *glmS* mRNA may also promote its degradation by RNase E (46). The fact that the *glmS* mRNA accumulates in the absence of PAP I at 30°C, 37°C but not at 44°C is consistent with the idea that the efficiency of its degradation by RNase E increases at high temperature. In contrast, this observation indicates that the contribution of the poly(A)-dependent pathway to the decay of this message increases at low temperature. It therefore appears

that RNase E exerts a predominant role in mRNA decay around the optimum temperature of growth and that the poly(A)-dependent degradation may substitute for RNase E at lower temperatures. The idea that poly(A)-dependent decay can substitute for RNase E is also supported by earlier data showing that this pathway becomes effective on mRNAs that are no longer degraded by RNase E (25–27).

GlmS catalyzes the conversion of fructose-6-phosphate into glucosamine-6-phosphate, which then undergoes sequential transformations leading to the formation of UDP-N-acetylglucosamine, the major intermediate in the biosynthesis of all amino-sugar-containing macromolecules in the cell (47). Regulation of *glmS* expression occurs both at the transcriptional level (36) with NagC acting as both an activator and a repressor of the *glmUS* operon in *E.coli* (35) and at the posttranscriptional level (present work). Interestingly, the expression of the *glmS* gene of *Bacillus subtilis* is also subject to a regulatory mechanism involving the processing and the degradation of its mRNA. However, in these bacteria lacking orthologs of both RNase E and PAP I (48,49), an alternate strategy has emerged. A new class of ribozyme that cleaves the messenger of the *glmS* mRNA was discovered, which is activated by GlcN6P (50). In spite of some sequence homology in the *glmU-glmS* intergenic region in bacterial species closely related to *E.coli*, in-line probing did not reveal any structural changes or self-cleavage activity when the entire *E.coli* intergenic *glmU-glmS* region was incubated with either Glc6P or GlcN6P (Elaine Lee and Ron Breaker, personal communication). It therefore appears that different strategies exist that allow sensitive control of *glmS* mRNA and hence protein levels despite the fact that the machineries of mRNA processing and degradation are very different. Such pathways may present selective advantages that could explain their conservation in Gram-positive and Gram-negative bacteria.

## ACKNOWLEDGEMENTS

We are grateful to J. Plumbridge for careful critical reading of the manuscript, discussions and her constant interest during this work. We thank M. P. Deutscher and S. D. Kushner for kindly supplying strains, M. A. Badet for antibodies and GlmS purified protein and E. Lee and R. Breaker for communication of unpublished results. Funding to pay the Open Access publication charge was provided by CNRS Departement des Sciences de la Vie.

*Conflict of interest statement.* None declared.

## REFERENCES

- Belasco, J.G., Nillson, G., von Gabain, A. and Cohen, S.N. (1986) The stability of *E.coli* gene transcripts is dependent on determinants localized to specific mRNA segments. *Cell*, **46**, 245–251.
- Bouvet, P. and Belasco, J. (1992) RNA degradation in *E.coli* by endonuclease RNase E: control by 5'-terminal base pairing. *Nature*, **360**, 488–491.
- Folichon, M., Arluisson, V., Pellegrini, O., Huntzinger, E., Regnier, P. and Hajnsdorf, E. (2003) The poly(A) binding protein Hfq protects RNA from RNase E and exoribonucleolytic degradation. *Nucleic Acids Res.*, **31**, 7302–7310.
- Feng, Y., Huang, H., Liao, J. and Cohen, S.N. (2001) Escherichia coli poly(A) binding proteins that interact with components of degradosomes or impede RNA decay mediated by polynucleotide phosphorylase and RNase E. *J. Biol. Chem.*, **276**, 31651–31656.
- Iost, I. and Dreyfus, M. (1994) mRNAs can be stabilized by DEAD-box proteins. *Nature*, **372**, 193–196.
- Braun, F., Le Derout, J. and Régnier, P. (1998) Ribosomes inhibit an RNase E cleavage which induces the decay of the rpsO mRNA of *Escherichia coli*. *EMBO J.*, **17**, 4790–4797.
- Hambraeus, G., Karhumaa, K. and Rutberg, B. (2002) A 5' stem-loop and ribosome binding but not translation are important for the stability of *Bacillus subtilis* aprE leader mRNA. *Microbiology*, **148**, 1795–1803.
- Bardwell, J.C.A., Régnier, P., Chen, S.M., Nakamura, Y., Grunberg-Manago, M. and Court, D.L. (1989) Autoregulation of RNase III operon by mRNA processing. *EMBO J.*, **8**, 3401–3407.
- Wachi, M., Kaga, N., Umitsuki, G., Clark, D.P. and Nagai, K. (2001) A novel RNase G mutant that is defective in degradation of adhe mRNA but proficient in the processing of 16S rRNA precursor. *Biochem. Biophys. Res. Com.*, **289**, 1301–1306.
- Deutscher, M.P. (2006) Degradation of RNA in bacteria: comparison of mRNA and stable RNA. *Nucleic Acids Res.*, **34**, 659–666.
- Haugel-Nielsen, J., Hajnsdorf, E. and Regnier, P. (1996) The rpsO mRNA of *Escherichia coli* is polyadenylated at multiple sites resulting from endonucleolytic processing and exonucleolytic degradation. *EMBO J.*, **15**, 3144–3152.
- Dreyfus, M. and Régnier, P. (2002) The poly(A) tail of mRNAs: bodyguard in eukaryotes, scavenger in bacteria. *Cell*, **111**, 611–613.
- Gagliardi, D., Stepien, P.P., Temperley, R.J., Lightowlers, R.N. and Chrzanoswska-Lightowlers, Z.M. (2004) Messenger RNA stability in mitochondria: different means to an end. *Trends Genet.*, **20**, 260–267.
- Komine, Y., Kwong, L., Anguera, M.C., Schuster, G. and Stern, D.B. (2000) Polyadenylation of three classes of chloroplast RNA in *Chlamydomonas reinhardtii*. *RNA*, **6**, 598–607.
- Anderson, J.T. (2005) RNA turnover: unexpected consequences of being tailed. *Curr. Biol.*, **15**, R635–638.
- West, S., Gromak, N., Norbury, C.J. and Proudfoot, N.J. (2006) Adenylation and exosome-mediated degradation of cotranscriptionally cleaved pre-messenger RNA in human cells. *Mol. Cell*, **21**, 437–443.
- Cao, G.-J. and Sarkar, N. (1992) Poly(A) RNA in *Escherichia coli*: nucleotide sequence at the junction of the lpp transcript and the polyadenylate moiety. *Proc. Natl. Acad. Sci. USA*, **89**, 7546–7550.
- He, L., Söderbom, F., Wagner, E.G.H., Binnie, U., Binns, N. and Masters, M. (1993) PcnB is required for the rapid degradation of RNAI, the antisense RNA that controls the copy number of ColE1-related plasmids. *Mol. Microbiol.*, **9**, 1131–1142.
- Xu, F., Lin-Chao, S. and Cohen, S.N. (1993) The *Escherichia coli* pcnB gene promotes adenylation of antisense RNAI of ColE1-type plasmids in vivo and degradation of RNAI decay intermediates. *Proc. Natl. Acad. Sci. USA*, **90**, 6756–6760.
- Deutscher, M.P. (2003) Degradation of stable RNA in bacteria. *J. Biol. Chem.*, **278**, 45041–45044.
- Kadaba, S., Krueger, A., Trice, T., Krecic, A.M., Hinnebusch, A.G. and Anderson, J. (2004) Nuclear surveillance and degradation of hypomodified initiator tRNAMet in *S. cerevisiae*. *Genes Dev.*, **18**, 1227–1240.
- Xu, F. and Cohen, S.N. (1995) RNA degradation in *Escherichia coli* regulated by 3' adenylation and 5' phosphorylation. *Nature*, **374**, 180–183.
- Mohanty, B.K. and Kushner, S.R. (1999) Analysis of the function of *Escherichia coli* poly(A) polymerase I in RNA metabolism. *Mol. Microbiol.*, **34**, 1094–1108.
- Aiso, T., Yoshida, H., Wada, A. and Ohki, R. (2005) Modulation of mRNA stability participates in stationary-phase-specific expression of ribosome modulation factor. *J. Bacteriol.*, **187**, 1951–1958.
- Hajnsdorf, E., Braun, F., Haugel-Nielsen, J. and Regnier, P. (1995) Polyadenylation destabilizes the rpsO mRNA of *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*, **92**, 3973–3977.
- O'Hara, E.B., Chekanova, J.A., Ingle, C.A., Kushner, Z.R., Peters, E. and Kushner, S.R. (1995) Polyadenylation helps regulate mRNA



- decay in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*, **92**, 1807–1811.
27. Marujo, P.E., Braun, F., Haugel-Nielsen, J., Le Derout, J., Arraiano, C.M. and Regnier, P. (2003) Inactivation of the decay pathway initiated at an internal site by RNase E promotes poly(A)-dependent degradation of the rpsO mRNA in *Escherichia coli*. *Mol. Microbiol.*, **50**, 1283–1294.
  28. Neidhardt, F.C., Bloch, P.L., Pedersen, S. and Reeh, S. (1977) Chemical measurement of steady-state levels of ten aminoacyl-transfer ribonucleic acid synthetases in *Escherichia coli*. *J. Bacteriol.*, **129**, 378–387.
  29. Neidhardt, F.C., Bloch, P.L. and Smith, D.F. (1974) Culture medium for enterobacteria. *J. Bacteriol.*, **119**, 736–747.
  30. Hajnsdorf, E. and Régnier, P. (2000) Host factor HFq of *Escherichia coli* stimulates elongation of poly(A) tails by poly(A) polymerase I. *Proc. Natl. Acad. Sci. USA*, **97**, 1501–1505.
  31. Hajnsdorf, E., Carpousis, A.J. and Régnier, P. (1994) Nucleolytic inactivation and degradation of the RNase III processed pnp message encoding polynucleotide phosphorylase of *Escherichia coli*. *J. Mol. Biol.*, **239**, 439–454.
  32. Hajnsdorf, E., Steier, O., Coscoy, L., Teyssset, L. and Régnier, P. (1994) Roles of RNase E, RNase II and PNPase in the degradation of the rpsO transcripts of *Escherichia coli*: stabilizing function of RNase II and evidence for efficient degradation in an ams-rnb-pnp mutant. *EMBO J.*, **13**, 3368–3377.
  33. Mohanty, B.K. and Kushner, S.R. (2002) Polyadenylation of *Escherichia coli* transcripts plays an integral role in regulating intracellular levels of polynucleotide phosphorylase and RNase E. *Mol. Microbiol.*, **45**, 1315–1324.
  34. Mengin-Lecreulx, D. and van Heijenoort, J. (1994) Copurification of glucosamine-1-phosphate acetyltransferase and N-acetylglucosamine-1-phosphate uridylyltransferase activities of *Escherichia coli*: characterization of the glmU gene product as a bifunctional enzyme catalyzing two subsequent steps in the pathway for UDP-N-acetylglucosamine synthesis. *J. Bacteriol.*, **176**, 5788–5795.
  35. Plumbridge, J. (1995) Co-ordinated regulation of amino sugar biosynthesis and degradation: the NagC repressor acts as both an activator and a repressor for the transcription of the glmUS operon and requires two separated NagC binding sites. *EMBO J.*, **14**, 3958–3965.
  36. Plumbridge, J.A., Cochet, O., Souza, J.M., Altamirano, M.M., Calcagno, M.L. and Badet, B. (1993) Coordinated regulation of amino sugar-synthesizing and -degrading enzymes in *Escherichia coli* K-12. *J. Bacteriol.*, **175**, 4951–4956.
  37. Li, Z., Reimers, S., Pandit, S. and Deutscher, M.P. (2002) RNA quality control: degradation of defective transfer RNA. *EMBO J.*, **21**, 1132–1138.
  38. Li, Z. and Deutscher, M.P. (2002) RNase E plays an essential role in the maturation of *Escherichia coli* tRNA precursors. *RNA*, **8**, 97–109.
  39. Kushner, S.R. (2002) mRNA decay in *Escherichia coli* comes of age. *J. Bacteriol.*, **184**, 4658–4665.
  40. Hajnsdorf, E. and Régnier, P. (1999) *E. coli* rpsO mRNA decay: RNase E processing at the beginning of the coding sequence stimulates poly(A)-dependent degradation of the mRNA. *J. Mol. Biol.*, **286**, 1033–1043.
  41. Mohanty, B.K. and Kushner, S.R. (2006) The majority of *Escherichia coli* mRNAs undergo post-transcriptional modification in exponentially growing cells. *Nucleic Acids Res.*, **34**, 5695–5704.
  42. Pedersen, K., Zavialov, A.V., Pavlov, M.Y., Elf, J., Gerdes, K. and Ehrenberg, M. (2003) The bacterial toxin relE displays codon-specific cleavage of mRNAs in the ribosomal A site. *Cell*, **112**, 131–140.
  43. Christensen, S.K. and Gerdes, K. (2003) RelE toxins from Bacteria and Archaea cleave mRNAs on translating ribosomes, which are rescued by tmRNA. *Mol. Microbiol.*, **48**, 1389–1400.
  44. Sunohara, T., Jojima, K., Tagami, H., Inada, T. and Aiba, H. (2004) Ribosome stalling during translation elongation induces cleavage of mRNA being translated in *Escherichia coli*. *J. Biol. Chem.*, **279**, 15368–15375.
  45. Yamanishi, H. and Yonesaki, T. (2005) RNA cleavage linked with ribosomal action. *Genetics*, **171**, 419–425.
  46. Mackie, G.A. (1998) Ribonuclease E is a 5'-end-dependent endonuclease. *Nature*, **395**, 720–723.
  47. Badet, B., Vermoote, P., Haumont, P.Y., Lederer, F. and LeGoffic, F. (1987) Glucosamine synthetase from *Escherichia coli*: purification, properties, and glutamine-utilizing site location. *Biochemistry*, **26**, 1940–1948.
  48. Raynal, L.C., Krisch, H.M. and Carpousis, A.J. (1998) The *Bacillus subtilis* nucleotidyltransferase is a tRNA CCA-adding enzyme. *J. Bacteriol.*, **180**, 6276–6282.
  49. Condon, C. (2003) RNA processing and degradation in *Bacillus subtilis*. *Microbiol. Mol. Biol. Rev.*, **67**, 157–174.
  50. Winkler, W.C., Nahvi, A., Roth, A., Collins, J.A. and Breaker, R.R. (2004) Control of gene expression by a natural metabolite-responsive ribozyme. *Nature*, **428**, 281–286.
  51. Le Derout, J., Régnier, P. and Hajnsdorf, E. (2002) Both temperature and medium composition regulate RNase E processing efficiency of the rpsO mRNA coding for ribosomal protein S15 of *Escherichia coli*. *J. Mol. Biol.*, **319**, 341–349.
  52. Goldblum, K. and Apirion, D. (1981) Inactivation of the ribonucleic acid-processing enzyme RNase E blocks cell division. *J. Bacteriol.*, **146**, 128–132.
  53. Reuven, N.B., Zhou, Z. and Deutscher, M.P. (1997) Functional overlap of tRNA nucleotidyltransferase, poly(A) polymerase I and polynucleotide phosphorylase. *J. Biol. Chem.*, **272**, 33255–33259.
  54. Régnier, P. and Hajnsdorf, E. (1991) Decay of mRNA encoding ribosomal protein S15 of *Escherichia coli* is initiated by an RNase E-dependent endonucleolytic cleavage that removes the 3' stabilizing stem and loop structure. *J. Mol. Biol.*, **217**, 283–292.
  55. Bardey, V., Vallet, C., Robas, N., Charpentier, B., Thouvenot, B., Mougou, A., Hajnsdorf, E., Regnier, P., Springer, M. *et al.* (2005) Characterization of the molecular mechanisms involved in the differential production of erythrose-4-phosphate dehydrogenase, 3-phosphoglycerate kinase and class II fructose-1,6-bisphosphate aldolase in *Escherichia coli*. *Mol. Microbiol.*, **57**, 1265–1287.
  56. Abe, H., Abo, T. and Aiba, H. (1999) Regulation of intrinsic terminator by translation in *Escherichia coli*: transcription termination at a distance downstream. *Genes to Cells*, **4**, 87–97.