



Draft Genome Sequence of a Novel *Rhizobium* Species Isolated from the Marine Macroalga *Codium fragile* (Oyster Thief)

Christopher H. Miller,^a Chanel Taylor,^a  Jeremy G. Owen,^a  Wayne M. Patrick,^a  Chelsea J. Vickers^a

^aCentre for Biodiscovery, School of Biological Sciences, Victoria University of Wellington, Wellington, New Zealand

Christopher H. Miller and Chanel Taylor contributed equally to this work. Author order was determined alphabetically.

ABSTRACT In the process of studying the relationship between marine macroalgae and their bacterial symbionts, we isolated a new species of *Rhizobium*, which we designated *Rhizobium* sp. nov. C1 (for “*Codium* 1”). Here, we report the complete genome sequence of *Rhizobium* sp. nov. C1.

Oyster Thief (*Codium fragile*) is a green macroalga that is established worldwide and is becoming of increasing interest to researchers due to the identification of the high-value bioactive compounds it produces (1–5). We isolated and sequenced the genome of a bacterium associated with *C. fragile* that we subsequently identified as the Gram-negative *Rhizobium* sp. nov. C1.

The bacterium was isolated from *C. fragile* samples collected from a shoreline in Tauranga, New Zealand. Marine minimal seawater medium [0.5 M NaCl, 17 mM Na₂HPO₄, 11 mM KH₂PO₄, 4 mM (NH₄)₂SO₄, 0.8 mM MgSO₄·7H₂O, and 0.3 mM CaCl₂·2H₂O] was supplemented with trace elements (6) and inoculated with *C. fragile* (10% [wt/vol]). The culture was grown in a shaking incubator (200 rpm) at 25°C. Cultures were subsequently subcultured every 2 to 5 days over a period of 2 weeks. Growth from liquid culture was spread onto BD Difco marine agar 2216. Repeated single-colony selection from marine agar was undertaken until a clonal strain was isolated, as determined by its uniform colony appearance and uniform growth rates upon restreaking. A stock of this strain was made by culturing the strain in BD Difco marine broth 2216 at 25°C, adding glycerol (25% [vol/vol]), and storing the strain at –80°C.

Rhizobium bacteria are of interest as important plant symbionts (7, 8), but their role as algal symbionts are less understood; therefore, we sequenced the genome of this new species. A single colony was used to inoculate 5 mL of Zobell marine broth, which was incubated for 15 h at 25°C with shaking. From this culture, genomic DNA was extracted using the Wizard genomic DNA purification kit (Promega) with the Gram-negative bacteria protocol. The library was prepared using in-house Tn5-mediated tagmentation (9), followed by PCR enrichment with KAPA HiFi HotStart ReadyMix (Roche). DNA fragments of 400 bp were size selected before 150-bp paired-end reads were generated on an Illumina HiSeq 4000 system by Genewiz (China). Other sequencing and assembly statistics are shown in Table 1.

Genome assembly was achieved using raw reads (no reads were trimmed or discarded on grounds of quality, which was verified with FastQC v.0.11.7 [10]). Reference assembly was carried out using SPAdes v.0.4.7 (11) with the *Rhizobium ipomoeae* reference genome ASM491216v1 (RefSeq assembly accession number [GCF_004912165.1](https://.ncbi.nlm.nih.gov/assembly/GCF_004912165.1)); this produced an assembly containing 22 contigs, with an *N*₅₀ value of 469,644 bp. Genome annotation occurred through the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) v.5.13 (12) as part of the GenBank submission process. All programs were run with default parameters unless otherwise stated. Genome features are summarized in Table 1.

The results of genome sequencing confirmed that we had isolated a species of *Rhizobium*, which we designated C1 (for “*Codium* 1”). The closest identified organism,

Editor Leighton Pritchard, SIPBS, University of Strathclyde

Copyright © 2022 Miller et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Chelsea J. Vickers, chelsea.vickers@vuw.ac.nz.

The authors declare no conflict of interest.

Received 13 January 2022

Accepted 11 April 2022

Published 2 May 2022

TABLE 1 Assembly statistics and genome information for *Rhizobium* sp. nov. C1

Parameter	Finding for <i>Rhizobium</i> sp. nov. C1
Sequencing and assembly	
No. of Illumina reads	711,961
Total no. of Illumina bases	106,794,150
Avg coverage (×)	25
Genome description	
Genome size (bp)	4,240,398
G+C content (%)	61.6
Total no. of genes	4,022
Total no. of coding sequences	3,971
No. of proteins	3,928
No. of pseudogenes	43
No. of rRNAs	3
No. of tRNAs	44
No. of noncoding RNAs	4

based on average nucleotide identity (87.72%) determined using GTDB-Tk v.1.7.0 (13), was the plant-growth-promoting bacterium *Rhizobium rhizophilum* 7209-2 (RefSeq assembly accession number [GCA_004912145.1](https://doi.org/10.1093/ncbi/taaa001)) isolated from rape (*Brassica napus* L.) (7). The close relationship between *Rhizobium* sp. nov. C1 and this plant-growth-promoting bacterium indicates that *Rhizobium* sp. nov. C1 could be a marine algal symbiont; however, further investigation is required to validate this hypothesis.

Data availability. All data can be found under the NCBI BioProject accession number [PRJNA773134](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA773134), including raw reads (SRA accession number [SRX13114500](https://www.ncbi.nlm.nih.gov/sra/SRX13114500)) and genome assembly information (GenBank accession number [NZ_JAJGPW01000000](https://www.ncbi.nlm.nih.gov/genbank/NZ_JAJGPW01000000)).

ACKNOWLEDGMENTS

This work was supported via strategic research funds from the School of Biological Sciences at Victoria University of Wellington and by a grant from the Marsden Fund (grant 18-VUW-050).

REFERENCES

- Lee J, Ohta Y, Hayashi K, Hayashi T. 2010. Immunostimulating effects of a sulfated galactan from *Codium fragile*. *Carbohydr Res* 345:1452–1454. <https://doi.org/10.1016/j.carres.2010.02.026>.
- Sabry DA, Cordeiro SL, Silva CHF, Farias EHC, Sasaki GL, Nader HB, Rocha HAO. 2019. Pharmacological prospection and structural characterization of two purified sulfated and pyruvylated homogalactans from green algae *Codium isthmocladum*. *Carbohydr Polymers* 222:115010. <https://doi.org/10.1016/j.carbpol.2019.115010>.
- Wang L, Oh JY, Je JG, Jayawardena TU, Kim Y, Ko JY, Fu X, Jeon Y. 2020. Protective effects of sulfated polysaccharides isolated from the enzymatic digest of *Codium fragile* against hydrogen peroxide-induced oxidative stress in *in vitro* and *in vivo* models. *Algal Res* 48:101891. <https://doi.org/10.1016/j.algal.2020.101891>.
- Li N, Mao W, Yan M, Liu X, Xia Z, Wang S, Xiao B, Chen C, Zhang L, Cao S. 2015. Structural characterization and anticoagulant activity of a sulfated polysaccharide from the green alga *Codium divaricatum*. *Carbohydr Polym* 121:175–182. <https://doi.org/10.1016/j.carbpol.2014.12.036>.
- Tabarsa M, Karnjanapratum S, Cho M, Kim J, You S. 2013. Molecular characteristics and biological activities of anionic macromolecules from *Codium fragile*. *Int J Biol Macromol* 59:1–12. <https://doi.org/10.1016/j.ijbiomac.2013.04.022>.
- Finster K, Tanimoto Y, Bak F. 1992. Fermentation of methanethiol and dimethylsulfide by a newly isolated methanogenic bacterium. *Arch Microbiol* 157:425–430. <https://doi.org/10.1007/BF00249099>.
- Gao J, Wang L, Xue J, Tong S, Peng G, Sun Y, Zhang X, Sun J. 2020. *Rhizobium rhizophilum* sp. nov., an indole acetic acid-producing bacterium isolated from rape (*Brassica napus* L.) rhizosphere soil. *Int J Syst Evol Microbiol* 70:5019–5025. <https://doi.org/10.1099/ijsem.0.004374>.
- Fisher RF, Long SR. 1992. Rhizobium-plant signal exchange. *Nature* 357:655–660. <https://doi.org/10.1038/357655a0>.
- Hennig BP, Velten L, Racke I, Tu CS, Thoms M, Rybin V, Besir H, Remans K, Steinmetz LM. 2018. Large-scale low-cost NGS library preparation using a robust Tn5 purification and tagmentation protocol. *G3 (Bethesda)* 8:79–89. <https://doi.org/10.1534/g3.117.300257>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Tatusova T, DiCuccio M, Badretdin A, Chetverin V, Nawrocki EP, Zaslavsky L, Lomsadze A, Pruitt KD, Borodovsky M, Ostell J. 2016. NCBI Prokaryotic Genome Annotation Pipeline. *Nucleic Acids Res* 44:6614–6624. <https://doi.org/10.1093/nar/gkw569>.
- Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. 2019. GTDB-Tk: a tool-kit to classify genomes with the Genome Taxonomy Database. *Bioinformatics* 36:1925–1927. <https://doi.org/10.1093/bioinformatics/btz848>.