

ARTICLE

Received 4 Nov 2015 | Accepted 25 May 2016 | Published 12 Jul 2016

DOI: 10.1038/ncomms12054

OPEN

Orthogonal photoswitching in a multifunctional molecular system

Michael M. Lerch¹, Mickel J. Hansen¹, Willem A. Velema¹, Wiktor Szymanski^{1,2} & Ben L. Feringa¹

The wavelength-selective, reversible photocontrol over various molecular processes in parallel remains an unsolved challenge. Overlapping ultraviolet-visible spectra of frequently employed photoswitches have prevented the development of orthogonally responsive systems, analogous to those that rely on wavelength-selective cleavage of photo-removable protecting groups. Here we report the orthogonal and reversible control of two distinct types of photoswitches in one solution, that is, a donor-acceptor Stenhouse adduct (DASA) and an azobenzene. The control is achieved by using three different wavelengths of irradiation and a thermal relaxation process. The reported combination tolerates a broad variety of differently substituted photoswitches. The presented system is also extended to an intramolecular combination of photoresponsive units. A model application for an intramolecular combination of switches is presented, in which the DASA component acts as a phase-transfer tag, while the azobenzene moiety independently controls the binding to α -cyclodextrin.

¹Centre for Systems Chemistry, Stratingh Institute for Chemistry, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands. ²Department of Radiology, University of Groningen, University Medical Center Groningen, Hanzeplein 1, 9713 GZ Groningen, The Netherlands. Correspondence and requests for materials should be addressed to W.S. (email: w.szymanski@umcg.nl) or to B.L.F. (email: b.l.feringa@rug.nl).

Light as an external stimulus has been extensively used in chemistry, biology and material sciences to control processes and properties in a non-invasive manner with high spatiotemporal precision^{1–7}. However, as systems grow more complex, this control often needs to be extended to processes that act simultaneously, using multiple sources of light at different wavelengths⁸ in an orthogonal manner. When Bochet and co-workers^{9,10} reported the proof-of-principle for the irreversible wavelength-selective removal of photolabile protecting groups (PPGs; Fig. 1a) in 2000, it paved the way for numerous impressive applications⁸. These include functionalized surfaces¹¹, hydrogels¹² and lithography¹³, control of nucleic acids¹⁴, gene expression^{15,16}, regulation of enzyme activity^{17,18} and interfering with neuronal processes¹⁹. Despite the successful use of this concept, light-mediated uncaging is an inherently irreversible process⁶, which limits future applications towards responsive systems both in material and life sciences.

On the other hand, the field of molecular photoswitches exploits the reversibility of the photoswitching process^{2,3,20–24}. However, the concept of addressing photoactive molecules in parallel with light of different wavelengths has not been translated from wavelength-selective uncaging of PPGs to independent photoswitching (Fig. 1b). Such orthogonal control has great potential for employment in various fields, as many of the above-mentioned applications used for wavelength-selective uncaging would benefit from the reversibility of activation. Furthermore, reversible orthogonal control would enable independent modulation of different material properties and investigation of complex networks of signalling pathways in biological systems.

The wavelength selectivity (orthogonality) of the photoswitching is essential and the lack thereof currently represents a major limiting factor for successful applications (N.B.: throughout this article we refer to orthogonality as a characteristic of a system, where each component of the system can be controlled independently and irrespective of the order of addressing)²⁵.

Although the idea of combining two different photoswitches in an intermolecular manner and addressing them independently may seem simple, its execution remains a major challenge²⁶. Wavelength-selective control was so far confined to irreversible processes likely because of the fact that many of the most abundantly used photoswitches, such as azobenzenes, diarylethenes and spiropyrans, have overlapping absorption bands in the ultraviolet-visible region^{2,8}. Thus, it is difficult to find spectral regions with large differences in photoswitching efficiency (defined, at a given wavelength, as a product of extinction coefficient, ϵ and quantum yield, ϕ). Moreover, the possibility of undesirable energy transfer between the chromophores²⁷ can lead to a major loss of selectivity. Herein, we present an orthogonal, intermolecular combination of two classes of photoswitches and their intramolecular combination as a first-generation orthogonally addressable, dual functional molecular system. The reported intermolecular combination of photoswitches is based on a donor–acceptor Stenhouse adduct (DASA, Fig. 1c) and an azobenzene (Fig. 1d). Both photoswitches can be independently controlled by irradiation with light of three different wavelengths and by taking advantage of a thermal isomerization process. The modular and experimentally robust combination allows for functionalization of each constituent. A model application for an




Figure 1 | Orthogonal reversible photoswitching. Comparison of concepts described in this manuscript: wavelength-selective uncaging of photolabile protecting groups (**a**) and orthogonal photoswitching (**b**). Molecular structures and switching of DASAs (DASA **1** and **2**) (**c**) and azobenzenes (**3–6**) (**d**) with a conceptual representation of the ultraviolet-visible spectra of compounds **1** and **3** (**e**).

intramolecular combination of switches is presented, in which the DASA component acts as a phase-transfer tag, while the azobenzene moiety independently controls the binding to α -cyclodextrin (α -CD).

Results

Selecting a compatible photoswitch pair. Defining a photochromically compatible pair of photoswitches would constitute the first example of wavelength-selective/orthogonal control of multiple photoswitches in one solution. For this reason, we aimed to find compatible classes of photoswitches (Fig. 1), especially those that would exhibit a high complementarity in the 300–800 nm spectral region where photoswitches normally operate (Fig. 1e). We chose azobenzene photoswitches as one class; they have been successfully applied in molecular photocontrol^{20–24,28–31}, are easily accessible and allow for custom modification, functionalization and spectral fine-tuning. Azobenzenes have their main absorption bands located between 300 and 500 nm (π - π^* and n - π^* transitions; Fig. 1d,e and Table 1)^{20,32}. We envisioned that the T-type² DASAs (DASA 1 and DASA 2, Fig. 1c) could be used as the complementary class^{33,34} since they show very little absorption between 300 and 500 nm (Fig. 1c,e and Table 1)^{35–37}. Owing to the fact that the reverse switching of DASAs is thermally driven^{33,34,36,37}, three different wavelengths would be sufficient for wavelength-selective control. Importantly, for the design of orthogonal systems, one can exploit the tunability of thermal half-lives of the thermodynamically unstable states of the photoswitches in use. By changing the substitution pattern of azobenzenes, both very long and very short thermal half-lives have been attained. Increased thermal stability could be achieved, for example, through *ortho*-substitution^{32,38–41}. On the other hand, compounds with shortened thermal half-lives can be obtained by tuning of the electron density, through incorporation of both electron-donating groups (for example, -NMe₂ and -OMe), electron-withdrawing groups and combinations of both (push-pull systems)⁴². For the DASAs a clear relation between half-life and structure has not been determined yet^{33,34}. However, it has been shown that changing the acceptor can influence the half-life to some extent^{33,34}. Methylbarbituric acid-based cyclized DASAs (Fig. 1c, X=NMe; Y=C=O) are generally less thermally stable in comparison with the ones based on Meldrum's acid (Fig. 1c, X=O; Y=C(Me)₂). Finally, tuning half-lives of switches is often relatively independent of structural features required for function. This leaves ample space to tune the thermal relaxation rate of switches to magnitudes needed for different applications^{20–24,28–31,43}.

Intermolecular combination of photoswitches. Initially, we studied push-pull azobenzene **3** in combination with DASA **1** (Fig. 1 and Table 1). The 4'-methoxy group in azobenzenes increases photostationary states⁴⁴ (PSS, defined as the percentage

of the thermodynamically unstable isomer that can be produced upon irradiation at a given wavelength), while the ester moiety represents a handle for convenient functionalization. The absorption spectrum of the mixture of **1** and **3** is, to a large degree, a linear combination of the spectra of the separate photoswitches, indicating a low level of intermolecular energy transfer (Supplementary Fig. 39)²⁶.

The combination of photoswitches **1** and **3** showed a remarkably high level of wavelength selectivity, apparent in ultraviolet-visible spectra and reversible photochromism plots (Fig. 2). Irradiation of azobenzene **3** led to a selective, nearly quantitative *trans*-*cis* isomerization ($\lambda = 370$ nm, irradiation 1, Fig. 2), which proceeds with a quantum yield of $\phi = 0.15$. The relatively long half-life of the *cis*-isomer of azobenzene **3** ($t_{1/2} > 8$ h; Table 2) makes it thermally stable on the timescale of the experiment (< 1 h). Irradiation of the *cis*-**3** isomer ($\lambda = 430$ nm, irradiation 2, Fig. 2) allowed selective *cis*-*trans* isomerization without affecting photoswitch **1**. DASA **1** could be selectively cyclized by longer wavelength irradiation ($\lambda = 590$ nm, irradiation 3, Fig. 2), without affecting the azobenzene, after which it showed a short thermal half-life for cycloreversion (37 s at room temperature; Table 2). DASA **1** can also be switched selectively (irradiation 5, Fig. 2) without affecting the azobenzene **3** after it has been switched to the *cis*-isomer (irradiation 4, Fig. 2). By using a broad-spectrum visible light source (white light, Supplementary Table 1), both switches could be addressed at the same time (Supplementary Fig. 41) and different combinations of PSS could be attained by adjusting the irradiation intensity and/or duration (Supplementary Fig. 41). Moreover, this intermolecular combination operates successfully throughout a large concentration range (Supplementary Figs 51–53).

In summary, in a mixture of photoswitches **1** and **3** (Fig. 2), azobenzene **3** could be selectively switched from *trans* to *cis* (irradiation 1) and back from *cis* to *trans* (irradiation 2). The same applies for DASA photoswitch **1**, which could be selectively cyclized (irradiation 3) and then relaxed back to its thermally stable state. It is important to note that full chromatic orthogonality was observed because both states of **1**, as well as the thermal relaxation process, were not affected by light pulses that address **3** (irradiations 1, 2 and 4). Similarly, both states of **3** were not affected by light pulses addressing **1** (irradiations 3 and 5).

Structural scope of photoswitches. Subsequently, we investigated whether other azobenzenes could be used in a two-switch system with compound **1**. Towards that end, azobenzenes **4–6** (Fig. 3 and Table 2) were selected. These photochromes are not only structurally diverse, but they also exhibit very different functional characteristics: the λ_{max} of **4** is bathochromically shifted^{20,38–42}, whereas λ_{max} of **5** and **6** are hypsochromically shifted (Fig. 3 and Table 2) as compared with compound **3**. Moreover, they differ markedly in the half-lives of their thermally unstable *cis*-isomers as compared with **3**, with a shortened half-life for **4**^{20,32,38–42}, comparable half-life for **6** and a longer half-life for **5** (ref. 20; Table 2; Supplementary Figs 16, 20 and 24). All combinations of DASA **1** with azobenzenes **3**, **4**, **5** or **6** showed chromatic orthogonality, thus showing the tolerance of this photoswitch combination towards structural modification of the azobenzene core with functionally different behaviour (Supplementary Figs 43–48).

We then investigated possibilities for structural diversity of DASAs that can be incorporated in an orthogonal switching system with azobenzenes. In principle, changing the acceptor part of this photochromic compound affects the absorption maximum and thermal half-life, while the donor moiety has little influence on λ_{max} affecting only its solubility^{33,34}. DASA **2** (Fig. 3), with a

Table 1 | Properties of photoswitches 1 and 3.

Entry	λ (nm)	Open-DASA 1 , ϵ ($10^2 \text{ M}^{-1} \text{ cm}^{-1}$)	Trans-azobenzene 3 , ϵ ($10^2 \text{ M}^{-1} \text{ cm}^{-1}$)
1	360	<3	356.5 \pm 4.1
2	365	<3	349.4 \pm 4.2
3	370	<3	324.8 \pm 4.1
4	430	<3	17.0 \pm 0.8
5	445	5.7 \pm 1.5	16.7 \pm 0.7
6	515	279.7 \pm 2.9	4.3 \pm 0.7
7	570	1758.0 \pm 15.3	<3

DASA, donor-acceptor Stenhouse adduct.
Overview of molar absorptivities of photoswitches **1** and **3** at relevant wavelengths.




Figure 2 | Orthogonal control of a mixture of photoswitches. Proof-of-concept for the reversible and orthogonal photocontrol in a mixture of DASA **1** (—) and azobenzene **3** (- - -). The orthogonal photoswitching of both compounds in one solution (**1**: $\sim 4 \mu\text{M}$; **3**: $\sim 20 \mu\text{M}$; toluene, room temperature) was monitored at characteristic wavelengths for each photoswitch ($\lambda = 360 \text{ nm}$ for **3** and $\lambda = 570 \text{ nm}$ for **1**). Steps 1–5 indicate distinct irradiation experiments. Coloured areas indicate different wavelengths of irradiation. The coloured bar below the figure indicates the different states of the photoswitches. The structure of photoswitches and their corresponding photostationary states (determined by $^1\text{H-NMR}$ and ultraviolet-visible spectroscopy) are given for the different states during photoswitching.

Meldrum's acid-derived acceptor moiety, shows a hypsochromically shifted absorption maximum (Fig. 3 and Table 2) and its cyclized form exhibits a longer half-life ($\sim 130 \text{ s}$; Supplementary Fig. 8), when compared with that of DASA **1** (refs 33,34). We selected azobenzene **6** (Fig. 3) to study the orthogonality of photoswitching in solution since it is compatible with DASA **2** because of its hypsochromically shifted $\pi\text{-}\pi^*$ absorption band (Fig. 3, Table 2 and Supplementary Fig. 22). Gratifyingly, both switches **2** and **6** can also be selectively addressed in parallel with light of different wavelengths (Supplementary Figs 49 and 50). The flexibility with regard to the DASA photoswitch highlights the robustness and modularity of the photoswitch combination and bodes well for versatile use in different applications.

Orthogonal photoswitching in an intramolecular system.

Apparently, loss of selectivity through energy transfer did not pose a problem for the intermolecular system described thus far, even at high concentrations up to $\sim 80 \mu\text{M}$ (**1**) and $\sim 400 \mu\text{M}$ (**3**; Supplementary Figs 51–53). Therefore, the reversible photocontrol in the intermolecular case is purely dependent on the spectral properties of the two photoswitches involved. For an intramolecular combination, on the other hand, linker design, geometry and distance of the photochromes become more important²⁶. While intramolecular energy transfer between chromophores can drastically reduce the switching selectivity^{8,26,27}, it can be partially controlled by adjusting the lifetime of the photo-excited state and/or by changing the design of the linker^{8,26,27}. Furthermore, properties that arise from energy

transfer and the interplay of the combination of two photoswitches can sometimes be desired. This is the case, for example, in molecular logics^{26,45–48}, where such hybrid multiswitches (or multiphotochromes) allow (sequential) access to multiple different states^{45,46}, usually in combination with external stimuli other than light^{49–56}.

To probe whether the level of selectivity of the intermolecular combination could be retained in the intramolecular system, or whether new properties could be observed, we combined an azobenzene and a DASA photoswitch into a single molecule. Towards that end, compounds **7** and **8**, which differ only in the length of the linker, were synthesized (Fig. 4) and the selectivity of their photoresponsiveness was evaluated (Fig. 5).

Importantly, the selective switching of the azobenzene moiety is retained. However, irradiation with $\lambda = 590 \text{ nm}$ not only switched the DASA moiety but also affected, to some extent, the azobenzene part in both **7** and **8** (Fig. 5). Generally, improved selectivities were observed with the elongated linker unit (from $-(\text{CH}_2)_4-$ to $-(\text{CH}_2)_{12}-$). In compound **7** (shorter linker), irradiation at $\lambda = 370 \text{ nm}$ (Fig. 5a) does partially affect the DASA moiety, which is not observed for compound **8** (longer linker). Interestingly, compound **8** shows a slight decrease in addressability of exclusively the DASA photoswitch with $\lambda = 590 \text{ nm}$, but not with white light (Fig. 5b). No such effect is observed for the compound with shorter linker (compound **7**). As a control experiment, the intermolecular switching of the corresponding azobenzenes in combination with the DASA photoswitch **1** was studied and orthogonality was observed (Supplementary Figs 54–58).

Table 2 | Overview of photoswitch characteristics.

Entry	Switch	$t_{1/2}$	λ_{\max} (nm)	PSS for irradiation at λ^{**}			
				365	430	590	White light
1	Azo 3	> 8 h	360 ($\pi-\pi^*$) 445 ($n-\pi^*$)	4:96	70:30	ND	77:23
2	Azo 4	180 s	434 ($\pi-\pi^*$) 374 ($n-\pi^*$)	ND	ND	ND	ND
3	Azo 5	> 8 h	320 ($\pi-\pi^*$) 436 ($n-\pi^*$)	32:68 [†]	82:18	ND	77:23
4	Azo 6	> 8 h	348 ($\pi-\pi^*$) 444 ($n-\pi^*$)	< 3:97	69:31	ND	76:24
5	DASA 1	37 s	570	ND	ND	< 20:80 [‡]	< 20:80 [‡]
6	DASA 2	130 s	545	ND	ND	< 20:80 ^{‡,§}	< 20:80 [‡]

DASA, donor-acceptor Stenhouse adduct; ND, not determined; PSS, photostationary state.

***Trans/cis*

[†]Same for 312 nm.

[‡]Open/cyclized.

[§]546 nm.




Figure 3 | Structural scope of photoswitches. Structural scope of DASAs (**1** and **2**) and azobenzenes (**3-6**). Overlay of the ultraviolet-visible absorption spectra of compounds **1-6** (compounds **1** and **2**: $\sim 4 \mu\text{M}$; compounds **3-6**: $\sim 20 \mu\text{M}$; toluene; room temperature) with their corresponding absorption maxima.

Modulation of phase transfer and supramolecular interactions.

We proceeded to apply this intramolecular system for controlling the location and the function of the molecule. Previously reported extraction experiments of DASA molecules^{33,34} established the potential of DASAs as a phase-transfer tag^{34,37}. Reversible photoswitching of DASAs is only observed in aromatic solvents. Polar protic solvents (as, for example, methanol or water) stabilize DASA's zwitterionic cyclized state, whereas halogenated solvents favour the elongated triene structure. Photoswitching results in formation of the hydrophilic zwitterionic cyclized form and phase transfer to the aqueous layer³⁴. We envisioned

exploiting this behaviour to transport photoswitchable cargo that is, an azobenzene. On an independent level of photocontrol, the isomerization of the azobenzene could be used to manipulate a certain function, for example, the well-known host-guest binding to CDs (especially α -CD)⁵⁷⁻⁶⁰. Therefore, we set off to evaluate a model system with dual functionality (Fig. 6) in which the DASA moiety of **8** controls the phase where the molecule resides (location), whereas the azobenzene moiety controls the supramolecular binding to α -CD (function). In toluene, the azobenzene moiety of compound **8** could be switched selectively between *trans*- and *cis*-forms (Fig. 5b). Upon cyclization of the




Figure 4 | Synthesis of compounds 7 and 8. Compounds 7 and 8 were used to probe the chromatic selectivity in intramolecular photoswitching.




Figure 5 | Selectivity in intramolecular photoswitching. Reversible photochromism of an intramolecular photoswitch combination of azobenzene and DASA with different linker lengths. Compound 7 (a) and compound 8 (b) ($\sim 4 \mu\text{M}$; toluene, room temperature). Steps 1–6 indicate distinct irradiation experiments.

DASA moiety, phase transfer of compound 8 to the aqueous layer is expected. In the aqueous layer, the photocontrol of the azobenzene moiety would allow control over binding to the water-soluble α -CD.

Towards that end, we established the phase-transfer behaviour of DASA in the intramolecular system 8. Inspired by traditional $\log P_{(\text{oct/wat})}^1$ measurements⁶¹, a functional phase-transfer system was designed to operate at pH 9 (using a saturated aqueous NaHCO_3 solution to adjust the pH Fig. 7). The extraction experiment described in Fig. 7 establishes the photoswitching and

light-controlled phase-transfer behaviour of compound 8. Compound 8 is dissolved in toluene and the organic layer is underlaid with water (pH 9). Upon irradiation of the biphasic mixture (stage I, Fig. 7) with $\lambda = 365 \text{ nm}$, the azobenzene part of 8 is selectively switched, as can be observed by ultraviolet-visible spectroscopy (stage II, Fig. 7). Subsequent irradiation with white light results in a successful phase transfer of 8 to the aqueous layer (stage III, Fig. 7). Separation of the biphasic mixture and addition of dichloromethane to the aqueous phase results in back-extraction and recolouration of the organic solution




Figure 6 | Design principle of switchable phase transfer and binding. Schematic outline of the application of light-controlled guest **8** with light-controlled phase-transfer properties. Compound **8** (*trans*-open) is soluble in toluene. Irradiation with $\lambda = 365$ nm (*trans*-*cis* isomerization) and $\lambda = 430$ nm (*cis*-*trans* isomerization) controls the state of the azobenzene. Cyclization of the DASA moiety with visible light results in phase transfer of compound **8**. In the aqueous layer ($\text{pH} \geq 9$), the azobenzene binds the water-soluble α -CD in its *trans*-state, but not in the *cis*-state.




Figure 7 | Demonstration of light-controlled phase transfer. Dynamic phase transfer of compound **8** in a toluene/aqueous NaHCO_3 ($\text{pH} = 9$) mixture. The independent switching and phase-transfer events can be monitored by phase-colour (**a**) and ultraviolet-visible spectroscopy (**b**). *Stage I*: initial conditions, compound **8** resides in the organic phase (purple colour). *Stage II*: the azobenzene part of compound **8** was switched with irradiation at $\lambda = 365$ nm. The DASA part remains untouched. *Stage III*: irradiation with white light results in the cyclization of the DASA moiety, decolouration and phase transfer of compound **8**. *Stage IV*: separating the layers and back-extraction of the aqueous layer with dichloromethane results in back-transfer of compound **8** to the organic phase, as can be observed by the recolouration of this phase (purple colour). The presence of compounds in the different phases is schematically depicted (**c**).

(stage IV, Fig. 7), taking advantage of the short half-life of the DASA 1-type moiety. Importantly, the geometry of the azobenzene moiety can be controlled without affecting the

DASA moiety in the organic phase (Fig. 7). Isolation of phase-transferred compound **8** and photoswitching in water ($\text{pH} \geq 9$, K_2CO_3) established reversibility with negligible fatigue




Figure 8 | Binding between azobenzene and α -CD. Binding studies on the host-guest binding of α -CD and compound **8**. Titration of α -CD to an aqueous solution (pH ≥ 9) of compound **8** ($11 \mu\text{M}$). A clear change of the absorption spectrum is observed, indicating the binding of the *trans*-azobenzene moiety of **8** (a). Binding is significantly reduced in a photostationary state reached under 365 nm irradiation (inducing *trans*-*cis* isomerization; b).

of the azobenzene moiety in the basic aqueous phase (Supplementary Figs 61, 62 and 69).

Having established the light-induced phase transfer of compound **8** through DASA cyclization (light-control of location), we studied the possibility of controlling host-guest binding of compound **8** in water (light control of function). CDs are well known as host molecules with a multitude of studied guest molecules^{57,58,60}. α -CD binds *trans*-azobenzenes but not *cis*-azobenzenes^{59,62}.

Compound **8** was isolated after phase transfer and was then subjected to binding studies with α -CD in water (pH ≥ 9 ; ref. 57). A clear change in the absorption spectrum upon titration of the host was observed (Supplementary Fig. 63) for the thermally adapted compound **8**, indicating the binding of *trans*-azobenzene moiety of **8** to α -CD. For irradiated samples, this effect was markedly reduced (Supplementary Figs 64–67), suggesting a much weaker binding of the *cis*-azobenzene moiety of **8** to α -CD. Photoswitching is retained even in the bound state, where interestingly different PSS values are observed (Supplementary Figs 68–70). Thus, by switching the azobenzene moiety, the affinity for the α -CD host can be tuned (Fig. 8).

Overall, the azobenzene moiety of compound **8** can be independently controlled by light in the organic phase with irradiation at $\lambda = 365 \text{ nm}$ (*trans*-*cis* isomerization; Figs 5b and 7) and $\lambda = 430 \text{ nm}$ (*cis*-*trans* isomerization; Fig. 5b). Upon switching of the DASA moiety with white light, compound **8** undergoes phase transfer to the basic aqueous layer. Herein, the *trans*-azobenzene moiety binds to α -CD more strongly than the *cis*-azobenzene one. Compound **8** can be recovered from the aqueous

layer by back-extraction to dichloromethane. In summary, compound **8** incorporates the two independently photocontrolled, functionally different capabilities for phase transfer, induced by DASA cyclization and host-guest binding, dependent on the geometry of the azobenzene moiety.

Discussion

In conclusion, an intermolecular two-switch system consisting of a DASA and an azobenzene was developed by a rational, spectra-guided design. The robustness of the photoswitch combination is illustrated by the compatibility of differently substituted azobenzenes, as well as DASAs. The flexibility in functional group substitution emphasizes the generality and modularity of this approach and may potentially also allow combination with other classes of photoswitches or PPGs. Furthermore, control of thermal relaxation allows specific tailoring of the switch to the needs of the system. Recent non-orthogonal applications of independent switches in the same solution highlight the great potential of our approach^{63–68}. The intramolecular photoswitching extends the applicability of this combination, which was exemplified by a dual functional molecule where both the binding to α -CD and phase transfer can be controlled with different wavelengths of light. The intramolecular combination shows interesting switching behaviour dependent on the linker length—an aspect that will be studied in-depth in the future. The presented study shows levels of orthogonal selectivity that are unparalleled both for wavelength-selective uncaging⁸ and multiphotochromes²⁶. It represents a major step towards the

development of future orthogonal and reversible photoswitchable tools that can be used for non-invasively interfering with and the manipulation of multifunctional molecular systems. Future research should focus on overcoming the strong solvent dependence of the photoswitching of the DASAs and orthogonal switching in the near infrared region of the spectrum^{38–41}.

Methods

Synthesis of photoswitches and combinations thereof. The synthesis of compounds 1–9 is detailed in Supplementary Methods. For NMR characterization of the described compounds, see Supplementary Figs 71–83.

Photochemical characterization. Irradiation of samples in toluene (Uvasol) was performed with different light sources and filters (Supplementary Tables 1 and 2). Photoswitching was observed with an Agilent 8453 UV-Visible Spectrophotometer in a 10 mm quartz cuvette using Uvasol grade solvents. Generally, the concentrations of the photoswitches were adjusted to result in a measured absorbance between 0.5 and 1.0 (a.u.). Half-lives of photoswitches were measured at λ_{\max} and fitted with single exponential process.

For the photochemical characterization of photoswitches 1–9, see Supplementary Figs 1–31. For the determination of photostationary states reached, see Supplementary Figs 32–38.

Determination of quantum yields. The quantum yield was determined by following initial photoswitching kinetics (percentage of switching ~10%) as decrease in absorbance at λ_{\max} and using the light intensity determined from standard ferrioxalate actinometry⁶⁹. A more detailed description can be found in Supplementary Methods.

Photoswitching experiments. Each mixture of compounds was analysed by measuring ultraviolet-visible spectra of the different accessible states and by kinetic measurement of the time-dependent reversible photochromism. Henceforth, each photoswitch was followed at its λ_{\max} . Importantly, samples were irradiated during measurements at an ~90° angle to the light path of the ultraviolet-visible spectrometer with distances from light source and their respective intensities at this distance detailed in Supplementary Tables 1 and 2. For the photochemical characterization of two-photoswitch mixtures, see Supplementary Figs 39–50. For photoswitching at higher concentrations, see Supplementary Figs 51–53. For the photochemical characterization of the intramolecular combinations of both switches, see Supplementary Figs 54–58.

Model application studies. For an analysis of the phase-transfer behaviour of compounds 3 and 9, see Supplementary Fig. 59. For the phase-transfer behaviour of compound 8, see Supplementary Fig. 60. Experimental details for the binding studies and switching behaviour of the azobenzene moiety of *cyclized-8* can be found in the Supplementary Methods section and Supplementary Figs 61–70 and Supplementary Tables 3–7.

Data availability. The authors declare that all data supporting the findings of this study are available within the article and its Supplementary Information Files.

References

- Griesbeck, A. G., Oelgemöller, M. & Ghetti, F. in *Handbook of Organic Photochemistry and Photobiology* (CRC Press, 2012).
- Feringa, B. L. & Browne, W. R. (eds). *Molecular Switches*, 2nd ed. (Wiley-VCH: Weinheim, 2011).
- Russek, M.-M. & Hecht, S. Photoswitches: from molecules to materials. *Adv. Mater.* **22**, 3348–3360 (2010).
- Klán, P. *et al.* Photoremovable protecting groups in chemistry and biology: reaction mechanisms and efficacy. *Chem. Rev.* **113**, 119–191 (2013).
- Lee, H.-M., Larson, D. R. & Lawrence, D. S. Illuminating the chemistry of life: design, synthesis, and applications of ‘caged’ and related photoresponsive compounds. *ACS Chem. Biol.* **4**, 409–427 (2009).
- Brieke, C., Rohrbach, F., Gottschalk, A., Mayer, G. & Heckel, A. Light-controlled tools. *Angew. Chem. Int. Ed.* **51**, 8446–8476 (2012).
- Szobota, S. & Isacoff, E. Y. Optical control of neuronal activity. *Annu. Rev. Biophys.* **39**, 329–348 (2010).
- Hansen, M. J., Velema, W. A., Lerch, M. M., Szymanski, W. & Feringa, B. L. Wavelength-selective cleavage of photoprotecting groups: Strategies and applications in dynamic systems. *Chem. Soc. Rev.* **44**, 3358–3377 (2015).
- Bochet, C. G. Wavelength-selective cleavage of photolabile protecting groups. *Tetrahedron Lett.* **41**, 6341–6346 (2000).
- Blanc, A. & Bochet, C. G. Wavelength-controlled orthogonal photolysis of protecting groups. *J. Org. Chem.* **67**, 5567–5577 (2002).
- Miguel, V. S., Bochet, C. G. & del Campo, A. Wavelength-selective caged surfaces: how many functional levels are possible? *J. Am. Chem. Soc.* **133**, 5380–5388 (2011).
- Azagarsamy, M. A. & Anseth, K. S. Wavelength-controlled photocleavage for the orthogonal and sequential release of multiple proteins. *Angew. Chem. Int. Ed.* **52**, 13803–13807 (2013).
- García-Fernández, L. *et al.* Dual photosensitive polymers with wavelength-selective photoresponse. *Adv. Mater.* **26**, 5012–5017 (2014).
- Rodrigues-Correia, A., Weyel, X. M. M. & Heckel, A. Four levels of wavelength-selective uncaging for oligonucleotides. *Org. Lett.* **15**, 5500–5503 (2013).
- Fournier, L. *et al.* A blue-absorbing photolabile protecting group for *in vivo* chromatically orthogonal photoactivation. *ACS Chem. Biol.* **8**, 1528–1536 (2013).
- Yamazoe, S., Liu, Q., McQuade, L. E., Deiters, A. & Chen, J. K. Sequential gene silencing using wavelength-selective caged morpholino oligonucleotides. *Angew. Chem. Int. Ed.* **53**, 10114–10118 (2014).
- Priestman, M. A., Sun, L. & Lawrence, D. S. Dual wavelength photoactivation of cAMP- and cGMP-dependent protein kinase signaling pathways. *ACS Chem. Biol.* **6**, 377–384 (2011).
- Velema, W. A., van der Berg, J. P., Szymanski, W., Driessen, A. J. M. & Feringa, B. L. Orthogonal control of antibacterial activity with light. *ACS Chem. Biol.* **9**, 1969–1974 (2014).
- Amatrudo, J. M., Olson, J. P., Agarwal, H. K. & Ellis-Davies, G. C. R. Caged compounds for multichromic optical interrogation of neural systems. *Eur. J. Neurosci.* **41**, 5–16 (2014).
- Beharry, A. A. & Woolley, G. A. Azobenzene photoswitches for biomolecules. *Chem. Soc. Rev.* **40**, 4422–4437 (2011).
- Szymański, W., Beierle, J. M., Kistemaker, H. A. V., Velema, W. A. & Feringa, B. L. Reversible photocontrol of biological systems by the incorporation of molecular photoswitches. *Chem. Rev.* **113**, 6114–6178 (2013).
- Fehrentz, T., Schönberger, M. & Trauner, D. Optochemical genetics. *Angew. Chem. Int. Ed.* **50**, 12156–12182 (2011).
- Velema, W. A., Szymanski, W. & Feringa, B. L. Photopharmacology: Beyond proof of principle. *J. Am. Chem. Soc.* **136**, 2178–2191 (2014).
- Broichhagen, J., Frank, J. A. & Trauner, D. A roadmap to success in photopharmacology. *Acc. Chem. Res.* **48**, 1947–1960 (2015).
- Barany, G. & Merrifield, R. B. A new amino protecting group removable by reduction. Chemistry of the dithiasuccinoyl (Dts) function. *J. Am. Chem. Soc.* **99**, 7363–7365 (1977).
- Fihey, A., Perrier, A., Browne, W. R. & Jacquemin, D. Multiphotochromic molecular systems. *Chem. Soc. Rev.* **44**, 3719–3759 (2015).
- Raymo, F. M. & Tomasulo, M. Electron and energy transfer modulation with photochromic switches. *Chem. Soc. Rev.* **34**, 327–336 (2005).
- Götl, R., Senf, A. & Hecht, S. Remote-controlling chemical reactions by light: towards chemistry with high spatio-temporal resolution. *Chem. Soc. Rev.* **43**, 1982–1996 (2014).
- Jamsaard, S. *et al.* Conversion of light into macroscopic helical motion. *Nat. Chem.* **6**, 229–235 (2014).
- Ware, T. H., McConney, M. E., Wie, J. J., Tondiglia, V. P. & White, T. J. Actuating materials. Voxellated liquid crystal elastomers. *Science* **347**, 982–984 (2015).
- Baroncini, M. *et al.* Photoinduced reversible switching of porosity in molecular crystals based on star-shaped azobenzene tetramers. *Nat. Chem.* **7**, 634–640 (2015).
- Bandara, H. M. D. & Burdette, S. C. Photoisomerization in different classes of azobenzene. *Chem. Soc. Rev.* **41**, 1809–1825 (2012).
- Helmy, S. *et al.* Photoswitching using visible light: a new class of organic photochromic molecules. *J. Am. Chem. Soc.* **136**, 8169–8172 (2014).
- Helmy, S., Oh, S., Leibfarth, F. A., Hawker, C. J. & Read de Alaniz, J. Design and synthesis of donor-acceptor Stenhouse adducts: a visible light photoswitch derived from furfural. *J. Org. Chem.* **79**, 11316–11329 (2014).
- Singh, S. *et al.* Spatiotemporal photopatterning on polycarbonate surface through visible light responsive polymer bound DASA compounds. *ACS Macro Lett.* **4**, 1273–1277 (2015).
- Mason, B. P. *et al.* A temperature-mapping molecular sensor for polyurethane-based elastomers. *Appl. Phys. Lett.* **108**, 041906 (2016).
- Helmy, S. & Read de Alaniz, J. Chapter three - photochromic and thermochromic heterocycles. *Adv. Heterocycl. Chem.* **117**, 131–177 (2015).
- Beharry, A. A., Sadowski, O. & Woolley, G. A. Azobenzene photoswitching without ultraviolet light. *J. Am. Chem. Soc.* **133**, 19684–19687 (2011).
- Dong, M., Babalhavaej, A., Samanta, S., Beharry, A. A. & Woolley, G. A. Red-shifting azobenzene photoswitches for *in vivo* use. *Acc. Chem. Res.* **48**, 2662–2670 (2015).
- Bléger, D., Schwarz, J., Brouwer, A. M. & Hecht, S. o-Fluoroazobenzenes as readily synthesized photoswitches offering nearly quantitative two-way isomerization with visible light. *J. Am. Chem. Soc.* **134**, 20597–20600 (2012).
- Bléger, D. & Hecht, S. Visible-light-activated molecular switches. *Angew. Chem. Int. Ed.* **54**, 11338–11349 (2015).

42. Nishimura, N. *et al.* Thermal cis-to-trans isomerization of substituted azobenzenes II. Substituent and solvent effects. *Bull. Chem. Soc. Jpn.* **49**, 1381–1387 (1976).
43. Kienzler, M. A. *et al.* A red-shifted, fast-relaxing azobenzene photoswitch for visible light control of an ionotropic glutamate receptor. *J. Am. Chem. Soc.* **135**, 17683–17686 (2013).
44. Szymański, W., Wu, B., Poloni, C., Janssen, D. B. & Feringa, B. L. Azobenzene photoswitches for Staudinger-Bertozzi ligation. *Angew. Chem. Int. Ed.* **52**, 2068–2072 (2013).
45. Frigoli, M. & Mehl, G. H. Multiple addressing in a hybrid biphotochromic system. *Angew. Chem. Int. Ed.* **44**, 5048–5052 (2005).
46. Szalóki, G., Sevez, G., Berthet, J., Pozzo, J.-L. & Delbaere, S. A simple molecule-based octastate switch. *J. Am. Chem. Soc.* **136**, 13510–13513 (2014).
47. de Silva, A. P. in *Molecular Logic-Based Computation, Monographs in Supramolecular Chemistry* (RSC Publishing, 2012).
48. de Silva, A. P., James, M. R., McKinney, B. O. F., Pears, D. A. & Weir, S. M. Molecular computational elements encode large populations of small objects. *Nat. Mater.* **5**, 787–790 (2006).
49. Straight, S. D. *et al.* All-photon molecular XOR and NOR logic gates based on photochemical control of fluorescence in a fulgimide–porphyrin–dithienylethene triad. *Adv. Funct. Mater.* **17**, 777–785 (2007).
50. Andréasson, J., Straight, S. D., Moore, T. A., Moore, A. L. & Gust, D. Molecular all-photon encoder–decoder. *J. Am. Chem. Soc.* **130**, 11122–11128 (2008).
51. Andréasson, J. *et al.* All-photon multifunctional molecular logic device. *J. Am. Chem. Soc.* **133**, 11641–11648 (2011).
52. Bälter, M., Li, S., Nilsson, J. R., Andréasson, J. & Pischel, U. An all-photon molecule-based parity generator/checker for error detection in data transmission. *J. Am. Chem. Soc.* **135**, 10230–10233 (2013).
53. Myles, A. J., Wigglesworth, T. J. & Branda, N. R. A multi-addressable photochromic 1,2-dithienylcyclopentene-phenoxynaphthacenequinone hybrid. *Adv. Mater.* **15**, 745–748 (2003).
54. Kinashi, K. & Ueda, Y. Synthesis and photochromic behavior of bi-functional photochromic compound. *Mol. Cryst. Liq. Cryst.* **445**, 223/[513]–230/[520] (2006).
55. Kinashi, K., Ono, Y., Naitoh, Y., Otomo, A. & Ueda, Y. Time-resolved fluorescence study on the photomerocyanine form of spiropyran and its derivative with azobenzene. *J. Photochem. Photobiol. A Chem.* **217**, 35–39 (2011).
56. He, Y., Zhu, Y., Chen, Z., He, W. & Wang, X. Remote-control photocycloreversion of dithienylethene driven by strong push-pull azo chromophores. *Chem. Commun.* **49**, 5556–5558 (2013).
57. Rekharsky, M. V. & Inoue, Y. Complexation thermodynamics of cyclodextrins. *Chem. Rev.* **98**, 1875–1918 (1998).
58. Connors, K. A. Population characteristics of cyclodextrin complex stabilities in aqueous solution. *J. Pharm. Sci.* **84**, 843–848 (1995).
59. Nepogodiev, S. A. & Stoddart, J. F. Cyclodextrin-based catenanes and rotaxanes. *Chem. Rev.* **98**, 1959–1976 (1998).
60. Crini, G. Review: a history of cyclodextrins. *Chem. Rev.* **114**, 10940–10975 (2014).
61. Leo, A., Hansch, C. & Elkins, D. Partition coefficients and their uses. *Chem. Rev.* **71**, 525–616 (1971).
62. Tomatsu, I., Hashidzume, A. & Harada, A. Photoresponsive hydrogel system using molecular recognition of α -cyclodextrin. *Macromolecules* **38**, 5223–5227 (2005).
63. Scott, T. F., Kowalski, B. A., Sullivan, A. C., Bowman, C. N. & McLeod, R. R. Two-color single-photon photoinitiation and photoinhibition for subdiffraction photolithography. *Science* **324**, 913–917 (2009).
64. Chen, W.-C., Lee, Y.-W. & Chen, C.-T. Diastereoselective, synergistic dual-mode optical switch with integrated chirochromic helicene and photochromic bis-azobenzene moieties. *Org. Lett.* **12**, 1472–1475 (2010).
65. Dube, H. & Rebek, J. Selective guest exchange in encapsulation complexes using light of different wavelengths. *Angew. Chem. Int. Ed.* **51**, 3207–3210 (2012).
66. Nishioka, H., Liang, X., Kato, T. & Asanuma, H. A photon-fueled DNA nanodevice that contains two different photoswitches. *Angew. Chem. Int. Ed.* **51**, 1165–1168 (2012).
67. Hildebrandt, K. *et al.* λ -Orthogonal pericyclic macromolecular photoligation. *Angew. Chem. Int. Ed.* **54**, 2838–2843 (2015).
68. Manna, D., Udayabhaskararao, T., Zhao, H. & Klajn, R. Orthogonal light-induced self-assembly of nanoparticles using differently substituted azobenzenes. *Angew. Chem. Int. Ed.* **54**, 12394–12397 (2015).
69. Hatchard, C. G. & Parker, C. A. A new sensitive chemical actinometer. II. Potassium ferrioxalate as a standard chemical actinometer. *Proc. R. Soc. Lond. A Math. Phys. Eng. Sci.* **235**, 518–536 (1956).

Acknowledgements

We gratefully acknowledge generous support from NanoNed, The Netherlands Organization for Scientific Research (NWO-CW, Top grant to B.L.F. and NWO VIDI grant no. 723.014.001 for W.S.), the Royal Netherlands Academy of Arts and Sciences (KNAW), the Ministry of Education, Culture and Science (Gravitation programme 024.001.035) and the European Research Council (Advanced Investigator Grant no. 227897 to B.L.F.). Furthermore, we thank the Swiss Study Foundation for a fellowship to M.M.L. We also thank T. Tiemersma-Wegman for ESI-MS analyses and we especially thank Professor Dr W.R. Browne, Dr S.J. Wezenberg and Juan Chen for technical assistance and discussions, and Professor Dr Denis Jacquemin for fruitful discussions.

Author contributions

B.L.F. and W.S. guided the research. B.L.F., W.S., M.J.H., W.A.V. and M.M.L. have designed the experiments. M.M.L. has conducted the experiments. M.M.L. has drafted the article, and all authors contributed to the final manuscript.

Additional information

Supplementary Information accompanies this paper at <http://www.nature.com/naturecommunications>

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Lerch, M. M. *et al.* Orthogonal photoswitching in a multi-functional molecular system. *Nat. Commun.* 7:12054 doi: 10.1038/ncomms12054 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016