

# Complete Genome Sequence of *Vibrio parahaemolyticus* Environmental Strain UCM-V493

S. S. Kalburge,<sup>a</sup> S. W. Polson,<sup>a</sup> K. Boyd Crotty,<sup>a</sup> L. Katz,<sup>b</sup> M. Turnsek,<sup>b</sup> C. L. Tarr,<sup>b</sup> J. Martinez-Urtaza,<sup>c</sup> E. F. Boyd<sup>a</sup>

Department of Biological Sciences, University of Delaware, Newark, Delaware, USA<sup>a</sup>; Centers for Disease Control and Prevention, Atlanta, Georgia, USA<sup>b</sup>; Department of Biology and Biochemistry, University of Bath, Bath, United Kingdom<sup>c</sup>

***Vibrio parahaemolyticus* is the leading bacterial cause of seafood-related gastroenteritis in the world. Here, we report the complete genome sequence and annotation of an environmental strain of *V. parahaemolyticus*, UCM-V493, with the aim of understanding the differences between the clinical and environmental isolates of the bacteria. We also make some preliminary sequence comparisons with the clinical strain RIMD2210633.**

Received 7 February 2014 Accepted 21 February 2014 Published 13 March 2014

**Citation** Kalburge SS, Polson SW, Boyd Crotty K, Katz L, Turnsek M, Tarr CL, Martinez-Urtaza J, Boyd EF. 2014. Complete genome sequence of *Vibrio parahaemolyticus* environmental strain UCM-V493. *Genome Announc.* 2(2):e00159-14. doi:10.1128/genomeA.00159-14.

**Copyright** © 2014 Kalburge et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to E. F. Boyd, [fboyd@udel.edu](mailto:fboyd@udel.edu).

*Vibrio parahaemolyticus* is a moderately halophilic Gram-negative bacterium found in marine environments in association with plankton, fish, and shellfish (1–4). *V. parahaemolyticus* is the leading bacterial cause of seafood-related gastroenteritis, with the CDC estimating 45,000 cases of infection yearly in the United States alone (4–7). Pathogenic strains of *V. parahaemolyticus* are characterized by the presence of *tdh* and *trh* genes coding for the thermostable direct hemolysin (TDH) and TDH-related hemolysin (TRH), respectively (1, 8–11). The first complete genome sequence of a *V. parahaemolyticus* strain was announced for the pandemic clinical isolate RIMD2210633 (12). Genomic analysis of this strain revealed the presence of 7 genomic islands, VP<sub>AI</sub>-1 to VP<sub>AI</sub>-7, ranging from 10 kb to 81 kb in size (13). An extensively studied environmental isolate, *V. parahaemolyticus* BB22OP, was recently sequenced and found to lack 5 of the 7 genomic islands present in the clinical isolate (14). Here, we report the complete genome sequence of *V. parahaemolyticus* UCM-V493. This strain is an O2:K28 serovar isolated in 2002 from a sediment sample in Spain. It is a *tdh*-negative and *trh*-negative strain and lacks all 7 genomic islands present in the clinical isolate (13, 15, 16).

*V. parahaemolyticus* UCM-V493 was grown at 37°C overnight in Luria-Bertani (LB) broth (Fisher Scientific, Fair Lawn, NJ) (pH 7) supplemented with streptomycin (200 µg/ml), with the final NaCl (Fisher Scientific) concentration adjusted to 3%. Genomic DNA was isolated using the GNOME DNA isolation kit (MP Biomedicals, Solon, OH).

Single-molecule real-time (SMRT) sequencing was performed on the PacBio RS2 platform (Pacific Biosciences, Menlo Park, CA). *De novo* assembly was performed using the hierarchical genome assembly process (HGAP) on the PacBio SMRT portal (17), and a coverage of 75× was obtained. Additional sequencing was performed using the Illumina MiSeq platform and Nextera technology (Illumina, San Diego, CA). The Illumina reads were assembled using CG-Pipeline (18) and CLC Genomics Workbench software package version 6.0.4 (CLC bio, Aarhus, Denmark), and

the resulting contigs were used to manually fill gaps with the MEGA5 software (19). The UCM-V493 genome is composed of two circular chromosomes and a circular plasmid. Annotation was performed using MAKER2 (20) and the RAST server (21). Chromosome 1 is 3.446 Mb and contains 3,187 coding sequences (CDSs), chromosome 2 is 1.698 Mb and contains 1,557 CDSs, and the plasmid is 88.5 kb and contains 116 CDSs. The G+C content is 45.3% for chromosome 1, 45.6% for chromosome 2, and 40.8% for the plasmid.

UCM-V493 shares a high homology with the clinical strain RIMD2210633; >80% of the CDSs are similar in the two strains. The UCM-V493 genome also shows a high level of gene synteny compared to RIMD2210633. Two novel prophage elements were identified in the UCM-V493 genome using the PHAST search tool (22). The prophage element on chromosome 1 shows homology to filamentous phage VCY-φ found in environmental *Vibrio cholerae* strains, and the prophage element on chromosome 2 shows homology to filamentous phage VFJ of *V. cholerae*. A detailed comparative analysis of the UCM-V493 and the RIMD2210633 genomes will be published elsewhere.

**Nucleotide sequence accession numbers.** The complete, annotated genome sequence for *V. parahaemolyticus* strain UCM-V493 was deposited at NCBI under accession no. [CP007004](https://.ncbi.nlm.nih.gov/nucl/CP007004) (UCM-V493\_Chromosome\_1), [CP007005](https://.ncbi.nlm.nih.gov/nucl/CP007005) (UCM-V493\_Chromosome\_2), and [CP007006](https://.ncbi.nlm.nih.gov/nucl/CP007006) (UCM-V493\_pVPUCMV\_plasmid).

## ACKNOWLEDGMENTS

We thank Brewster Kingham and Olga Shevchenko at the Sequencing and Genotyping Center at the Delaware Biotechnology Institute (DBI) for assistance with the genome sequencing. Computational infrastructure and other support provided by the University of Delaware Center for Bioinformatics and Computational Biology Core Facility and DBI was made possible through Delaware INBRE (NIH NIGMS 8 P20 GM103446-12) and Delaware EPSCoR (NSF EPS-081425).

This research was supported by a National Science Foundation CAREER award DEB-0844409 to E.F.B.

## REFERENCES

1. Johnson CN, Bowers JC, Griffitt KJ, Molina V, Clostio RW, Pei S, Laws E, Paranjpye RN, Strom MS, Chen A, Hasan NA, Huq A, Noriega NF III, Grimes DJ, Colwell RR. 2012. Ecology of *Vibrio parahaemolyticus* and *Vibrio vulnificus* in the coastal and estuarine waters of Louisiana, Maryland, Mississippi, and Washington (United States). *Appl. Environ. Microbiol.* 78:7249–7257. <http://dx.doi.org/10.1128/AEM.01296-12>.
2. Joseph SW, Colwell RR, Kaper JB. 1982. *Vibrio parahaemolyticus* and related halophilic vibrios. *Crit. Rev. Microbiol.* 10:77–124. <http://dx.doi.org/10.3109/10408418209113506>.
3. Kaneko T, Colwell RR. 1973. Ecology of *Vibrio parahaemolyticus* in Chesapeake Bay. *J. Bacteriol.* 113:24–32.
4. Krantz GE, Colwell RR, Lovelace E. 1969. *Vibrio parahaemolyticus* from the blue crab *Callinectes sapidus* in Chesapeake Bay. *Science* 164:1286–1287. <http://dx.doi.org/10.1126/science.164.3885.1286>.
5. Daniels NA, MacKinnon L, Bishop R, Altekruze S, Ray B, Hammond RM, Thompson S, Wilson S, Bean NH, Griffin PM, Slutsker L. 2000. *Vibrio parahaemolyticus* infections in the United States, 1973–1998. *J. Infect. Dis.* 181:1661–1666. <http://dx.doi.org/10.1086/315459>.
6. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, Jones JL, Griffin PM. 2011. Foodborne illness acquired in the United States—major pathogens. *Emerg. Infect. Dis.* 17:7–15. <http://dx.doi.org/10.3201/eid1701.P11101>.
7. Su YC, Liu C. 2007. *Vibrio parahaemolyticus*: a concern of seafood safety. *Food Microbiol.* 24:549–558. <http://dx.doi.org/10.1016/j.fm.2007.01.005>.
8. DePaola A, Ulaszek J, Kaysner CA, Tenge BJ, Nordstrom JL, Wells J, Puhr N, Gendel SM. 2003. Molecular, serological, and virulence characteristics of *Vibrio parahaemolyticus* isolated from environmental, food, and clinical sources in North America and Asia. *Appl. Environ. Microbiol.* 69:3999–4005. <http://dx.doi.org/10.1128/AEM.69.7.3999-4005.2003>.
9. Johnson CN, Flowers AR, Young VC, Gonzalez-Escalona N, DePaola A, Noriega NF III, Grimes DJ. 2009. Genetic relatedness among *tdh+* and *trh+* *Vibrio parahaemolyticus* cultured from Gulf of Mexico oysters (*Crassostrea virginica*) and surrounding water and sediment. *Microb. Ecol.* 57:437–443. <http://dx.doi.org/10.1007/s00248-008-9418-3>.
10. Paranjpye R, Hamel OS, Stojanovski A, Liermann M. 2012. Genetic diversity of clinical and environmental *Vibrio parahaemolyticus* strains from the Pacific Northwest. *Appl. Environ. Microbiol.* 78:8631–8638. <http://dx.doi.org/10.1128/AEM.01531-12>.
11. Parvathi A, Kumar HS, Bhanumathi A, Ishibashi M, Nishibuchi M, Karunasagar I, Karunasagar I. 2006. Molecular characterization of thermostable direct haemolysin-related haemolysin (TRH)-positive *Vibrio parahaemolyticus* from oysters in Mangalore, India. *Environ. Microbiol.* 8:997–1004. <http://dx.doi.org/10.1111/j.1462-2920.2006.00990.x>.
12. Makino K, Oshima K, Kurokawa K, Yokoyama K, Uda T, Tagomori K, Iijima Y, Najima M, Nakano M, Yamashita A, Kubota Y, Kimura S, Yasunaga T, Honda T, Shinagawa H, Hattori M, Iida T. 2003. Genome sequence of *Vibrio parahaemolyticus*: a pathogenic mechanism distinct from that of *V. cholerae*. *Lancet* 361:743–749. [http://dx.doi.org/10.1016/S0140-6736\(03\)12659-1](http://dx.doi.org/10.1016/S0140-6736(03)12659-1).
13. Hurlley CC, Quirke A, Reen FJ, Boyd EF. 2006. Four genomic islands that mark post-1995 pandemic *Vibrio parahaemolyticus* isolates. *BMC Genomics* 7:104. <http://dx.doi.org/10.1186/1471-2164-7-104>.
14. Jensen RV, Depasquale SM, Harbolick EA, Hong T, Kernell AL, Kruchko DH, Modise T, Smith CE, McCarter LL, Stevens AM. 2013. Complete genome sequence of pre-pandemic *Vibrio parahaemolyticus* BB22OP. *Genome Announc.* 1(1):e00002-12.
15. Boyd EF, Cohen AL, Naughton LM, Ussery DW, Binnewies TT, Stine OC, Parent MA. 2008. Molecular analysis of the emergence of pandemic *Vibrio parahaemolyticus*. *BMC Microbiol.* 8:110. <http://dx.doi.org/10.1186/1471-2180-8-110>.
16. Martinez-Urtaza J, Lozano-Leon A, DePaola A, Ishibashi M, Shimada K, Nishibuchi M, Liebana E. 2004. Characterization of pathogenic *Vibrio parahaemolyticus* isolates from clinical sources in Spain and comparison with Asian and North American pandemic isolates. *J. Clin. Microbiol.* 42:4672–4678. <http://dx.doi.org/10.1128/JCM.42.10.4672-4678.2004>.
17. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat. Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
18. Kislyuk AO, Katz LS, Agrawal S, Hagen MS, Conley AB, Jayaraman P, Nelakuditi V, Humphrey JC, Sammons SA, Govil D, Mair RD, Tatti KM, Tondella ML, Harcourt BH, Mayer LW, Jordan IK. 2010. A computational genomics pipeline for prokaryotic sequencing projects. *Bioinformatics* 26:1819–1826. <http://dx.doi.org/10.1093/bioinformatics/btq284>.
19. Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. 2011. MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28:2731–2739. <http://dx.doi.org/10.1093/molbev/msr121>.
20. Holt C, Yandell M. 2011. MAKER2: an annotation pipeline and genome-database management tool for second-generation genome projects. *BMC Bioinformatics* 12:491. <http://dx.doi.org/10.1186/1471-2105-12-491>.
21. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
22. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res.* 39:W347–W352. <http://dx.doi.org/10.1093/nar/gkr485>.