

Contribution of the Proximal Nerve Stump in End-to-side Nerve Repair: In a Rat Model

Jun Mo Jung, MD, Moon Sang Chung, MD*, Min Bom Kim, MD*, Goo Hyun Baek, MD*

*Department of Orthopedic Surgery, Yeseon Hospital, Bucheon, *Department of Orthopedic Surgery, Seoul National University College of Medicine, Seoul, Korea*

Background: The aim of this study was to evaluate the contribution of the proximal nerve stump, in end-to-side nerve repair, to functional recovery, by modifying the classic end-to-side neurorrhaphy and suturing the proximal nerve stump to a donor nerve in a rat model of a severed median nerve.

Methods: Three experimental groups were studied: a modified end-to-side neurorrhaphy with suturing of the proximal nerve stump (double end-to-side neurorrhaphy, Group I), a classic end-to-side neurorrhaphy (Group II) and a control group without neurorrhaphy (Group III). Twenty weeks after surgery, grasping testing, muscle contractility testing, and histological studies were performed.

Results: The grasping strength, muscle contraction force and nerve fiber count were significantly higher in group I than in group II, and there was no evidence of nerve recovery in group III.

Conclusions: The contribution from the proximal nerve stump in double end-to-side nerve repair might improve axonal sprouting from the donor nerve and help achieve a better functional recovery in an end-to-side coaptation model.

Keywords: Median nerve, Proximal stump, Peripheral nerve injury, End-to-side neurorrhaphy, Nerve recovery

End-to-end neurorrhaphy has had relatively satisfying results with regard to nerve regeneration since the introduction of microsurgery. The microsurgical end-to-end neurorrhaphy is currently regarded as the gold standard for nerve reconstruction.^{1,2)} Even so, this procedure is not always applicable. In cases with old or extensive nerve injuries, nerve grafting or direct muscle neurotization has been used as an alternative procedure; however, these techniques do not produce as successful results as the end-to-end procedure. The end-to-side (ETS) neurorrhaphy^{3,4)} was developed for these situations; it is a procedure where the distal stump of a transected nerve is sutured to an adjacent healthy nerve. The recovered axonal

passage to the recipient nerve, through the coaptation site, has been demonstrated in animal experiments.⁵⁻⁸⁾

The hypothesis of our study was that the original neurons of severed nerves contribute to axonal regeneration after ETS neurorrhaphy. If this is the case, then a double ETS neurorrhaphy, which requires the additional coaptation of the proximal end of the severed nerve to the donor nerve, might be more effective in nerve repair than a single ETS neurorrhaphy. The grasping strength and the muscle contraction force of the finger flexor muscle of rats and the number of myelinated axons of the injured median nerve were measured to evaluate the contribution from the proximal nerve stump after ETS nerve repair on functional recovery.

METHODS

Materials

Healthy male mature Sprague-Dawley rats (average weight, 300 g) were used in the experiments. We used the median nerve as the recipient nerve and the ulnar nerve

Received June 18, 2008; Accepted September 11, 2008

Correspondence to: Goo Hyun Baek, MD

Department of Orthopedic Surgery, Seoul National University College of Medicine, 28 Yeongeon-dong, Jongno-gu, Seoul 110-774, Korea

Tel: +82-2-2072-3787, Fax: +82-2-764-2718

E-mail: ghbaek@snu.ac.kr

as the donor nerve. As the median nerve innervates solely the finger flexor muscle of the rat,⁹⁾ this design facilitates functional assessment of the finger flexor muscle. In addition, as the ulnar nerve was kept intact, some sensation of the upper limb remained unaffected, and self mutilation or self-biting, due to the loss of sensation among the experimental animals, could be prevented.

Methods

A total of 45 Sprague-Dawley rats were randomly divided into three groups, with 15 rats in each group. All surgery was performed under general anesthesia using sodium pentobarbital (50 mg/kg body weight) administered intraperitoneally. The median and ulnar nerves were exposed using a 2 cm incision in the brachial area. A 1.5 cm segment of the right median nerve was removed at a point 1 cm proximal to the elbow joint. The distal stump of the median nerve was coapted to the ulnar nerve through an epineurial window (approximately 2 mm²) in an end-to-side fashion with 11-0 nylon sutures using an operating microscope (Zeiss S 22, OPMI 6-SFR, Zeiss, Munich, Germany). In group I, the proximal stump of the median nerve was coapted in an end-to-side fashion to the ulnar nerve, 2 cm proximal to the above mentioned coaptation site. In group II, the proximal stump end was pulled more proximally, ligated and buried beneath the pectoralis major to prevent the proximal stump from involvement in the reinnervation process. In group III, the transected median nerve was not anastomosed. The wound was closed with 4-0 nylon skin sutures.

Functional recovery was measured using the grasping test and the finger flexor muscle contraction force test. Morphological changes were assessed by comparing the number of regenerated myelinated axons.

Grasping test

The grasping strength was measured¹⁰⁾ preoperatively, 10 and 20 weeks after the surgical procedure. Animals were gently lifted by the tail and allowed to grasp the grid of a cage with their forepaws. Finger flexion then was observed. For the assessment of the grasping strength, a 15 × 15 cm square wire grid, 1.5 mm in diameter, was connected to an electronic balance [pull and push gauge (model AK-1, Algol, Nagano, Japan)] using a clamp and rod designed for the grid. The rats were allowed to grip the grid, while being pulled vertically upward by the tail at a constant rate of 10 mm/s, until the grip loosened. At the precise moment of release, the maximal negative value registered on the gauge was recorded (Fig. 1). Each rat was tested three times and the average value was documented. When

the grid was raised by the wrist or the nail, it was ignored. At the 10th post-surgical week, each 2 cm segment of the median and ulnar nerve, in the contralateral forepaws, was resected to prevent gripping the grid⁹⁻¹¹⁾ and to measure the number of myelinated axons in the normal nerve. At 20 weeks postoperatively, after the grasping test, the nerve coapted site was exposed and the muscle contraction force test and a nerve biopsy were performed under anesthesia.

Contraction strength test for finger flexor muscle

For the assessment of the finger flexor muscle contraction strength, the volar aspect of the forearm was incised and then the tendon and muscle belly of the flexor were exposed. The finger flexor tendon was connected to a tension transducer by 3-0 non-absorbable sutures. The forearm and wrist were fixated with an 18 gauge needle. The electrical stimuli (60 Hz, 1 ms in duration and 25 V) from a Harvard stimulator (Harvard Apparatus Ltd, Edenbridge, UK) were delivered via bipolar electrodes to the mid-belly of finger flexor muscle. The isometric muscle contraction strength was recorded using a Harvard tension transducer and was displayed on an oscilloscope.^{6,12,13)}

Morphologic evaluation

After completing the measurement of force, a nerve biopsy was performed at A, A', B, C and D in group I (double ETS neurorrhaphy) while at A, B and C in group II (ETS neurorrhaphy)(Fig. 2). In group I, if the proximal




Fig. 1. A rat underwent a grasping strength test. The animal was gently lifted by the tail and allowed to grasp the grid connected to an electronic balance. While grasping, the rat continued to be lifted by the tail with increasing firmness until it loosened its grip. At that precise moment, the negative value demonstrated by the balance was recorded.

coaptation point was x, the distal neurorrhaphy site was y. A was between x and y in the ulnar nerve, B was distal to y in the median nerve, C was distal to y in the ulnar nerve, D was proximal to x in the median nerve, and A' was proximal to x in the ulnar nerve. In group II, when the coaptation site was z, A was proximal to z in the ulnar nerve, B was distal to z in the median nerve, and C was distal to z in the ulnar nerve. Since the proximal stump of the median nerve was not coapted to the proximal ulnar nerve in group II, the site corresponding to A' in group I was similar to A in group II. Therefore, we ignored the test at A' in group II.

The nerve tissues were prepared for optical microscopy.^{7,8,12,14} Axonal counts were performed by a single laboratory technician, blinded to the specimen identity.

Data analysis

All data were analyzed using the SPSS ver. 11.0 (SPSS Inc., Chicago, IL, USA). The Kruskal-Wallis test was used for the differences in grasping strength and flexor muscle contraction force and compared between the two groups. The differences in the number of myelinated axons were evaluated using the Wilcoxon signed-rank test. The power of the test was 81% with a 0.05 significance level.

RESULTS

Grasping Test

The normal grasping strength in the forepaw before surgery was 581 ± 103 g. In group I (double ETS neurorrhaphy), the average was 137 ± 30 g at 10 weeks postoperatively and 155 ± 40 g at 20 weeks postoperatively. In group II (single ETS neurorrhaphy), it was 45 ± 19 g

and 97 ± 31 g, respectively (Table 1). In group III, it was 0 g at both testing times. The differences between the groups were statistically significant both at postoperative week 10 ($p = 0.001$) and 20 ($p = 0.002$).

Finger Flexor Muscle Contraction Strength

At postoperative week 20, the muscle contraction force was 45.4 ± 5.6 g in group I and 30.2 ± 5.6 g in group II (Table 1). The differences between the groups were statistically significant ($p = 0.004$).

The Number of Myelinated Axons

The number of myelinated axons in the median and ulnar nerve of a normal rat was 2283 ± 223 and 1495 ± 268 respectively. In group I, the number of myelinated nerve fibers was 1760 ± 310 at A (Fig. 3A), 1205 ± 233 at B (Fig. 3B), 1713 ± 341 at C (Fig. 3C), 1643 ± 215 at D (Fig. 3D), and 1958 ± 209 at A' (Fig. 3E). In group II, it was 1895 ± 213 at A (Fig. 4A), 536 ± 184 at B (Fig. 4B), and 1800 ± 264 at C (Fig. 4C, Table 2).

No statistical significance was observed in the number of axons between the two groups at A ($p = 0.175$) and at C ($p = 0.441$), but the number of axons was significant in comparisons with B ($p = 0.002$). The difference in the number of axons at site A and C was not statistically significant for each group.

DISCUSSION

It is widely accepted that spontaneous regeneration of nerves occurs relatively often in small animals such as the rat. Therefore, care was taken to prevent the proximal stump of the transected median nerve from reattaching spontaneously to the ulnar nerve, or to the distal portion of the injured median nerve. Such reconnection was not observed in groups II and III.

As in humans, anatomical variations between the median and ulnar nerve in rats must be taken into consideration. While Martin-Gruber anastomosis, an anomalous




Fig. 2. Schematic drawing of the end-to-side neurorrhaphy in Group I and Group II. MN: Median nerve, UN: Ulnar nerve.

Table 1. Comparison of Grasping Strengths and Flexor Muscle Contraction Forces in Group I and II

	Grasping strength (g)		Flexor muscle contraction force (g)
	10 weeks	20 weeks	
Group I	137 ± 30	155 ± 40	45.4 ± 5.6
Group II	45 ± 19	97 ± 31	30.2 ± 5.6
<i>p</i> value	0.001	0.002	0.004




Fig. 3. Photomicrographs of the cross-section in Group I. (A) Site A. Average number of the myelinated nerve fibers was 1760 ± 310 . (B) Site B. Average number of the myelinated nerve fibers was 1205 ± 233 . Regenerated nerve fibers were more compact with increased axonal size than in Group II-site B. (C) Site C. Average number of the myelinated nerve fibers was 1713 ± 341 . (D) Site D. Average number of the myelinated nerve fibers was 1643 ± 215 . (E) Site A'. Average number of the myelinated nerve fibers was 1958 ± 209 (Toluidine blue, $\times 200$).




Fig. 4. Photomicrographs of the cross-section in Group II. (A) Site A. Average number of the myelinated nerve fibers was 1895 ± 213 . (B) Site B. Average number of the myelinated nerve fibers was 536 ± 184 . (C) Site C. Average number of the myelinated nerve fibers was 1800 ± 264 (Toluidine blue, $\times 200$).

median-to-ulnar nerve communication in the forearm, is known to be present in 17% of the population,¹⁵⁾ its prevalence in rats has not been studied. Bertelli et al.⁹⁾ reported that transection of the ulnar nerve would be a safe procedure in studies involving the median nerve in a rat model. In this study, there was no choice but to preserve the ulnar nerve as it was used as the donor nerve for the ETS neuroorrhaphy. Since the grasping strength was 0 g in group III, the above mentioned anomaly was thought to not interfere with our results. The measurement of grasping strength had the technical problem of flexion

of the wrist being affected by the flexor carpi ulnaris innervated by the ulnar nerve; this could have interfered with accurate measurement of the grasping strength of the fingers. Therefore, the grid space was made narrow enough to prevent the wrist from being trapped in the wire.

The ETS neuroorrhaphy offers two theoretical benefits. First, it does not sacrifice the donor nerve that replaces a nerve defect. Second, suturing the distal stump of the injured nerve to a healthy donor nerve, in an end-to-side fashion, promotes recovery and prevents motor end-plate atrophy that could occur during the reinnervation period

Table 2. Comparison of Numbers of Myelinated Nerve Fibers in Group I and II

	Numbers of myelinated nerve fibers				
	Site A	Site B	Site C	Site D	Site A'
Group I	1760 ± 310	1205 ± 233	1713 ± 341	1643 ± 215	1958 ± 209
Group II	1895 ± 213	536 ± 184	1800 ± 264	NA	NA
<i>p</i> value	0.175	0.002	0.441	NA	NA

NA: Not available.

through a nerve graft for end-to-end neurorrhaphy in cases with a long segmental defect.

In recent studies regarding ETS neurorrhaphy, concerns about the analysis of factors and variables that enhance axonal sprouting after the procedure have been raised; these include the cross section size of the epineurial window, the number of Schwann cells at the coaptation site, the angle between the donor nerve and the recipient nerve, the helicoid technique, timing of coaptation, the growth factors at the coaptation site, and the type of reinnervated muscle.^{5,16,17} Indeed, promoting axonal sprouting directly correlates with reinnervation through the distal stump in the ETS neurorrhaphy. Therefore, increasing axonal sprouting is the most important factor for the clinical outcome of this technique.^{5-8,13,14,16}

The reinnervation mechanism after ETS neurorrhaphy has not been clearly identified. Experiments on the origin of new formed axons have yielded different results. For convenience, the current theories can be broadly categorized into three explanations.^{8,18-23}

First, the axons regenerated from the original neurons can reconnect to the original target organ by chance. This can be assumed especially when a nerve stump is anastomosed to the trunk of the spinal cord at a similar level. This theory is supported by experiments that showed that regeneration cannot occur if the recipient and the donor nerve originate from different spinal cord levels,²¹ and the number of myelinated axons increased in the donor nerve at the proximal site of the ETS neurorrhaphy.²² In the present study, the number of myelinated axons at the proximal site of the ETS coaptation of the ulnar nerve was 1760 in group I and 1895 in group II, while it was 1495 in the normal rats. These findings suggest that the proximal stump of the severed nerve contributed to the axonal regeneration.

Second, collateral axonal sprouting from a donor nerve can connect with a recipient nerve.^{8,19,23} According to some investigators, a donor nerve appears to cut off

its nerve branch connected to the original target organ in order to create a functional connection to a recipient nerve.¹² This idea is based on experiments showing the downgrading of the donor nerve distal to the coaptation site with no functional connection between the reinnervated muscle and the one innervated by the donor nerve. The results of the present study indicated that collateral axonal sprouting played an essential role in nerve repair. There was no significant difference in the number of myelinated axons in the ulnar nerve (donor nerve) at site A of each group, but the number of myelinated axons at site B in group I was almost two times the number in group II, 1205 and 536 respectively. However, downgrading of the donor nerve distal to the coaptation site was not observed.

Third, terminal sprouting from the injured donor nerve during the construction of an epineurial window or performing the ETS neurorrhaphy can reinnervate the target organ. However, this theory is not widely accepted because degenerative changes of the donor nerve distal to the coaptation site occur rarely. The difference in the number of myelinated axons between the proximal and distal sites in relation to the ETS neurorrhaphy donor nerve was not statistically significant in the present study.

Various models designed to prevent the proximal stump from interfering with the evaluation of the efficacy of the ETS neurorrhaphy have been suggested. However, no prior studies have investigated whether the proximal stump can facilitate the regeneration process. Prior studies have focused on the findings that support regeneration after an ETS neurorrhaphy. By contrast, we sought to use the proximal stump to determine whether it facilitated the nerve repair process after an ETS neurorrhaphy in cases with long segmental nerve defects where the proximal stump is available.

We performed experiments to assess the efficacy of a double ETS neurorrhaphy in a rat model. With regard to the grasping force, the flexor muscle contraction strength and the axonal regeneration, our findings showed that the method used was more effective for peripheral nerve reconstruction than the conventional ETS neurorrhaphy. These results imply that the contribution from the proximal nerve stump in the double end-to-side nerve repair facilitated axonal sprouting from the donor nerve and aided in improved functional recovery in an end-to-side coaptation model.

ACKNOWLEDGEMENTS

This study was supported by grant no. 04-2003-035-0 from the SNUH General Research Fund.

REFERENCES

1. Zhao Q, Dahlin LB, Kanje M, Lundborg G. Specificity of muscle reinnervation following repair of the transected sciatic nerve: a comparative study of different repair techniques in the rat. *J Hand Surg Br.* 1992;17(3):257-61.
2. Millesi H. Peripheral nerve surgery today: turning point or continuous development? *J Hand Surg Br.* 1990;15(3):281-7.
3. Viterbo F, Trindade JC, Hoshino K, Mazzoni A. Two end-to-side neurorrhaphies and nerve graft with removal of the epineurial sheath: experimental study in rats. *Br J Plast Surg.* 1994;47(2):75-80.
4. Zhao JZ, Chen ZW, Chen TY. Nerve regeneration after terminolateral neurorrhaphy: experimental study in rats. *J Reconstr Microsurg.* 1997;13(1):31-7.
5. Fortes WM, Noah EM, Liuzzi FJ, Terzis JK. End-to-side neurorrhaphy: evaluation of axonal response and upregulation of IGF-I and IGF-II in a non-injury model. *J Reconstr Microsurg.* 1999;15(6):449-57.
6. Liu K, Chen LE, Seaber AV, Goldner RV, Urbaniak JR. Motor functional and morphological findings following end-to-side neurorrhaphy in the rat model. *J Orthop Res.* 1999;17(2):293-300.
7. McCallister WV, Tang P, Smith J, Trumble TE. Axonal regeneration stimulated by the combination of nerve growth factor and ciliary neurotrophic factor in an end-to-side model. *J Hand Surg Am.* 2001;26(3):478-88.
8. Noah EM, Williams A, Jorgenson C, Skoulis TG, Terzis JK. End-to-side neurorrhaphy: a histologic and morphometric study of axonal sprouting into an end-to-side nerve graft. *J Reconstr Microsurg.* 1997;13(2):99-106.
9. Bertelli JA, Taleb M, Saadi A, Mira JC, Pecot-Dechavassine M. The rat brachial plexus and its terminal branches: an experimental model for the study of peripheral nerve regeneration. *Microsurgery.* 1995;16(2):77-85.
10. Bertelli JA, Mira JC. The grasping test: a simple behavioral method for objective quantitative assessment of peripheral nerve regeneration in the rat. *J Neurosci Methods.* 1995; 58(1-2):151-5.
11. Bertelli JA, dos Santos AR, Calixto JB. Is axonal sprouting able to traverse the conjunctival layers of the peripheral nerve? A behavioral, motor, and sensory study of end-to-side nerve anastomosis. *J Reconstr Microsurg.* 1996;12(8): 559-63.
12. Rovak JM, Cederna PS, Kuzon WM Jr. Terminolateral neurorrhaphy: a review of the literature. *J Reconstr Microsurg.* 2001;17(8):615-24.
13. Yan JG, Matloub HS, Sanger JR, Zhang LL, Riley DA, Jaradeh SS. A modified end-to-side method for peripheral nerve repair: large epineurial window helicoid technique versus small epineurial window standard end-to-side technique. *J Hand Surg Am.* 2002;27(3):484-92.
14. Karacaoglu E, Yuksel F, Peker F, Guler MM. Nerve regeneration through an epineurial sheath: its functional aspect compared with nerve and vein grafts. *Microsurgery.* 2001; 21(5):196-201.
15. Leibovic SJ, Hastings H 2nd. Martin-Gruber revisited. *J Hand Surg Am.* 1992;17(1):47-53.
16. Bryan DJ, Wang KK, Chakalis-Haley DP. Effect of Schwann cells in the enhancement of peripheral-nerve regeneration. *J Reconstr Microsurg.* 1996;12(7):439-6.
17. Bryan DJ, Wang KK, Summerhayes C. Migration of schwann cells in peripheral-nerve regeneration. *J Reconstr Microsurg.* 1999;15(8):591-6.
18. Alvarez J, Moreno RD, Llanos O, et al. Axonal sprouting induced in the sciatic nerve by the amyloid precursor protein (APP) and other antiproteases. *Neurosci Lett.* 1992; 144(1-2):130-4.
19. Lundborg G, Zhao Q, Kanje M, Danielsen N, Kerns JM. Can sensory and motor collateral sprouting be induced from intact peripheral nerve by end-to-side anastomosis? *J Hand Surg Br.* 1994;19(3):277-82.
20. McCallister WV, Tang P, Trumble TE. Is end-to-side neurorrhaphy effective? A study of axonal sprouting stimulated from intact nerves. *J Reconstr Microsurg.* 1999;15(8):597-603.
21. Zhang F, Cheng C, Chin BT, et al. Results of termino-lateral neurorrhaphy to original and adjacent nerves. *Microsurgery.* 1998;18(4):276-81.
22. Giovanoli P, Koller R, Meuli-Simmen C, et al. Functional and morphometric evaluation of end-to-side neurorrhaphy for muscle reinnervation. *Plast Reconstr Surg.* 2000;106(2): 383-92.
23. Chen YG, Brushart TM. The effect of denervated muscle and Schwann cells on axon collateral sprouting. *J Hand Surg Am.* 1998;23(6):1025-33.