



Published in final edited form as:

*Neuroimage*. 2017 June ; 153: 382–398. doi:10.1016/j.neuroimage.2016.11.069.

## Time to wake up: Studying neurovascular coupling and brain-wide circuit function in the un-anesthetized animal

Yu-Rong Gao<sup>a,b</sup>, Yuncong Ma<sup>d</sup>, Qingguang Zhang<sup>b</sup>, Aaron T. Winder<sup>b</sup>, Zhifeng Liang<sup>d</sup>, Lilith Antinori<sup>d</sup>, Patrick J. Drew<sup>a,b,c,d,\*</sup>, and Nanyin Zhang<sup>a,d,\*\*</sup>

<sup>a</sup>Neuroscience Graduate Program, Pennsylvania State University, University Park, PA 16802, United States

<sup>b</sup>Department of Engineering Science and Mechanics, Pennsylvania State University, University Park, PA 16802, United States

<sup>c</sup>Department of Neurosurgery, Pennsylvania State University, University Park, PA 16802, United States

<sup>d</sup>Department of Biomedical Engineering, Pennsylvania State University, University Park, PA 16802, United States

### Abstract

Functional magnetic resonance imaging (fMRI) has allowed the noninvasive study of task-based and resting-state brain dynamics in humans by inferring neural activity from blood-oxygenation-level dependent (BOLD) signal changes. An accurate interpretation of the hemodynamic changes that underlie fMRI signals depends on the understanding of the quantitative relationship between changes in neural activity and changes in cerebral blood flow, oxygenation and volume. While there has been extensive study of neurovascular coupling in anesthetized animal models, anesthesia causes large disruptions of brain metabolism, neural responsiveness and cardiovascular function. Here, we review work showing that neurovascular coupling and brain circuit function in the awake animal are profoundly different from those in the anesthetized state. We argue that the time is right to study neurovascular coupling and brain circuit function in the awake animal to bridge the physiological mechanisms that underlie animal and human neuroimaging signals, and to interpret them in light of underlying neural mechanisms. Lastly, we discuss recent experimental innovations that have enabled the study of neurovascular coupling and brain-wide circuit function in un-anesthetized and behaving animal models.

### Keywords

Neurovascular coupling; Awake animal; Imaging; Hemodynamics

---

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\*\*Correspondence to: Department of Biomedical Engineering, W-341 Millennium Science Complex, Pennsylvania State University, University Park, PA 16802, United States. \*Correspondence to: Department of Engineering Science & Mechanics, Department of Neurosurgery, W-317 Millennium Science Complex, Pennsylvania State University, University Park, PA 16802, United States. [pjd17@psu.edu](mailto:pjd17@psu.edu) (P.J. Drew), [nuz2@psu.edu](mailto:nuz2@psu.edu) (N. Zhang).

### Conflict of interest

None.

## Introduction

The discovery of anesthetics has been a boon to mankind, allowing invasive surgical procedures to be performed with little pain. In the history of neurophysiology, anesthetics were widely used to immobilize animals and to reduce variability in neural responses due to behavioral and attentional changes. Neurophysiological studies using the anesthetized preparation has tremendously advanced our understanding of brain function. Anesthetics are also an important tool for investigating the phenomenon of consciousness and theory of mind (Alkire et al., 2008; Brown et al., 2010, 2011), given their remarkable ability to manipulate consciousness level.

Despite the significant role that anesthesia plays in neuroscience research, it has become increasingly clear that anesthetics produced a neurological ‘state’ unlike any natural physiological condition, but rather, anesthesia is more akin to a ‘temporary, reversible coma’ (Brown, 2010). As a consequence of this better understanding of the nature of the anesthetized brain, and improvements in experimental methodology, there has been a push in the neuroscience community to utilize awake animal models in neurophysiological experiments (Ferenczi et al., 2016). Such models can be particularly valuable to the neurovascular coupling and animal neuroimaging communities, because the interpretation of both task-based and resting-state functional magnetic resonance imaging (fMRI) data collected in un-anesthetized humans depends on our understanding of neurovascular coupling under normal physiological conditions, and consequently, it is critical for translational animal experiments to be done without anesthesia as well. Furthermore, the vast majority of animal models of brain disorders rely on behavioral assessment conducted in the awake condition. Therefore, it is essential to image un-anesthetized animals if the imaging data will be used to interpret behavioral measurements (Ferenczi et al., 2016; Liang et al., 2014).

In this review, we summarize a large body of literature showing that anesthetics produce profound changes in cerebral hemodynamics, brain metabolism, neural activity, neurovascular coupling, and functional connectivity relative to the awake or sleeping states. Because of the large physiological differences between anesthetized and un-anesthetized animals, it is inappropriate to generalize physiological experimental data from anesthetized animals to physiological observations in awake humans. Given that technical advances have made it possible to study neurovascular coupling and brain circuit function in awake, behaving animals, the time is right for the community to move towards paradigms using awake animals.

### Physiological basis of the BOLD signal, and its sensitivity to anesthetics

The changes in oxygenation that underlie the blood-oxygenation-level dependent (BOLD) fMRI signal are due to an ‘oversupply’ of oxygenated blood to active areas of the brain (Buxton, 2012; Kim and Ogawa, 2012) (Fig. 1). Neural activity directly and indirectly leads to the release of vasoactive substances (Attwell et al., 2010; Cauli and Hamel, 2010), which relax contractile tissue around arteries (Kim and Ogawa, 2012) and potentially capillaries (Hall et al., 2014; Hartmann et al., 2015; Stefanovic et al., 2008, Fernández-Klett et al., 2010; Hill et al., 2015). Because the resistance of a blood vessel depends on its diameter,

these dilations cause a sharp reduction of vascular resistance, which increases blood velocity and flux. This increase in arterial diameter and blood flow is relatively rapid, occurring within less than a second following the stimulation (Chen et al., 2011; Drew et al., 2011; Gao et al., 2015; Kim et al., 2013). Arterial dilation and increased blood flow result in an elevated influx of oxygenated hemoglobin that exceeds the oxygen demand of the surrounding neural tissue (Fox and Raichle, 1986). Once the oxygenated blood transits through the capillary bed and into the draining veins, the increase in venous blood oxygenation can be detected using BOLD fMRI (Kim and Ogawa, 2012) (Fig. 1). The BOLD signal depends on an interaction between local neural metabolic activity (which may not perfectly match up with the spike rate of neurons), increases in blood flow, volume and oxygenation. Because of the transit time, the BOLD signal is delayed relative to the increase in blood volume (Hirano et al., 2011; Silva et al., 2007a). A slow dilation of the veins underlies the “balloon” or “windkessel” models of the BOLD responses (Buxton et al., 2004; 1998; Mandeville et al., 1999; Miller et al., 2001). If the stimulus is sustained for tens of seconds, a slow dilation of veins is observed (Drew et al., 2011; Huo et al., 2015a; Kim and Kim, 2011; Mandeville et al., 1999), though the venous dilation seems to fail under some anesthetics (Drew et al., 2011; Hillman et al., 2007; Lee et al., 2001). Studying the relationship between neural activity and vascular dynamics (neurovascular coupling) is crucial for interpreting brain functional imaging signals. Substantial amount of previous work using anesthetized animals has significantly contributed to our understanding of neurovascular coupling and mechanisms of neuroimaging signals. However, as all of the processes that underlie the BOLD response are affected by anesthesia (Fig. 1) (Aksenov et al., 2015; Martin et al., 2006b; PISAURO et al., 2013; Alkire et al., 2000; Lyons et al., 2016; Sellers et al., 2015; Ferezou et al., 2007; Nimmerjahn et al., 2009; Thrane et al., 2012; Chapin and Woodward, 1981; Crane et al., 1978; Cazakoff et al., 2014; Constantinople and Bruno, 2011; de Kock and Sakmann, 2009; Dudley et al., 1982; Ueki et al., 1992; Ferenczi et al., 2016), measurements made in the awake animal are expected to improve the quantification of neurovascular coupling and facilitate the translation over into the awake human studies.

### **Anesthetics attenuate the amplitude and increase the lag of the hemodynamic responses**

Importantly, though anesthesia reduces neural activity, it will disproportionately suppress the measured hemodynamic (BOLD, cerebral blood volume (CBV), and cerebral blood flow (CBF)) responses (Logothetis et al., 2001; Goense and Logothetis, 2008; PISAURO et al., 2013; Aksenov et al., 2015) relative to neural activity. The different components of the hemodynamic and neural responses that lead to the BOLD signal (CBV, cerebral metabolic rate of oxygen (CMRO<sub>2</sub>), etc.) can be differentially affected by anesthesia. While CBF, CBV, and CMRO<sub>2</sub> are all affected by anesthesia, this does not mean that the changes cancel out, and that the BOLD impulse response function will not be affected by anesthesia. In fact, as we will discuss below, the BOLD impulse response is greatly affected by anesthesia. For instance, in the somatosensory cortex, fentanyl and/or isoflurane decrease multiunit neural activity and the BOLD response, however, the decrease in the BOLD response is substantially larger than the decrease in multi-unit or LFP neural activity (Aksenov et al., 2015), implying a reduction in BOLD hemodynamic response function (HRF) amplitude. In addition, PISAURO et al. directly measured multi-unit activity (MUA) and CBV responses in

awake and anesthetized mice (Pisauro et al., 2013), and found that the spike rates to visual stimulation in awake animals were slightly increased relative to anesthetized animals (~20%), but the CBV response was increased to a much larger extent (~100%). This disproportional change in CBV and neural responses again means that the amplitude of the CBV HRF is reduced. A similar effect of anesthesia on the amplitude of the CBV and CBF HRFs is also seen when neural activity is measured using the local field potential (LFP) (Martin et al., 2006b).

Anesthesia also causes a *slowing* in the BOLD, CBV, and CBF HRFs. Pisauro et al., observed a 2 s delay in the CBV HRF in anesthetized mice relative to awake animals (Pisauro et al., 2013). This delay cannot be accounted for by a slight slowing in neural dynamics (~100 ms) caused by anesthesia. The anesthetic-induced lag in the neural response is too short (~100 ms) to account for the increase in the lag of the CBV change (~2 s) without a change in the *shape* of impulse response function that relates neural activity to CBV changes. As two impulse response functions with different shapes are by definition not the same, these neural recordings and CBV measurements clearly show a change in the HRF relating MUA to CBV by anesthesia. The same group of researcher also reanalyzed the BOLD data of Logothetis et al. (2001) and Goense and Logothetis, 2008 and found a similar slowing of the BOLD hemodynamic response in the anesthetized monkey relative to the awake monkey (Pisauro et al., 2013), which is a clear demonstration that anesthesia alters the shape of the BOLD HRF relative to the awake condition. Further, Martin and colleagues observed a similar slowing in the CBV, CBF and oxygenated hemoglobin responses in anesthetized rats relative to awake rats (Martin et al., 2006b). To our knowledge, every published comparison of neurovascular coupling between the awake and anesthetized condition has found substantial differences in every aspect of the hemodynamic response considered (CBV, BOLD, etc.). Taken together, these literature studies have found the BOLD, CBV and CBF HRFs are slowed in speed and decreased in amplitude by anesthesia (Fig. 2).

Critically, an accurate, quantitative understanding of the relationship between neural activity and hemodynamic signals is critical for interpreting neuroimaging data. Handwerker et al. (2004) showed that slight variations in the time course of the hemodynamic response (1 s) can lead to erroneous fMRI statistical results. This difference is smaller than the 2 s difference in delays found between awake and anesthetized animals (Pisauro et al., 2013). Therefore, to faithfully infer neuronal activity from hemodynamic signals, a quantitatively accurate HRF is essential, and anesthesia can represent a critical confounder in this aspect.

Current dominant working models for the vascular basis of neurovascular coupling (Buxton et al., 2004; 1998; Mandeville et al., 1999; Miller et al., 2001) lump the vasculature into 'arterial' and 'postarterial' (capillaries and veins) components, assuming that changes in arterial volume are small, and venous distention dominate the bulk of CBV changes. However, using optical approaches, experiments with anesthetized animals had failed to detect any venous dilations (Hillman et al., 2007; Lee et al., 2001) and typically found small (~10%) average changes in arterial diameter in response to prolonged sensory stimulation (Devor et al., 2007; Nizar et al., 2013). In another two-photon imaging study, urethane blocked the dilation of veins (Drew et al., 2011). The lack of venous dilation under some

anesthetic regimens is likely due to the reduced blood pressure caused by anesthesia, as decreasing blood pressure in the awake animal blocks the venous dilation (Huo et al., 2015b). Interestingly, there are examples of putative slow venous dilations being detected using MRI (Kim and Kim, 2010; Silva et al., 2007a), though in both cases low doses of anesthetics were used.

Early optical imaging experiments revealed that in the awake animal, hemodynamic signals are substantially larger and more widespread in response to whisker stimulation (Martin et al., 2006a) or limb stimulation (Lahti et al., 1999; Peeters et al., 2001) than in anesthetized animals. Similarly, in the visual cortex, hemodynamic signals during wakefulness are approximately twice as large as those in the anesthetized animal (Goense and Logothetis, 2008; Liu et al., 2013a; Shtoyerman et al., 2000). Two-photon microscopy in awake, head-fixed mice has shown that arterial dilations in response to sensory stimulation are much larger (two-fold or more) than those in the anesthetized animal (Fig. 3). Because the resistance of a vessel is inversely proportional to diameter to the fourth power, a small change in vessel diameter can lead to a large change in blood flow. Our data (Fig. 3) showed a 20% dilation in the arteries in awake animals, will cause a 100% increase in flow, while in anesthetized animals an 8% dilation was observed, which will cause a 36% increase in flow. The dilation of arteries is accompanied by a slow distention of veins, which can build to levels of 10% or more for prolonged stimuli (Drew et al., 2011; Huo et al., 2015a; Gao and Drew, 2016). Voluntary locomotion drives nearly identical dynamics and amplitudes of arterial and venous dilations as those evoked by passive stimulation (Huo et al., 2015a), suggesting that this pattern of arterial and venous dilation is an invariant feature of cerebral hemodynamics found across different behavioral and cardiovascular states when the animal is awake.

The anesthetic-induced reduction of hemodynamic response is at least in part caused by direct actions of anesthetics on ion channels of the vasculature. Like neurons, endothelial and smooth muscle cells have a rich complement of ion channels, and many anesthetics have direct actions on these ion channels. The perhaps most well-studied anesthetic is the widely used vasodilatory anesthetic isoflurane. Isoflurane induces the relaxation of blood vessels via its action on ATP-sensitive potassium channels (Kokita et al., 1999), as well as reducing the calcium signals (Flynn et al., 1991) by blocking calcium channels (Buljubasic et al., 1992; Akata et al., 2003) in vascular smooth muscle. Some anesthetics, like halothane, cause vasodilation via an interaction with the endothelial cells to trigger the release of relaxing factor (Muldoon et al., 1988), or via nitric oxide, like propofol (Park et al., 1995). These direct actions of anesthetics on ion channels present in the vasculature tend to have a shunting effect on the neurally induced currents, which can reduce the amplitude of the evoked dilation (Hill, 2012).

Lastly, the amplitude and dynamics of the hemodynamic response and its relationship to neural activity are greatly affected by the type and dosage of the anesthetic used (Masamoto and Kanno, 2012; Schlegel et al., 2015; Masamoto et al., 2009). For example, increasing isoflurane concentration (from 1% to 2%) does not measurably alter the neural responses to electrical stimulation, but it does change the stimulation-evoked blood flow response (Masamoto et al., 2009). Importantly, disproportionate influences of anesthesia on neuronal

and vascular activities can lead to an altered neurovascular coupling relationship. Using optical imaging spectroscopy in awake rats, the neural–hemodynamic coupling relationship was found to be linear (Martin et al., 2006b). This result is consistent with an approximately linear neurovascular coupling relationship found using fMRI in the microvasculature of awake humans (Zhang et al., 2008a; Zhang et al., 2009; Zhang et al., 2008b). However, this linear neurovascular coupling relationship was not present in anesthetized animals (Martin et al., 2006b). Consequently, the results obtained under one anesthetic agent or dose are often not comparable to those obtained under another anesthetic agent or dose (Masamoto and Kanno, 2012; Schlegel et al., 2015; Masamoto et al., 2009). The issue as to which is the ‘correct’ anesthetic to use can be completely bypassed by working in the awake animal.

### **Anesthetics cause profound decreases in baseline metabolism of the brain**

The brain has an intense metabolic demand (Harris and Attwell, 2012), which is thought to constrain the computations neurons can perform (Attwell and Gibb, 2005; Howarth et al., 2012; Niven and Laughlin, 2008). Much of this energy is likely spent on recycling synaptic vesicles (Harris and Attwell, 2012; Rangaraju et al., 2014), with spiking activity and restoring the membrane gradient contributing less to energy consumption. Because glucose is the primary energy source for the brain (Siesjö, 1978), the most basic comparison across brain states is how the rate of glucose utilization ( $CMR_{glc}$ ) changes.  $CMR_{glc}$  can be measured in animal models using the 2-deoxyglucose (2-DG) technique (Sokoloff et al., 1977), or in humans using positron emission tomography (PET) (Reivich et al., 1979) or  $^{13}C$  MR spectroscopy techniques (Rothman et al., 2011).

Anesthetics cause a substantial depression of the resting metabolic rate in the brain (Fig. 4), with the decreases in the metabolic rates in the cortex being larger than those in other brain structures. For example, alpha-chloralose, which is widely used in neurovascular coupling studies, causes a decrease of 50% or more in the metabolic rate of the cortex compared to the awake or lightly anesthetized state (Dudley et al., 1982; Hyder et al., 2002) with regional and layer variations in sensitivity (Dudley et al., 1982; Ueki et al., 1992). A dose dependent reduction of  $CMR_{glc}$  by pentobarbital is also observed, with a depression of 56% when cerebral pentobarbital levels exceed 50  $\mu g$  per g of tissue (Crane et al., 1978). In humans, halothane reduces global brain  $CMR_{glc}$  by ~40% (Alkire et al., 2000), and anesthesia with propofol (Alkire et al., 1995; Kaisti et al., 2003) or isoflurane (Alkire et al., 1997) cuts the cortical metabolic rate by 50% or more. To put this in context, one can compare anesthetic-induced decreases in brain metabolism to those observed in humans during sleep and those measured from patients in comas. Under normal physiological conditions, non-REM sleep decreases the brain metabolic rate by ~30% (Buchsbau et al., 1989; Kennedy et al., 1982), while REM sleep decreases metabolic activity by ~10% (Buchsbau et al., 1989), consistent with blood flow changes associated with sleep state (Braun et al., 1997). Patients in ‘locked-in’ and vegetative states showed decreases of 25% and > 50% of brain metabolism rate, respectively (Levy et al., 1987). The large changes in basal metabolism will likely affect blood flow, as astrocytes, which play a key role in brain metabolism, exert control over tonic blood flow in the brain (Rosenegger et al., 2015). Although these measurements were done in a wide range of species (humans, non-human primate, rodent, etc.), as the effects of all the anesthetics (whether used for human surgery or reported here) generalize across all

species and anesthetics, it seems reasonable to assume that the profound metabolic depression caused by anesthetics is universal across species.

Anesthesia-related brain metabolism changes can also affect oxygenation. Microscopic oxygen measurements using polarographic electrodes have shown great heterogeneity in tissue oxygen concentration in anesthetized animals (Ndubuizu and La Manna, 2007), making direct comparisons between the awake and anesthetized state difficult. Recent technical developments combining phosphorescent oxygen-sensitive dyes with two-photon microscopy have enabled the visualization of oxygen gradients at the micrometer scale (Sakadzic et al., 2010, 2014), showing that tissue oxygen tension varies widely depending on the distance to the nearest arteriole. Two-photon phosphorescence point measurements of oxygenation have found that the oxygen tension in the brains of awake mice are lower than in anesthetized mice, and have also failed to find laminar differences in oxygen levels which are seen in anesthetized animals (Lyons et al., 2016). Though further studies are warranted, these results suggest that the metabolic changes caused by anesthesia could alter normal brain oxygenation, which will have implications on BOLD signals measured in anesthetized animals.

### **Anesthetics directly alter the responsiveness and dynamics of neurons and glia**

There is a long history of using anesthetized animal models in sensory neurophysiology studies (Hubel and Wiesel, 1959), allowing the state of the animal to be controlled and the presentation of stimuli unaffected by eye movements, attention and other sources of variability present in awake animals. Gradually, it became apparent that neural dynamics during anesthesia were very different from those in the awake animal in several important ways. First, anesthetics, in general, tend to hyperpolarize neurons by increasing inhibition and/or decreasing excitation (Alkire et al., 2008; Ries and Puil, 1999), which usually leads to a suppression in the average firing rate of neurons (Cazakoff et al., 2014; Constantinople and Bruno, 2011; de Kock and Sakmann, 2009). Electroencephalography (EEG) recordings show that baseline neuronal activity is suppressed by various anesthetics (Antunes et al., 2003; Austin et al., 2005; Bonvento et al., 1994), while the evoked neural response is different in shape and is significantly delayed under anesthesia (Martin et al., 2006b; Sellers et al., 2015). Higher order aspects of neural activity are also profoundly affected, as correlations between neurons are increased during anesthesia (Greenberg et al., 2008).

The receptive field of neurons in sensory areas is also altered by anesthesia. A common but not universal finding is that the receptive field of a neuron is reduced by anesthesia (Armstrong-James and George, 1988; Chapin and Woodward, 1981; Friedberg et al., 1999). This distortion will have implications for neuroimaging-based mapping experiments since the activated area following sensory stimulation may be much larger in the awake animal (Ferezou et al., 2007). However, in the visual cortex, anesthesia seems to increase the spatial spread and prolong the duration of the response to visual input (Haider et al., 2013; Vaiceliunaite et al., 2013). In the olfactory bulb, odor evoked responses are larger in the anesthetized animal (Kato et al., 2013; Rinberg et al., 2006). In higher order sensory and sensorimotor areas, anesthesia can have large effects on sensory gating and responsiveness (Cardin and Schmidt, 2004, 2003; Schmidt and Konishi, 1998). These complicated, and

brain-region specific, changes make translation of imaging results obtained under anesthesia to the awake brain difficult.

Although anesthesia affects neural activities in many ways, these effects are not uniform across the brain, or even within the cortex. Frontal areas tend to be more affected by anesthesia than sensory areas (Sellers et al., 2013; Liang et al., 2015a), and there are laminar differences as well (Raz et al., 2014; Sellers et al., 2013). Anesthesia is also not a temporally uniform state. Many anesthetics, including urethane and isoflurane, are known to make the cortex alternate between a depolarized up-state and a hyperpolarized down-state (Castro-Alamancos, 2009; Steriade et al., 2001). In the 'up' state, neurons are more responsive to external input (Curto et al., 2009; Shu et al., 2003), similar to the sustained depolarization of wakefulness, while the synaptic activity is reduced during the 'down' state (Castro-Alamancos, 2009). Some effects of anesthesia may be due to changes in the modulatory state of the brain, as stimulation of modulatory nuclei can induce 'awake like' behavior. Stimulation of the nucleus basalis, which send cholinergic projections to cortex, can facilitate motor behavior (Berg et al., 2005) and alter sensory responses in anesthetized animals to a more 'awake' like state (Castro-Alamancos and Gulati, 2014; Goard and Dan, 2009). Anesthesia also increases the cortical adaptation to repetitive sensory stimulation relative to the awake state (Castro-Alamancos, 2004; Masamoto et al., 2009).

In addition to the depressant effects of anesthesia on neural activity, anesthetics decrease the frequency of calcium signals in astrocytes. Astrocytes are essential for the supply of energy metabolites to neurons, and provide a physical link to the vasculature (Simard et al., 2003; Takano et al., 2006). Astrocytes are also thought to have a role in mediating vasodilation in response to increased neural activity (Hirase, 2005; Takano et al., 2006), although this issue is contentious (see (Bonder and McCarthy, 2014; Lind et al., 2013; Nizar et al., 2013; Otsu et al., 2014)). Using two-photon microscopy, clear decreases in the spontaneous calcium signals in astrocytes have been observed under isoflurane, ketamine and urethane (Nimmerjahn et al., 2009; Thrane et al., 2012). In both studies, the anesthetic-induced astrocyte activity decrease was larger than the decrease induced by local pharmacological blocker of sodium channels or glutamate receptors, implying that the anesthetics have direct action on astrocytes. These effects of anesthesia on neural and glial activity are further complicated by the possibility that anesthesia could disrupt processes that relate neural activity changes to the hemodynamic response. For example, a study investigating the role of neuronally-generated nitric oxide (NO) in neurovascular coupling found different effects of NO synthase inhibitor on CBV in awake and anesthetized animals (Nakao et al., 2001), suggesting differential involvements of neurovascular coupling pathways depending on the state of the animal.

Lastly, due to the decreased responsiveness of neurons during anesthesia, many groups use electric shocks, rather than tactile or visual stimuli, to elicit hemodynamic responses. Electric shock is an unpleasant experience that bears little resemblance to normal sensory experience that indiscriminately activate proprioceptive, pain and other fibers, and can also induce large systemic changes in blood pressure that obscure neuronally-generated hemodynamic signals (Schroeter et al., 2014). As much larger (two-fold or greater) hemodynamic responses can be elicited in the awake animal (Drew et al., 2011; Pisauero et



al., 2013) using stimuli that are more of the type that might be presented to a human during an fMRI experiment, the advantages of the awake preparation are clear.

### **Anesthetics decrease brain temperature and affect other brain physiological processes**

While neurophysiologists focus on electrical measures of neural activity, there are other physiological variables that are greatly perturbed by anesthesia that could potentially influence the neurovascular coupling. Anesthetics decrease brain temperature by several degrees (Hayward, 1968; Shirey et al., 2015; Kalmbach and Waters, 2012). The decrease in brain temperature persists even when the core body temperature is kept in the physiological range (Shirey et al., 2015; Kalmbach and Waters, 2012; Podgorski and Ranganathan, 2016). The decrease in brain temperature is the largest at the surface of the brain (Kalmbach and Waters, 2012; Podgorski and Ranganathan, 2016), potentially altering the dynamics of the large pial vessels and supra-granular neurons more than the intraparenchymal vessels. Since neural activity warms the brain (Kiyatkin et al., 2002), the brain cooling associated with anesthesia is partially due to the anesthesia-evoked decreases in brain metabolism. Alternations of temperature, either up or down, affect the baseline diameter, myogenic tone, and spontaneous vasomotion of an arteriole (Osol and Halpern, 1988; Ogura et al., 1991), which will change evoked vascular dynamics under anesthesia (Pisauro et al., 2013). Brain temperature changes of one degree or more are large enough to produce changes in animal behavior (Long and Fee, 2008; Aronov and Fee, 2012).

Another physiological change induced by anesthesia is an increase in the volume fraction of extracellular space. The brain is composed of approximately 20% extracellular space (Nicholson, 2001; Korogod et al., 2015). The extracellular volume fraction has been shown to increase relative to the awake condition under ketamine (Xie et al., 2013). Because the extracellular space exchanges ions with active neurons, changes in the volume of this extracellular space should impact neural excitability (Somjen, 2002) and cerebrospinal fluid flow (Iloff et al., 2013). Besides brain temperature and extracellular space, intracranial pressure (ICP) is another important physiological variable that can be altered by various anesthetics (Artru, 1984; Gao and Drew, 2016). Elevating ICP alters heart beat and respiratory rhythm (Matsuura et al., 1984), decreases CBF in the cortex (Sadoshima et al., 1981), and changes neural activity (Seigo Nagao et al., 2009), all of which could impact neurovascular coupling. Lastly, isoflurane can open the blood-brain barrier, leading to ionic changes in the brain (Chever et al., 2008). While these physiological changes by anesthesia may not be as commonly appreciated as neural or cardiovascular changes, they are likely very disruptive.

### **Anesthesia alters neural circuit function**

Investigating neural circuit function plays a fundamental role in neuroscience. The brain is a highly inter-connected system that gives rise to complex, diverse and integrated information processing (Alkire et al., 2008; Tononi et al., 1998). The proper function of neural circuits and networks is essential to maintain meaningful information exchange and integration, and thus to support normal brain function. Elucidating the functional properties of neural networks, and their dysfunction in the context of pathology, is a critically important avenue of investigation in neuroscience.

Neural circuit function can be studied in both resting and activated states using fMRI techniques. In the resting state, fMRI (termed resting-state fMRI or rsfMRI) measures resting-state functional connectivity (RSFC) between brain regions by quantifying the temporal synchronization of spontaneously fluctuating rsfMRI signals (Biswal et al., 1995; Fox and Raichle, 2007). This technique offers a convenient tool to study neural circuit function without relying on any external stimulation, and thus is particularly suitable in some clinical settings where patients have difficulty to perform certain tasks inside the MRI scanner. Indeed, RSFC in awake humans has been extensively used to reveal neural circuit function in both normal (Albert et al., 2009; Horowitz et al., 2009) and pathological conditions (Anand et al., 2005; Carter et al., 2012; Greicius et al., 2007; Greicius et al., 2004; Hunter et al., 2012; Kennedy et al., 2006; Lowe et al., 2002; Lustig et al., 2003; Mayer et al., 2011; Tian et al., 2006; van Meer et al., 2012; Whitfield-Gabrieli et al., 2009).

Anesthesia has been shown to profoundly alter RSFC measurements of neural circuit function. A number of animal studies indicate that RSFC is tremendously weakened by anesthesia. Lu *et al.* demonstrated a dose-dependent decrease of cross-hemispheric RSFC in alpha-chloralose-anesthetized rats (Lu et al., 2007). This finding is in agreement with the study by Liu and colleagues who found that intrinsic BOLD fluctuations and functional connectivity in the resting rat are strongly dependent on anesthesia depth (Liu et al., 2011). In addition, several studies have shown that, relative to the awake state, thalamo-cortical connectivity is significantly reduced by anesthesia in the rat (Liang et al., 2012b; Liang et al., 2013). This result is consistent with extensive literature demonstrating a pivotal role of the thalamus in the systems-level mechanisms of anesthetic-induced unconsciousness (Nallasamy and Tsao, 2011). Interestingly, there is evidence indicating that thalamo-cortical connectivity in higher-tier associative networks is more strongly affected than that in lower-level sensory-motor networks by anesthesia, highlighting the importance of consciousness on integrative multimodal processing (Boveroux et al., 2010; Liang et al., 2012b; Liang et al., 2013). Furthermore, dynamic rsfMRI approaches reveal that anesthesia diminishes RSFC between medial prefrontal cortex (mPFC) and the hippocampus (Liang et al., 2015a), as well as RSFC between mPFC and amygdala (Liang et al., 2012b) as compared to the awake state, suggesting that anesthesia can profoundly impact neural circuits subserving cognitive and emotional function. This result is consistent with the notion that cognitive and emotional regulations are overall disrupted in the anesthetized state.

Similar effects of anesthesia on RSFC have also been observed in numerous human studies. RSFC in the cortex is ablated in deeply anesthetized patients (Peltier et al., 2005). In particular, like animal studies, anesthesia disrupts RSFC within the thalamocortical and frontoparietal networks in humans (Breshears et al., 2010; Lee et al., 2009a, 2009b; Velly et al., 2007; White and Alkire, 2003). Like anesthesia, sleep has similar effects on RSFC (Tomasi et al., 2009). Spontaneous fluctuations of the BOLD signal and RSFC are significantly reduced in sleep compared to wakefulness (Wilson et al., 2015).

Anesthesia can also lead to considerable changes in anticorrelated RSFC. Although the majority of rsfMRI studies use positive temporal correlations between BOLD time series as a measure of RSFC, negative correlations (i.e. anticorrelated RSFC) also contains important information of neural circuit function (e.g. inhibitory interactions between regions). For

instance, anticorrelated RSFC has been shown to play a critical role in mediating cognitive function in humans (Fox et al., 2005; Hampson et al., 2010; Kelly et al., 2008). In animal studies, robust anticorrelated RSFC in a well-characterized frontolimbic circuit between the mPFC and amygdala exhibited in the awake rat, whereas this anticorrelated RSFC was abolished in anesthetized rats (Liang et al., 2012a) (Fig. 5). These results are consistent with a recent study reporting both positive and negative RSFC in the awake monkey, but only positive RSFC when the monkey was anesthetized (Barttfeld et al., 2015).

In addition to the resting state, neural circuit function can be studied in activated states. The emergence of optogenetics has opened a new avenue for interrogating the neural circuit function that mediates many behaviors (Deisseroth et al., 2006). Such capacity has been highlighted by a series of recent studies that combine optogenetic control and fMRI readout (Abe et al., 2012; Kahn et al., 2013; Lee et al., 2010; Li et al., 2011; Liang et al., 2015b; Weitz et al., 2015). In an optogenetic approach, light-activated ion channels are introduced to a specifically targeted neuron population through adeno-associated viral (AAV) injection. This allows researchers to specifically activate/inhibit the infected neurons through optical stimulation (Deisseroth et al., 2006). The scientific value of this technique is tremendously expanded by combining it with fMRI (opto-fMRI), because the signaling impact of photo-activating a specific circuit element on the rest of circuit can be monitored across the whole brain (Lee et al., 2010). Compared to conventional electrical stimulation approaches, opto-fMRI does not suffer from electromagnetic-related artifacts that electrical approaches commonly encounter.

Anesthesia also exerts significant effects on neural circuit function in the activated state. Opto-fMRI conducted in awake mice (Desai et al., 2011) and rats (Ferenczi et al., 2016; Liang et al., 2015b) have consistently indicated that BOLD activations accompanying optogenetic stimulation in both injected regions and postsynaptic targets are strongly suppressed by anesthesia. In addition, the neural circuit connectivity pattern is significantly altered in the anesthetized state (Liang et al., 2015b) (Fig. 6). Taken together, all these studies strongly suggest that anesthesia can significantly alter neural circuit function. In fact, RSFC changes in anesthetized states have been detected using rsfMRI and have provided important insight into the system-level neural mechanism underlying anesthetic-induced unconsciousness (Liang et al., 2012b).

### **Anesthesia causes changes in brain network organization**

The intrinsic functional connectivity architecture is a crucial component of the governing principles of brain organization. This connectivity architecture has typically been studied by combining rsfMRI (or diffusion tensor imaging) with *graph theory*. Brain graphs are composed of a number of *nodes* interconnected by a set of *edges*. Anatomically or functionally defined region of interests (ROIs) are defined as nodes in the brain graph, and functional or structural connectivity between ROIs is used to define edges. Connectional architecture of brain networks can be quantitatively evaluated using a set of topological parameters of the brain graph, such as *degree*, *clustering coefficient*, *shortest path length* and *modularity*.

Relative to the awake state, brain networks are topologically altered to support different patterns of information transfer in the anesthetized state. Our data indicate that neural networks are significantly reorganized and exhibit different network properties (Liang et al., 2012b). For instance, several brain regions demonstrate pronounced changes in local topological metrics such as *clustering coefficient* and *betweenness centrality*. In particular, relative to the awake condition, regions of the basal ganglia, including nucleus accumbens and septal nuclei, show significantly reduced local clustering coefficients in the isoflurane-induced anesthetized condition, which suggests that the communication between those regions and their neighboring regions is reduced in the anesthetized animal. Interestingly, the same two regions can enhance anesthetic effects when they are pharmacologically inactivated (Ma and Leung, 2006; Ma et al., 2002). These results collectively highlight the susceptibility of the basal ganglia to anesthesia. In addition, thalamic nuclei show decreased betweenness centrality, again indicating impaired information relay in the thalamus in the anesthetized brain. Furthermore, the community structure between the awake and anesthetized states considerably differs in ways with stark functional implications (Fig. 7). Brain modules in the awake brain are more likely to contain both cortical and sub-cortical regions (Liang et al., 2011), whereas brain modules in the anesthetized brain tend to include only cortical or only subcortical regions, implying compromised communication between the cortex and subcortex in the anesthetized state (Liang et al., 2012b). The impact of anesthesia on the topological organization of brain networks has also been reported in several EEG studies. Lee et al. demonstrated reconfiguration of network hub structure after propofol-induced unconsciousness (Lee et al., 2013), and a disruption of network topology during the transition to loss of consciousness (Lee et al., 2011). Taken together, these studies strongly indicate that brain networks are topologically reorganized to support new patterns of information integration in the anesthetized condition.

### **Technical advances enable the study of neurovascular coupling and neural circuit function in the awake animal**

In the systems neuroscience community, the realization that anesthesia profoundly disrupts normal neural circuit function has driven neurophysiologists to embrace the use of awake animals, and many of these approaches and tools can be translated to the study of neurovascular coupling. A key advance has been the widespread adoption of head-fixation methods, first developed in the primate neurophysiology community (Wurtz, 1969) and then adapted the use in rodents (Dombeck et al., 2007) (see also Kleinfeld and Griesbeck (2005) and references therein). Both mice (Murphy et al., 2016) and rats (Scott et al., 2013; Scott et al., 2015) can be trained to voluntarily engage in head fixation, indicating that head fixation is not an aversive experience.

Though hemodynamic signals can be imaged from awake, head-fixed rodents using fMRI or optical imaging, there are methodological differences due to technical considerations. Additionally, there are species differences, with mice acclimating to head-fixation more rapidly than rats, which in turn acclimate more rapidly than marmosets (Belcher et al., 2013b; Liu et al., 2013a) and other primates. Each species has its own unique advantages and limitations, and the choice of species should be driven by the scientific questions

addressed (Hall et al., 2014; Hill et al., 2015; Hung et al., 2015a). Below we discuss the approaches used in mice and rats.

### Habituation to head fixation

The habituation procedures depend on species (mice or rats) and imaging modality (MRI or optical). Optical imaging requires a very still head relative to the optical imaging apparatus, requiring a metal headbar, but this type of fixation is avoided in MRI due to the issue of devices' MRI compatibility. Anecdotally, mice are easier to acclimate to head fixation, with some groups being able to bypass acclimation all together (Dombeck et al., 2009; Sofroniew et al., 2014).

For optical imaging experiments, prior to habituation a headbar is surgically attached to the skull to hold the head still. A cranial window may also be implanted at this time. After allowing adequate time for surgical recovery, we begin the habituation to head-fixation, which can be done on treadmill that allows locomotion (Dombeck et al., 2010; Gao et al., 2015; Gao and Drew, 2014; Huo et al., 2014; Rosenegger et al., 2015; Tran and Gordon 2015), or body restraining cylinders or slings (Berwick et al., 2002; Drew et al., 2011) (Fig. 8). We typically habituate mice to being on the treadmill or head-fixing in a tube by acclimating them over two to five days (Drew et al., 2010a; Drew et al., 2011; Huo et al., 2015a; Gao and Drew, 2016) in several 15-min to ½-h sessions. Mice are briefly (< 1 min) anesthetized with isoflurane before being head fixed to minimize handling stress. This habituation procedure and schedule is based on those used in neurophysiology experiments (Dombeck et al., 2007; Zagha et al., 2013). With a stable preparation, the noise in the intrinsic signal due to animal motion is negligible, as it is several orders of magnitude below the amplitude of the sensory-evoked response (Huo et al., 2014; 2015a; 2015b). Two-photon microscopy is sensitive to animal movement, but movement during voluntary locomotion are typically small enough to be corrected using motion correction software (Dombeck et al., 2007; Gao and Drew, 2016). Head-fixation and optical imaging can be longitudinally done on very young animals, starting as early as postnatal day 3 (P3) (Letourneur et al., 2014). Using two-photon microscopy, we are able to image locomotion-induced vasodilation in the somatosensory cortex (Fig. 9). Note that even small movements can drive 5–10% increases in arterial diameter, and prolonged motion can lead to arterial dilation of 40% or more, much larger than what is observed in the anesthetized animal. Large windows allow chronic measurement of hemodynamic signals over the entire dorsal surface of the cortex. The arterial dilation is localized to the somatosensory and visual cortex (Fig. 10), with little arterial dilation seen in the frontal cortex (Huo et al., 2014, 2015a, 2015b, Gao and Drew, 2016).

Technical challenges involved in imaging awake animals in MR scanners include controlling for motion artifacts and minimizing the stress induced by scanning noise and environment. To resolve these issues, an acclimation procedure is typically applied to habituate the animal to MRI scanning environment. Through this acclimation paradigm, animal's motion and stress during MRI scanning are substantially diminished (Ferenczi et al., 2016; King et al., 2005). This method has been extensively used in a number of awake animal fMRI studies involving multiple species including mice (Desai et al., 2011; Harris et al., 2015), rats

(Becerra et al., 2011; Ferris et al., 2006; Huang et al., 2011; Lahti et al., 1998; Liang et al., 2011; Martin et al., 2006; Zhang et al., 2010), monkeys (Belcher et al., 2013a; Hung et al., 2015b; Liu et al., 2013b), rabbits (Schroeder et al., 2016) and dogs (Andics et al., 2016; Berns et al., 2012).

Although acclimation paradigms vary across labs, they typically use a noninvasive restraint system and a routine habituation procedure over several days (see review (Febo, 2011)). A rat head restrainer is usually composed to a bite bar, a pair of adjustable ear bars or ear pads, a nose clamp and a shoulder pad. In terms of acclimation procedure, using our lab as an example (Fig. 11), the rat is first briefly anesthetized (2–3% isoflurane given for 3–5 min) to enable secure placement of the animal's head into a head holder with the canines secured over a bite bar, the nose fixed with a clamp, and ears positioned inside the head holder with adjustable ear pads or ear bars. Topical application of lidocaine (2%) or EMLA cream is used to relieve any discomfort associated with head-fixation. The animal's forepaws and hindpaws are loosely taped to prevent any self-injurious behavior (i.e. scratching). The body is then placed in a Plexiglas body tube that allows unrestricted respiration and movement of the trunk and limbs. The entire unit is subsequently secured onto a firm base, and then placed in a black opaque tube "mock scanner". An audio-recording of acoustic sounds from various imaging pulse sequences are played throughout the acclimation procedure. The animal is exposed to these conditions for 7 days before imaging. The time for exposure during acclimation is increased from 15 min on the first day to 60 min on days 4, 5, 6, and 7, with an increment of 15 min per day.

### Minimizing stress during awake imaging

One concern with the use of awake animals is that they might be overly stressed by head-fixation or imaging. While we cannot say that there is no stress associated with awake imaging procedures, this stress is most likely minimal.

For optical imaging in mice, habituation for several days is likely on the conservative side, as some groups successfully use as short as one day of acclimation (Sofroniew et al., 2014), animals become proficient in using a treadmill in 10–15 min (Niell and Stryker, 2010), and habituation does not change neural responses, as neural activity in the motor cortex is the same in habituated and un-habituated mice (Dombeck et al., 2009). Mice rapidly (usually within one session) acclimate to head-fixation, and learn to 'operate' the treadmill. After acclimation, resting heart rates in head-fixed mice are comparable to those from recorded telemetrically from mice in their home cage (Gehrmann et al., 2000, Huo et al., 2015b) (Fig. 10), indicating that that head fixation after acclimation is not stressful. In our hands, the pattern of evoked hemodynamic signals are stable up to 10 months of continuous imaging (Huo et al., 2015a, 2015b), which would not be the case if the animals were stressed. Many other groups have used head-fixed rodents, and have shown that they can perform complex, cognitively demanding tasks (Harvey et al., 2009; Komiyama et al., 2010; Mayrhofer et al., 2013; Mehta, 2007; Smear et al., 2011) that use memory or discrimination of stimuli at very high levels, which would not be possible if the animals were excessively stressed.

In awake animals, invasive measurements of physiological variables (arterial pressure, blood gasses, etc.) are more difficult to measure. However, other physiological parameters, such as

heart rate (Fig. 10), can be easily extracted from optically signals (Drew et al., 2011; Huo et al., 2015a, 2015b). Other indicators of state, such as whisker or body motion (Figs. 8–10) can also be used as moment-to-moment measures of animal's state.

Likewise, although restraint is involved in the MRI acclimation paradigm, this procedure is not likely to induce chronic stress in animals. Unlike studies investigating the chronic stress effects of immobilization, the restraint for acclimation and imaging requires considerably shorter restraining period per day and only involves restraint of the head as oppose to the whole body. Indeed, King et al. reported that the level of stress hormone (corticosterone) and other physiologic parameters of rats return to pre-stress baseline following 5–8 days of acclimation (King et al., 2005). The data are consistent with reports that heart rate and blood pressure normalized after initial exposure to restraint in rats, indicating that the animal well adapted to the restrainer (Parry and McElligott, 1993). Rats that were acclimated and imaged up to 4 times did not show significantly altered anxiety level compared to intact rats as measured by elevated plus maze test (Fig. 12).

Importantly, acclimation does not compound effects of other stressors on animals. In our recent study, the stress attributed to acclimation was minimal and significantly less than the stress introduced by a different stressor (i.e. single episode of predator odor exposure) (Liang et al., 2014). Notably, we found that there was no interaction between the acclimation-related stress and predator odor-related stress, suggesting that any possible residual effects from acclimation- and/or imaging-related stress can be subtracted out in testing animals with appropriate controls, which undergo the identical acclimation and/or imaging procedures. These results allow us to specifically examine the effect predator odor stress on neural circuit function in awake rodents (Liang et al., 2014). Taken together, these data suggest that stress induced by acclimation and imaging is very mild in animals and can be well controlled.

It is important to keep in mind that anesthesia is not 'stress-free', and in some cases can induce stress-related hormonal changes substantially larger than behavioral stressors. Urethane increases circulating adrenocorticotrophic hormone by a factor of four, substantially higher than open-field stress (Donnerer and Lembeck, 1988). Studies in human patients have shown that anesthesia increases circulating cytokines (Brand et al., 1997), and can increase circulating epinephrine, norepinephrine, and cortisol (Kono et al., 1981) by up to a factor of two or more, though fentanyl can cause a substantial decrease in circulating cortisol relative to control conditions (Kono et al., 1981). Above and beyond the effects of anesthetic, surgical procedures can cause enormous hormonal changes (Desborough, 2000), making it preferable to temporally separate any surgical procedure (such as cranial window implantation) from the imaging session.

### **Neurovascular coupling and functional connectivity are robust and stable in the awake animal**

Historically, anesthesia has been used as an important experimental tool for perturbing the state of the brain in a reproducible manner. For awake animal imaging approaches to be successfully applied, it is essential to validate the reproducibility of data collected.

Neuroimaging data collected in awake animals using aforementioned techniques demonstrate a high degree of reproducibility. Our opto-fMRI data acquired in two sessions separated by 2–7 days show highly consistent BOLD activation patterns in all rats (Fig. 13a). Besides, BOLD amplitude of the corresponding voxels in the two maps were highly correlated ( $r=0.79$ ,  $p < 10^{-20}$ ) (Liang et al., 2015b). In addition, rsfMRI data collected in two subgroups of awake rodents show highly reproducible connectivity profile, as well as a high correlation in voxel-wise connectivity strength (Fig. 13b) (Liang et al., 2012a). Furthermore, whole-brain RSFC profiles can be used to produce connectivity-based functional parcellations of the rat brain, which exhibit high reproducibility across animals (Ma et al., 2016). Similarly, when locomotion-induced changes in CBV are fit with linear models, these fits are very reproducible across days (Huo et al., 2015a) (Fig. 13c and d), indicating that even with spontaneous, volitional behaviors, the hemodynamic response is stable in head-fixed animals.

One factor that may confound the reproducibility of neuroimaging data in awake animals is spontaneous behaviors. Any awake animal (or human) will engage in small spontaneous behaviors in the scanner, and these spontaneous behaviors could affect hemodynamic signals. Several behaviors known to cause or correlate with measurable changes in hemodynamic signals and functional connectivity include eye state (closed, open or fixating) (Patriat et al. 2013; Chang et al., 2016), blink rate (Guipponi et al., 2014), and variations in respiration (Birn et al., 2009). In rodents, behaviors like whisking and sniffing (Moore et al., 2013) are analogous to eye movements in humans and primates (Kleinfeld et al., 2006; Crapse and Sommer, 2008; Ahissar and Assa, 2016), and these behaviors would be expected to have similar impacts on functional connectivity. These behavioral effects can be regressed out, as is currently done with physiological noise (Birn 2012; Murphy et al., 2013). Going forward, monitoring of the relevant behavior(s) (Fig. 8 and (Guipponi et al., 2014; Chang et al., 2016; McGinley et al., 2015; O'Connor et al., 2010; Yang et al., 2016)) in animals is likely to play an important role in animal fMRI experiments, just as it has in human experiments (Richlan et al., 2014; Wang et al., 2016). The development of new techniques for analyzing behavioral data in other paradigms have undergone enormous advances in recent years (Gomez-Marín et al. 2014), and it is likely that some of these approaches can be applied to the studies of more subtle behaviors.

Another concern is the normal fluctuations in blood pressure and heart rate that are found in the awake state (Goldberger et al., 2002) might affect neurovascular coupling. However, sensory-evoked changes in CBF and arterial dilation are not affected by large changes in heart rate in the awake animal. CBF measurements in mice where heart rate was pharmacologically elevated with glycopyrrolate, or decreased with atenolol showed that sensory-evoked CBF increase was not affected by these large manipulations of heart rate, and neither was arterial dilation (Huo et al., 2015b). However, pharmacological depression of heart rate reduced venous dilation in response to sensory stimulation, similar to that in the anesthetized animal (Drew et al., 2011). These results indicate that in the awake animal, the hemodynamic response is largely un-affected by cardiovascular fluctuations as long as the heart rate is not depressed. Further work is still needed to understand what contributions normal cardiovascular fluctuations might have on neurovascular coupling.



## Conclusions

Anesthetics have been widely used for studying neurovascular coupling in *in-vivo* animal models, and these models have added tremendously to our understanding of neurovascular coupling and associated physiological processes (Logothetis et al., 2001; Vincent et al., 2007; Shulman et al., 1999). However, anesthesia has profound actions on vascular, electrical, metabolic, and other physiological processes, and alters the coupling relationship between neural and vascular responses. Therefore, it needs to be cautious when one aims to translate functional neuroimaging data acquired in anesthetized animals to awake humans. Recent technical developments have enabled the use of optical and MR imaging in awake and behaving animals, allowing scientists to address critical questions about neurovascular coupling and brain-wide circuit function in a preparation similar to the awake human. Certainly, these methods are still in a relatively early stage, and will likely undergo further improvement in the future. Nevertheless, we believe the time is right for the community to move towards using awake animal paradigms when addressing questions involving neurovascular coupling and circuit function.

## Acknowledgments

PJD is supported by award 12SDG9130022 from the American Heart Association, a Scholar Award from the McKnight Endowment Fund for Neuroscience, and R01NS078168, R01EB021703, and R01NS079737 from the NIH. NZ is supported by R01MH098003 from the National Institute of Mental Health and R01NS085200 from the National Institute of Neurological Disorders and Stroke.

## References

- Abe Y, Sekino M, Terazono Y, Ohsaki H, Fukazawa Y, Sakai S, Yawo H, Hisatsune T. Opto-fMRI analysis for exploring the neuronal connectivity of the hippocampal formation in rats. *Neurosci Res*. 2012; 74:248–255. [PubMed: 22982343]
- Ahissar, E., Assa, E. Perception as a closed-loop convergence process; *eLife*. 2016. p. 5 <http://dx.doi.org/10.7554/eLife.12830>
- Akata T, Kanna T, Yoshino J, Takahashi S. Mechanisms of direct inhibitory action of isoflurane on vascular smooth muscle of mesenteric resistance arteries. *Anesthesiology*. 2003; 99:666–677. [PubMed: 12960552]
- Aksenov DP, Li L, Miller MJ, Iordanescu G, Wyrwicz AM. Effects of anesthesia on BOLD signal and neuronal activity in the somatosensory cortex. *J Cereb Blood Flow Metab*. 2015; 35(11):1819. <http://dx.doi.org/10.1038/jcbfm.2015.130>. [PubMed: 26104288]
- Albert NB, Robertson EM, Miall RC. The resting human brain and motor learning. *Curr Biol*. 2009; 19:1023–1027. [PubMed: 19427210]
- Alkire MT, Haier RJ, Barker SJ, Shah NK, Wu JC, Kao YJ. Cerebral metabolism during propofol anesthesia in humans studied with positron emission tomography. *Anesthesiology*. 1995; 82:393–403. (discussion 27A). [PubMed: 7856898]
- Alkire MT, Haier RJ, Fallon JH. Toward a unified theory of narcosis: brain imaging evidence for a thalamocortical switch as the neurophysiologic basis of anesthetic-induced unconsciousness. *Conscious Cogn*. 2000; 9:370–386. <http://dx.doi.org/10.1006/ccog.1999.0423>. [PubMed: 10993665]
- Alkire MT, Haier RJ, Shah NK, Anderson CT. Positron emission tomography study of regional cerebral metabolism in humans during isoflurane anesthesia. *Anesthesiology*. 1997; 86:549–557. [PubMed: 9066320]
- Alkire MT, Hudetz AG, Tononi G. Consciousness and anesthesia. *Science*. 2008; 322:876–880. <http://dx.doi.org/10.1126/science.1149213>. [PubMed: 18988836]

- Anand A, Li Y, Wang Y, Wu J, Gao S, Bukhari L, Mathews VP, Kalnin A, Lowe MJ. Activity and connectivity of brain mood regulating circuit in depression: a functional magnetic resonance study. *Biol Psychiatry*. 2005; 57:1079–1088. [PubMed: 15866546]
- Andics A, Gabor A, Gacsi M, Farago T, Szabo D, Miklosi A. Neural mechanisms for lexical processing in dogs. *Science*. 2016; 353:1030–1032. [PubMed: 27576923]
- Antunes LM, Golledge HDR, Roughan JV, Flecknell PA. Comparison of electroencephalogram activity and auditory evoked responses during isoflurane and halothane anaesthesia in the rat. *Vet Anaesth Analg*. 2003; 30:15–23. [PubMed: 14498913]
- Armstrong-James M, George MJ. Influence of anesthesia on spontaneous activity and receptive field size of single units in rat SmI neocortex. *Exp Neurol*. 1988; 99:369–387. [PubMed: 3338529]
- Aronov D, Fee MS. Natural changes in brain temperature underlie variations in song tempo during a mating behavior. *PLoS One*. 2012; 7:e47856. [PubMed: 23112858]
- Artru AA. Relationship between cerebral blood volume and CSF pressure during anesthesia with isoflurane or fentanyl in dogs. *Anesthesiology*. 1984; 60:575–579. [PubMed: 6731913]
- Attwell D, Buchan AM, Charpak S, Lauritzen M, Macvicar BA, Newman EA. Glial and neuronal control of brain blood flow. *Nature*. 2010; 468:232–243. <http://dx.doi.org/10.1038/nature09613>. [PubMed: 21068832]
- Attwell D, Gibb A. Neuroenergetics and the kinetic design of excitatory synapses. *Nat Rev Neurosci*. 2005; 6:841–849. <http://dx.doi.org/10.1038/nrn1784>. [PubMed: 16261178]
- Austin VC, Blamire AM, Allers KA, Sharp T, Styles P, Matthews PM, Sibson NR. Confounding effects of anesthesia on functional activation in rodent brain: a study of halothane and  $\alpha$ -chloralose anesthesia. *NeuroImage*. 2005; 24:92–100. <http://dx.doi.org/10.1016/j.neuroimage.2004.08.011>. [PubMed: 15588600]
- Barttfeld P, Uhrig L, Sitt JD, Sigman M, Jarraya B, Dehaene S. Signature of consciousness in the dynamics of resting-state brain activity. *Proc Natl Acad Sci USA*. 2015; 112:887–892. [PubMed: 25561541]
- Becerra L, Pendse G, Chang PC, Bishop J, Borsook D. Robust reproducible resting state networks in the awake rodent brain. *PLoS One*. 2011; 6:e25701. [PubMed: 22028788]
- Belcher AM, Yen CC, Stepp H, Gu H, Lu H, Yang Y, Silva AC, Stein EA. Large-scale brain networks in the awake, truly resting marmoset monkey. *J Neurosci*. 2013a; 33:16796–16804. [PubMed: 24133280]
- Belcher AM, Yen CC, Stepp H, Gu H, Lu H, Yang Y, Silva AC, Stein EA. Large-scale brain networks in the awake, truly resting marmoset monkey. *J Neurosci*. 2013b; 33:16796–16804. [PubMed: 24133280]
- Berg RW, Friedman B, Schroeder LF, Kleinfeld D. Activation of nucleus basalis facilitates cortical control of a brain stem motor program. *J Neurophysiol*. 2005; 94:699–711. <http://dx.doi.org/10.1152/jn.01125.2004>. [PubMed: 15728764]
- Berns GS, Brooks AM, Spivak M. Functional MRI in awake unrestrained dogs. *PLoS One*. 2012; 7:e38027. [PubMed: 22606363]
- Berwick J, Martin C, Martindale J, Jones M, Johnston D, Zheng Y, Redgrave P, Mayhew J. Hemodynamic response in the unanesthetized rat: intrinsic optical imaging and spectroscopy of the barrel cortex. *J Cereb Blood Flow Metab*. 2002; 22:670–679. [PubMed: 12045665]
- Birn RM. The role of physiological noise in resting-state functional connectivity. *NeuroImage*. 2012; 62(2):864–870. <http://dx.doi.org/10.1016/j.neuroimage.2012.01.016>. [PubMed: 22245341]
- Birn RM, Smith Ma, Jones TB, Bandettini Pa. The Respiration Response Function: the temporal dynamics of fMRI signal fluctuations related to changes in respiration. *NeuroImage*. 2009; 40(2): 644–654.
- Biswal B, Yetkin FZ, Haughton VM, Hyde JS. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn Reson Med*. 1995; 34:537–541. [PubMed: 8524021]
- Bonder DE, McCarthy KD. Astrocytic Gq-GPCR-linked IP3R-dependent  $Ca^{2+}$  signaling does not mediate neurovascular coupling in mouse visual cortex in vivo. *J Neurosci*. 2014; 34:13139–13150. [PubMed: 25253859]

- Bonvento G, Charbonné R, Corrèze JL, Borredon J, Seylaz J, Lacombe P. Is alpha-chloralose plus halothane induction a suitable anesthetic regimen for cerebrovascular research? *Brain Res.* 1994; 665:213–221. [PubMed: 7895056]
- Boveroux P, Vanhauzenhuysse A, Bruno MA, Noirhomme Q, Lauwick S, Luxen A, Degueldre C, Plenevaux A, Schnakers C, Phillips C, Brichant JF, Bonhomme V, Maquet P, Greicius MD, Laureys S, Boly M. Breakdown of within- and between-network resting state functional magnetic resonance imaging connectivity during propofol-induced loss of consciousness. *Anesthesiology.* 2010; 113:1038–1053. [PubMed: 20885292]
- Brand JM, Kirchner H, Poppe C, Schmucker P. The effects of general anesthesia on human peripheral immune cell distribution and cytokine production. *Clin Immunol Immunopathol.* 1997; 83:190–194. [PubMed: 9143381]
- Braun AR, Balkin TJ, Wesenten NJ, Carson RE, Varga M, Baldwin P, Selbie S, Belenky G, Herscovitch P. Regional cerebral blood flow throughout the sleep-wake cycle. An H2(15)O PET study. *Brain.* 1997; 120:1173–1197. [PubMed: 9236630]
- Breshears JD, Roland JL, Sharma M, Gaona CM, Freudenburg ZV, Tempelhoff R, Avidan MS, Leuthardt EC. Stable and dynamic cortical electrophysiology of induction and emergence with propofol anesthesia. *Proc Natl Acad Sci USA.* 2010; 107:21170–21175. [PubMed: 21078987]
- Brown, DL. *Atlas of Regional Anesthesia.* Elsevier Health Sciences; Philadelphia: 2010.
- Brown EN, Lydic R, Schiff ND. General anesthesia, sleep, and coma. *N Engl J Med.* 2010; 363:2638–2650. [PubMed: 21190458]
- Brown EN, Purdon PL, Van Dort CJ. General anesthesia and altered states of arousal: a systems neuroscience analysis. *Annu Rev Neurosci.* 2011; 34:601–628. [PubMed: 21513454]
- Buchsbaum MS, Gillin JC, Wu J, Hazlett E, Sicotte N, Dupont RM, Bunney WE. Regional cerebral glucose metabolic rate in human sleep assessed by positron emission tomography. *Life Sci.* 1989; 45:1349–1356. [PubMed: 2796606]
- Buljubasic N, Rusch NJ, Marijic J, Kampine JP, Bosnjak ZJ. Effects of halothane and isoflurane on calcium and potassium channel currents in canine coronary arterial cells. *Anesthesiology.* 1992; 76:990–998. [PubMed: 1318010]
- Buxton RB. Dynamic models of BOLD contrast. *NeuroImage.* 2012; 62:953–961. <http://dx.doi.org/10.1016/j.neuroimage.2012.01.012>. [PubMed: 22245339]
- Buxton RB, Uluda K, Dubowitz DJ, Liu TT. Modeling the hemodynamic response to brain activation. *NeuroImage.* 2004; 23(Suppl. 1):S220–S233. <http://dx.doi.org/10.1016/j.neuroimage.2004.07.013>. [PubMed: 15501093]
- Buxton RB, Wong EC, Frank LR. Dynamics of blood flow and oxygenation changes during brain activation: the balloon model. *Magn Reson Med.* 1998; 39:855–864. <http://dx.doi.org/10.1002/mrm.1910390602/pdf>. [PubMed: 9621908]
- Cardin JA, Schmidt MF. Auditory responses in multiple sensorimotor song system nuclei are co-modulated by behavioral state. *J Neurophysiol.* 2004; 91:2148–2163. <http://dx.doi.org/10.1152/jn.00918.2003>. [PubMed: 14724261]
- Cardin JA, Schmidt MF. Song system auditory responses are stable and highly tuned during sedation, rapidly modulated and unselective during wakefulness, and suppressed by arousal. *J Neurophysiol.* 2003; 90:2884–2899. <http://dx.doi.org/10.1152/jn.00391.2003>. [PubMed: 12878713]
- Carter AR, Shulman GL, Corbetta M. Why use a connectivity-based approach to study stroke and recovery of function? *Neuroimage.* 2012; 62:2271–2280. [PubMed: 22414990]
- Castro-Alamancos MA. Cortical up and activated states: implications for sensory information processing. *Neuroscientist.* 2009; 15:625–634. <http://dx.doi.org/10.1177/1073858409333074>. [PubMed: 19321459]
- Castro-Alamancos MA. Absence of rapid sensory adaptation in neocortex during information processing states. *Neuron.* 2004; 41:455–464. [PubMed: 14766183]
- Castro-Alamancos MA, Gulati T. Neuromodulators produce distinct activated states in neocortex. *J Neurosci.* 2014; 34:12353–12367. <http://dx.doi.org/10.1523/JNEUROSCI.1858-14.2014>. [PubMed: 25209276]
- Cauli B, Hamel E. Revisiting the role of neurons in neurovascular coupling. *Front Neuroenergetics.* 2010; 2:9. <http://dx.doi.org/10.3389/fnene.2010.00009>. [PubMed: 20616884]

- Cazakoff BN, Lau BYB, Crump KL, Demmer HS, Shea SD. Broadly tuned and respiration-independent inhibition in the olfactory bulb of awake mice. *Nat Neurosci.* 2014; 17:569–576. <http://dx.doi.org/10.1038/nn.3669>. [PubMed: 24584050]
- Chang C, Leopold DA, Scholvinck ML, Mandelkow H, Picchioni D, Liu X, Ye FQ, Turchi JN, Duyn JH. Tracking brain arousal fluctuations with fMRI. *Proc Natl Acad Sci USA.* 2016; 113(16):4518–4523. <http://dx.doi.org/10.1073/pnas.1520613113>. [PubMed: 27051064]
- Chapin JK, Woodward DJ. Modulation of sensory responsiveness of single somatosensory cortical cells during movement and arousal behaviors. *Exp Neurol.* 1981; 72:164–178. [PubMed: 7202621]
- Chen BR, Bouchard MB, McCaslin AFH, Burgess SA, Hillman EMC. High-speed vascular dynamics of the hemodynamic response. *NeuroImage.* 2011; 54:1021–1030. <http://dx.doi.org/10.1016/j.neuroimage.2010.09.036>. [PubMed: 20858545]
- Chever O, Sik A, Amzica F, Te S. Opening of the blood–brain barrier during isoflurane anaesthesia. *Eur J Neurosci.* 2008; 28:1330–1341. [PubMed: 18973560]
- Constantinople CM, Bruno RM. Effects and mechanisms of wakefulness on local cortical networks. *Neuron.* 2011; 69:1061–1068. [PubMed: 21435553]
- Crane PD, Braun LD, Cornford EM, Cremer JE, Glass JM, Oldendorf WH. Dose dependent reduction of glucose utilization by pentobarbital in rat brain. *Stroke.* 1978; 9:12–18. <http://dx.doi.org/10.1161/01.STR.9.1.12>. [PubMed: 622738]
- Crapse TB, Sommer MA. Corollary discharge across the animal kingdom. *Nat Rev Neurosci.* 2008; 9(8):587–600. <http://dx.doi.org/10.1038/nrn2457>. [PubMed: 18641666]
- Curto C, Sakata S, Marguet S, Itskov V. A simple model of cortical dynamics explains variability and state dependence of sensory responses in urethane-anesthetized auditory cortex. *J Neurosci.* 2009; 29:10600–10612. [PubMed: 19710313]
- de Kock CPJ, Sakmann B. Spiking in primary somatosensory cortex during natural whisking in awake head-restrained rats is cell-type specific. *Proc Natl Acad Sci USA.* 2009; 106:16446–16450. <http://dx.doi.org/10.1073/pnas.0904143106>. [PubMed: 19805318]
- Deisseroth K, Feng G, Majewska AK, Miesenbock G, Ting A, Schnitzer MJ. Next-generation optical technologies for illuminating genetically targeted brain circuits. *J Neurosci.* 2006; 26:10380–10386. [PubMed: 17035522]
- Desborough JP. The stress response to trauma and surgery. *Br J Anaesth.* 2000; 109–117. [PubMed: 10927999]
- Desai M, Kahn I, Knoblich U, Bernstein J, Atallah H, Yang A, Kopell N, Buckner RL, Graybiel AM, Moore CI, Boyden ES. Mapping brain networks in awake mice using combined optical neural control and fMRI. *J Neurophysiol.* 2011; 105:1393–1405. [PubMed: 21160013]
- Devor A, Tian P, Nishimura N, Teng IC, Hillman EMC, Narayanan SN, Ulbert I, Boas DA, Kleinfeld D, Dale AM. Suppressed neuronal activity and concurrent arteriolar vasoconstriction may explain negative blood oxygenation level-dependent signal. *J Neurosci: Off J Soc Neurosci.* 2007; 27:4452–4459. <http://dx.doi.org/10.1523/JNEUROSCI.0134-07.2007>.
- Dombeck DA, Graziano MS, Tank DW. Functional clustering of neurons in motor cortex determined by cellular resolution imaging in awake behaving mice. *J Neurosci.* 2009; 29:13751–13760. [PubMed: 19889987]
- Dombeck DA, Harvey CD, Tian L, Looger LL, Tank DW. Functional imaging of hippocampal place cells at cellular resolution during virtual navigation. *Nat Neurosci.* 2010; 13:1433–1440. <http://dx.doi.org/10.1038/nn.2648>. [PubMed: 20890294]
- Dombeck DA, Khabbaz A, Collman F, Adelman T, Tank DW. Imaging large-scale neural activity with cellular resolution in awake, mobile mice. *Neuron.* 2007; 56:43–57. <http://dx.doi.org/10.1016/j.neuron.2007.08.003>. [PubMed: 17920014]
- Donnerer J, Lembeck F. Neonatal capsaicin treatment of rats reduces ACTH secretion in response to peripheral neuronal stimuli but not to centrally acting stressors. *Br J Pharmacol.* 1988; 94:647–652. [PubMed: 2846107]
- Drew PJ, Shih A, Driscoll J, Knutsen P, Blinder P, Davalos D, Akassoglou K, Tsai P, Kleinfeld D. Chronic optical access through a polished and reinforced thinned skull. *Nat Methods.* 2010a; 7:981–984. <http://dx.doi.org/10.1038/nmeth.1530>. [PubMed: 20966916]

- Drew PJ, Blinder P, Cauwenberghs G, Shih AY, Kleinfeld D. Rapid determination of particle velocity from space-time images using the Radon transform. *J Comput Neurosci*. 2010b; 29:5–11. [PubMed: 19459038]
- Drew PJ, Shih A, Kleinfeld D. Fluctuating and sensory-induced vasodynamics in rodent cortex extend arteriole capacity. *Proc Natl Acad Sci USA*. 2011; 108:8473–8478. [PubMed: 21536897]
- Dudley RE, Nelson SR, Samson F. Influence of chloralose on brain regional glucose utilization. *Brain Res*. 1982; 233:173–180. [http://dx.doi.org/10.1016/0006-8993\(82\)90938-6](http://dx.doi.org/10.1016/0006-8993(82)90938-6). [PubMed: 7059799]
- Febo M. Technical and conceptual considerations for performing and interpreting functional MRI studies in awake rats. *Front Psychiatry*. 2011; 2:43. [PubMed: 21808625]
- Ferenczi EA, Zalocusky KA, Liston C, Grosenick L, Warden MR, Amatya D, Katovich K, Mehta H, Patenaude B, Ramakrishnan C, Kalanithi P, Etkin A, Knutson B, Glover GH, Deisseroth K. Prefrontal cortical regulation of brainwide circuit dynamics and reward-related behavior. *Science*. 2016; 351:aac9698. [PubMed: 26722001]
- Ferezou I, Haiss F, Gentet LJ, Aronoff R, Weber B, Petersen CCH. Spatiotemporal dynamics of cortical sensorimotor integration in behaving mice. *Neuron*. 2007; 56:907–923. <http://dx.doi.org/10.1016/j.neuron.2007.10.007>. [PubMed: 18054865]
- Fernández-Klett F, Offenhauser N, Dirnagl U, Priller J, Lindauer U. Pericytes in capillaries are contractile in vivo, but arterioles mediate functional hyperemia in the mouse brain. *Proc Natl Acad Sci USA*. 2010; 107:22290–22295. <http://dx.doi.org/10.1073/pnas.1011321108>. [PubMed: 21135230]
- Ferris CF, Febo M, Luo F, Schmidt K, Brevard M, Harder JA, Kulkarni P, Messenger T, King JA. Functional magnetic resonance imaging in conscious animals: a new tool in behavioural neuroscience research. *J Neuroendocrinol*. 2006; 18:307–318. [PubMed: 16629829]
- Flynn N, Buljubasic N, Bosnjak ZJ, Kampine JP. Cerebral vascular responses to anesthetics. *Adv Exp Med Biol*. 1991; 301:237–246. [PubMed: 1662464]
- Fox PT, Raichle ME. Focal physiological uncoupling of cerebral blood flow and oxidative metabolism during somatosensory stimulation in human subjects. *Proc Natl Acad Sci USA*. 1986; 83:1140–1144. [PubMed: 3485282]
- Fox MD, Raichle ME. Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nat Rev Neurosci*. 2007; 8:700–711. [PubMed: 17704812]
- Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci USA*. 2005; 102:9673–9678. [PubMed: 15976020]
- Friedberg MH, Lee SM, Ebner FF. Modulation of receptive field properties of thalamic somatosensory neurons by the depth of anesthesia. *J Neurophysiol*. 1999; 81:2243–2252. <http://dx.doi.org/10.1002/cne.903030204>. [PubMed: 10322063]
- Gao YR, Drew PJ. Effects of voluntary locomotion and calcitonin gene-related peptide on the dynamics of single dural vessels in awake mice. *J Neurosci*. 2016; 36:2503–2516. <http://dx.doi.org/10.1523/JNEUROSCI.3665-15.2016>. [PubMed: 26911696]
- Gao YR, Drew PJ. Determination of vessel cross-sectional area by thresholding in Radon space. *J Cereb Blood Flow Metab*. 2014; 34:1180–1187. <http://dx.doi.org/10.1038/jcbfm.2014.67>. [PubMed: 24736890]
- Gao YR, Greene SE, Drew PJ. Mechanical restriction of intracortical vessel dilation by brain tissue sculpts the hemodynamic response. *NeuroImage*. 2015; 115:162–176. <http://dx.doi.org/10.1016/j.neuroimage.2015.04.054>. [PubMed: 25953632]
- Gehrmann J, Hammer PE, Maguire CT, Wakimoto H, Triedman JK, Berul CI. Phenotypic screening for heart rate variability in the mouse. *Am J Physiol - Heart Circ Physiol*. 2000; 279:H733–H740. [PubMed: 10924073]
- Goard M, Dan Y. Basal forebrain activation enhances cortical coding of natural scenes. *Nat Neurosci*. 2009; 12:1444–1449. <http://dx.doi.org/10.1038/nn.2402>. [PubMed: 19801988]
- Goense J, Logothetis NK. Positive and Negative BOLD-Signals From Blood Vessels in Monkey Visual Cortex. 2008
- Goldberger AL, Amaral LAN, Hausdorff JM, Ivanov PC, Peng CK, Stanley HE. Fractal dynamics in physiology: alterations with disease and aging. *Proceedings of the National Academy of Sciences*

- of the United States of America. 2002; 99:2466–2472. <http://dx.doi.org/10.1073/pnas.012579499>. [PubMed: 11875196]
- Gomez-Marin A, Paton JJ, Kampff AR, Costa RM, Mainen ZF. Big behavioral data: psychology, ethology and the foundations of neuroscience. *Nat Neurosci*. 2014; 17(11):1455–1462. <http://dx.doi.org/10.1038/nn.3812>. [PubMed: 25349912]
- Greenberg DS, Houweling AR, Kerr JND. Population imaging of ongoing neuronal activity in the visual cortex of awake rats. *Nat Neurosci*. 2008; 11:749–751. <http://dx.doi.org/10.1038/nn.2140>. [PubMed: 18552841]
- Greicius MD, Flores BH, Menon V, Glover GH, Solvason HB, Kenna H, Reiss AL, Schlaggar BL, Tootell RB, Mendola JD, Huxley-Hamill C, Drevets WC, Hyndy G, Ungerleider LG, Lavenex P, Rosen BR, Yeo BT, Polk S, D'Esposito M. Resting-state functional connectivity in major depression: abnormally increased contributions from subgenual cingulate cortex and thalamus. *Biol Psychiatry*. 2007; 62:429–437. [PubMed: 17210143]
- Greicius MD, Srivastava G, Reiss AL, Menon V. Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc Natl Acad Sci USA*. 2004; 101:4637–4642. [PubMed: 15070770]
- Guipponi O, Odouard S, Pinede S, Wardak C, Ben Hamed S. fMRI cortical correlates of spontaneous eye blinks in the nonhuman primate. *Cereb Cortex*. 2014
- Haider B, Häusser M, Carandini M. Inhibition dominates sensory responses in the awake cortex. *Nature*. 2013; 493:97–100. <http://dx.doi.org/10.1038/nature11665>. [PubMed: 23172139]
- Hall CN, Reynell C, Gesslein B, Hamilton NB, Mishra A, Sutherland BA, O'Farrell FM, Buchan AM, Lauritzen M, Attwell D. Capillary pericytes regulate cerebral blood flow in health and disease. *Nature*. 2014; 508:55–60. <http://dx.doi.org/10.1038/nature13165>. [PubMed: 24670647]
- Hampson M, Driesen N, Roth JK, Gore JC, Constable RT. Functional connectivity between task-positive and task-negative brain areas and its relation to working memory performance. *Magn Reson Imaging*. 2010; 28:1051–1057. [PubMed: 20409665]
- Handwerker DA, Ollinger JM, D'Esposito M. Variation of BOLD hemodynamic responses across subjects and brain regions and their effects on statistical analyses. *Neuroimage*. 2004; 21:1639–1651. [PubMed: 15050587]
- Harris AP, Lennen RJ, Marshall I, Jansen MA, Pernet CR, Brydges NM, Duguid IC, Holmes MC. Imaging learned fear circuitry in awake mice using fMRI. *Eur J Neurosci*. 2015; 42:2125–2134. [PubMed: 25943794]
- Harris JJ, Attwell D. The energetics of CNS white matter. *J Neurosci*. 2012; 32:356–371. <http://dx.doi.org/10.1523/JNEUROSCI.3430-11.2012>. [PubMed: 22219296]
- Hartmann DA, Underly RG, Grant RI, Watson AN, Lindner V, Shih AY. Pericyte structure and distribution in the cerebral cortex revealed by high-resolution imaging of transgenic mice. *Neurophotonics*. 2015; 2:041402. <http://dx.doi.org/10.1117/1.NPh.2.4.041402>. [PubMed: 26158016]
- Harvey CD, Collman F, Dombeck DA, Tank DW. Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature*. 2009; 461:941–946. <http://dx.doi.org/10.1038/nature08499>. [PubMed: 19829374]
- Hayward JN. Brain temperature regulation during sleep and arousal in the dog. *Exp Neurol*. 1968; 21:201–212. [PubMed: 5675327]
- Hill CE. Long Distance Conduction of Vasodilation: A Passive or Regenerative Process? *Microcirculation*. 2012; 19(5):379–390. <http://dx.doi.org/10.1111/j.1549-8719.2012.00169.x>. [PubMed: 22313498]
- Hill RA, Tong L, Yuan P, Murikinati S, Gupta S, Grutzendler J. Regional blood flow in the normal and ischemic brain is controlled by arteriolar smooth muscle cell contractility and not by capillary pericytes. *Neuron*. 2015; 87:95–110. <http://dx.doi.org/10.1016/j.neuron.2015.06.001>. [PubMed: 26119027]
- Hillman EMC, Devor A, Bouchard MB, Dunn AK, Krauss GW, Skoch J, Bacskaï BJ, Dale AM, Boas DA. Depth-resolved optical imaging and microscopy of vascular compartment dynamics during somatosensory stimulation. *Neuroimage*. 2007; 35:89–104. <http://dx.doi.org/10.1016/j.neuroimage.2006.11.032>. [PubMed: 17222567]

- Hirano Y, Stefanovic B, Silva AC. Spatiotemporal evolution of the functional magnetic resonance imaging response to ultrashort stimuli. *J Neurosci: Off J Soc Neurosci*. 2011; 31:1440–1447. <http://dx.doi.org/10.1523/JNEUROSCI.3986-10.2011>.
- Hirase H. A multi-photon window onto neuronal-glia-vascular communication. *Trends Neurosci*. 2005; 28:217–219. <http://dx.doi.org/10.1016/j.tins.2005.03.002>. [PubMed: 15866194]
- Horowitz SG, Braun AR, Carr WS, Picchioni D, Balkin TJ, Fukunaga M, Duyn JH. Decoupling of the brain's default mode network during deep sleep. *Proc Natl Acad Sci USA*. 2009; 106:11376–11381. [PubMed: 19549821]
- Howarth C, Gleeson P, Attwell D. Updated energy budgets for neural computation in the neocortex and cerebellum. *J Cereb Blood Flow Metab*. 2012; 32:1222–1232. <http://dx.doi.org/10.1038/jcbfm.2012.35>. [PubMed: 22434069]
- Huang W, Heffernan ME, Li Z, Zhang N, Overstreet DH, King JA. Fear induced neuronal alterations in a genetic model of depression: an fMRI study on awake animals. *Neurosci Lett*. 2011; 489:74–78. [PubMed: 21134416]
- Hubel DH, Wiesel TN. Receptive fields of single neurones in the cat's striate cortex. *J Physiol*. 1959; 148:574–591. [PubMed: 14403679]
- Hung CC, Yen CC, Ciuchta JL, Papoti D, Bock NA, Leopold DA, Silva AC. Functional mapping of face-selective regions in the extrastriate visual cortex of the marmoset. *J Neurosci*. 2015a; 35:1160–1172. [PubMed: 25609630]
- Hung CC, Yen CC, Ciuchta JL, Papoti D, Bock NA, Leopold DA, Silva AC. Functional MRI of visual responses in the awake, behaving marmoset. *Neuroimage*. 2015b; 120:1–11. [PubMed: 26149609]
- Hunter JV, Wilde EA, Tong KA, Holshouser BA. Emerging imaging tools for use with traumatic brain injury research. *J Neurotrauma*. 2012; 29:654–671. [PubMed: 21787167]
- Huo BX, Gao YR, Drew PJ. Quantitative separation of arterial and venous cerebral blood volume increases during voluntary locomotion. *Neuroimage*. 2015a; 105:369–379. <http://dx.doi.org/10.1016/j.neuroimage.2014.10.030>. [PubMed: 25467301]
- Huo BX, Greene SE, Drew PJ. Venous cerebral blood volume increase during voluntary locomotion reflects cardiovascular changes. *Neuroimage*. 2015b; 118:301–312. <http://dx.doi.org/10.1016/j.neuroimage.2015.06.011>. [PubMed: 26057593]
- Huo BX, Smith JB, Drew PJ. Neurovascular coupling and decoupling in the cortex during voluntary locomotion. *J Neurosci: Off J Soc Neurosci*. 2014; 34:10975–10981. <http://dx.doi.org/10.1523/JNEUROSCI.1369-14.2014>.
- Hyder F, Rothman DL, Shulman RG. Total neuroenergetics support localized brain activity: implications for the interpretation of fMRI. *PNAS*. 2002; 99:10771–10776. <http://dx.doi.org/10.1073/pnas.132272299>. [PubMed: 12134057]
- Illiff JJ, Wang M, Zeppenfeld D, Venkataraman A, Plog B, Liao Y, Deane R, Nedergaard M. Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J Neurosci: Off J Soc Neurosci*. 2013; 33:18190–18199. <http://dx.doi.org/10.1523/JNEUROSCI.1592-13.2013>.
- Kahn I, Knoblich U, Desai M, Bernstein J, Graybiel AM, Boyden ES, Buckner RL, Moore CI. Optogenetic drive of neocortical pyramidal neurons generates fMRI signals that are correlated with spiking activity. *Brain Res*. 2013; 1511:33–45. [PubMed: 23523914]
- Kaisti KK, Långsjö JW, Aalto S, Oikonen V, Sipilä H, Teräs M, Hinkka S, Metsähonkala L, Scheinin H. Effects of sevoflurane, propofol, and adjunct nitrous oxide on regional cerebral blood flow, oxygen consumption, and blood volume in humans. *Anesthesiology*. 2003; 99:603–613. [PubMed: 12960544]
- Kalmbach AS, Waters J. Brain surface temperature under a craniotomy. *J Neurophysiol*. 2012; 108:3138–3146. [PubMed: 22972953]
- Kato HK, Gillet SN, Peters AJ, Isaacson JS, Komiyama T. Parvalbumin-expressing interneurons linearly control olfactory bulb output. *Neuron*. 2013; 80:1218–1231. <http://dx.doi.org/10.1016/j.neuron.2013.08.036>. [PubMed: 24239124]
- Kelly AM, Uddin LQ, Biswal BB, Castellanos FX, Milham MP. Competition between functional brain networks mediates behavioral variability. *Neuroimage*. 2008; 39:527–537. [PubMed: 17919929]

- Kennedy C, Gillin JC, Mendelson W, Suda S, Miyaoka M, Ito M, Nakamura RK, Storch FI, Pettigrew K, Mishkin M, Sokoloff L. Local cerebral glucose utilization in non-rapid eye movement sleep. *Nature*. 1982; 297:325–327. [PubMed: 7078644]
- Kennedy DP, Redcay E, Courchesne E. Failing to deactivate: resting functional abnormalities in autism. *Proc Natl Acad Sci USA*. 2006; 103:8275–8280. [PubMed: 16702548]
- Kim SG, Harel N, Jin T, Kim T, Lee P, Zhao F. Cerebral blood volume MRI with intravascular superparamagnetic iron oxide nanoparticles. *NMR Biomed*. 2013; 26:949–962. <http://dx.doi.org/10.1002/nbm.2885>. [PubMed: 23208650]
- Kim SG, Ogawa S. Biophysical and physiological origins of blood oxygenation level-dependent fMRI signals. *J Cereb Blood Flow Metab*. 2012; 32:1188–1206. <http://dx.doi.org/10.1038/jcbfm.2012.23>. [PubMed: 22395207]
- Kim T, Kim SG. Temporal dynamics and spatial specificity of arterial and venous blood volume changes during visual stimulation: implication for BOLD quantification. *J Cereb Blood Flow Metab*. 2011; 31:1211–1222. <http://dx.doi.org/10.1038/jcbfm.2010.226>. [PubMed: 21179068]
- Kim T, Kim SG. Cortical layer-dependent arterial blood volume changes: improved spatial specificity relative to BOLD fMRI. *NeuroImage*. 2010; 49:1340–1349. <http://dx.doi.org/10.1016/j.neuroimage.2009.09.061>. [PubMed: 19800013]
- King JA, Garelick TS, Brevard ME, Chen W, Messenger TL, Duong TQ, Ferris CF. Procedure for minimizing stress for fMRI studies in conscious rats. *J Neurosci Methods*. 2005; 148:154–160. [PubMed: 15964078]
- Kiyatkin EA, Brown PL, Wise RA. Brain temperature fluctuation: a reflection of functional neural activation. *Eur J Neurosci*. 2002; 16:164–168. [PubMed: 12153543]
- Kleinfeld D, Ahissar E, Diamond ME. Active sensation: insights from the rodent vibrissa sensorimotor system. *Curr Opin Neurobiol*. 2006; 16(4):435–444. [PubMed: 16837190]
- Kleinfeld D, Griesbeck O. From art to engineering? The rise of in vivo mammalian electrophysiology via genetically targeted labeling and nonlinear imaging. *Plos Biol*. 2005; 3:e355. <http://dx.doi.org/10.1371/journal.pbio.0030355>. [PubMed: 16207078]
- Kokita N, Stekiel TA, Yamazaki M, Bosnjak ZJ, Kampine JP, Stekiel WJ. Potassium channel-mediated hyperpolarization of mesenteric vascular smooth muscle by isoflurane. *Anesthesiology*. 1999; 90:779–788. [PubMed: 10078680]
- Komiyama T, Sato TR, O'Connor DH, Zhang YX, Huber D, Hooks BM, Gabitto M, Svoboda K. Learning-related fine-scale specificity imaged in motor cortex circuits of behaving mice. *Nature*. 2010; 464:1182–1186. <http://dx.doi.org/10.1038/nature08897>. [PubMed: 20376005]
- Kono K, Philbin DM, Coggins CH, Moss J, Rosow CE, Schneider RC, Slater EE. Renal function and stress response during halothane or fentanyl anesthesia. *Anesth Analg*. 1981; 60:552–556. [PubMed: 7020486]
- Korogod N, Petersen CC, Knott GW. Ultrastructural analysis of adult mouse neocortex comparing aldehyde perfusion with cryo fixation. *eLife*. 2015; 4:1–17.
- Lahti KM, Ferris CF, Li F, Sotak CH. Comparison of evoked cortical activity in conscious and propofol-anesthetized rats using functional MRI. *Magn Reson*. 1999
- Lahti KM, Ferris CF, Li F, Sotak CH, King JA. Imaging brain activity in conscious animals using functional MRI. *J Neurosci Methods*. 1998; 82:75–83. [PubMed: 10223517]
- Lee H, Mashour GA, Noh GJ, Kim S, Lee U. Reconfiguration of network hub structure after propofol-induced unconsciousness. *Anesthesiology*. 2013; 119:1347–1359. [PubMed: 24013572]
- Lee JH, Durand R, Gradinaru V, Zhang F, Goshen I, Kim DS, Fenno LE, Ramakrishnan C, Deisseroth K. Global and local fMRI signals driven by neurons defined optogenetically by type and wiring. *Nature*. 2010; 465:788–792. [PubMed: 20473285]
- Lee SP, Duong TQ, Yang G, Iadecola C, Kim SG. Relative changes of cerebral arterial and venous blood volumes during increased cerebral blood flow: implications for BOLD fMRI. *Magn Reson Med: Off J Soc Magn Reson Med/Soc Magn Reson Med*. 2001; 45:791–800. <http://dx.doi.org/10.1002/mrm.1107>.
- Lee U, Kim S, Noh GJ, Choi BM, Hwang E, Mashour GA. The directionality and functional organization of frontoparietal connectivity during consciousness and anesthesia in humans. *Conscious Cogn*. 2009a; 18:1069–1078. [PubMed: 19443244]



- Lee U, Mashour GA, Kim S, Noh GJ, Choi BM. Propofol induction reduces the capacity for neural information integration: implications for the mechanism of consciousness and general anesthesia. *Conscious Cogn.* 2009b; 18:56–64. [PubMed: 19054696]
- Lee U, Muller M, Noh GJ, Choi B, Mashour GA. Dissociable network properties of anesthetic state transitions. *Anesthesiology.* 2011; 114:872–881. [PubMed: 21383615]
- Letourneur A, Chen V, Waterman G, Drew PJ. A method for longitudinal, transcranial imaging of blood flow and remodeling of the cerebral vasculature in postnatal mice. *Physiol Rep.* 2014; 2:e12238. (e12238). [PubMed: 25524276]
- Levy DE, Sidtis JJ, Rottenberg DA, Jarden JO, Strother SC, Dhawan V, Ginos JZ, Tramo MJ, Evans AC, Plum F. Differences in cerebral blood flow and glucose utilization in vegetative versus locked-in patients. *Ann Neurol.* 1987; 22:673–682. <http://dx.doi.org/10.1002/ana.410220602>. [PubMed: 3501694]
- Li N, Downey JE, Bar-Shir A, Gilad AA, Walczak P, Kim H, Joel SE, Pekar JJ, Thakor NV, Pelled G. Optogenetic-guided cortical plasticity after nerve injury. *Proc Natl Acad Sci USA.* 2011; 108:8838–8843. [PubMed: 21555573]
- Liang Z, King J, Zhang N. Uncovering intrinsic connective architecture of functional networks in awake rat brain. *J Neurosci.* 2011; 31:3776–3783. [PubMed: 21389232]
- Liang Z, King J, Zhang N. Anticorrelated resting-state functional connectivity in awake rat brain. *NeuroImage.* 2012a; 59:1190–1199. [PubMed: 21864689]
- Liang Z, King J, Zhang N. Intrinsic organization of the anesthetized brain. *J Neurosci.* 2012b; 32:10183–10191. [PubMed: 22836253]
- Liang Z, King J, Zhang N. Neuroplasticity to a single-episode traumatic stress revealed by resting-state fMRI in awake rats. *NeuroImage.* 2014; 103:485–491. [PubMed: 25193500]
- Liang Z, Li T, King J, Zhang N. Mapping thalamocortical networks in rat brain using resting-state functional connectivity. *Neuroimage.* 2013; 83:237–244. [PubMed: 23777756]
- Liang Z, Liu X, Zhang N. Dynamic resting state functional connectivity in awake and anesthetized rodents. *NeuroImage.* 2015a; 104:89–99. [PubMed: 25315787]
- Liang Z, Watson GD, Alloway KD, Lee G, Neuberger T, Zhang N. Mapping the functional network of medial prefrontal cortex by combining optogenetics and fMRI in awake rats. *Neuroimage.* 2015b; 117:114–123. [PubMed: 26002727]
- Lind BL, Brazhe AR, Jessen SB, Tan FCC, Lauritzen MJ. Rapid stimulus-evoked astrocyte Ca<sup>2+</sup> elevations and hemodynamic responses in mouse somatosensory cortex in vivo. *Proc Natl Acad Sci USA.* 2013; 110:E4678–E4687. [PubMed: 24218625]
- Liu JV, Hirano Y, Nascimento GC, Stefanovic B, Leopold DA, Silva AC. fMRI in the awake marmoset: somatosensory-evoked responses, functional connectivity, and comparison with propofol anesthesia. *Neuroimage.* 2013a; 78:186–195. [PubMed: 23571417]
- Liu JV, Hirano Y, Nascimento GC, Stefanovic B, Leopold DA, Silva AC. fMRI in the awake marmoset: somatosensory-evoked responses, functional connectivity, and comparison with propofol anesthesia. *Neuroimage.* 2013b; 78:186–195. [PubMed: 23571417]
- Liu X, Zhu XH, Zhang Y, Chen W. Neural origin of spontaneous hemodynamic fluctuations in rats under burst-suppression anesthesia condition. *Cereb Cortex.* 2011; 21:374–384. [PubMed: 20530220]
- Logothetis NK, Pauls J, Augath M, Trinath T, Oeltermann A. Neurophysiological investigation of the basis of the fMRI signal. *Nature.* 2001; 412:150–157. <http://dx.doi.org/10.1038/35084005>. [PubMed: 11449264]
- Long MA, Fee MS. Using temperature to analyse temporal dynamics in the songbird motor pathway. *Nature.* 2008; 456:189–194. <http://dx.doi.org/10.1038/nature07448>. [PubMed: 19005546]
- Lowe MJ, Phillips MD, Lurito JT, Mattson D, Dziedzic M, Mathews VP. Multiple sclerosis: low-frequency temporal blood oxygen level-dependent fluctuations indicate reduced functional connectivity initial results. *Radiology.* 2002; 224:184–192. [PubMed: 12091681]
- Lu H, Zuo Y, Gu H, Waltz JA, Zhan W, Scholl CA, Rea W, Yang Y, Stein EA. Synchronized delta oscillations correlate with the resting-state functional MRI signal. *Proc Natl Acad Sci USA.* 2007; 104:18265–18269. [PubMed: 17991778]


- Lustig C, Snyder AZ, Bhakta M, O'Brien KC, McAvoy M, Raichle ME, Morris JC, Buckner RL. Functional deactivations: change with age and dementia of the Alzheimer type. *Proc Natl Acad Sci USA*. 2003; 100:14504–14509. [PubMed: 14608034]
- Lyons DG, Parpaleix A, Roche M, Charpak S. Mapping oxygen concentration in the awake mouse brain. *Elife*. 2016; 5
- Ma J, Leung LS. Limbic system participates in mediating the effects of general anesthetics. *Neuropsychopharmacology*. 2006; 31:1177–1192. [PubMed: 16205783]
- Ma J, Shen B, Stewart LS, Herrick IA, Leung LS. The septohippocampal system participates in general anesthesia. *J Neurosci*. 2002; 22:RC200. [PubMed: 11784812]
- Ma Z, Perez P, Ma Z, Liu Y, Hamilton C, Liang Z, Zhang N. Functional atlas of the awake rat brain: a neuroimaging study of rat brain specialization and integration. *Neuroimage*. 2016
- Mandeville JB, Marota JJ, Ayata C, Zaharchuk G, Moskowitz MA, Rosen BR, Weisskoff RM. Evidence of a cerebrovascular postarteriole windkessel with delayed compliance. *J Cereb Blood Flow Metab*. 1999; 19:679–689. <http://dx.doi.org/10.1097/00004647-199906000-00012>. [PubMed: 10366199]
- Martin C, Jones M, Martindale J, Mayhew J. Haemodynamic and neural responses to hypercapnia in the awake rat. *Eur J Neurosci*. 2006a; 24:2601–2610. <http://dx.doi.org/10.1111/j.1460-9568.2006.05135.x>. [PubMed: 17100848]
- Martin C, Martindale J, Berwick J, Mayhew J. Investigating neural-hemodynamic coupling and the hemodynamic response function in the awake rat. *NeuroImage*. 2006b; 32:33–48. <http://dx.doi.org/10.1016/j.neuroimage.2006.02.021>. [PubMed: 16725349]
- Masamoto K, Fukuda M, Vazquez A, Kim SG. Dose-dependent effect of isoflurane on neurovascular coupling in rat cerebral cortex. *Eur J Neurosci*. 2009; 30:242–250. <http://dx.doi.org/10.1111/j.1460-9568.2009.06812.x>. [PubMed: 19659924]
- Masamoto K, Kanno I. Anesthesia and the quantitative evaluation of neurovascular coupling. *J Cereb Blood Flow Metab*. 2012; 32:1233–1247. [PubMed: 22510601]
- Matsuura S, Sakamoto H, Hayashida Y, Kuno M. Efferent discharges of sympathetic and parasympathetic nerve fibers during increased intracranial pressure in anesthetized cats in the absence and presence of pressor response. *Brain Res*. 1984; 305:291–301. [PubMed: 6744068]
- Mayer AR, Mannell MV, Ling J, Gasparovic C, Yeo RA. Functional connectivity in mild traumatic brain injury. *Hum Brain Mapp*. 2011; 32:1825–1835. [PubMed: 21259381]
- Mayrhofer JM, Skreb V, von der Behrens W, Musall S, Weber B, Haiss F. Novel two-alternative forced choice paradigm for bilateral vibrotactile whisker frequency discrimination in head-fixed mice and rats. *J Neurophysiol*. 2013; 109:273–284. <http://dx.doi.org/10.1152/jn.00488.2012>. [PubMed: 23054598]
- McGinley MJ, Vinck M, Reimer J, Batista-Brito R, Zagha E, Cadwell CR, Tolia AS, Cardin JA, McCormick DA. Waking state: rapid variations modulate neural and behavioral responses. *Neuron*. 2015; 87(6):1143–1161. <http://dx.doi.org/10.1016/j.neuron.2015.09.012>. [PubMed: 26402600]
- Mehta MR. Cortico-hippocampal interaction during up-down states and memory consolidation. *Nat Neurosci*. 2007; 10:13–15. <http://dx.doi.org/10.1038/nn0107-13>. [PubMed: 17189946]
- Miller KL, Luh WM, Liu TT, Martinez A, Obata T, Wong EC, Frank LR, Buxton RB. Nonlinear temporal dynamics of the cerebral blood flow response. *Hum Brain Mapp*. 2001; 13:1–12. [PubMed: 11284042]
- Moore JD, Deschênes M, Furuta T, Huber D, Smear MC, Demers M, Kleinfeld D. Hierarchy of orofacial rhythms revealed through whisking and breathing. *Nature*. 2013; 497:205–210. <http://dx.doi.org/10.1038/nature12076>. [PubMed: 23624373]
- Muldoon SM, Hart JL, Bowen KA, Freas W. Attenuation of endothelium-mediated vasodilation by halothane. *Anesthesiology*. 1988; 68:31–37. [PubMed: 3257364]
- Murphy K, Birn RM, Bandettini PA. Resting-state fMRI confounds and cleanup. *NeuroImage*. 2013; 80:349–359. [PubMed: 23571418]
- Murphy TH, Boyd JD, Bolanos F, Vanni MP, Silasi G, Haupt D, LeDue JM. High-throughput automated home-cage mesoscopic functional imaging of mouse cortex. *Nat Commun*. 2016; 7:11611. [PubMed: 27291514]

- Nakao Y, Itoh Y, Kuang TY, Cook M, Jehle J, Sokoloff L. Effects of anesthesia on functional activation of cerebral blood flow and metabolism. *PNAS*. 2001; 98:7593–7598. <http://dx.doi.org/10.1073/pnas.121179898>. [PubMed: 11390971]
- Nallasamy N, Tsao DY. Functional connectivity in the brain: effects of anesthesia. *Neuroscientist*. 2011; 17:94–106. [PubMed: 21343409]
- Ndubuizu O, LaManna JC. Brain tissue oxygen concentration measurements. *Antioxid Redox Signal*. 2007; 9(8):1207–1219. DOI: 10.1089/ars.2007.1634 [PubMed: 17536959]
- Nicholson C. Diffusion and related transport mechanisms in brain tissue. *Rep Prog Phys*. 2001; 64:815–884. <http://dx.doi.org/10.1088/0034-4885/64/7/202>.
- Niell CM, Stryker MP. Modulation of visual responses by behavioral state in mouse visual cortex. *Neuron*. 2010; 65:472–479. [PubMed: 20188652]
- Nimmerjahn A, Mukamel EA, Schnitzer MJ. Motor behavior activates bergmann glial networks. *Neuron*. 2009; 62:400–412. <http://dx.doi.org/10.1016/j.neuron.2009.03.019>. [PubMed: 19447095]
- Niven JE, Laughlin SB. Energy limitation as a selective pressure on the evolution of sensory systems. *J Exp Biol*. 2008; 211:1792–1804. <http://dx.doi.org/10.1242/jeb.017574>. [PubMed: 18490395]
- Nizar K, Uhlirova H, Tian P, Saisan PA, Cheng Q, Reznichenko L, Weldy KL, Steed TC, Sridhar VB, MacDonald CL, Cui J, Gratiy SL, Sakadzi S, Boas DA, Beka TI, Einevoll GT, Chen J, Masliah E, Dale AM, Silva GA, Devor A. In vivo stimulus-induced vasodilation occurs without IP3 receptor activation and may precede astrocytic calcium increase. *J Neurosci: Off J Soc Neurosci*. 2013; 33:8411–8422. <http://dx.doi.org/10.1523/JNEUROSCI.3285-12.2013>.
- O'Connor DH, Clack NG, Huber D, Komiyama T, Myers EW, Svoboda K. Vibrissa-based object localization in head-fixed mice. *J Neurosci*. 2010; 30(5):1947–1967. [PubMed: 20130203]
- Ogura K, Takayasu M, Dacey RG. Effects of hypothermia and hyperthermia on the reactivity of rat intracerebral arterioles in vitro. *J Neurosurg*. 1991; 75:433–439. <http://dx.doi.org/10.3171/jns.1991.75.3.0433>. [PubMed: 1869945]
- Osol G, Halpern W. Spontaneous vasomotion in pressurized cerebral arteries from genetically hypertensive rats. *Am J Physiol*. 1988; 254:H28–33. [PubMed: 3337256]
- Otsu Y, Couchman K, Lyons DG, Collot M, Agarwal A, Mallet JM, Pfrieger FW, Bergles DE, Charpak S. Calcium dynamics in astrocyte processes during neurovascular coupling. *Nat Neurosci*. 2014; 1–11. [PubMed: 24369367]
- Patriat R, Molloy EK, Meier TB, Kirk GR, Nair VA, Meyerand ME, Prabhakaran V, Birn RM. The effect of resting condition on resting-state fMRI reliability and consistency: a comparison between resting with eyes open, closed, and fixated. *Neuroimage*. 2013; 78:463–473. <http://dx.doi.org/10.1016/j.neuroimage.2013.04.013>. [PubMed: 23597935]
- Park KW, Dai HB, Lowenstein E, Sellke FW. Propofol-associated dilation of rat distal coronary arteries is mediated by multiple substances, including endothelium-derived nitric oxide. *Anesthesia Analg*. 1995; 81:1191.
- Parry TJ, McElligott JG. A method for restraining awake rats using head immobilization. *Physiol Behav*. 1993; 53:1011–1015. [PubMed: 8099749]
- Peeters RR, Tindemans I, De Schutter E, Van der Linden A. Comparing BOLD fMRI signal changes in the awake and anesthetized rat during electrical forepaw stimulation. *Magn Reson Imaging*. 2001; 19:821–826. [http://dx.doi.org/10.1016/S0730-725X\(01\)00391-5](http://dx.doi.org/10.1016/S0730-725X(01)00391-5). [PubMed: 11551722]
- Peltier SJ, Kerssens C, Hamann SB, Sebel PS, Byas-Smith M, Hu X. Functional connectivity changes with concentration of sevoflurane anesthesia. *Neuroreport*. 2005; 16:285–288. [PubMed: 15706237]
- Pisauro MA, Dhruv NT, Carandini M, Benucci A. Fast hemodynamic responses in the visual cortex of the awake mouse. *J Neurosci: Off J Soc Neurosci*. 2013; 33:18343–18351. <http://dx.doi.org/10.1523/JNEUROSCI.2130-13.2013>.
- Podgorski K, Ranganathan GN. Brain heating induced by near infrared lasers during multi-photon microscopy. *J Neurophysiol*. 2016:00275, 02016.
- Rangaraju V, Calloway N, Ryan TA. Activity-driven local ATP synthesis is required for synaptic function. *Cell*. 2014; 156:825–835. <http://dx.doi.org/10.1016/j.cell.2013.12.042>. [PubMed: 24529383]


- Raz A, Grady SM, Krause BM. Preferential effect of isoflurane on top-down vs. bottom-up pathways in sensory cortex. *Front Syst.* 2014
- Reivich M, Kuhl D, Wolf A, Greenberg J, Phelps M, Ido T, Casella V, Fowler J, Hoffman E, Alavi A, Som P, Sokoloff L. The [<sup>18</sup>F]fluorodeoxyglucose method for the measurement of local cerebral glucose utilization in man. *Circ Res.* 1979; 44:127–137. [PubMed: 363301]
- Richlan F, Gagl B, Hawelka S, Braun M, Schurz M, Kronbichler M, Hutzler F. Fixation-related fMRI analysis in the domain of reading research: using self-paced eye movements as markers for hemodynamic brain responses during visual letter string processing. *Cereb Cortex.* 2014; 24(10): 2647–2656. <http://dx.doi.org/10.1093/cercor/bht117>. [PubMed: 23645718]
- Ries CR, Puil E. Mechanism of anesthesia revealed by shunting actions of isoflurane on thalamocortical neurons. *J Neurophysiol.* 1999; 81:1795–1801. [PubMed: 10200213]
- Rinberg D, Koulakov A, Gelperin A. Sparse odor coding in awake behaving mice. *J Neurosci.* 2006; 26:8857–8865. <http://dx.doi.org/10.1523/JNEUROSCI.0884-06.2006>. [PubMed: 16928875]
- Rosenegger DG, Tran CH, Wamsteeker Cusulin JL, Gordon GR. Tonic Local Brain Blood Flow Control by Astrocytes Independent of Phasic Neurovascular Coupling. *J Neurosci.* 2015; 35:13463–13474. [PubMed: 26424891]
- Rothman DL, De Feyter HM, de Graaf RA, Mason GF, Behar KL. <sup>13</sup>C MRS studies of neuroenergetics and neurotransmitter cycling in humans. *NMR Biomed.* 2011; 24:943–957. [PubMed: 21882281]
- Sadoshima S, Thames M, Heistad D. Cerebral blood flow during elevation of intracranial pressure: role of sympathetic nerves. *Am J Physiol.* 1981; 241:H78–H84. [PubMed: 7246792]
- Sakadzic S, Mandeville ET, Gagnon L, Musacchia JJ, Yaseen MA, Yucel MA, Lefebvre J, Lesage F, Dale AM, Eikermann-Haerter K, Ayata C, Srinivasan VJ, Lo EH, Devor A, Boas DA. Large arteriolar component of oxygen delivery implies a safe margin of oxygen supply to cerebral tissue. *Nat Commun.* 2014; 5:5734. [PubMed: 25483924]
- Sakadzic S, Roussakis E, Yaseen MA, Mandeville ET, Srinivasan VJ, Arai K, Ruvinskaya S, Devor A, Lo EH, Vinogradov SA, Boas DA. Two-photon high-resolution measurement of partial pressure of oxygen in cerebral vasculature and tissue. *Nat Methods.* 2010; 7:755–759. [PubMed: 20693997]
- Schmidt MF, Konishi M. Gating of auditory responses in the vocal control system of awake songbirds. *Nat Neurosci.* 1998; 1:513–518. <http://dx.doi.org/10.1038/2232>. [PubMed: 10196550]
- Schroeder MP, Weiss C, Proccisi D, Disterhoft JF, Wang L. Intrinsic connectivity of neural networks in the awake rabbit. *Neuroimage.* 2016; 129:260–267. [PubMed: 26774609]
- Schroeter A, Schlegel F, Seuwen A, Grandjean J, Rudin M. Specificity of stimulus-evoked fMRI responses in the mouse: the influence of systemic physiological changes associated with innocuous stimulation under four different anesthetics. *NeuroImage.* 2014; 94:372–384. [PubMed: 24495809]
- Scott BB, Brody CD, Tank DW. Cellular resolution functional imaging in behaving rats using voluntary head restraint. *Neuron.* 2013; 80:371–384. <http://dx.doi.org/10.1016/j.neuron.2013.08.002>. [PubMed: 24055015]
- Scott BB, Constantinople CM, Erlich JC, Tank DW, Brody CD. Sources of noise during accumulation of evidence in unrestrained and voluntarily head-restrained rats. *eLife.* 2015; 4:e11308. [PubMed: 26673896]
- Shtoyerman E, Arieli A, Slovlin H, Vanzetta I, Grinvald A. Long-term optical imaging and spectroscopy reveal mechanisms underlying the intrinsic signal and stability of cortical maps in V1 of behaving monkeys. *J Neurosci.* 2000; 20:8111–8121. [PubMed: 11050133]
- Nagao, Seigo, Roccaforte, Peter, Moody, Robert A. Acute intracranial hypertension and auditory brain-stem responses. *J Neurosurg.* 2009; 51:669–676. <http://dx.doi.org/10.3171/jns.1979.51.5.0669><http://dx.doi.org/10.3171/jns.1979.51.5.0669>.
- Sellers KK, Bennett DV, Hutt A, Fröhlich F. Anesthesia differentially modulates spontaneous network dynamics by cortical area and layer. *J Neurophysiol.* 2013; 110:2739–2751. <http://dx.doi.org/10.1152/jn.00404.2013>. [PubMed: 24047911]
- Sellers KK, Bennett DV, Hutt A, Williams JH, Fröhlich F. Awake vs anesthetized: layer-specific sensory processing in visual cortex and functional connectivity between cortical areas. *J*

- Neurophysiol. 2015; 113:3798–3815. <http://dx.doi.org/10.1152/jn.00923.2014>. [PubMed: 25833839]
- Shirey MJ, Smith JB, Kudlik DE, Huo BX, Greene SE, Drew PJ. Brief anesthesia, but not voluntary locomotion, significantly alters cortical temperature. *J Neurophysiol.* 2015; 114:3009–3022. <http://dx.doi.org/10.1152/jn.00046.2015>.
- Shu Y, Hasenstaub A, McCormick DA. Turning on and off recurrent balanced cortical activity. *Nature.* 2003; 423:288–293. <http://dx.doi.org/10.1038/nature01616>. [PubMed: 12748642]
- Shulman RG, Rothman DL, Hyder F. Stimulated changes in localized cerebral energy consumption under anesthesia. *PNAS.* 1999; 96:3245–3250. <http://dx.doi.org/10.1073/pnas.96.6.3245>. [PubMed: 10077669]
- Siesjö, BK. *Brain Energy Metabolism.* John Wiley & Sons; New York: 1978.
- Silva AC, Koretsky AP, Duyn JH. Functional MRI impulse response for BOLD and CBV contrast in rat somatosensory cortex. *Magnetic resonance in medicine: official journal of the Society of Magnetic Resonance in Med/Soc Magn Reson Med.* 2007a; 57:1110–1118. <http://dx.doi.org/10.1002/mrm.21246>.
- Simard M, Arcuino G, Takano T, Liu QS, Nedergaard M. Signaling at the gliovascular interface. *J Neurosci.* 2003; 23:9254–9262. [PubMed: 14534260]
- Smear M, Shusterman R, O'Connor R, Bozza T, Rinberg D. Perception of sniff phase in mouse olfaction. *Nature.* 2011; 479:397–400. <http://dx.doi.org/10.1038/nature10521>. [PubMed: 21993623]
- Sofroniew NJ, Cohen JD, Lee AK, Svoboda K. Natural whisker-guided behavior by head-fixed mice in tactile virtual reality. *J Neurosci.* 2014; 34:9537–9550. [PubMed: 25031397]
- Sokoloff L, Reivich M, Kennedy C, Rosiers Des, Patlak MH, Pettigrew CS, Sakurada KD, Shinohara OM. The [<sup>14</sup>C]deoxyglucose method for the measurement of local cerebral glucose utilization: theory, procedure, and normal values in the conscious and anesthetized albino rat. *J Neurochem.* 1977; 28:897–916. [PubMed: 864466]
- Somjen GG. Ion regulation in the brain: implications for pathophysiology. *Neuroscientist.* 2002; 8:254–267. [PubMed: 12061505]
- Stefanovic B, Hutchinson E, Yakovleva V, Schram V, Russell JT, Belluscio L, Koretsky AP, Silva AC. Functional reactivity of cerebral capillaries. *J Cereb Blood Flow Metab.* 2008; 28:961–972. <http://dx.doi.org/10.1038/sj.jcbfm.9600590>. [PubMed: 18059431]
- Steriade M, Timofeev I, Grenier F. Natural waking and sleep states: a view from inside neocortical neurons. *J Neurophysiol.* 2001; 85:1969–1985. [http://dx.doi.org/10.1016/0306-4522\(89\)90423-5](http://dx.doi.org/10.1016/0306-4522(89)90423-5). [PubMed: 11353014]
- Takano T, Tian GF, Peng W, Lou N, Libionka W, Han X, Nedergaard M. Astrocyte-mediated control of cerebral blood flow. *Nat Neurosci.* 2006; 9:260–267. <http://dx.doi.org/10.1038/nn1623>. [PubMed: 16388306]
- Thrane AS, Thrane VR, Zeppenfeld D, Lou N, Xu Q, Nagelhus EA, Nedergaard M. General anesthesia selectively disrupts astrocyte calcium signaling in the awake mouse cortex. *PNAS.* 2012; 109:18974–18979. <http://dx.doi.org/10.1073/pnas.1209448109>. [PubMed: 23112168]
- Tian L, Jiang T, Wang Y, Zang Y, He Y, Liang M, Sui M, Cao Q, Hu S, Peng M, Zhuo Y. Altered resting-state functional connectivity patterns of anterior cingulate cortex in adolescents with attention deficit hyperactivity disorder. *Neurosci Lett.* 2006; 400:39–43. [PubMed: 16510242]
- Tomasi D, Wang RL, Telang F, Boronikolas V, Jayne MC, Wang GJ, Fowler JS, Volkow ND. Impairment of attentional networks after 1 night of sleep deprivation. *Cereb Cortex.* 2009; 19:233–240. [PubMed: 18483003]
- Tononi G, Edelman GM, Sporns O. Complexity and coherency: integrating information in the brain. *Trends Cogn Sci.* 1998; 2:474–484. [PubMed: 21227298]
- Tran CH, Gordon GR. Acute two-photon imaging of the neurovascular unit in the cortex of active mice. *Front Cell Neurosci.* 2015; 9:11. [PubMed: 25698926]
- Ueki M, Mies G, Hossmann KA. Effect of alpha-chloralose, halothane, pentobarbital and nitrous oxide anesthesia on metabolic coupling in somatosensory cortex of rat. *Acta Anaesthesiol Scand.* 1992; 36:318–322. [PubMed: 1595336]

- Vaiceliunaite A, Eriskien S, Franzen F, Katzner S, Busse L. Spatial integration in mouse primary visual cortex. *J Neurophysiol.* 2013; 110:964–972. <http://dx.doi.org/10.1152/jn.00138.2013>. [PubMed: 23719206]
- van Meer MP, Otte WM, van der Marel K, Nijboer CH, Kavelaars A, van der Sprekel JW, Viergever MA, Dijkhuizen RM. Extent of bilateral neuronal network reorganization and functional recovery in relation to stroke severity. *J Neurosci.* 2012; 32:4495–4507. [PubMed: 22457497]
- Velly LJ, Rey MF, Bruder NJ, Gouvitsos FA, Witjas T, Regis JM, Peragut JC, Gouin FM. Differential dynamic of action on cortical and subcortical structures of anesthetic agents during induction of anesthesia. *Anesthesiology.* 2007; 107:202–212. [PubMed: 17667563]
- Vincent JL, Patel GH, Fox MD, Snyder AZ, Baker JT, Van Essen DC, Zempel JM, Snyder LH, Corbetta M, Raichle ME. Intrinsic functional architecture in the anaesthetized monkey brain. *Nature.* 2007; 447:83–86. [PubMed: 17476267]
- Wang C, Ong JL, Patanaik A, Zhou J, Chee MW. Spontaneous eyelid closures link vigilance fluctuation with fMRI dynamic connectivity states. *Proc Natl Acad Sci USA.* 2016; 113(34): 9653–9658. <http://dx.doi.org/10.1073/pnas.1523980113>. [PubMed: 27512040]
- Weitz AJ, Fang Z, Lee HJ, Fisher RS, Smith WC, Choy M, Liu J, Lin P, Rosenberg M, Lee JH. Optogenetic fMRI reveals distinct, frequency-dependent networks recruited by dorsal and intermediate hippocampus stimulations. *Neuroimage.* 2015; 107:229–241. [PubMed: 25462689]
- White NS, Alkire MT. Impaired thalamocortical connectivity in humans during general-anesthetic-induced unconsciousness. *Neuroimage.* 2003; 19:402–411. [PubMed: 12814589]
- Whitfield-Gabrieli S, Thermenos HW, Milanovic S, Tsuang MT, Faraone SV, McCarley RW, Shenton ME, Green AI, Nieto-Castanon A, LaViolette P, Wojcik J, Gabrieli JD, Seidman LJ. Hyperactivity and hyperconnectivity of the default network in schizophrenia and in first-degree relatives of persons with schizophrenia. *Proc Natl Acad Sci USA.* 2009; 106:1279–1284. [PubMed: 19164577]
- Wilson RS, Mayhew SD, Rollings DT, Goldstone A, Przewdzik I, Arvanitis TN, Bagshaw AP. Influence of epoch length on measurement of dynamic functional connectivity in wakefulness and behavioural validation in sleep. *NeuroImage.* 2015; 112:169–179. [PubMed: 25765256]
- Wurtz RH. Visual receptive fields of striate cortex neurons in awake monkeys. *J Neurophysiol.* 1969
- Xie L, Kang H, Xu Q, Chen MJ, Liao Y, Thiyagarajan M, O'Donnell J, Christensen DJ, Nicholson C, Iliff JJ, Takano T, Deane R, Nedergaard M. Sleep drives metabolite clearance from the adult brain. *Science.* 2013; 342:373–377. <http://dx.doi.org/10.1126/science.1241224>. [PubMed: 24136970]
- Yang H, Kwon SE, Severson KS, O'Connor DH. Origins of choice-related activity in mouse somatosensory cortex. *Nat Neurosci.* 2016; 19(1):127–134. <http://dx.doi.org/10.1038/nn.4183>. [PubMed: 26642088]
- Zagha E, Casale AE, Sachdev RNS, McGinley MJ, McCormick Da. Motor cortex feedback influences sensory processing by modulating network state. *Neuron.* 2013; 79:567–578. [PubMed: 23850595]
- Zhang N, Liu Z, He B, Chen W. Noninvasive study of neurovascular coupling during graded neuronal suppression. *J Cereb Blood Flow Metab.* 2008a; 28:280–290. [PubMed: 17700632]
- Zhang N, Rane P, Huang W, Liang Z, Kennedy D, Frazier JA, King J. Mapping resting-state brain networks in conscious animals. *J Neurosci Methods.* 2010; 189:186–196. [PubMed: 20382183]
- Zhang N, Yacoub E, Zhu XH, Ugurbil K, Chen W. Linearity of blood-oxygenation-level dependent signal at microvasculature. *Neuroimage.* 2009; 48:313–318. [PubMed: 19580875]
- Zhang N, Zhu XH, Chen W. Investigating the source of BOLD nonlinearity in human visual cortex in response to paired visual stimuli. *Neuroimage.* 2008b; 43:204–212. [PubMed: 18657623]




**Fig. 1.** Effects of anesthetics on hemodynamic responses. Schematic showing how a stimulus-generated changes in neural activity are converted into a detectable BOLD signal. Physiological processes affected by anesthesia are outlined in purple.




**Fig. 2.** Schematic showing net effects of anesthesia on (BOLD) hemodynamic response function. Anesthesia slows and attenuates the HRF. The awake hemodynamic response is approximately twice as large and has a faster onset compared to the HRF in anesthetized animals (Goense and Logothetis, 2008; PISAURO et al., 2013; AKSENOV et al., 2015).






**Fig. 3.** Comparison of dilation of individual arteries and veins measured using two-photon microscopy in the somatosensory cortex of awake and anesthetized mice. The sensory stimulus is puffs of air aimed at the vibrissae. Urethane drastically decreases the arterial dilation, and abolishes the venous dilation. Adapted from Drew et al. (2011).




**Fig. 4.**


Brain metabolic rate is significantly decreased during anesthesia and sleep. Summary plot of measured brain metabolic rates under different anesthetics and conditions. References, species and measurement technique are as follows: REM sleep: Buchsbaum et al. (1989), Human, 18-fluorodeoxyglucose PET. Non-REM sleep: Buchsbaum et al. (1989), Human, 18-fluorodeoxyglucose PET, Kennedy et al. (1982), Rhesus monkey, 2-deoxyglucose. Halothane: Alkire et al. (1999), Human, 18-fluorodeoxyglucose PET. Propofol: Alkire et al. (1995), Human, 18-fluorodeoxyglucose PET. Propofol: Alkire et al. (1995), Human, 18-fluorodeoxyglucose PET. Alpha-chloralose: Dudley et al. (1982), Rat, 2-deoxyglucose. Isoflurane: Alkire et al. (1997), Humans, 18-fluorodeoxyglucose PET. Pentobarbital: Crane et al. (1978), Rat, 2-deoxyglucose. Vegetative coma: Levy et al. (1987), Human, 18-fluorodeoxyglucose PET. Note that these are changes in brain metabolism measured over the entire brain. Anesthesia-induced decreases in metabolic rate in gray matter will be larger than that in white matter because the metabolic rate of white matter is not appreciably affected by neural activity (Harris and Attwell, 2012).




**Fig. 5.** Infralimbic cortex (IL) resting-state functional connectivity maps in awake and anesthetized rats. (a) The IL RSFC map in the awake rat. (b) The IL RSFC map in the anesthetized rat. Anticorrelated RSFC between IL and amygdala was completely abolished by anesthesia. All maps were overlaid on anatomical images. Distances to Bregma (mm) are labeled for each slice. Adapted from (Liang et al., 2012a).




**Fig. 6.** Optogenetic activation of brain networks in the awake and anesthetized animal. Averaged activation maps and time courses in response to infralimbic (IL) optogenetic stimulation in awake (left) and isoflurane-anesthetized (right) states. (a) The averaged activation map in the awake state. (b) The averaged activation map in the anesthetized state. Green arrow, implanted optical fiber. Distance to Bregma is labeled in each slice. (c) BOLD time courses of anterior cingulate cortex in the awake state. (d) BOLD time courses of anterior cingulate cortex in the anesthetized state. Note the difference in the vertical scale. Red lines indicate the stimulation periods. Blue shades shows SEM. AC, anterior cingulate cortex; AGm, medial agranular (frontal) cortex; Aid, dorsal agranular insular cortex; AON, anterior olfactory nucleus; BST, Bed nucleus of stria terminalis; CL, claustrum; EN, endopiriform nucleus; CP, caudate-putamen; Hb, habenula; IL, infralimbic; IP, interpeduncular nucleus; LHyp, lateral hypothalamus; LS, lateral septum; MD, mediodorsal nucleus; MS, medial septum; NAcc, nucleus accumbens; OC, orbital cortex; PMd, dorsal preammillary nucleus; PrL, prelimbic; SNr, substantia nigra; SUB, subiculum; RSC, retrosplenial cortex. Adapted from (Liang et al., 2015b).



**Fig. 7.** Anesthesia disrupts large-scale brain organization. Community structures in the awake (a) and isoflurane-anesthetized (b) states. Brain regions with the same color are within the same module. Brain regions without the annotation of L or R imply bilateral sides. L: left, R: right, A: anterior, P: posterior, V: ventral, D: dorsal. Adapted from (Liang et al., 2012b).




**Fig. 8.** Imaging hemodynamic signals in awake mice. (A) Mouse head-fixed in a plexiglass cylinder. The whiskers are illuminated from below to allow tracking of movements. (B) Mouse head-fixed on a spherical treadmill with one degree of rotational freedom. The mouse is free to run or stand still. An electrical connector allows measurement of simultaneous ECoG or LFP from implanted electrodes. (C) Head fixed mice exhibit active sensing (whisking) behavior. Top, camera images taken of whiskers of a head-fixed mouse at various time points and protraction angles. Bottom, whisker angle versus time tracked using the Radon transform (Drew et al., 2010b; Gao and Drew, 2014). (D) The entire dorsal cortex can be imaged using optical imaging. Photo of a large, thinned skull window showing locations of forelimb/hindlimb representations (green, FL/HL), frontal cortex (red), visual cortex (purple, V1), and whisker representation (dark blue) reconstructed from the cytochrome oxidase stained cortex. Red dot denotes the location of bregma. Rostral is up, medial to the left. Scale bar 1 mm.



**Fig. 9.**


Cerebral arterial and venous responses during voluntary locomotion in head-fixed awake mice measured with two-photon microscopy. (A) Top left, photograph of the cortical vasculature taken through a polished and reinforced thinned skull (PoRTS) window. The window was implanted over the parietal cortex, which includes the hindlimb region (HL) and vibrissa cortex (Vi). The black box denotes the region where the two-photon images were obtained. Top right, schematic drawing of the major surface vasculatures on the PoRTS window in A. Arterioles are in red, while venules are in blue. Bottom, one second averaged two-photon images of an arteriole (in red box) and a venule (in blue box) enclosed in the black box in A during resting (right, marked by purple asterisk) and locomoting (left, marked by magenta asterisk). The arteriole dilates strongly during locomotion. (B) The responses of the arteriole (top) and venule (bottom) shown in C during voluntary locomotion. Black tick marks on top are the binarized locomotion events. Locomoting and resting periods where the two-photon images shown in A are marked by magenta and purple asterisks, respectively. Arteriole dilation is rapid and large, while venule diameter changes are smaller and slower. The dilations of both vessels can be captured by convolving the binarized locomotion events with an exponential decay (Gao and Drew, 2016; Huo et al., 2015a, 2015b). The fitted responses are shown in light red and light blue for arteriole and venule respectively. Insets are the fitted exponential kernels.




**Fig. 10.**

Intrinsic optical signal imaging of hemodynamic signals from the cortex during locomotion. (A) Location of the frontal, forelimb/hindlimb whisker and visual cortex regions of interest (ROI). All but the frontal area are determined from cytochrome oxidase staining. Scale bar denotes 1 mm. (B) Changes in reflectance obtained at 530 nm over the entire cranial window at different time points during the imaging session. Decreases in reflectance correspond to increases in blood volume, caused by vessel dilation. (C) Top, change in reflectance in each of the four ROIs versus time. The black ticks show locomotion events. There is a large decrease in reflectance, corresponding to vasodilation, in all ROIs but the frontal cortex during locomotion. Note the spontaneous fluctuations in blood volume at rest. Middle, spectrogram of the intrinsic optical signal over the entire hemisphere showing heart-rate fluctuations. The heart-rate is visible as a peak in the spectrogram, and increases during the locomotion bout. The blue line shows the peak of the heart-rate. Bottom, spectrogram of the electrocortigram (ECoG) versus time. Locomotion is accompanied by increases in high frequency (gamma-band) power. Gamma band power, changes versus resting, in dB, are plotted in green. (D) Locomotion-triggered average from the entire 30 min trial. Note the strong responses in the FL/HL, whisker and visual cortex, but lack of response in frontal cortex. (E) The arterial and venous components of the blood volume response to locomotion in each pixel can be separated using a linear fitting procedure into an arterial component and a venous component (Huo et al., 2015a). The amplitude of the arterial component for each pixel is plotted at the top (in red), and the venous amplitude (in blue) for each pixel is plotted versus the bottom. The brighter the color, the larger the amplitude of the fit to the pixel. The venous dilation is widely spread across the cortex, while the arterial response is more localized. Scale bar is 1 mm.






**Fig. 11.** Procedure for imaging an awake rat using MRI. Schematic illustration of the animal setup and imaging procedures.



**Fig. 12.**

Stress induced by imaging is minimal. The elevated-plus maze (EPM) score (open/open +closed arm) was separately evaluated in three groups of rats: rats without any manipulation (n=8), rats imaged four times and rats exposed to a single episode of predator odor (i.e. traumatized) but did not undergo any acclimation and imaging procedures (n=8). EPM in the last two groups was measured 7 days after either the trauma exposure or the last imaging. Two sample t-tests showed that there was no significant difference between the no manipulation group and four-time imaging group ( $p=0.8$ ), while the predator odor exposed group showed a significantly higher stress level relative to the imaging group ( $p=0.03$ ). These results suggest that predator odor significantly increased the stress level while multiple imaging acquisitions had no significant effect. Bars show SEM.



**Fig. 13.** Functional networks and hemodynamic responses are reproducible and stable over time. (A) BOLD activation maps in response to optogenetic stimulation of IL in awake rats on two separate days. Left panel: BOLD activation map acquired on one day; middle panel: BOLD activation map acquired in the same group of awake animals on another day; right panel: correlation of voxel-wise BOLD amplitude between the two days. Adapted from (Liang et al., 2015). (B) IL RSFC maps in two separate subgroups of animals. Left panel: IL RSFC map in subgroup 1; middle panel: IL RSFC map in subgroup 2; right panel: correlation of voxel-wise RSFC strength between the two subgroups. Adapted from (Liang et al., 2011). (C) Top, the correlation coefficient for each pixel in the window between the predicted hemodynamic response to locomotion (using a fitted hemodynamic response function) during one 30-min IOS imaging session. The blue rectangles denote histologically-identified forelimb/hindlimb representations. The fit has a high correlation in these regions, and not in

the frontal cortex because there is no hemodynamic response in the frontal cortex to locomotion. Bottom shows the correlation between the predicted hemodynamic response and actual response plotted for each pixel for a different 30-min imaging session, using the fits same in the top panel, demonstrating a high reproducibility of the response. (D) Matrix of cross-correlations between the predicted and actual hemodynamic response to voluntary locomotion during a 30-min trial for the histologically defined SI FL/HL area. Fits from the trials numbered in the rows were used to predicted data from the trials in the columns. Imaging sessions were done on seven different days with 17 days between the first and last day, showing that the hemodynamic responses are stable. C and D adapted from (Huo et al., 2015a).