

xCyp26c* Induced by Inhibition of BMP Signaling Is Involved in Anterior-Posterior Neural Patterning of *Xenopus laevis

Saet-Byeol Yu¹, Zobia Umair¹, Shiv Kumar¹, Unjoo Lee², Seung-Hwan Lee¹, Jong-Il Kim³, SungChan Kim¹, Jae-Bong Park¹, Jae-Yong Lee¹, and Jaebong Kim^{1,*}

Vertebrate neurogenesis requires inhibition of endogenous bone morphogenetic protein (BMP) signals in the ectoderm. Blocking of BMPs in animal cap explants causes the formation of anterior neural tissues as a default fate. To identify genes involved in the anterior neural specification, we analyzed gene expression profiles using a *Xenopus* Affymetrix Gene Chip after BMP-4 inhibition in animal cap explants. We found that the *xCyp26c* gene, encoding a retinoic acid (RA) degradation enzyme, was upregulated following inhibition of BMP signaling in early neuroectodermal cells. Whole-mount *in situ* hybridization analysis showed that *xCyp26c* expression started in the anterior region during the early neurula stage. Overexpression of *xCyp26c* weakly induced neural genes in animal cap explants. *xCyp26c* abolished the expression of all trans-*l*-cis-RA-induced posterior genes, but not basic FGF-induced posterior genes. Depletion of *xCyp26c* by morpholino-oligonucleotides suppressed the normal formation of the axis and head, indicating that *xCyp26c* plays a critical role in the specification of anterior neural tissue in whole embryos. In animal cap explants, however, *xCyp26c* morpholinos did not alter anterior-to-posterior neural tissue formation. Together, these results suggest that *xCyp26c* plays a specific role in anterior-posterior (A-P) neural patterning of *Xenopus* embryos.

INTRODUCTION

Induction of three germ layers (ectoderm, mesoderm, and endoderm) and axis formation including the dorsoventral and anteroposterior posterior regions are major events during ver-

tebrate embryo development. Signals from bone morphogenetic protein-4 (BMP-4), fibroblast growth factor (FGF), and Wnt play crucial roles during early embryo development and early patterning (Harland and Gerhart, 1997; Hemmati-Brivanlou and Thomsen, 1995; Kessler and Melton, 1994; Lee et al., 2011a; Schmidt et al., 1995).

In *Xenopus*, BMP-4 is expressed in the ventral mesoderm and ectoderm and acts as a ventral morphogen in a direct and long-range fashion to establish its gradient and specific cell fate (Dosch et al., 1997; Xu et al., 1999). When the BMP-4 gradient is established, responding cells activate different genes at distinct threshold levels, resulting in dorsoventral patterning of the mesoderm as well as ectoderm during early *Xenopus* development (Jones and Smith, 1998; Knecht and Harland, 1997). If BMP-4 mRNA is injected into *Xenopus* embryos, the dorsal mesoderm is converted into the ventral mesoderm (Dale et al., 1992; Hwang et al., 2002; 2003; Jones et al., 1992). On the other hand, over-expression of a dominant negative BMP-4 receptor (DNBR) into ventral territories results in the formation of a secondary body axis (Graff et al., 1994; Suzuki et al., 1994). BMP-4 is a strong candidate as an epidermal, ventral mesodermal inducer as well as a neural inhibitor (Xu et al., 1995). Previously, Wawersik et al. (2005) reported that *in-vivo* BMP inhibition triggered at stage 5 not only expands the neural plate but also represses the neural crest marker.

The development of the central nervous system (CNS) begins when the ectoderm gives rise to the neural plate. Early neurogenesis is initiated by inhibition of BMP signaling in the ectoderm by BMP antagonists expressed in Spemann's organizer region, resulting in anterior neural tissue formation (Hemmati-Brivanlou and Melton, 1994). According to a generally accepted mechanism, inhibition of BMP signaling generates anterior neural tissue as a default pathway (Nieuwkoop, 1952). Wnts, FGF and retinoic acids (RA) act as modifiers of anterior to posterior neural specification (Blumberg et al., 1997; Kolm et al., 1997; Ruiz i Altaba and Jessell, 1991; Xu et al., 1997). However, it is unclear whether anterior neural tissue produced by inhibition of BMP is simply generated by a default pathway of pre-existing intracellular molecules or by the newly expressed posterior modifier inhibition molecules.

The intracellular level of active RA is determined by the balance between RA synthesis by retinaldehyde dehydrogenases (RALDHs) and its degradation by *Cyp26* enzymes, the latter of which constitute a group of P450 enzymes that metabolize RA

¹Department of Biochemistry, Institute of Cell Differentiation and Aging, College of Medicine, Hallym University, Kangwon 200-702, Korea, ²Department of Electrical Engineering, Hallym University, Kangwon200-702, Korea, ³Department of Biochemistry and Molecular Biology, Seoul National University College of Medicine, Seoul 110-799, Korea
*Correspondence: jbkim@hallym.ac.kr

Received 6 January, 2016; revised 15 January, 2016; accepted 18 January, 2016; published online 29 February, 2016

Keywords: A-P neural formation, BMP-4, *xCyp26c*, *Xenopus laevis*

eISSN: 0219-1032

© The Korean Society for Molecular and Cellular Biology. All rights reserved.

© This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>.

to its inactive forms (Fujii et al., 1997; Ray et al., 1997; White et al., 1996). *Cyp26* enzymes are thought to play a central role in the appropriate regulation of the RA signal as a posteriorizing factor in CNS development (Abu-Abed et al., 2001; Sakai et al., 2001; Sirbu et al., 2005). Mice and humans possess three *Cyp26* genes: *Cyp26a1*, *Cyp26b1* and *Cyp26c1* (MacLean et al., 2001; Nebert and Russell, 2002; Tahayato et al., 2003). *Cyp26* genes have been previously characterized in various species, but their function in *Xenopus leavis* have not yet been fully identified. The role of *Cyp26* during RA utilization and the anteriorization of the neuroectoderm during the early embryonic stages of *Xenopus* is also unclear.

Wnt and FGF are also known as anterior to posterior modifiers during early neurogenesis. *Cyp26* is regulated by both Wnt and FGF signaling (Kudoh et al., 2002; Lee et al., 2011b). Wnt is reportedly responsible for (A-P) neural patterning, as an injection of a truncated form of BMP receptor into the ventral side of embryos results in the formation of a secondary axis without a head. Inhibition of XWnt8 with dn-XWnt8 induces the formation of a complete axis with a head, indicating that inhibition of Wnt signaling is important for head formation (Glinka et al., 1997). *xCyp26c* is negatively regulated by Wnt signaling and is different from *xCyp26a*. *Cyp26c* is expressed in the anterior region, and the expression of *Cyp26c* is positively regulated by inhibition of canonical Wnt signaling and also by a high-dose RA treatment in the neurula (Tanibe et al., 2008).

To examine whether RA metabolism is involved in the anterior neurogenesis caused by the inhibition of BMP, we sought to identify the genes that are regulated during early neurogenesis and anterior neural patterning. The gene expression profiles were analyzed using *Xenopus* Affymetrix gene chips. We found that the RA-degrading enzyme *xCyp26c* was upregulated and the RA-synthesizing enzyme RALDH was downregulated following BMP-4 inhibition with DNBR in *Xenopus* animal cap explants. We found that *xCyp26c* played a critical role in the specification of anterior neural tissue in animal cap explants and whole embryos. In summary, we concluded that *xCyp26c* upregulated by DNBR is required for anterior neural development via RA degradation during early *Xenopus* development. We suggest that anterior neural tissue is not simply generated by a default pathway of pre-existing intracellular molecules but by the posterior modifier inhibition molecules, including *xCyp26c*, which is newly expressed after inhibition of BMP signaling.

MATERIALS AND METHODS

Embryo injection and explant culture

Xenopus leavis embryos were obtained by artificial fertilization (Smith and Slack, 1983). Developmental stages were determined according to Nieuwkoop and Faber (Nieuwkoop and Faber, 1956). The vitelline membranes were removed by immersing the embryos in 2.5 % thioglycolic acid. Embryos at the one-cell stage or two-cell stage were injected in the animal pole with messenger RNA, as described in the figure legends. Animal caps were dissected from the injected embryos at stage 8-9 and cultured to various stages in 67% Leibovitz's L-15 medium (GIBCO/BRL) with BSA (1 mg/ml), 7 mM Tris-HCl (pH 7.5) and gentamicin (50 µg/ml).

In vitro transcription

All of the synthetic mRNAs used for microinjection were produced by *in vitro* transcription. The *xCyp26c* cDNA was inserted in the pCS2 vector. The cDNA was linearized and used for in

synthesis of capped mRNA using an *in vitro* transcription kit (Ambion) in accordance with the manufacturer's instructions. The synthetic RNA was quantified by ethidium bromide staining in comparison with a standard RNA (Yoon et al., 2014a).

RNA isolation and reverse transcription-polymerase chain reaction (RT-PCR)

Total RNA was extracted from whole embryos or cultured animal explants with TRIzol reagent (Tel-Test Inc.) following the manufacturer's instruction. RT-PCR was performed with a Superscript preamplification system (Invitrogen). PCR was performed as follows: first, a denaturation step of 94°C for 5 min; second, 94°C for 1 min; third, annealing temperature appropriate for each primer pair for 1 min; fourth, 72°C for 1 min; and fifth, repeat the second, third and fourth steps for 19-30 cycles of amplification as described by the *Xenopus* Molecular Marker Resource (XMMR; University of Texas).

Morpholino oligos

Morpholino oligos (Gene tools LLC) for antisense oligodeoxynucleotides were used for loss-of-function studies. The base composition of the antisense oligodeoxynucleotide was a 25-mer morpholino 5' TAC AAG ATG TTC CTC CTT GAG ATC A 3' (MO-*xCyp26c*). Morpholino oligos substitute riboside moieties with nitrogen-containing morpholine moieties and are phosphorodiamidate-linked (Summerton and Weller, 1997). Oligos were re-suspended in sterile water and injected at doses of 40 ng per embryo (Yoon et al., 2014b).

Whole-mount *in situ* hybridization

Whole-mount *in situ* hybridization was completed as described (Harland, 1991) with modifications as reported (Holleman et al., 1998). Embryos were fixed in MEMFA and processed for the whole-mount *in situ* hybridization using digoxigenin-labeled antisense RNA probes. To produce the *xCyp26c* antisense probe, the cDNA was linearized by *EcoRI* and transcribed with T7 RNA polymerase (Ambion).

RESULTS

xCyp26c is induced by BMP inhibition and expressed in an anterior specific pattern in *Xenopus* embryos

To identify the genes involved during early neurogenesis, homogeneous animal cap explant samples were obtained from embryos with or without BMP signaling inhibition. Two nanograms of synthetic mRNA of DNBR was injected into one- or two-cell stage embryos. Animal cap explants isolated from blastula stage (stage 8-9) embryos were allowed to develop until the early gastrula stage (stage 11.5-12). The gene expression profiles were analyzed using *Xenopus* Affymetrix Gene chips containing 14,400 gene transcripts. Expression of various genes during early neurogenesis was induced by BMP signaling inhibition. We found 'Xl.1946.1.A1_' to be a newly expressed gene in early neuroectodermal cells (Fig. 1A). We isolated the full-length cDNA of this gene and identified it as a member of the *Xenopus* cytochrome P450 subfamily 26c (*xCyp26c*) through homology analysis of amino acid sequences in NCBI protein databases. Alignment of multiple sequences and phylogenetic analysis were used to determine the *Cyp26* subfamilies (Fig. 1B).

Using RT-PCR, we analyzed the temporal expression of *xCyp26c* mRNA throughout embryogenesis (Fig. 1C). Temporal-*xCyp26c* was not expressed maternally, but was detected in the gastrula stages (stage 12), and expression was sustained




Fig. 1. Induction of *xCyp26c* by blocking BMP in animal cap explants and the expression pattern of *xCyp26c* in whole embryos. (A) Selection of the 'Xl.1946.1.A1_at' gene induced by DNBR in a cDNA microarray. The expression of 'Xl.1946.1.A1_at' was confirmed by RT-PCR. (B) Phylogenetic relationships among the xCYP26 proteins. (C) Developmental expression of *xCyp26c*. RT-PCR analysis was performed at various stages, as indicated (Nieuwkoop and Faber, 1956). Histone 4 or *EF1 α* serve as an RNA loading control. (D) Spatial expression patterns of *xCyp26c*. Whole-mount *in situ* hybridization analysis of *xCyp26c* was performed during oocyte to tail bud stages. Views of whole embryos at the stages are indicated on top of each column. *xCyp26c* was expressed in the presumptive anterior neural ectoderm, and the expression continued to the tail bud stage.

to the tail bud stage. To examine the spatial expression pattern of *xCyp26c*, we performed whole-mount *in situ* hybridization during the embryonic stages (Fig. 1D). *xCyp26c* shows a restricted expression pattern in the anterior region from the early neural to tail bud stage (Fig. 1D). Zygotic expression of *xCyp26c* was detected as a semi-circular shape during the late gastrula stage in the dorsal region (Fig. 1D) and at a tail-bud stage in the presumptive anterior neural region (Fig. 1D).

Ectopic expression of *xCyp26c* induces neural genes in animal cap explants

To investigate the function of *xCyp26c* during early *Xenopus* development, gain-of-function experiments were performed. For effective *xCyp26c* expression in *Xenopus* embryos, *xCyp26c* mRNA was injected into one- or two-cell stage embryos. Animal caps were dissected from the injected embryos at stage 8-9 and cultured to stages 12 and 24, as described in the Materials and Methods.

At stage 12, *xCyp26c* induced the expression of the early neural marker *Zic3*, *Ngnr1* (Fig. 2A). However, *xCyp26c* had no effect on the expression of ventral (*Vent1*) or mesodermal markers (*Xbra*). At stage 24, the expression of the anterior neural marker *Otx2* was slightly increased by *xCyp26c* (Fig. 2B). Otherwise, the posterior neural marker *HoxB9* and the mesoderm marker *actin* were not induced by *xCyp26c*. These results provided evidence for a specific role for *xCyp26c* in anterior neural development, but not in mesoderm and posterior formation at the molecular level.

xCyp26c inhibits all-trans and 9-cis RA-induced posteriorization of anterior neural tissue in animal cap explants

To investigate whether *xCyp26c* was involved in A-P patterning, we examined the effect of *xCyp26c* in combination with DNBR and RA (all-trans and 9-cis RA), which induced both anterior and posterior neural tissue in animal cap explants (Tanibe et al., 2008). Animal caps were isolated from embryos injected with *xCyp26c* mRNA or DNBR and cultured in the presence or ab-




Fig. 2. Induction of neural genes by overexpression of *xCyp26c* in animal cap explants. *Xenopus* embryos at the one cell stage or two-cell stage were injected in the animal pole with *xCyp26c* mRNA. Animal caps were dissected from the injected embryos at stages 8-9 and cultured to stage 12 (A) or 24 (B). RT-PCR was then performed to detect expression of the indicated genes. *EF1 α* serves as a RNA loading control.

sence of RA (all-trans and 9-cis RA). RT-PCR analysis showed that *xCyp26c* enhanced anterior neural marker (*Otx2*, *BF1*) and inhibited posterior neural marker expression (*HoxB9*) (Figs. 3A and 3B). However, *xCyp26c* did not affect the expression of the anterior neural marker in A-P patterning induced by a combination of DNBR and basic FGF (bFGF), whereas it enhanced posterior neural markers (Fig. 3C). As results, we suggest that *xCyp26c* blocks the posteriorization caused by RA, but not by bFGF.

Depletion of *xCyp26c* inhibits the formation of normal axis and head structures in whole embryos

To further verify the function of *xCyp26c* during normal development, we performed a loss-of-function study using the anti-




Fig. 3. Inhibition of all-trans and 9-cis retinoic acid-induced posteriorization by overexpression of *xCyp26c* in animal cap explants. *xCyp26c* was injected with DNBR, and animal caps were treated with all-trans retinoic acid (AT-RA) (A), 9-cis retinoic acid (9cRA) (B), and basic FGF (bFGF) (C) at stage 8 and then cultured to stage 24. The expression pattern of marker genes were analyzed by RT-PCR.

sense morpholino directed against *xCyp26c* (MO-*xCyp26c*). We generated a MO-*xCyp26c* that targets the first 25 nucleotides of the untranslated region, including the AUG translational start site in *Xenopus xCyp26c* mRNA. MO-*xCyp26c* was effective in specifically reducing the level of the 3'-terminal end HA- or Flag-tagged *xCyp26c* protein (*xCyp26c*-HA or *xCyp26c*-flag) (Fig. 4A). Injection of MO-*xCyp26c* into the animal region at the one- or two-cell stage caused inhibition of axis formation and the head structure of tadpole stage embryos (Fig. 4B).

To investigate whether depletion of *xCyp26c* could inhibit DNBR-induced anterior neural marker expression, animal caps isolated from embryos injected with MO-*xCyp26c* alone or together with DNBR were incubated until stage 24. MO-*xCyp26c* did not suppress the expression of *Otx2* induced by DNBR (Fig 4C).

DISCUSSION

In this study, we examined whether RA metabolism was involved in the anterior neurogenesis caused by the inhibition of BMP. We found that the RA-degrading enzyme *xCyp26c* was upregulated and that the RA-synthesizing enzyme RALDH was downregulated after BMP signaling blockade in animal cap




Fig. 4. Inhibition of normal head and axis formation by depletion of endogenous *xCyp26c*. Morpholino activity was confirmed by Western blot. (B) Embryos injected with 40 ng of MO-*xCyp26c* did not have head structures and had short axis formation. (C) *xCyp26c* was injected with DNBR and animal caps were cultured to stage 24. Expression patterns of *Otx2* and *HoxB9* were analyzed by RT-PCR.

explants from *Xenopus* embryos. We also observed that ectopic expression of *xCyp26c* induced neural gene expression and inhibited all-trans and 9-cis RA-induced posteriorization of neural tissue in animal cap explants. In addition, depletion of *xCyp26c* inhibited the formation of normal axis and head structures in whole embryos. Taken together, *xCyp26c* upregulation by DNBR is necessary for anterior neural development in *Xenopus*. We suggest that anterior neurogenesis may not be the default pathway for pre-existing intracellular molecules, but requires newly synthesized posterior modifier inhibitors, including *xCyp26c*, after inhibition of BMP signaling.

We examined whether the inhibition of BMP directly elicited anterior neurogenesis without the involvement of RA metabolism or with RA-metabolizing enzymes, which were newly up-regulated after BMP signaling blockade. To identify the genes involved in early neurogenesis, we used a *Xenopus* Affymetrix gene chip containing 14400 gene transcripts and obtained 107 ESTs that was upregulated after inhibition of BMP signaling. Among these ESTs, we found that *X1.19461A1-at* was upregulated. This was further characterized as a member the *Xenopus* cytochrome P450 subfamily 26, *xCyp26c*, which is a metabolic enzyme that converts active RA to its inactive form. In this study, we found that BMP inhibition upregulated the expression of the RA degradation enzyme *xCyp26c* and downregulated the RA-synthesizing enzyme RALDH during neurogenesis in animal cap explants. A-P neural patterning plays a key role in the formation of normal neural development in embryos. Antagonists of Wnt or BMPs are expressed as anteriorizing factors (Gamse and Sive, 2000), whereas Wnts induce neuroectoderm posteriorization in *Xenopus* (Kuhl, 2003). Other

signaling molecules, such as FGFs and RA, are also involved in embryonic A-P specification as modifiers (Doniach, 1995; Gavalas and Krumlauf, 2000; Mason, 1996). Previously, it has been reported that *xCyp26c* is expressed in the anterior region during early neural development (Hollemann et al., 1998). Its expression is specifically restricted to the anterior neural region during the early neurula to tailbud stages. We also found that *xCyp26c* is not expressed maternally. Most importantly, components of RA signaling, RA receptor, and RA converting enzyme (RALDH), which are expressed in dorsolateral ectoderm and more posteriorly, lead to a reduction in anterior structures and induction of positional genes (Gamse and Sive, 2000). Alternatively, RA hydroxylase (*Cyp26*), which targets RA for degradation, leads to the expansion of anterior structures (de Roos et al., 1999; Hollemann et al., 1998). To examine the expression profiles of posterior modifier modulating genes in early neurogenesis (stage 11), we analyzed 98 genes that were the most upregulated by BMP inhibition in animal cap explants. At least, 3 genes were RA-related enzymes, and 4 genes were Wnt signal related molecules, including secreted frizzled protein 2, frizzled 4 and 7. In fact, *xCyp26c* has been reported as one of the genes that are negatively regulated by canonical Wnt signaling through the anterior neuroectoderm of early neural embryos (Tanibe et al., 2008). A more detailed study of the relationship between Wnt and RA and their mutual regulation mechanisms during anterior neurogenesis is needed. We suggest that anterior neurogenesis caused by BMP inhibition may not be the default pathway of pre-existing molecules but may require the expression of new molecules that inhibit anterior to posterior modifier molecules such as RA and Wnt.

In this study, we found that *xCyp26c* was expressed during the late gastrula stage, and the expression remained higher in later developmental stages. Whole-mount *in situ* hybridization revealed that it is anteriorly expressed, similar to previous reports that also mentioned that temporal expression of *xCyp26c* in the anterior embryonic region was 40-fold higher than in the posterior region. *xCyp26c* is expressed in only a subset of cranial nerves during late embryonic development (Tanibe et al., 2008). Because inhibition of BMP with DNBR upregulated the expression *xCyp26c* in animal explants, we examined the roles of *xCyp26c* in neurogenesis. We found that overexpression of *xCyp26c* upregulated the expression of neural markers *zic3*, *otx2*, and *ngnr1* in animal explants. However, the expression of ventral markers *vent-1*, *vent-2*, *GATA-2*, *xbra* and *actin* (mesodermal marker) were not affected much. It is interesting how the overexpression of the RA degradation enzyme *xCyp26c* led to the upregulation of neural genes, including *zic3*, *otx2* and *ngnr1*. The neurogenesis mechanism caused by *xCyp26c* remains unclear.

Previously, Kolm et al. (1997) reported that the *Cyp26* enzyme utilizes RA as its main substrate within an expected range to a limit of RA-mediated posteriorization of embryos. *Cyp26* plays a crucial role in the A-P patterning of the neural ectoderm in the late blastula stage, whereas *Hoxb1* is expressed later as an earlier posterior marker (Kudoh et al., 2002). FGF signaling is also involved in posterior neural formation (Xu et al., 1997). We examined whether *xCyp26c* was specific for A-P patterning including RA or is also involved in FGF-mediated patterning. *xCyp26c* was specifically involved in A-P patterning by blocking the posteriorization caused by RA, but did not affect the posteriorization induced by FGF. The results showed that *xCyp26c* induced anterior neural marker (*otx2* and *BF1*) and inhibited posterior marker expression (*HoxB9*), but did not affect the expression of anterior neural markers in A-P

patterning in the presence of FGF and DNBR. Rather, *xCyp26c* upregulated the posterior neural marker expression (Fig. 3c). In a previous report in *zebrafish* embryos (Kudoh et al., 2002), FGF3 and Wnt signaling were reported to independently suppress *Cyp26*, and FGF or Wnt signals initiate posterior gene expression only in the presence of an intact RA signaling pathway. Because FGF over-activation may lead to an expansion of posterior genes more than that due to abrogation of *Cyp26*, FGF-mediated posteriorization may include other factors and/or different FGFs may use a different set of RA-metabolizing enzymes. Wnt facilitates the A-P patterning and requires *Cyp26c* mediated RA metabolism (Tanibe et al., 2008). In our study, the results indicated that bFGF mediated A-P patterning was not fully dependent on RA metabolism by *xCyp26c*.

We also found that the loss of function assay of *xCyp26c* by using *Mo-xCypc26* led to anterior truncation in whole embryos. However, DNBR-induced anterior neural tissue was not suppressed by the depletion of *xCyp26c* in animal cap explants. These results indicate that *xCyp26c* is not an essential factor for anterior neurogenesis caused by BMP inhibition. Instead, we suggest that other factors, such as Wnt signal inhibitors, must contribute to the anterior neurogenesis in animal cap explants. Taken together, all of these results proved our proposed hypothesis that *xCyp26c* is involved in anterior neural tissue development during early embryonic stages of *Xenopus laevis*. However, the detailed interaction mechanism among FGF, Wnt and RA signals in neurogenesis and the mechanism for anterior neurogenesis by BMP inhibition require further study.

ACKNOWLEDGMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Education, Science, and Technology (NRF-2013R1A1A2008541) and Hallym University Research Fund (HRF-G-2015-3).

REFERENCES

- Abu-Abed, S., Dollé, P., Metzger, D., Beckett, B., Chambon, P. and Petkovich, M. (2001). The retinoic acid-metabolizing enzyme, *Cyp26A1*, is essential for normal hindbrain patterning, vertebral identity, and development of posterior structures. *Genes Dev.* 15, 226-240.
- Blumberg, B., Bolado, J., Moreno, T.A., Kintner, C., Evans, R.M. and Papalopulu, N. (1997). An essential role for retinoid signaling in anteroposterior neural patterning. *Development* 124, 373-379.
- Dale, L., Howes, G., Price, B. and Smith, J. (1992). Bone morphogenetic protein 4: a ventralizing factor in early *Xenopus* development. *Development* 115, 573-585.
- de Roos, K., Sonneveld, E., Compaaan, B., ten Berge, D., Durston, A.J. and van der Saag, P.T. (1999). Expression of retinoic acid 4-hydroxylase (*Cyp26*), during mouse and *Xenopus laevis* embryogenesis. *Mech. Dev.* 82, 205-211.
- Doniach, T. (1995). Basic FGF as an inducer of anteroposterior neural pattern. *Cell* 83, 1067-1070.
- Dosch, R., Gawantka, V., Delius, H., Blumenstock, C. and Niehrs, C. (1997). *Bmp-4* acts as a morphogen in dorsoventral mesoderm patterning in *Xenopus*. *Development* 124, 2325-2334.
- Fujii, H., Sato, T., Kaneko, S., Gotoh, O., Fujii-Kuriyama, Y., Osawa, K., Kato, S. and Hamada, H. (1997). Metabolic inactivation of retinoic acid by a novel P450 differentially expressed in developing mouse embryos. *EMBO J.* 16, 4163-4173.
- Gamse, J. and Sive, H. (2000). Vertebrate anteroposterior patterning: the *Xenopus* neuroectoderm as a paradigm. *BioEssays* 22, 976-986.
- Gavalas, A. and Krumlauf, R. (2000). Retinoid signalling and hindbrain patterning. *Curr. Opin. Genet. Dev.* 10, 380-386.
- Glinka, A., Wu, W., Onichtchouk, D., Blumenstock, C. and Niehrs,

- C. (1997). Head induction by simultaneous repression of Bmp and Wnt signalling in *Xenopus*. *Nature* 389, 517-519.
- Graff, J.M., Thies, R.S., Song, J.J., Celeste, A.J. and Melton, D.A. (1994). Studies with a *Xenopus* BMP receptor suggest that ventral mesoderm-inducing signals override dorsal signals *in vivo*. *Cell* 79, 169-179.
- Harland, R.M. (1991). *In situ* hybridization: an improved whole-mount method for *Xenopus* embryos. *Methods Cell Biol.* 36, 685.
- Harland, R. and Gerhart, J. (1997). Formation and function of Spemann's organizer. *Annu. Rev. Cell Dev. Biol.* 13, 611-667.
- Hemmati-Briuanlou, A. and Melton, D.A. (1994). Inhibition of activin receptor signaling promotes neuralization in *Xenopus*. *Cell* 77, 273-281.
- Hemmati-Briuanlou, A. and Thomsen, G.H. (1995). Ventral mesodermal patterning in *Xenopus* embryos: expression patterns and activities of BMP-2 and BMP-4. *Dev. Genet.* 17, 78-89.
- Holleman, T., Chen, Y., Grunz, H. and Pieler, T. (1998). Regionalized metabolic activity establishes boundaries of retinoic acid signalling. *EMBO J.* 17, 7361-7372.
- Hwang, Y.-S., Seo, J.-J., Cha, S.-W., Lee, H.-S., Lee, S.-Y., Roh, D.-H., Kung, H.-f., Kim, J., and Park, M.J. (2002). Antimorphic PV. 1 causes secondary axis by inducing ectopic organizer. *Biochem. Biophys. Res. Commun.* 292, 1081-1086.
- Hwang, Y.-S., Lee, H.-S., Roh, D.-H., Cha, S.-W., Lee, S.-Y., Seo, J.-J., Kim, J. and Park, M.J. (2003). Active repression of organizer genes by C-terminal domain of PV. 1. *Biochem. Biophys. Res. Commun.* 308, 79-86.
- Jones, C.M., and Smith, J. (1998). Establishment of a BMP-4 morphogen gradient by long-range inhibition. *Dev. Biol.* 194, 12-17.
- Jones, C.M., Lyons, K.M., Lapan, P., Wright, C., and Hogan, B. (1992). DVR-4 (bone morphogenetic protein-4). as a posterior-ventralizing factor in *Xenopus* mesoderm induction. *Development* 115, 639-647.
- Kessler, D.S., and Melton, D.A. (1994). Vertebrate embryonic induction: mesodermal and neural patterning. *Science* 266, 596-604.
- Knecht, A.K., and Harland, R.M. (1997). Mechanisms of dorsal-ventral patterning in noggin-induced neural tissue. *Development* 124, 2477-2488.
- Kolm, P.J., Apekin, V., and Sive, H. (1997). *Xenopus* hindbrain patterning requires retinoid signaling. *Dev. Biol.* 192, 1-16.
- Kudoh, T., Wilson, S.W., and Dawid, I.B. (2002). Distinct roles for Fgf, Wnt and retinoic acid in posteriorizing the neural ectoderm. *Development* 129, 4335-4346.
- Kuhl, M. (2003). Wnt Signaling in Development.
- Lee, H.-S., Lee, S.-Y., Lee, H., Hwang, Y.-S., Cha, S.-W., Park, S., Lee, J.-Y., Park, J.-B., Kim, S., and Park, M.J. (2011a). Direct response elements of BMP within the PV. 1A promoter are essential for its transcriptional regulation during early *Xenopus* development. *PLoS One* 6, e22621.
- Lee, S.-Y., Yoon, J., Lee, H.-S., Hwang, Y.-S., Cha, S.-W., Jeong, C.-H., Kim, J.-I., Park, J.-B., Lee, J.-Y., and Kim, S. (2011b). The function of heterodimeric AP-1 comprised of c-Jun and c-Fos in activin mediated Spemann organizer gene expression. *PLoS One* 6, e21796.
- MacLean, G., Abu-Abed, S., Dollé, P., Tahayato, A., Chambon, P., and Petkovich, M. (2001). Cloning of a novel retinoic-acid metabolizing cytochrome P450, Cyp26B1, and comparative expression analysis with Cyp26A1 during early murine development. *Mech. Dev.* 107, 195-201.
- Mason, I. (1996). Neural induction: Do fibroblast growth factors strike a cord? *Curr. Biol.* 6, 672-675.
- Nebert, D.W., and Russell, D.W. (2002). Clinical importance of the cytochromes P450. *The Lancet* 360, 1155-1162.
- Nieuwkoop, P. (1952). Activation and organization of the central nervous system in amphibians. Part III. Synthesis of a new working hypothesis. *J. Exp. Zool.* 120, 83-108.
- Nieuwkoop, P.D., and Faber, J. (1956). Normal table of *Xenopus laevis* (Daudin). A systematical and chronological survey of the development from the fertilized egg till the end of metamorphosis. (Amsterdam: North-Holland Publishing Company. Guilders).
- Ray, W.J., Bain, G., Yao, M., and Gottlieb, D.I. (1997). Cyp26, a novel mammalian cytochrome P450, is induced by retinoic acid and defines a new family. *J. Biol. Chem.* 272, 18702-18708.
- Ruiz i Altaba, A., and Jessell, T.M (1991). Retinoic acid modifies the pattern of cell differentiation in the central nervous system of neurula stage *Xenopus* embryos. *Development* 112, 945-958.
- Sakai, Y., Meno, C., Fujii, H., Nishino, J., Shiratori, H., Saijoh, Y., Rossant, J., and Hamada, H. (2001). The retinoic acid-inactivating enzyme Cyp26 is essential for establishing an uneven distribution of retinoic acid along the anterior-posterior axis within the mouse embryo. *Genes Dev.* 15, 213-225.
- Schmidt, J., Francois, V., Bier, E., and Kimelman, D. (1995). *Drosophila* short gastrulation induces an ectopic axis in *Xenopus*: evidence for conserved mechanisms of dorsal-ventral patterning. *Development* 121, 4319-4328.
- Sirbu, I.O., Gresh, L., Barra, J., and Duester, G. (2005). Shifting boundaries of retinoic acid activity control hindbrain segmental gene expression. *Development* 132, 2611-2622.
- Smith, J., and Slack, J. (1983). Dorsalization and neural induction: properties of the organizer in *Xenopus laevis*. *J. Embryol. Exp. Morphol.* 78, 299-317.
- Summerton, J., and Weller, D. (1997). Morpholino antisense oligomers: design, preparation, and properties. *Antisense Nucleic Acid Drug Dev.* 7, 187-195.
- Suzuki, A., Thies, R.S., Yamaji, N., Song, J.J., Wozney, J.M., Murakami, K., and Ueno, N. (1994). A truncated bone morphogenetic protein receptor affects dorsal-ventral patterning in the early *Xenopus* embryo. *Proc. Natl. Acad. Sci. USA* 91, 10255-10259.
- Tahayato, A., Dollé, P., and Petkovich, M. (2003). Cyp26C1 encodes a novel retinoic acid-metabolizing enzyme expressed in the hindbrain, inner ear, first branchial arch and tooth buds during murine development. *Gene Expr. Patterns* 3, 449-454.
- Tanibe, M., Michiue, T., Yukita, A., Danno, H., Ikuzawa, M., Ishiura, S., and Asashima, M. (2008). Retinoic acid metabolizing factor *xCyp26c* is specifically expressed in neuroectoderm and regulates anterior neural patterning in *Xenopus laevis*. *Int. J. Dev. Biol.* 52, 893-901.
- Wawersik, S., Evola, C., and Whitman, M. (2005). Conditional BMP inhibition in *Xenopus* reveals stage-specific roles for BMPs in neural and neural crest induction. *Dev. Biol.* 277, 425-442.
- White, J.A., Guo, Y.-D., Baetz, K., Beckett-Jones, B., Bonasoro, J., Hsu, K.E., Dilworth, F.J., Jones, G., and Petkovich, M. (1996). Identification of the retinoic acid-inducible all-trans-retinoic acid 4-hydroxylase. *J. Biol. Chem.* 271, 29922-29927.
- Xu, R.-H., Kim, J., Taira, M., Zhan, S., Sredni, D., and Kung, H. (1995). A dominant negative bone morphogenetic protein 4 receptor causes neuralization in *Xenopus* ectoderm. *Biochem. Biophys. Res. Commun.* 212, 212-219.
- Xu, R.H., Kim, J., Taira, M., Sredni, D., and Kung, H. (1997). Studies on the role of fibroblast growth factor signaling in neurogenesis using conjugated/aged animal caps and dorsal ectoderm-grafted embryos. *J. Neurosci.* 17, 6892-6898.
- Xu, R.-H., Ault, K.T., Kim, J., Park, M.-J., Hwang, Y.-S., Peng, Y., Sredni, D., and Kung, H.-f. (1999). Opposite effects of FGF and BMP-4 on embryonic blood formation: roles of PV. 1 and GATA-2. *Dev. Biol.* 208, 352-361.
- Yoon, J., Kim, J.H., Kim, S.C., Park, J.B., Lee, J.Y., and Kim, J. (2014a). PV. 1 suppresses the expression of FoxD5b during neural induction in *Xenopus* embryos. *Mol. Cells* 37, 220-225.
- Yoon, J., Kim, J.-H., Lee, S.-Y., Kim, S., Park, J.-B., Lee, J.-Y., and Kim, J. (2014b). PV. 1 induced by FGF-Xbra functions as a repressor of neurogenesis in *Xenopus* embryos. *BMB Rep.* 47, 673.