

Bowel Obstruction after Liver Transplant: A Rare Cause

Tiago Ribeiro^{a,b} Miguel Mascarenhas^{a,b,c} Hélder Cardoso^{a,b,c}
Guilherme Macedo^{a,b,c}

^aDepartment of Gastroenterology, Centro Hospitalar Universitário de São João, Alameda Professor Hernâni Monteiro, Porto, Portugal; ^bWorld Gastroenterology Organization Gastroenterology and Hepatology Training Center, Porto, Portugal; ^cFaculty of Medicine of the University of Porto, Alameda Professor Hernâni Monteiro, Porto, Portugal

Keywords

Liver transplantation · Cirrhosis · Bowel obstruction · Gallstone ileus · Double-balloon enteroscopy

Obstrução Intestinal Após Transplante Hepático: Uma Causa Rara

Palavras Chave

Transplante hepático · Obstrução intestinal · Íleo biliar · Enteroscopia de duplo balão

We present the case of a 67-year-old man with a medical history significant for orthotopic liver transplant 20 years earlier due to alcohol-related liver cirrhosis. The posttransplant period was complicated with posttransplant lymphoproliferative disease, treated with adjustment of immunosuppressive therapy and systemic chemotherapy, and is currently on remission. This patient developed a biliary stenosis at the level of the biliary anastomosis (choledococholedocostomy), leading to secondary graft cirrhosis. Cross-sectional magnetic resonance cholangiopancreatography demonstrated the anastomotic stricture with a concomitant dilation of the distal common bile duct (native bile duct segment), with recurrent identification of large biliary stones in this dilated section. During the post-transplant follow-up period, he had multiple hospital admissions due to acute cholangitis (last episode in 2018), requiring multiple endoscopic retrograde cholangiopancreatographies for



Fig. 1. Axial abdominal CT scan image showing endoluminal content within a jejunal loop (arrow) with upstream small bowel dilation. CT, computed tomography.

stone extraction. The patient presented to the emergency department with right upper quadrant pain, jaundice, and fever. Laboratory workup revealed elevated total bilirubin (10 mg/dL) and C-reactive protein (77 mg/L). An abdominal ultrasound was performed and revealed a discrete dilation of intrahepatic bile ducts. The patient was hospitalized under antibiotic therapy for acute cholangitis. Given the prior episodes of acute cholangitis and lithiasis and the current clinical presentation, the patient underwent an endoscopic retrograde cholangiopancreatography which showed no evidence of dilated

intrahepatic bile ducts or biliary calculi but reported abundant gastric and duodenal stasis content. Over the following days, the patient developed recurrent biliary vomiting and abdominal distension, complicating with prerenal acute kidney injury and portosystemic hepatic encephalopathy. A contrast-enhanced abdominal computed tomography revealed dilation of the jejunal loops, with a maximum diameter of 40 mm, and the presence of obstructing endoluminal content (Fig. 1). Distal to this point, no dilation was evident. Due to his poor physical condition and underlying comorbidities, surgery was considered to carry an unacceptably high risk. A double-balloon enteroscopy (DBE) was performed revealing an impacted biliary stone with approximately 25 mm in the distal jejunum (Fig. 2). The stone was mobilized to the stomach using a polypectomy snare and ultimately removed using a retrieving

basket (RothNet[®], STERIS, Dublin, Ireland) (Fig. 3). The patient had resolution of obstructive symptoms and resumed oral feeding with tolerance.

Bowel obstruction is a rare late complication after liver transplantation, occurring in approximately in 1–2% of cases [1]. The most common etiologies include intra-abdominal adhesions and internal hernias [2]. Gallstone ileus is an uncommon etiology of mechanical small bowel obstruction, accounting for up to 4% of all cases, being more frequent in women [3]. Contrast-enhanced computed tomography is the *gold standard* for the diagnosis of this entity and can be of value for decision-making regarding the therapeutic strategy. To date, cases of gallstone ileus have scarcely been described in orthotopic liver transplant patients. Recently, 2 cases of gallstone ileus have been reported after liver transplantation [4, 5]. In both cases, the patients had been submitted to liver transplantation due to biliary atresia and had a bilioenteric anastomosis, favoring stone migration, and an altered gastrointestinal anatomy (Y-en-Roux jejunojunostomy). Both cases were managed with enterotomy for stone extraction. Opposite to these reports, in this case, the patient did not have a bilioenteric anastomosis and was managed using an endoscopic technique. Indeed, the location of stone impaction combined with the accessibility to DBE allowed a safe and efficient stone extraction in a patient with a poor surgical condition.

The resolution of gallstone ileus most often requires aggressive fluid and electrolyte followed by early intervention to relieve bowel obstruction [3]. Surgical enterotomy remains the standard of care for stone extraction and resolution of bowel obstruction. Moreover, when the presence of a bilioenteric fistula is responsible for the passage of a large stone to the small bowel, surgery allows the closure of the defect [3]. The use of DBE for resolution of bowel obstruction has scarcely been reported. Nevertheless, DBE can be a valid alternative to the standard surgical approach in patients with a poor preoperative condition. Endoscopic treatment options include electrohydraulic and mechanical lithotripsy, as well as retrieving baskets [6, 7].




Fig. 2. Gallstone and alimentary debris impacted in the distal jejunum.




Fig. 3. a Gallstone from the stomach before being removed with a retrieving basket. **b** Subsequent measurement of the gallstone confirmed its estimated size of 25 mm (photo taken after the specimen had been several hours in a refrigerator for preservation).

Statement of Ethics

All rules of the Local Ethics Committee (“Comissão de Ética para a Saúde do Centro Hospitalar de São João/Faculdade de Medicina da Universidade do Porto, Portugal”) were followed, preserving patient identity and confidentiality. Informed consent was obtained from the patient for the publication of his case. Written informed consent to share the medical case and images was obtained from the patient.

Conflict of Interest Statement

The authors declare that there was no source of funding.

Funding Sources

The authors have no funding sources to declare.

References

- 1 Porrett PM, Hsu J, Shaked A. Late surgical complications following liver transplantation. *Liver Transpl.* 2009;15(Suppl 2):S12–8.
- 2 Blachar A, Federle MP. Bowel obstruction following liver transplantation: clinical and ct findings in 48 cases with emphasis on internal hernia. *Radiology.* 2001;218(2):384–8.
- 3 Ayantunde AA, Agrawal A. Gallstone ileus: diagnosis and management. *World J Surg.* 2007;31(6):1292–7.
- 4 Vanerio P, Morgade P, Rodriguez F, San Martin G, Abelleira M. Gallstone ileus after hepaticojejunostomy in liver transplant recipient: case report. *HPB.* 2021;23: S581.
- 5 Komine R, Sakamoto S, Uchida H, Nakao T, Kodama T, Okada N, et al. Gallstone ileus at 17 years after living donor liver transplantation: a case report. *Pediatr Transplant.* 2023; 27(4):e14517.
- 6 Heinzow HS, Meister T, Wessling J, Domschke W, Ullerich H. Ileal gallstone obstruction: single-balloon enteroscopic removal. *World J Gastrointest Endosc.* 2010;2(9):321–4.
- 7 Bourke MJ, Schneider DM, Haber GB. Electrohydraulic lithotripsy of a gallstone causing gallstone ileus. *Gastrointest Endosc.* 1997; 45(6):521–3.

Author Contributions

Tiago Ribeiro: clinical management of the patient, collection of clinical and endoscopic data, bibliographic review, and drafting and critical revision of the manuscript. Miguel Mascarenhas: collection of clinical data, bibliographic review, drafting of the manuscript, and critical revision of the manuscript. Hélder Cardoso: performance of the endoscopic procedure and critical revision of the manuscript. Guilherme Macedo: critical revision of the manuscript and final approval of the manuscript. All authors agree to be accountable for all aspects of the manuscript.

Data Availability Statement

Data are stored in the hospital’s electronic patient database and will be provided upon reasonable request.