



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

- in patients with abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 2020;**60**:365–73.
- 2 Miller K, Joldes G, Qian J, Patel AP, Jung MS, Tavner A, et al. Maximum principal AAA wall stress is proportional to wall thickness. In: Miller K, Nielsen PMF, editors. *Computational Biomechanics for Medicine*. New York: Springer Nature; 2018. p. 43–53.
 - 3 Joldes GR, Noble C, Polzer S, Taylor ZA, Wittek A, Miller K. A simple method of incorporating the effect of the Uniform Stress Hypothesis in arterial wall stress computations. *Acta Bioeng Biomech* 2018;**20**:59–67.

Karol Miller*, Adam Wittek, Grand Joldes

Intelligent Systems for Medicine Laboratory, The University of Western Australia, Perth, WA, Australia

*Corresponding author.

Email-address: karol.miller@uwa.edu.au (Karol Miller)

Available online 20 July 2020

© 2020 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2020.05.017>

DOI of original article: <https://doi.org/10.1016/j.ejvs.2020.02.023>

Response to “Re Biomechanical Assessment Predicts Aneurysm Related Events in Patients with Abdominal Aortic Aneurysm”

We thank Miller *et al.* for their interest in our recent publication.¹ Indeed, it was always the intention to test the developed methods^{2,3} in the MA³RS Cohort,⁴ with our analyses beginning in 2017.

In response to the three points raised: (1) magnetic resonance imaging is often used to measure aortic wall thickness, with good repeatability reported.^{5–7} We found mean wall thickness measurements to differ by $\pm 12\%$ between three users. This equates to a 6% difference in maximum principal stress and <5% difference in peak aneurysm biomechanical ratio (ABR). (2) Including residual stress into wall stress simulations typically reduces the magnitude of the maximum principal stress. Therefore, all cases would have lower wall stress and lower ABR. It is unlikely that this would change the conclusions of the paper. We finalised our methods and began the ABR computation before Miller *et al.* published their residual stress approach in 2018.⁸ Furthermore, their publication does not state that the method is included in new releases of BioPARR. (3) One could argue that no model is truly “patient specific” as assumptions are almost always required. We followed the common use of the term “patient specific” in abdominal aortic aneurysm (AAA) studies whereby the aneurysm geometry (including wall thickness in our work) and pressure loading are patient specific. Also, the wall strength model uses geometric information (local ratio of diameter; local thickness of thrombus), gender, and family history,⁹ all of which are unique to that patient, to generate the pointwise strength data. Interestingly, Miller *et al.* refer to their AAA simulations as “patient specific”,^{9,10} despite

using only simplified uniform wall thickness AAA geometries and blood pressure loading, which are less patient specific data than those used in the models of Doyle *et al.*¹

REFERENCES

- 1 Doyle BJ, Bappoo N, Syed MJB, Forsythe RO, Powell JT, Conlisk N, et al. Biomechanical assessment predicts aneurysm related events in patients with abdominal aortic aneurysm. *Eur J Vasc Endovasc Surg* 2020;**60**:365–73.
- 2 Joldes G, Miller K, Wittek A, Forsythe R, Newby D, Doyle B. BioPARR: a software system for estimating the rupture potential index for abdominal aortic aneurysms. *Sci Rep* 2017;**7**:4641.
- 3 Joldes GR, Miller K, Wittek A, Doyle B. A simple, effective and clinically applicable method to compute abdominal aortic aneurysm wall stress. *J Mech Behav Biomed Mater* 2016;**58**:139–48.
- 4 MA³RS Study Investigators. Aortic wall inflammation predicts abdominal aortic aneurysm expansion, rupture, and need for surgical repair. *Circulation* 2017;**136**:787–97.
- 5 Mensel B, Kühn J, Schneider T, Quadrat A, Hegenscheid K. Mean thoracic aortic wall thickness determination by cine MRI with steady-state free precession: validation with dark blood imaging. *Acad Radiol* 2013;**20**:1004–8.
- 6 Li AE, Kamel I, Rando F, Anderson M, Kumbasar B, Lima JA, et al. Using MRI to assess aortic wall thickness in the Multi-Ethnic Study of Atherosclerosis (MESA): distribution by race, sex, and age. *AJR Am J Roentgenol* 2004;**182**:593–7.
- 7 Liu C-Y, Chen D, Bluemke DA, Wu CO, Teixido-Tura G, Chugh AR, et al. Evolution of aortic wall thickness and stiffness with atherosclerosis: long-term follow up from the Multi-Ethnic Study of Atherosclerosis (MESA). *Hypertension* 2015;**65**:1015–9.
- 8 Joldes GR, Noble C, Polzer S, Taylor Z, Wittek A, Miller K. A simple method of incorporating the effect of uniform stress hypothesis in arterial wall stress computations. *Acta Bioeng Biomech* 2018;**20**:59–67.
- 9 Vande Geest J, Wang D, Wisniewski S, Makaroun M, Vorp D. Towards a non-invasive method for determination of patient-specific wall strength distribution in abdominal aortic aneurysms. *Ann Biomed Eng* 2006;**34**:1098–106.
- 10 Miller K, Mufty H, Catlin A, Rogers C, Saunders B, Sciarra R, et al. Is there a relationship between stress in walls of abdominal aortic aneurysm and symptoms? *J Surg Res* 2020;**252**:37–46.

Barry Doyle on behalf of All The Authors

The University of Western Australia, Perth, Australia

Email-address: barry.doyle@uwa.edu.au

Available online 20 July 2020

© 2020 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2020.06.017>

DOI of original articles: <https://doi.org/10.1016/j.ejvs.2020.05.017>

<https://doi.org/10.1016/j.ejvs.2020.02.023>

Re “The Elephant in the Operating Room”

I read the work by Massiot *et al.*¹ and the commentary by Gonçalves and Chakfé² with great interest. Technology has provided us with high resolution imaging equipment allowing us to perform demanding procedures at a fraction of the exposure of older devices. In most centres, low dose protocols and radiation safety training are already in place. However,

daily exposure to radiation should not be underestimated as its cumulative effects could take decades to be evaluated. Because of the poly-aetiological nature of some pathologies (e.g. neoplasms), it may be impossible to link final health effects of radiation on medical personnel to exposure sustained by those personnel many years previously.

During a procedure, there are three to six individuals in the room. Any radiation dose received is dependent on an individual's distance from the radiation source, but continuous work in high volume centres results in accumulated exposure. Some of these professionals might be of reproductive age. Is it ethically acceptable to irradiate a number of young individuals daily and repeatedly for treatment of patients who are usually older and burdened with significant comorbidities?

This is a question that will raise the eyebrows of colleagues ready to cite the Hippocratic Oath. Yet, the situation described is not similar to putting a medical professional's health in jeopardy under extraordinary circumstances, e.g. intensive care unit specialists fighting COVID-19 or trauma surgeons operating in a war zone. The systematic exposure of young professionals to a silent hazard like radiation is the daily working environment of thousands. What happens when these individuals get sick or give birth to children with catastrophic genetic disorders? What are the social and long term financial burdens? Further research is required to find answers to these questions.

If we wish to discuss the details of the "Elephant in the Operating Room", we must also discuss the ethics of working with "Elephants".

REFERENCES

- 1 Massiot N, Abdallah IB, Duprey A, Leygnac S, Corcos O, Cadtier Y, et al. Multicentre evaluation of an extra low dose protocol to reduce radiation exposure in superior mesenteric artery stenting. *Eur J Vasc Endovasc Surg* 2020;60:925–31.
- 2 Bastos Goncalves F, Chakfe N. The elephant in the operating Room: radiation safety and vascular practice. *Eur J Vasc Endovasc Surg* 2020;60:932.

Nikolaos Patelis

3rd Department of Vascular Surgery, Athens Medical Center, Athens, Greece
Email-address: patelisin@gmail.com

Available online 9 October 2020

© 2020 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2020.09.008>

DOI of original article: <https://doi.org/10.1016/j.ejvs.2020.09.007>

Response to "Re The Elephant in the Operating Room"

We thank Dr Patelis for his thought provoking letter. The risk of working with radiation is evident but still largely

uncertain. There is no "radiation free" option for most specialists, certainly not for residents who are young and require great exposure in a short period of time, but also for nurses and anaesthetists. In short, we are compelled to put ourselves in harm's way to do our job. With the endovascular revolution, our practice now takes place in a dangerous environment, a fact that must be acknowledged.

This is one big mammoth we can hardly control, let alone ignore.

One cannot rely solely on ethics for radiation safety, we should strive for stricter regulation to protect professionals. This entails substantial investment in safety and monitoring equipment, improved risk assessment, enforcement of risk reduction measures, and obligatory (re)certification in radiation management. Administrators and physicians who do not comply need to be liable. Scientific societies and other professional associations, political stakeholders, and scientific publications like that of Massiot *et al.*¹ all play a role.

This elephant is not leaving the room. We must learn to tame it.

REFERENCE

- 1 Massiot N, Abdallah IB, Duprey A, Leygnac S, Corcos O, Cadtier Y, et al. Multicentre evaluation of an extra low dose protocol to reduce radiation exposure in superior mesenteric artery stenting. *Eur J Vasc Endovasc Surg* 2020;60:925–31.

Frederico Bastos Gonçalves*

Serviço de Angiologia e Cirurgia Vascular, Centro Hospitalar Universitário de Lisboa Central/NOVA Medical School, Rua de Santa Marta 50, Lisbon, 1150-299, Portugal

Nabil Chakfé

Department of Vascular Surgery and Kidney Transplantation and GEPROVAS, University Hospital of Strasbourg, France

*Corresponding author.

Email-address: f.bastosgoncalves@nms.unl.pt (Frederico Bastos Gonçalves)

Available online 13 October 2020

© 2020 European Society for Vascular Surgery. Published by Elsevier B.V. All rights reserved.

<https://doi.org/10.1016/j.ejvs.2020.09.007>

DOI of original article: <https://doi.org/10.1016/j.ejvs.2020.09.008>

Ethics in Occupational Exposure Needs a Green New Deal

We would like to thank Dr Patelis for his relevant comments regarding the worrying issue of long term occupational exposure.¹ We definitely agree that it is the real question lying behind all the recent literature regarding radiation. By law in France, staff are supposed to be trained, monitored, and adequately equipped against radiation. This is the institution's responsibility. On an individual basis, every operator has the moral responsibility of justification and optimisation.