

Article

Tetrel Bonds with π -Electrons Acting as Lewis Bases—Theoretical Results and Experimental Evidences

Sławomir J. Grabowski ^{1,2} 

¹ Faculty of Chemistry, University of the Basque Country and Donostia International Physics Center (DIPC), P.K. 1072, 20080 Donostia, Spain; s.grabowski@ikerbasque.org; Tel.: +34-943-01-5477

² IKERBASQUE, Basque Foundation for Science, 48011 Bilbao, Spain

Academic Editor: Steve Scheiner

Received: 30 April 2018; Accepted: 11 May 2018; Published: 15 May 2018



Abstract: MP2/aug-cc-pVTZ calculations were carried out for the ZFH₃-B complexes (Z = C, Si, Ge, Sn and Pb; B = C₂H₂, C₂H₄, C₆H₆ and C₅H₅⁻; relativistic effects were taken into account for Ge, Sn and Pb elements). These calculations are supported by other approaches; the decomposition of the energy of interaction, Quantum Theory of Atoms in Molecules (QTAIM) and Natural Bond Orbital (NBO) method. The results show that tetrel bonds with π -electrons as Lewis bases are classified as Z...C links between single centers (C is an atom of the π -electron system) or as Z... π interactions where F-Z bond is directed to the mid-point (or nearly so) of the CC bond of the Lewis base. The analogous systems with Z...C/ π interactions were found in the Cambridge Structural Database (CSD). It was found that the strength of interaction increases with the increase of the atomic number of the tetrel element and that for heavier tetrel elements the ZFH₃ tetrahedral structure is more deformed towards the structure with the planar ZH₃ fragment. The results of calculations show that the tetrel bond is sometimes accompanied by the Z-H...C hydrogen bond or even sometimes the ZFH₃-B complexes are linked only by the hydrogen bond interaction.

Keywords: electron charge shifts; tetrel bond; hydrogen bond; π -electrons as Lewis bases; σ -hole

1. Introduction

The tetrel bond is a Lewis acid—Lewis base interaction that may play an important role in some chemical and biological processes [1]; for example, it may be considered as a preliminary stage of the S_N2 reaction [2]. This interaction was classified as the σ -hole bond by Politzer and coworkers since it may be defined as an interaction between the 14th Group element acting as the Lewis acid centre through its σ -hole and a region that is rich of the electron density by a lone electron pair, π -electron system etc. [3,4]. The σ -hole is usually located in this case in the elongation of the covalent bond to the tetrel centre and it is often characterized by the positive electrostatic potential (EP) [3,4]. It seems that first time the tetrel bond was analyzed in terms of the σ -hole concept in the SiF₄ complexes with amines [5]. In spite of the fact that the term “tetrel bond” appeared recently [6] and also this interaction was classified as the σ -hole bond in the last decade [3–5] it was analyzed in earlier studies. For example, the SiF₄...NH₃ and SiF₄...(NH₃)₂ complexes were analyzed theoretically since ab initio MO calculations were performed with the use of STO-3G and STO-6G basis sets [7], another study that is more complex concerns a large sample of complexes of silicon derivatives with electron-rich groups [8]. Both latter studies were performed before the proposition of the σ -hole concept [9,10] and the introduction of the tetrel bond term [6].

One can mention other studies on tetrel bonds as for example that one where this interaction was analyzed as a preliminary stage of the S_N2 reaction [11], the theoretical analysis of structural and

energetic properties of acetonitrile complexes with the 14 Group tetrahalides [12]; a study on the Lewis acid carbon center, the corresponding interaction was labeled as the carbon bond and it was compared with the hydrogen bond [13]; it is worth mentioning that the carbon bond is a sub-class of the tetrel bond interactions [2,11]. The tetrel-hydride interaction is another sub-class of tetrel bonds where the negatively charged H-atom plays the role of Lewis base center [14].

There are other, more recent studies on this kind of interaction; only few are mentioned here; the analysis of factors which influence the strength of tetrel bonds [15], the analysis of mechanisms of S_N2 reactions, among them at the C centre [16], the role of tetrel bonds in the crystal structures' stabilization [17], the theoretical analysis of the H-Si...N and F-Si...N linear or nearly so arrangements [18], comparison of neutral and charge assisted tetrel bonds [19], the geometry deformations of monomers linked by tetrel bond [20] or the balance between the attractive forces of tetrel interactions and the steric repulsions in crystal structures [21].

One may cite numerous other examples since the number of studies on this kind of interaction has increased rapidly. However, it seems that there are no systematic and extensive studies on the tetrel bonds with π -electrons playing a role of Lewis bases or at most they are very rare and they are not a main goal of investigations. For example, very recently, various σ -hole bonds were analyzed with the use of few theoretical approaches: halogen, chalcogen, pnictogen and tetrel bonds were compared [22]; three different types of Lewis bases were considered there, neutral species (NH_3), anion (Cl^-) and the π -electron system (C_2H_2). Hence two tetrel bonded complexes with acetylene molecule playing a role of the Lewis base were analyzed there among various other complexes. These are the $SiFH_3 \cdots C_2H_2$ and $GeFH_3 \cdots C_2H_2$ complexes [22].

The other issue that is not analyzed so frequently concerns the tetrel bond interactions with the heavier tetrel elements, such analyses are rather rare and mainly concern germanium species. There are more experimental studies on tetrel bonds with heavier tetrel elements playing a role of the Lewis acid centers, however only sometimes such experimental analyses are supported by theoretical results [23]. One of examples where heavier tetrel elements were considered in tetrel bonds is a study on the SnF_4 and PbF_4 complexes with NH_3 and HCN that play a role of Lewis bases through the nitrogen centre [24]. A theoretical study analogous to the latter one was performed on the lighter tetrel species since the complexes of CF_4 , SiF_4 and GeF_4 Lewis acid units with NH_3 and AsH_3 Lewis bases were analyzed [25].

Returning to the π -electron species—numerous theoretical and experimental studies on interactions where such systems play a role of Lewis bases may be mentioned. These are mainly those studies that concern hydrogen bonded systems [26]. However other Lewis acid—Lewis base interactions with π -electron donors were analyzed very often [27–30]. One can even mention the triel bonds between the boron or aluminium Lewis acid center and acetylene or ethylene [31,32] or the recent study where the multivalent halogen centers act as the Lewis acids [33].

The aim of this study is an analysis of the tetrel bonds in complexes of ZFH_3 species, where Z labels the following centers; C, Si, Ge, Sn and Pb, thus light and heavy tetrrels are taken into account; the acetylene, ethylene, benzene and cyclopentadienyl anion were chosen as the π -electron moieties acting as the Lewis bases. Different theoretical techniques are applied here to deepen the understanding of the nature of these tetrel bond interactions; i.e., the Quantum Theory of Atoms in Molecules (QTAIM) [34], Natural Bond Orbital (NBO) approach [35], the decomposition of the energy of interaction [36,37] as well as the analysis of the electrostatic potential (EP) distribution [38]. The short descriptions of the theoretical approaches applied here are included in the section that concerns the computational details.

2. Results and Discussion

2.1. Energetic and Geometric Parameters

Figure 1 presents examples of complexes analyzed here. All kinds of Lewis bases that are considered are shown in selected examples of the figure. The molecular graphs are presented since they reflect geometry of species analyzed. However these graphs are discussed further here in the section on QTAIM results.




Figure 1. The molecular graphs of the selected complexes analyzed here; big circles— attractors, small green circles—BCPs, the nonnuclear attractor (NNA) is located (small red circle) between two BCPs of the CC bond in a case of the $\text{SnFH}_3\text{-C}_2\text{H}_2$ complex.

The energetic parameters of analyzed complexes, among them, the binding and interaction energies corrected for BSSE, E_{bin} and E_{int} , respectively, are included in Table 1. One can see much stronger interactions, i.e., greater $-E_{\text{int}}$ and $-E_{\text{bin}}$ values, for complexes with the cyclopentadienyl

anion, than for complexes with the other Lewis base units. This may be explained since the complexes of $C_5H_5^-$ anion that are linked through the tetrel bonds are assisted by negative charge; the latter anion is much stronger base than the remaining species chosen here. The following tendencies are also observed here, and it does not depend on the choice of $-E_{int}$ or $-E_{bin}$ value for the discussion; the strength of interaction for the same Lewis base increases in the following order of the tetrel center $C < Si < Ge < Sn \cong Pb$. It was observed earlier for tetrel bond interactions [3,4] and it was explained by the increase of the electrostatic part of the energy of interaction since the electrostatic potential (EP) at the tetrel σ -hole increases with the increase of the atomic number [3,4]. The calculations performed here show the EP of tetrel σ -hole of the ZFH_3 species equal to 0.033; 0.062; 0.068; 0.081; 0.080 au for C, Si, Ge, Sn and Pb centers, respectively (0.001 au electron density surfaces were chosen). The EP values for tin and lead centers are almost equal one to each other. This is why for complexes analyzed here similar Lewis acid properties are observed for these centers; however the strongest interaction is observed for the $SnFH_3-C_5H_5^-$ complex if the interaction energy is considered while if the binding energy is taken into account thus it is the $PbFH_3-C_5H_5^-$ complex.

Table 1. The energetic parameters of complexes analyzed (in kcal/mol); interaction energy, E_{int} , binding energy, E_{bin} , deformation energy, E_{def} and BSSE correction. The distance between Lewis base and Lewis acid units is included—the shortest $Z \cdots C$ distance was chosen, the values in parentheses show if this distance is greater than the corresponding sum of van der Waals radii (positive values) or if it is lower (negative ones), distances in Å.

Complex	Distance	E_{int}	E_{bin}	E_{def}	BSSE
$CFH_3-C_2H_2$	3.428 (+0.53)	−1.3	−1.2	0.0	0.3
$CFH_3-C_2H_4$	3.458 (+0.56)	−1.3	−1.3	0.0	0.4
$CFH_3-C_6H_6$	3.398 (+0.50)	−2.7	−2.7	0.0	1.0
$CFH_3-C_5H_5^-$	3.241 (+0.34)	−10.8	−10.5	0.3	1.2
$SiFH_3-C_2H_2$	3.344 (+0.04)	−2.4	−2.3	0.0	0.6
$SiFH_3-C_2H_4$	3.315 (−0.02)	−2.7	−2.6	0.1	0.8
$SiFH_3-C_6H_6$	3.253 (−0.05)	−3.7	−3.7	0.1	1.3
$SiFH_3-C_5H_5^-$	2.477 (−0.82)	−27.5	−18.1	9.4	2.0
$GeFH_3-C_2H_2$	3.285 (−0.02)	−2.6	−2.6	0.1	1.1
$GeFH_3-C_2H_4$	3.253 (−0.05)	−2.9	−2.9	0.1	1.5
$GeFH_3-C_6H_6$	3.203 (−0.10)	−4.3	−4.2	0.1	2.7
$GeFH_3-C_5H_5^-$	2.525 (−0.78)	−29.3	−21.0	8.3	4.4
$SnFH_3-C_2H_2$	3.325 (−0.13)	−3.4	−3.3	0.1	1.3
$SnFH_3-C_2H_4$	3.280 (−0.17)	−3.8	−3.7	0.2	1.8
$SnFH_3-C_6H_6$	3.183 (−0.27)	−5.5	−5.2	0.3	3.2
$SnFH_3-C_5H_5^-$	2.519 (−0.93)	−41.9	−30.6	11.4	5.2
$PbFH_3-C_2H_2$	3.323 (−0.18)	−3.5	−3.4	0.1	2.2
$PbFH_3-C_2H_4$	3.267 (−0.23)	−3.8	−3.7	0.1	3.1
$PbFH_3-C_6H_6$	3.148 (−0.35)	−5.8	−5.6	0.3	5.7
$PbFH_3-C_5H_5^-$	2.624 (−0.88)	−39.3	−31.8	7.6	8.3

If the Lewis acid unit is the same thus the interaction strength increases in the following order $C_2H_2 < C_2H_4 < C_6H_6 < C_5H_5^-$. One can also see that the $-E_{int}$ or $-E_{bin}$ values do not exceed 6 kcal/mol for all complexes of acetylene, ethylene and benzene while they are much greater in a case of complexes with cyclopentadienyl, especially large values are observed for the above-mentioned tin and lead complexes.

The BSSE corrections are greater for stronger interactions, especially large values are observed for interactions in cyclopentadienyl complexes. The deformation energy, E_{def} , is a parameter that is related to geometrical changes of the interacting systems. For example, in a case of the strong $A-H \cdots B$ hydrogen bonds, the complexation often leads to the meaningful elongation of the $A-H$ proton donating bond that results in the greater E_{def} values [39]. Steric effects are very important for tetrel bonded species [11,21] since the tetrel center, often characterized by the sp^3 hybridization and surrounded by four substituents (like for the systems considered here) is hardly available for the Lewis base (nucleophilic attack). Thus the tetrel-base link should cause greater deformations connected

with the increase of the availability of the tetrel center. In other words the ZFH₃ tetrahedral system should be closer to the trigonal bipyramid in the ZFH₃·B complex with the ZH₃ part being closer to planarity. For complexes of acetylene, ethylene and benzene E_{def} does not exceed 0.3 kcal/mol indicating negligible changes of geometry resulting from complexation. However for the C₅H₅[−] complexes, if one excludes the CFH₃-C₅H₅[−] complex with this energy amounting only 0.3 kcal/mol, E_{def} is close to 10 kcal/mol or even exceeds this value. This is in agreement with changes of geometry; one can see (Figure 1) the ZH₃ part close to planarity and the F-Z···C arrangement close to linearity for two complexes presented; SnFH₃-C₅H₅[−] and PbFH₃-C₅H₅[−].

The above-presented EP values at the Z-tetrel centre concern the σ-hole that occurs in the extension of F-Z bonds. For the ZFH₃ species analyzed here, similarly as for other sp³ hybridized tetrel centers four σ-holes located in extensions of covalent bonds to Z-center occur. However the electronegative F-substituent enhances F-Z σ-hole [3,4] that results in greater positive EP values than those of other H-Z σ-holes. For example, for the SnFH₃ molecule the EP value at the F-Sn σ-hole is equal to +0.081 au while this value for the H-Sn σ-hole amounts +0.048 au. For the clarity of the results' presentation only interactions of the F-Z σ-hole are considered here; it means that the F-Z σ-hole is directed to the π-electron system in the configurations analyzed.

Table 1 presents the Lewis acid—Lewis base distances, for each of complexes the shortest Z···C contact was chosen. One can see that these distances are usually greater than 3 Å, only for the C₅H₅[−] complexes where stronger interactions are observed such distances amount ~2.5 Å (except of the CFH₃-C₅H₅[−] complex where this distance is equal to ~3.2 Å). It was pointed out in numerous studies that the distance between interacting units is roughly related to the strength of interaction, this was observed for the hydrogen bonded complexes [40] but it seems that such dependence occurs also for other types of interactions [2]. It is often stated in various studies that the sum of van der Waals radii of two atoms being in contact roughly indicates at which distance a significant so-called noncovalent interaction begins [41]. Table 1 presents how the Z···C distances are related to the corresponding sum of Z and C van der Waals radii. These distances are greater than the corresponding sum for carbon complexes (Z = C) and for the SiFH₃-C₂H₂ complex while for the remaining ones these distances are lower than the van der Waals sum; the following van der Waals radii were applied here, H—1.2 Å [42], C—1.7 Å, Si—2.1 Å, Ge—2.1 Å, Sn—2.25 Å and Pb—2.3 Å [43]. One can see that for the strongest interactions in the C₅H₅[−] complexes the Z···C distance is almost by 1 Å lower than the sum of van der Waals radii. If one considers only the Z···C distances as a measure of the strength of interaction thus the interactions in the CFH₃ complexes are very weak and one may contest even their stabilizing nature.

Table 2 presents geometrical parameters related to the changes resulting from complexation; one of them is a percentage elongation of the Z-F bond related to the corresponding isolated ZFH₃ species that is not involved in the tetrel bond. One can see that these elongations correspond to the deformation energies, the greatest values are observed for the cyclopentadienyl complexes. For the species analyzed here three F-Z-H angles in the Lewis acid unit are very close one to each other; however for each complex considered its average value is considered in the further discussions; the latter angle is defined in Figure 2. For the isolated ZFH₃ species this tetrahedral angle, labeled as α_{iso}, amounts from 101.4° for PbFH₃ to 108.8° for CFH₃. It decreases in the complex to α_{comp} (up to 90° corresponding to the planar ZH₃ system in the trigonal bipyramid structure). The angle decrease values, [(α_{iso} − α_{comp})/α_{iso}] × 100%, are presented in Table 2. They correspond to the deformation energies discussed earlier here as well as to the elongations of the Z-F bonds since the greatest decreases are observed for the strongest interactions. The Z-F bond elongations result from the π_{CC} → σ_{ZF}^{*} and σ_{CH} → σ_{ZF}^{*} overlaps; Table 2 shows the orbital-orbital NBO energies corresponding to these overlaps; E_{NBO}¹ is a sum of all energies of such interactions for the F—C bond considered. Similarly the E_{NBO}² energy summarizes all π_{CC} → σ_{ZH}^{*} and σ_{CH} → σ_{ZH}^{*} overlaps, however in this case the whole ZFH₃ species is considered for one E_{NBO}² value (i.e., three C-H bonds). One can see (Table 2) that both E_{NBO}¹ and E_{NBO}² values are much greater for the C₅H₅[−] complexes than for other ones.

Table 2. The characteristics of complexes analyzed; ZF% and Angle% are the percentage increase of the Z-F distance and the percentage decrease of the F-Z-H angle, respectively; E_{NBO}^1 and E_{NBO}^2 are the NBO energies defined in the text (in kcal/mol); El-trans (au) is the electron charge transfer from the Lewis base to the Lewis acid while Z-charge is the charge of the Z-center in the complex considered (both charges in au calculated within NBO approach).

Complex	ZF%	Angle%	E_{NBO}^1	E_{NBO}^2	El-Trans	Z-Charge ^a
CFH ₃ -C ₂ H ₂	0.14	0.00	0.5	0.0	-0.001	-0.158
CFH ₃ -C ₂ H ₄	0.14	0.00	0.6	0.0	-0.002	-0.158
CFH ₃ -C ₆ H ₆	0.29	0.09	0.4	0.8	-0.004	-0.156
CFH ₃ -C ₅ H ₅ ⁻	1.95	0.28	1.5	2.4	-0.030	-0.378
SiFH ₃ -C ₂ H ₂	0.25	0.65	1.7	0.3	-0.014	1.227
SiFH ₃ -C ₂ H ₄	0.31	0.74	2.3	0.4	-0.021	1.218
SiFH ₃ -C ₆ H ₆	0.31	0.74	1.8	1.0	-0.019	1.221
SiFH ₃ -C ₅ H ₅ ⁻	4.46	9.06	24.5	20.9	-0.250	1.086
GeFH ₃ -C ₂ H ₂	0.40	0.85	3.1	0.6	-0.019	1.025
GeFH ₃ -C ₂ H ₄	0.52	0.85	4.1	0.7	-0.027	1.015
GeFH ₃ -C ₆ H ₆	0.52	0.85	3.2	1.6	-0.025	1.021
GeFH ₃ -C ₅ H ₅ ⁻	5.18	8.95	33.4	22.1	-0.250	0.906
SnFH ₃ -C ₂ H ₂	0.46	1.34	4.1	1.2	-0.024	1.246
SnFH ₃ -C ₂ H ₄	0.57	1.43	5.4	1.9	-0.035	1.232
SnFH ₃ -C ₆ H ₆	0.62	1.72	4.8	2.9	-0.035	1.240
SnFH ₃ -C ₅ H ₅ ⁻	5.11	11.45	37.9	40.5	-0.289	1.148
PbFH ₃ -C ₂ H ₂	0.59	0.99	4.9	1.0	-0.027	1.085
PbFH ₃ -C ₂ H ₄	0.78	1.28	7.0	2.4	-0.039	1.070
PbFH ₃ -C ₆ H ₆	0.83	1.68	6.1	3.9	-0.044	1.080
PbFH ₃ -C ₅ H ₅ ⁻	5.44	8.78	36.5	31.0	-0.289	1.013

^a Z-charge for CFH₃: -0.157, SiFH₃: +1.240, GeFH₃: +1.042, SnFH₃: +1.267, PbFH₃: +1.106 (all in au).




Figure 2. The definition of the angle expressing the change of the tetrahedral structure into the structure being closer to the trigonal bipyramid.

Table 2 shows the electron charge transfer values from the Lewis base to the Lewis acid unit, these transfers are especially great for the ZFH₃-C₅H₅⁻ complexes, except of the CFH₃-C₅H₅⁻ one. Such electron charge redistributions resulting from complexation are usually great for those complexes where the geometry deformations are important [2]; this is observed also here. The charge of the Z-central tetrel atom is shown in Table 2; one can see that this charge decreases (is “more negative”) in complexes in comparison with the corresponding isolated ZFH₃ species. This is in opposite to the A-H...B hydrogen bonded systems where the complexation usually results in the increase of the positive charge of the central H-atom [35].

2.2. Nature of Interactions—Decomposition of Interaction Energy

Table 3 presents terms of the energy of interaction resulting from the Ziegler and Rauk decomposition scheme [36,37] (see the Computational Details section).

Table 3. The terms of the energy of interaction (kcal/mol); Pauli repulsion, ΔE_{Pauli} , electrostatic, ΔE_{elstat} , orbital, ΔE_{orb} , dispersion, ΔE_{disp} , and the total interaction energy, ΔE_{int} .

Complex	ΔE_{Pauli}	ΔE_{elstat}	ΔE_{orb}	ΔE_{disp}	ΔE_{int}
CFH ₃ -C ₂ H ₂	2.4	−1.3	−0.8	−1.4	−1.1
CFH ₃ -C ₂ H ₄	2.7	−1.4	−0.9	−1.6	−1.2
CFH ₃ -C ₆ H ₆	4.7	−2.4	−1.5	−3.6	−2.8
CFH ₃ -C ₅ H ₅ [−]	8.9	−11.0	−5.6	−4.2	−11.9
SiFH ₃ -C ₂ H ₂	5.6	−3.4	−2.7	−2.2	−2.6
SiFH ₃ -C ₂ H ₄	7.3	−4.3	−3.5	−2.8	−3.2
SiFH ₃ -C ₆ H ₆	8.2	−4.1	−3.3	−4.5	−3.8
SiFH ₃ -C ₅ H ₅ [−]	54.2	−43.4	−34.9	−5.6	−29.8
GeFH ₃ -C ₂ H ₂	7.1	−4.4	−3.2	−2.7	−3.2
GeFH ₃ -C ₂ H ₄	9.1	−5.5	−4.1	−3.4	−3.8
GeFH ₃ -C ₆ H ₆	10.6	−5.6	−4.1	−5.8	−5.0
GeFH ₃ -C ₅ H ₅ [−]	56.1	−47.7	−32.8	−6.3	−30.7
SnFH ₃ -C ₂ H ₂	8.7	−5.8	−3.8	−3.0	−3.9
SnFH ₃ -C ₂ H ₄	11.6	−7.3	−5.0	−3.9	−4.6
SnFH ₃ -C ₆ H ₆	14.0	−7.8	−5.6	−6.9	−6.2
SnFH ₃ -C ₅ H ₅ [−]	75.3	−68.1	−41.8	−6.6	−41.2
PbFH ₃ -C ₂ H ₂	9.2	−6.3	−3.8	−2.8	−3.6
PbFH ₃ -C ₂ H ₄	12.9	−8.3	−5.1	−3.7	−4.2
PbFH ₃ -C ₆ H ₆	17.9	−10.0	−6.6	−7.4	−6.2
PbFH ₃ -C ₅ H ₅ [−]	71.4	−65.9	−37.2	−6.6	−38.4

One can see that only for the CFH₃ complexes with acetylene, ethylene and benzene the dispersion term, ΔE_{disp} , is the most important attractive one. This is typical for weak van der Waals interactions where attractive interaction energy terms related to charge distributions and to electron charge shifts, ΔE_{elstat} and ΔE_{orb} , are less important [2]. For the above-mentioned three complexes, $-\Delta E_{\text{int}}$ does not exceed 3 kcal/mol, and in two cases it is close to 1 kcal/mol. The electron charge shifts for these complexes (Table 2) do not exceed 4 millielectrons! The latter is connected with the practically unchanged carbon charge in the CFH₃ unit in those complexes in comparison with the isolated CFH₃ molecule. For the remaining complexes electrostatic interaction energy is the most important attractive term, only in a case of the SiFH₃-C₆H₆ and GeFH₃-C₆H₆ complexes the dispersive term is slightly “less negative” than the electrostatic one. If one excludes the above-mentioned three CFH₃ complexes, thus for majority of remaining complexes the orbital interaction, ΔE_{orb} , is the next most important attractive term, after electrostatic interaction.

It was discussed in recent studies on hydrogen bonds and on other σ -hole bonds that these interactions are accompanied by effects that are a response for the Pauli repulsion [2,44]. The latter was also discussed for halogen bonds where multivalent halogen center plays a role of the Lewis acid while the π -electrons are the Lewis base [33]. For such interactions correlations were found between the repulsion interaction energy and different terms of the attractive interaction. It was found in earlier studies that the orbital interaction energy (if one refers to the decomposition scheme applied here) well correlates with the repulsion term, correlations for other interaction energy terms are not so good. However, in general, the sum of all attractive terms correlates with the Pauli repulsion term [2,33,44]. Figure 3 presents such a correlation for the complexes analyzed here. Thus the attractive interaction which is related to various effects related to complexation, among them to the electron charge redistribution, is a response for the Pauli repulsion.




Figure 3. The linear correlation between the repulsion interaction energy and the sum of attractive terms (both in kcal/mol) for the ZFH₃-B complexes analyzed here.

The orbital interaction reflecting electron shifts corresponds to energy terms which are named in the other way in other decomposition schemes; most often they are labeled as the delocalization interaction energy, induction, charge transfer, polarization and others [2]. Figure 4 shows, for the complexes analyzed here, the correlation between the orbital energy, ΔE_{orb} , and the electron charge shift resulting from complexation.




Figure 4. The linear correlation between the orbital interaction energy, ΔE_{orb} , and the electron charge shift from the Lewis base unit to the Lewis acid (au).

2.3. Quantum Theory of Atoms in Molecules Parameters

Table 4 presents characteristics of the bond critical point (BCP) of the bond path that connects the Lewis acid and Lewis base units of the complex. It is a link between the tetrel center (tetrel attractor) or the hydrogen center (hydrogen attractor) and the critical point of the Lewis base species. This critical point may correspond to the carbon atom attractor, to the bond critical point (BCP) of the CC bond or to the non-nuclear attractor (NNA) located on the CC bond path. Hence one can see that there are various topologies of complexes analyzed here.

Table 4. The QTAIM parameters (in au) of BCP of the Lewis acid—Lewis base bond path; electron density at BCP, ρ_{BCP} , its laplacian, $\nabla^2\rho_{\text{BCP}}$, and the total electron energy density at BCP, H_{BCP} . The bond path type is also indicated.

Complex	ρ_{BCP}	$\nabla^2\rho_{\text{BCP}}$	H_{BCP}	BP-Type
CFH ₃ -C ₂ H ₂	0.005	0.020	0.001	C...C
CFH ₃ -C ₂ H ₄	0.005	0.018	0.001	C...C
CFH ₃ -C ₆ H ₆	0.007	0.022	0.001	(C)H...C
CFH ₃ -C ₅ H ₅ [−]	0.011	0.038	0.001	(C)H...C
SiFH ₃ -C ₂ H ₂	0.007	0.022	0.001	Si...C
SiFH ₃ -C ₂ H ₄	0.008	0.023	0.001	(Si)H...C
SiFH ₃ -C ₆ H ₆	0.008	0.025	0.001	(Si)H...C
SiFH ₃ -C ₅ H ₅ [−]	0.036	0.012	−0.012	Si...C
GeFH ₃ -C ₂ H ₂	0.009	0.027	0.001	Ge...NNA(CC)
GeFH ₃ -C ₂ H ₄	0.010	0.028	0.001	Ge...BCP(CC)
GeFH ₃ -C ₆ H ₆	0.010	0.028	0.001	Ge...C
-	0.007	0.024	0.001	(Ge)H...C
GeFH ₃ -C ₅ H ₅ [−]	0.037	0.045	−0.008	Ge...C
SnFH ₃ -C ₂ H ₂	0.010	0.025	0.001	Sn...NNA(CC)
SnFH ₃ -C ₂ H ₄	0.011	0.029	0.001	Sn...BCP(CC)
SnFH ₃ -C ₆ H ₆	0.012	0.029	0.001	Sn...C
-	0.007	0.022	0.001	(Sn)H...C
SnFH ₃ -C ₅ H ₅ [−]	0.045	0.066	−0.011	Sn...C
PbFH ₃ -C ₂ H ₂	0.011	0.034	0.001	Pb...NNA(CC)
PbFH ₃ -C ₂ H ₄	0.013	0.036	0.001	Pb...BCP(CC)
PbFH ₃ -C ₆ H ₆	0.014	0.039	0.001	Pb...C
-	0.009	0.026	0.001	(Pb)H...C
PbFH ₃ -C ₅ H ₅ [−]	0.042	0.074	−0.008	Pb...C
-	0.015	0.041	0.001	(Pb)H...C

The above-mentioned bond path may concern the tetrel bond if the Z-center of the Lewis acid unit is linked with the Lewis base critical point or it may concern the hydrogen bond if the H-atom attractor of the Lewis acid is linked with the Lewis base critical point. One may expect the (Z)H...C bond paths show some “artificial interactions”, especially since the meaning of the bond path and its usefulness to analyze interactions is often a subject of controversies [45] and disputes [46,47]. The presented here preliminary results on tetrel bonds where π -electrons play a role of Lewis bases need additional extended studies. However few arguments that the accompanying (Z)H...C bond paths observed for some benzene and cyclopentadienyl complexes may correspond to weak hydrogen bonds are listed here. The electron densities at the (Z)H...C bond critical points (BCPs) are not meaningless and they are comparable sometimes with such values for the Z...C BCPs; see the GeFH₃-C₆H₆ complex for example (Table 4). The PbFH₃-C₅H₅[−] complex is an example where the greatest electron density at the H...C BCP is observed since it amounts 0.015 au; note that for the medium in strength hydrogen bond in the water dimer the electron density at the H...O BCP is equal to 0.023 au (MP2/6-311++G(d,p) results [48]). Additionally the H...C intermolecular contacts correspond to the attractive electrostatic interactions since the carbon centers of the C₆H₆ and C₅H₅[−] moieties are characterized by the negative electrostatic potentials (EPs) while the H-centers of the ZFH₃ species by the positive EPs.

Particularly the following cases of bond paths are observed for complexes analyzed here. For the CFH₃-C₂H₂ and CFH₃-C₂H₄ complexes the irregular and nonlinear carbon-carbon bond paths are observed that may result from weak interactions (Figure 5); formally according to the QTAIM approach, they may be attributed to the tetrel bonds. For the CFH₃-C₆H₆ (Figure 1) and CFH₃-C₅H₅[−] complexes the H...C intermolecular bond paths are observed which may be attributed to the C-H...C hydrogen bonds! For the SiFH₃-C₂H₂, SiFH₃-C₂H₄ and CFH₃-C₆H₆ complexes the non-linear bond paths are detected, similarly as for the CFH₃-C₂H₂ and CFH₃-C₂H₄ complexes, which are attributed to the Si...C or H...C intermolecular links (Table 4).

In a case of the $\text{SiFH}_3\text{-C}_5\text{H}_5^-$ complex the clear almost linear $\text{Si}\cdots\text{C}$ bond path corresponding to the strong tetrel bond is observed, similarly as for the other $\text{ZFH}_3\text{-C}_5\text{H}_5^-$ complexes for $\text{Z} = \text{Ge}$, Sn and Pb . In a case of the $\text{PbFH}_3\text{-C}_5\text{H}_5^-$ complex the additional $\text{H}\cdots\text{C}$ bond path corresponding to the $\text{Pb-H}\cdots\text{C}$ hydrogen bond is observed (Figure 1). For the $\text{ZFH}_3\text{-C}_6\text{H}_6$ complexes ($\text{Z} = \text{Ge}$, Sn , Pb) the tetrel and hydrogen bonds are observed with the corresponding bond paths, the $\text{SnFH}_3\text{-C}_6\text{H}_6$ complex representing such a situation is presented in Figure 1. Similarly the $\text{SnFH}_3\text{-C}_2\text{H}_2$, $\text{SnFH}_3\text{-C}_2\text{H}_4$ complexes in Figure 1 reflect the same situation in analogues tin and lead complexes; in the case of acetylene Lewis base the $\text{Z}\cdots\text{NNA}$ bond path is observed while in the case of ethylene Lewis base this is the $\text{Z}\cdots\text{BCP}$ bond path.




Figure 5. The molecular graphs of the $\text{CFH}_3\text{-C}_2\text{H}_2$ and $\text{CFH}_3\text{-C}_2\text{H}_4$ complexes; big circles—attractors, small green circles—BCPs, the nonnuclear attractor (NNA) is located (small red circle) between two BCPs in a case of the $\text{CFH}_3\text{-C}_2\text{H}_2$ complex.

The characteristics of bond critical points presented in Table 4 reflect the strength of interaction. It was discussed in various studies that these characteristics may be often treated as measures of the strength of interaction [49]; especially for homogeneous samples of complexes. Numerous relationships were found between characteristics of the $\text{H}\cdots\text{B}$ BCP and the strength of interaction for the $\text{A-H}\cdots\text{B}$ hydrogen bonds. For complexes analyzed here greater values of the electron density at the bond critical point, ρ_{BCP} 's, are observed for the C_5H_5^- complexes. The Laplacian of the electron density at BCP, $\nabla^2\rho_{\text{BCP}}$, is positive for all complexes analyzed which may show these are not covalent interactions; the H_{BCP} values are positive and close to zero for all complexes of acetylene, ethylene and benzene as well as for the $\text{CFH}_3\text{-C}_5\text{H}_5^-$ complex. For the remaining complexes of the C_5H_5^- anion the negative H_{BCP} values are observed that may indicate these are partly covalent in nature interactions.

One may ask what is the difference between the $\text{Z}\cdots\pi$ and $\text{Z}\cdots\text{C}$ tetrel bonds that are presented here. These “two kinds” of connections correspond to the types of bond paths. For the majority of acetylene and ethylene complexes former connections are observed while for the benzene and cyclopentadienyl complexes the latter ones. The $\text{Z}\cdots\pi$ bond path is a link between Z -attractor that corresponds to the nucleus and BCP or NNA located at the CC bond of acetylene or ethylene (see Figure 1). The $\text{Z}\cdots\text{C}$ bond path is a link between Z and C attractors corresponding to nuclei. This difference occurs within the Quantum Theory of Atoms in Molecules (QTAIM) scheme but it seems it is not observed in other approaches; for example in both cases the same orbital-orbital overlaps occur that correspond to the $\text{Z}\cdots\pi$ interactions; i.e., $\pi_{\text{CC}} \rightarrow \sigma_{\text{ZF}}^*$ ones. All other accompanying overlaps specified earlier here are the same in both cases of contacts. The similar situations were observed earlier for the hydrogen bonded complexes with the π -electron systems playing a role of Lewis bases [50].

3. Computational Details

The calculations were performed with the Gaussian16 set of codes [51] using the second-order Møller-Plesset perturbation theory method (MP2) [52], and the aug-cc-pVTZ basis set [53].

The relativistic effects for the heavier Ge, Sn and Pb atoms were taken into account. The calculations for these elements were done with quasi-relativistic small-core effective core potentials: ECP10MDF, ECP28MDF and ECP60MDF, for Ge, Sn and Pb, respectively [54]. For the latter elements the basis sets corresponding to aug-cc-pVTZ were applied, i.e., ECP10MDF_AVTZ, ECP28MDF_AVTZ and ECP60MDF_AVTZ, respectively [55]. Frequency calculations were performed for the complexes analyzed and their monomers to confirm that the optimized structures correspond to energetic minima. The binding energy, E_{bin} , was calculated as difference between the energy of the complex and the sum of energies of monomers optimized separately while the interaction energy, E_{int} , is a difference between the energy of the complex and the sum of energies of monomers which geometries come from the geometry of the complex considered [56]. The binding and interaction energies are negative but their difference—the deformation energy, $E_{\text{def}} = E_{\text{bin}} - E_{\text{int}}$, is positive and it is connected with the change of geometries of monomers resulting from the complexation [39]. The Counterpoise (CP) correction was applied to calculate the basis set superposition error BSSE [57]; hence the E_{bin} and E_{int} values corrected for BSSE are analyzed in this study.

The Quantum Theory of ‘Atoms in Molecules’ (QTAIM) was also applied to characterize critical points (BCPs) in terms of the electron density (ρ_{BCP}), its Laplacian ($\nabla^2\rho_{\text{BCP}}$) and the total electron energy density at BCP (H_{BCP}) which is the sum of the potential electron energy density (V_{BCP}) and the kinetic electron energy density (G_{BCP}) [34]. The AIMAll program was used to carry out the QTAIM calculations [58].

The Natural Bond Orbital (NBO) method [35] was applied to calculate atomic charges, the electron charge shifts from the Lewis bases to the Lewis acids as well as the orbital-orbital interactions. The $n_{\text{B}} \rightarrow \sigma_{\text{AH}}^*$ orbital-orbital interaction is characteristic for the A-H...B hydrogen bond; n_{B} labels the lone electron pair of the B Lewis base center and σ_{AH}^* is the antibonding orbital of the A-H Lewis acid bond [35]. In a case of the hydrogen bonds where π -electrons and σ -electrons play a role of the Lewis bases, A-H... π and A-H... σ systems, the $\pi_{\text{B}} \rightarrow \sigma_{\text{AH}}^*$ and $\sigma_{\text{B}} \rightarrow \sigma_{\text{AH}}^*$ overlaps, respectively, are the most important orbital-orbital interactions [59]. The similar situation occurs for the tetrel bonds analyzed here, they may be classified as the Z... π or Z...C interactions (Z labels the tetrel centre). The $\pi_{\text{CC}} \rightarrow \sigma_{\text{ZF}}^*$ and $\pi_{\text{CC}} \rightarrow \sigma_{\text{ZH}}^*$ overlaps are observed here as the most important interactions; besides the $\sigma_{\text{CH}} \rightarrow \sigma_{\text{ZF}}^*$ and $\sigma_{\text{CH}} \rightarrow \sigma_{\text{ZH}}^*$ overlaps are also detected but they are characterized by lower energies than the former interactions. For example, the $\pi_{\text{CC}} \rightarrow \sigma_{\text{ZF}}^*$ interaction is calculated as the second-order perturbation theory energy (Equation (1)):

$$\Delta E (\pi_{\text{CC}} \rightarrow \sigma_{\text{ZF}}^*) = -2 \langle \pi_{\text{CC}} | F | \sigma_{\text{ZF}}^* \rangle^2 / (\varepsilon(\sigma_{\text{ZF}}^*) - \varepsilon(\pi_{\text{CC}})), \quad (1)$$

$\langle \pi_{\text{CC}} | F | \sigma_{\text{ZF}}^* \rangle$ designates the Fock matrix element and $(\varepsilon(\sigma_{\text{ZF}}^*) - \varepsilon(\pi_{\text{CC}}))$ is the orbital energy difference. The similar equations (to Equation (1)) for the remaining above-mentioned orbital-orbital interactions may be given.

The energy decomposition analysis (EDA) [36,37] was carried out with the BP86 functional [60,61] in conjunction with the Grimme dispersion corrections (BP86-D3) [62] using uncontracted Slater-type orbitals (STOs) as basis functions for all elements with triple- ζ quality (ADF-basis set TZP). The energy decomposition analysis (EDA) was performed with the use of the ADF2013.01 program [63] for all complexes analyzed here and characterized by geometries resulting from the MP2/aug-cc-pVTZ optimizations. The EDA method follows the energy partition of Morokuma [36,37]. The interaction energy, ΔE_{int} , between two fragments (A and B) in the A-B link, in the particular electronic reference state and in the frozen geometry of AB is considered in this approach. The ΔE_{int} interaction energy is divided into three components and the additional dispersion term, ΔE_{disp} (Equation (2)):

$$\Delta E_{\text{int}} = \Delta E_{\text{elstat}} + \Delta E_{\text{Pauli}} + \Delta E_{\text{orb}} + \Delta E_{\text{disp}}, \quad (2)$$

The ΔE_{elstat} term corresponds to the electrostatic interaction between the unperturbed charge distributions of atoms and is usually attractive. The Pauli repulsion, ΔE_{Pauli} , is the energy change

associated with the transformation from the superposition of the unperturbed electron densities of the isolated fragments to the wavefunction which properly obeys the Pauli principle through explicit antisymmetrization and renormalization of the product wavefunction; it comprises the destabilizing interactions between electrons of the same spin on either fragment. The orbital interaction, ΔE_{orb} , accounts for charge transfer and polarization effects.

Figure 6 presents the correlation between the interaction energy calculated within the MP2/aug-cc-pVTZ approach (Table 1), thus at the level corresponding to the systems' optimizations, and ΔE_{int} DFT energy calculated with the use of ADF codes. The excellent correlation observed here partly justifies the use of DFT calculations for the previously optimized MP2 geometries.




Figure 6. The linear correlation between the MP2 E_{int} interaction energy and the ΔE_{int} energy calculated within the DFT approach; both energies in kcal/mol.

4. Conclusions and Perspectives

The tetrel bonds in complexes where the π -electron system plays a role of the Lewis base were analyzed here. Practically for all complexes considered the NBO approach shows the existence of the $\pi_{\text{CC}} \rightarrow \sigma_{\text{EZ}}^*$ overlaps, however in a case of complexes of CFH_3 with acetylene, ethylene and benzene the corresponding energies are negligible thus the existence of tetrel bonds is problematic. On the other hand for the remaining complexes the above-mentioned interactions are significant that may indicate the existence of the tetrel bonds. The QTAIM approach often shows the complicated topology, sometimes the additional bond paths corresponding to the hydrogen bonds are observed, or like for the $\text{CFH}_3\text{-C}_5\text{H}_5^-$ complex, only $\text{C-H}\cdots\text{C}$ intermolecular link is observed that may indicate the existence of the hydrogen bond and not of the tetrel bond. However for the other cyclopentadienyl complexes the interactions are very strong and the $\text{Z}\cdots\text{C}$ bond paths exist there. Hence there is no doubt that these complexes are linked by the tetrel bonds; all theoretical approaches applied in this study support the existence of such interactions in these complexes.

Only for some of acetylene and ethylene complexes one may observe the link between tetrel center and the site corresponding to π -electrons, NNA or BCP of the CC bond of the Lewis base unit (see the $\text{SnFH}_3\text{-C}_2\text{H}_2$ and $\text{SnFH}_3\text{-C}_2\text{H}_4$ complexes in Figure 1 as examples). In a case of the C_6H_6 and C_5H_5^- aromatic systems, the $\text{Z}\cdots\text{C}$ bond path is observed that suggest the one-atom Lewis base center and not the π -electron system. It means that the existence of two types of tetrel bonds may be considered within the QTAIM approach, $\text{Z}\cdots\pi$ and $\text{Z}\cdots\text{C}$. However other approaches applied here do not distinguish rather between these types. Such a situation was earlier observed for the hydrogen bonded complexes [50].

The question arises if the interactions analyzed theoretically here really exist. This is why the Cambridge Structural Database (CSD) [64] search was performed. The following search criteria were taken into account; non-disordered structures, R less than 10%, 3D coordinated determined, non polymeric structures, single crystal structures and no errors (CSD updates up to February of 2018 were taken into account). The additional condition was that the Z tetrel center (C, Si, Ge, Sn and Pb) has to form two intermolecular Z···C contacts within corresponding sum of van der Waals radii. Two Z···C contacts were required since one may expect that in a case of double and triple CC bonds, or if CC bond concerns delocalized aromatic system; at least two Z···C distances within the van der Waals sum should be observed. 218 systems of crystal structures fulfilling those requirements were found in CSD. However only in 10 cases the clear tetrel-CC bond contacts with the tetrahedral (sp^3 hybridized) tetrel center were observed which suggest the existence of the tetrel··· π -electrons interactions. Figure 7 shows an example where one can observe the F-Si···CC contact (CC bond of the aromatic phenyl ring). This issue requires additional studies on the experimental crystal structures however. It seems that the search criteria could be also improved. More detailed study on experimental crystal structures' results is in the progress.




Figure 7. The fragment of the crystal structure (refcode: BAKZOF) where the Si···CC tetrel bond is observed.

Funding: Financial support comes from Eusko Jaurlaritz (GIC IT-588-13) and the Spanish Government MINECO/FEDER (CTQ2016-80955).

Acknowledgments: Technical and human support provided by Informatikako Zerbitzu Orokora—Servicio General de Informática de la Universidad del País Vasco (SGI/IZO-SGIker UPV/EHU), Ministerio de Ciencia e Innovación (MICINN), Gobierno Vasco Eusko Jaurlaritz (GV/EJ), European Social Fund (ESF) is gratefully acknowledged.

Conflicts of Interest: The author declares no conflict of interest.

References

- García-Llinás, X.; Bauzá, A.; Seth, S.K.; Frontera, A. Importance of R-CF₃···O Tetrel Bonding Interactions in Biological Systems. *J. Phys. Chem. A* **2017**, *121*, 5371–5376. [[CrossRef](#)] [[PubMed](#)]
- Grabowski, S.J. Hydrogen bonds, and σ -hole and π -hole bonds—mechanisms protecting doublet and octet electron structures. *Phys. Chem. Chem. Phys.* **2017**, *19*, 29742–29759. [[CrossRef](#)] [[PubMed](#)]
- Politzer, P.; Murray, J.S.; Clark, T. Halogen bonding and other σ -hole interactions: A perspective. *Phys. Chem. Chem. Phys.* **2013**, *15*, 11178–11189. [[CrossRef](#)] [[PubMed](#)]
- Bundhun, A.; Ramasami, P.; Murray, J.S.; Politzer, P. Trends in σ -hole strengths and interactions of F₃MX molecules (M = C, Si, Ge and X = F, Cl, Br, I). *J. Mol. Model.* **2013**, *19*, 2739–2746. [[CrossRef](#)] [[PubMed](#)]

5. Politzer, P.; Murray, J.S.; Lane, P.; Concha, M.C. Electrostatically Driven Complexes of SiF₄ with Amines. *Int. J. Quantum Chem.* **2009**, *109*, 3773–3780. [[CrossRef](#)]
6. Bauzá, A.; Mooibroek, T.J.; Frontera, A. Tetrel-Bonding Interaction: Rediscovered Supramolecular Force? *Angew. Chem. Int. Ed.* **2013**, *52*, 12317–12321. [[CrossRef](#)] [[PubMed](#)]
7. Chehayber, J.M.; Nagy, S.T.; Lin, C.S. Ab initio studies of complexes between SiF₄ and ammonia. *Can. J. Chem.* **1984**, *62*, 27–31. [[CrossRef](#)]
8. Alkorta, I.; Rozas, I.; Elguero, J. Molecular Complexes between Silicon Derivatives and Electron-Rich Groups. *J. Phys. Chem. A* **2001**, *105*, 743–749. [[CrossRef](#)]
9. Clark, T.; Hennemann, M.; Murray, J.S.; Politzer, P. Halogen bonding: The σ -hole. *J. Mol. Model.* **2007**, *13*, 291–296. [[CrossRef](#)] [[PubMed](#)]
10. Politzer, P.; Lane, P.; Concha, M.C.; Ma, Y.; Murray, J.S. An overview of halogen bonding. *J. Mol. Model.* **2007**, *13*, 305–311. [[CrossRef](#)] [[PubMed](#)]
11. Grabowski, S.J. Tetrel bond- σ -hole bond as a preliminary stage of the S_N2 reaction. *Phys. Chem. Chem. Phys.* **2014**, *16*, 1824–1834. [[CrossRef](#)] [[PubMed](#)]
12. Helminiak, H.M.; Knauf, R.R.; Danforth, S.J.; Phillips, J.A. Structural and Energetic Properties of Acetonitrile-Group IV (A & B) Halide Complexes. *J. Phys. Chem. A* **2014**, *118*, 4266–4277. [[PubMed](#)]
13. Mani, D.; Arunan, E. The X-C...Y (X = O/F, Y = O/S/F/Cl/Br/N/P) ‘carbon bond’ and hydrophobic interactions. *Phys. Chem. Chem. Phys.* **2013**, *15*, 14377–14383. [[CrossRef](#)] [[PubMed](#)]
14. Li, Q.-Z.; Zhuo, H.-Y.; Li, H.-B.; Liu, Z.-B.; Li, W.-Z.; Cheng, J.-B. Tetrel-Hydride Interaction between XH₃F (X = C, Si, Ge, Sn) and HM (M = Li, Na, BeH, MgH). *J. Phys. Chem. A* **2015**, *119*, 2217–2224. [[CrossRef](#)] [[PubMed](#)]
15. Scheiner, S. Systematic Elucidation of Factors That Influence the Strength of Tetrel Bonds. *J. Phys. Chem. A* **2017**, *121*, 5561–5568. [[CrossRef](#)] [[PubMed](#)]
16. Kubelka, J.; Bickelhaupt, F.M. Activation Strain Analysis of S_N2 Reactions at C, N, O, and F centers. *J. Phys. Chem. A* **2017**, *121*, 885–891. [[CrossRef](#)] [[PubMed](#)]
17. George, J.; Dronskowski, R. Tetrel Bonds in Infinite Molecular Chains by Electronic Structure Theory and Their Role for Crystal Stabilization. *J. Phys. Chem. A* **2017**, *121*, 1381–1387. [[CrossRef](#)] [[PubMed](#)]
18. Marín-Luna, M.; Alkorta, I.; Elguero, J. A theoretical study of the H_nF_{4-n}Si:N-base (n = 1–4) tetrel-bonded complexes. *Theor. Chem. Acc.* **2017**, *136*, 41. [[CrossRef](#)]
19. Liu, M.; Li, Q.; Scheiner, S. Comparison of tetrel bonds in neutral and protonated complexes of pyridine TF₃ and furan TF₃ (T = C, Si, and Ge) with NH₃. *Phys. Chem. Chem. Phys.* **2017**, *19*, 5550–5559. [[CrossRef](#)] [[PubMed](#)]
20. Zierkiewicz, W.; Michalczyk, M.; Scheiner, S. Implications of monomer deformations for tetrel and pnictogen bonds. *Phys. Chem. Chem. Phys.* **2018**, *20*, 8832–8841. [[CrossRef](#)] [[PubMed](#)]
21. Scheiner, S. Steric Crowding in tetrel Bonds. *J. Phys. Chem. A* **2018**, *122*, 2550–2562. [[CrossRef](#)] [[PubMed](#)]
22. Grabowski, S.J.; Sokalski, W.A. Are Various σ -Hole Bonds Steered by the Same Mechanisms? *ChemPhysChem* **2017**, *18*, 1569–1577. [[CrossRef](#)] [[PubMed](#)]
23. Roy, S.; Drew, M.G.B.; Bauzá, A.; Frontera, A.; Chattopadhyay, S. Non-covalent tetrel bonding interactions in hemidirectional lead (II) complexes with nickel(II)–salen type metalloligands. *New J. Chem.* **2018**, *42*, 6062–6076.
24. Grabowski, S.J. Tetrel bonds, penta- and hexa-coordinated tin and lead centres. *Appl. Organomet. Chem.* **2017**, *31*, e3727. [[CrossRef](#)]
25. Grabowski, S.J. Lewis Acid Properties of Tetrel tetrafluorides—The Coincidence of the σ -Hole Concept with the QTAIM Approach. *Crystals* **2017**, *7*, 43. [[CrossRef](#)]
26. Nishio, M.; Hirota, M.; Umezawa, Y. *The CH/ π Interaction, Evidence, Nature, and Consequences*; Wiley-VCH: New York, NY, USA, 1998.
27. Vasilyev, A.V.; Lindeman, S.V.; Kochi, J.K. Noncovalent binding of the halogens to aromatic donors. Discrete structures of labile Br₂ complexes with benzene and toluene. *Chem. Commun.* **2001**, *10*, 909–910. [[CrossRef](#)]
28. Duarte, D.J.R.; de las Vallejos, M.M.; Peruchena, N.M. Topological analysis of aromatic halogen/hydrogen bonds by electron charge density and electrostatic potentials. *J. Mol. Model.* **2010**, *16*, 737–748. [[CrossRef](#)] [[PubMed](#)]
29. Zhuo, H.; Li, Q.; Li, W.; Cheng, J. Is π halogen bonding or lone pair... π interaction formed between borazine and some halogenated compounds. *Phys. Chem. Chem. Phys.* **2014**, *16*, 159–165. [[CrossRef](#)] [[PubMed](#)]

30. Liu, C.; Zeng, Y.; Li, X.; Meng, L.; Zhang, X. A comprehensive analysis of P \cdots π pnictogen bonds: Substitution effects and comparison with Br \cdots π halogen bonds. *J. Mol. Model.* **2015**, *21*, 143. [[CrossRef](#)] [[PubMed](#)]
31. Fau, S.; Frenking, G. Theoretical investigation of the weakly bonded donor-acceptor complexes X₃B-H₂, X₃B-C₂H₄, and X₃B-C₂H₂ (X = H, F, Cl). *Mol. Phys.* **1999**, *96*, 519–527.
32. Grabowski, S.J. Triel Bonds, π -Hole- π -Electrons Interactions in Complexes of Boron and Aluminium Trihalides and Trihydrides with Acetylene and Ethylene. *Molecules* **2015**, *20*, 11297–11316. [[CrossRef](#)] [[PubMed](#)]
33. Grabowski, S.J. New Type of Halogen Bond: Multivalent Halogen Interacting with π - and σ -Electrons. *Molecules* **2017**, *22*, 2150. [[CrossRef](#)] [[PubMed](#)]
34. Bader, R.F.W. *Atoms in Molecules, a Quantum Theory*; Oxford University Press: Oxford, UK, 1990.
35. Weinhold, F.; Landis, C. *Valency and Bonding, a Natural Bond Orbital Donor—Acceptor Perspective*; Cambridge University Press: Cambridge, UK, 2005.
36. Morokuma, K. Molecular Orbital Studies of Hydrogen Bonds. III. C=O \cdots H-O Hydrogen Bond in H₂CO \cdots H₂O and H₂CO \cdots 2H₂O. *J. Chem. Phys.* **1971**, *55*, 1236–1244. [[CrossRef](#)]
37. Ziegler, T.; Rauk, A. On the calculation of bonding energies by the Hartree–Fock Slater method. *Theor. Chim. Acta* **1977**, *46*, 1–10. [[CrossRef](#)]
38. Murray, J.S.; Politzer, P. Molecular electrostatic potentials and noncovalent interactions. *WIREs Comput. Mol. Sci.* **2017**, *7*, e1326. [[CrossRef](#)]
39. Grabowski, S.J.; Sokalski, W.A. Different types of hydrogen bonds: Correlation analysis of interaction energy components. *J. Phys. Org. Chem.* **2005**, *18*, 779–784. [[CrossRef](#)]
40. Gilli, P.; Bertolasi, V.; Ferretti, V.; Gilli, G. Evidence for resonance-assisted hydrogen bonding. 4. Covalent nature of the strong homonuclear hydrogen bond. Study of the O-H \cdots O system by crystal structure correlation methods. *J. Am. Chem. Soc.* **1994**, *116*, 909–915. [[CrossRef](#)]
41. Murray, J.S.; Politzer, P. Molecular Surfaces, van der Waals Radii and Electrostatic Potentials in Relation to Noncovalent Interactions. *Croat. Chem. Acta* **2009**, *82*, 267–275.
42. Pauling, L. *The Nature of the Chemical Bond*, 3rd ed.; Cornell University Press: New York, NY, USA, 1960.
43. Batsanov, S.S. Van der Waals Radii of Elements. *Inorg. Mater.* **2001**, *37*, 871–885. [[CrossRef](#)]
44. Grabowski, S.J. Hydrogen bonds and other interactions as a response to protect doublet/octet electron structure. *J. Mol. Model.* **2018**, *24*, 38. [[CrossRef](#)] [[PubMed](#)]
45. Poater, J.; Solà, M.; Bickelhaupt, F.M. Hydrogen-Hydrogen Bonding in Planar Biphenyl, Predicted by Atoms-In-Molecules Theory, Does Not Exist. *Chem. Eur. J.* **2006**, *12*, 2889–2895. [[CrossRef](#)] [[PubMed](#)]
46. Bader, R.F.W. Pauli Repulsions Exist Only in the Eye of the Beholder. *Chem. Eur. J.* **2006**, *12*, 2896–2901. [[CrossRef](#)] [[PubMed](#)]
47. Poater, J.; Solà, M.; Bickelhaupt, F.M. A Model of the Chemical Bond Must Be Rooted in Quantum Mechanics, Provide Insight, and Possess Predictive Power. *Chem. Eur. J.* **2006**, *12*, 2902–2905. [[CrossRef](#)] [[PubMed](#)]
48. Grabowski, S.J. Ab Initio Calculations on Conventional and Unconventional Hydrogen Bonds—Study of the Hydrogen Bond Strength. *J. Phys. Chem. A* **2001**, *105*, 10739–10746. [[CrossRef](#)]
49. Espinosa, E.; Molins, E.; Lecomte, C. Hydrogen bond strengths revealed by topological analyses of experimentally observed electron densities. *Chem. Phys. Lett.* **1998**, *285*, 170–173. [[CrossRef](#)]
50. Grabowski, S.J.; Ugalde, J.M. Bond Paths Show Preferable Interactions: Ab Initio and QTAIM Studies on the X-H \cdots π Hydrogen Bond. *J. Phys. Chem. A* **2010**, *114*, 7223–7229. [[CrossRef](#)] [[PubMed](#)]
51. Frisch, M.J.; Trucks, G.W.; Schlegel, H.B.; Scuseria, G.E.; Robb, M.A.; Cheeseman, J.R.; Scalmani, G.; Barone, V.; Petersson, G.A.; Nakatsuji, H.; et al. *Gaussian 16, Revision A.03*, Gaussian, Inc.: Wallingford, CT, USA, 2016.
52. Møller, C.; Plesset, M.S. Note on an Approximation Treatment for Many-Electron Systems. *Phys. Rev.* **1934**, *46*, 618–622. [[CrossRef](#)]
53. Woon, D.E.; Dunning, T.H., Jr. Gaussian Basis Sets for Use in Correlated Molecular Calculations. III. The second row atoms, Al–Ar. *J. Chem. Phys.* **1993**, *98*, 1358–1371. [[CrossRef](#)]
54. Metz, B.; Stoll, H.; Dolg, M. Small-core multiconfiguration-Dirac-Hartree-Fock-adjusted pseudopotentials for post-d main group elements: Application to PbH and PbO. *J. Chem. Phys.* **2000**, *113*, 2563–2569. [[CrossRef](#)]
55. Peterson, K.A. Systematically convergent basis sets with relativistic pseudopotentials. I. Correlation consistent basis sets for the post-d group 13–15 elements. *J. Chem. Phys.* **2003**, *119*, 11099–11112. [[CrossRef](#)]
56. Piela, L. *Ideas of Quantum Chemistry*; Elsevier Science Publishers: Amsterdam, The Netherlands, 2007; pp. 684–691.

57. Boys, S.F.; Bernardi, F. The calculation of small molecular interactions by the differences of separate total energies. Some procedures with reduced errors. *Mol. Phys.* **1970**, *19*, 553–561. [[CrossRef](#)]
58. Keith, T.A. *AIMAll (Version 11.08.23)*; TK Gristmill Software: Overland Park, KS, USA, 2011. Available online: aim.tkgristmill.com.
59. Grabowski, S.J. Dihydrogen bond and X-H... σ interaction as sub-classes of hydrogen bond. *J. Phys. Org. Chem.* **2013**, *26*, 452–459. [[CrossRef](#)]
60. Becke, A.D. Density-functional exchange-energy approximation with correct asymptotic behaviour. *Phys. Rev. A* **1988**, *38*, 3098–3100. [[CrossRef](#)]
61. Perdew, J.P. Density-functional approximation for the correlation energy of the inhomogeneous electron gas. *Phys. Rev. B* **1986**, *33*, 8822–8824. [[CrossRef](#)]
62. Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. A consistent and accurate ab initio parametrization of density functional dispersion correction (DFT-D) for the 94 elements H-Pu. *J. Chem. Phys.* **2010**, *132*, 154104. [[CrossRef](#)] [[PubMed](#)]
63. Velde, G.T.E.; Bickelhaupt, F.M.; Baerends, E.J.; Guerra, C.F.; van Gisbergen, S.J.A.; Snijders, J.G.; Ziegler, T. Chemistry with ADF. *J. Comput. Chem.* **2001**, *22*, 931–967. [[CrossRef](#)]
64. Wong, R.; Allen, F.H.; Willett, P. The scientific impact of the Cambridge Structural Database: A citation-based study. *J. Appl. Crystallogr.* **2010**, *43*, 811–824. [[CrossRef](#)]

Sample Availability: Samples of the compounds are not available from the author.



© 2018 by the author. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).