



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Correction

Hyperglycemia in acute COVID-19 is characterized by insulin resistance and adipose tissue infectivity by SARS-CoV-2

Moritz Reiterer, Mangala Rajan, Nicolás Gómez-Banoy, Jennifer D. Lau, Luis G. Gomez-Escobar, Lunkun Ma, Ankit Gilani, Sergio Alvarez-Mulett, Evan T. Sholle, Vasuretha Chandar, Yaron Bram, Katherine Hoffman, Priya Bhardwaj, Phoebe Piloco, Alfonso Rubio-Navarro, Skyler Uhl, Lucia Carrau, Sean Houhgton, David Redmond, Alpana P. Shukla, Parag Goyal, Kristy A. Brown, Benjamin R. tenOever, Laura C. Alonso, Robert E. Schwartz, Edward J. Schenck, Monika M. Safford, and James C. Lo*

*Correspondence: jlo@med.cornell.edu

<https://doi.org/10.1016/j.cmet.2021.10.014>

(Cell Metabolism 33, 2174–2188.e1–e5, November 2, 2021)

It was brought to the authors' attention by readers that ARDS was not adequately defined. The STAR Methods have now been updated with specific clinical (ventilatory and oxygenation) parameters used to qualify for ARDS along with accompanying references.