


# Client perspectives on an outreach approach for HIV prevention targeting Indonesian MSM and transwomen

Adi Nugroho <sup>1,2,3,\*</sup>, Vicki Erasmus<sup>1</sup>, Sarah E. Krier<sup>4,5</sup>,  
Kusnindyah P. Reviagana<sup>2</sup>, Pipiet A. Laksmono<sup>3</sup>, Asti Widiastuti<sup>3</sup>, and  
Jan Hendrik Richardus<sup>1</sup>

<sup>1</sup>Department of Public Health, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The Netherlands, <sup>2</sup>School of Public Health, Faculty of Medicine, Lambung Mangkurat University, Banjarbaru, Indonesia, <sup>3</sup>GWL-INA Network, Jakarta, Indonesia, <sup>4</sup>Department of Infectious Diseases and Microbiology, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA and <sup>5</sup>Center for LGBT Health Research, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA, USA

\*Corresponding author. E-mail: adie.noegroho@gmail.com

## Summary

This study explored clients' perspective on an outreach approach to promote HIV testing in Indonesia targeting men who have sex with men (MSM) and transgender women (transwomen or waria). Semi-structured qualitative interviews were conducted with 32 individuals (21 MSM and 11 waria) who had received services from outreach workers (OWs) in five cities in Indonesia. Participants in this study reported positive experiences with the outreach approach and perceived OWs as their motivators in accessing HIV testing as well as HIV care and treatment. OWs provided easy-to-understand HIV information. Clients expected OWs to be well-trained and more creative in performing outreach. They perceived that the Internet and social media have helped them considerably to stay in touch with OWs. Yet, they expressed that such virtual contacts could not simply replace the face-to-face contact, especially for waria. Furthermore, clients suggested outreach to be delivered in a more appealing manner, for example through activities that may facilitate clients learning professional or life skills. They also asserted that as an HIV prevention approach, outreach needs to use more positive framing and go beyond HIV and health contents, chiefly for the youth. Future outreach programmes should facilitate OWs in providing tailored services based on the level and type of support that the clients need, and in applying varied proportion and levels of sophistication in the use of online and virtual platforms for outreach.

**Key words:** HIV prevention, outreach, MSM, transgender, qualitative study, Indonesia

## INTRODUCTION

Studies have suggested that HIV prevention strategies should apply combined approach that includes behavioural

change, biomedical strategies, antiretroviral treatment and social justice and human rights components (Coates *et al.*, 2008). Consequently, much research strives to meet

biomedical or behavioural objectives, such as reducing sexually transmitted infection (STIs), increasing consistent condom use, and improving HIV testing uptake (Hankins and de Zalduondo, 2010). In Indonesia, the national HIV prevention strategy focuses on the prevention of the sexual transmission of HIV. The strategy includes four key components: (i) mobilizing relevant stakeholders; (ii) implementing behaviour change communication; (iii) improving management and distribution of condom and lubricant and (iv) providing comprehensive treatment and care for STIs. One of the crucial elements of HIV prevention effort in Indonesia involves the provision of stigma free HIV testing services as well as promoting and improving access to routine HIV testing for populations at risk for HIV infection (National AIDS Commission of Indonesia, 2012).

Accessing routine HIV testing allows key populations to know their HIV status, and engages them in medical care as soon as they are diagnosed with HIV (Walensky *et al.*, 2007; Jain and Deeks, 2010). However, raising awareness about the importance of engaging in immediate and routine HIV testing among high-risk individuals in Indonesia remains a challenge. Men who have sex with men (MSM) and transgender women (waria in Bahasa Indonesia) are groups least reached by HIV prevention programmes. In 2013, only 38% (vs. 25% in 2009) of MSM were reported to have an HIV test (with result), 24% of MSM were reported to take an HIV test in the past 12 months. And 54% of transwomen reported ever having an HIV test (with result) (Ministry of Health Republic of Indonesia, 2013).

HIV programmes targeting these populations have been consistently conducted by an outreach approach (van Griensven and de Lind van Wijngaarden, 2010). This approach uses outreach workers (OWs) at the frontline to engage individuals who are at risk for HIV. The main task of the OWs is to connect members from the community to public health services, so that they can be reached by HIV programmes. OWs disseminate HIV prevention packages, provide safe sex education materials and support for behaviour changes, and most importantly, refer the community for HIV testing and counselling (Agüero *et al.*, 1996). The success of this approach is generally measured using two parameters, i.e. the number/percentage of individuals who are reached, and the number/percentage of individuals who receive testing of HIV (UNAIDS, 2013).

HIV prevention interventions for MSM and waria in Indonesia are implemented primarily through civil society organizations (CSOs) (WHO SEARO, 2010). When tailoring HIV intervention approaches to the characteristics of MSM and waria, every effort is made by the organizations to hire OWs who are members of these

communities as this facilitates the reaching of their peers. Several reviews found that an outreach approach is commonly used among peer-based interventions for MSM and transwomen (Simoni *et al.*, 2011; Higa *et al.*, 2013). Peer outreach facilitates the detection of those who are at risk for HIV but may not be discovered through regular health services. This, however, is labour-intensive and costly. Therefore, efforts are made to increase the effectiveness of outreach programmes (Platteau *et al.*, 2012).

The performance of outreach programmes in Indonesia is still relatively low. In 2013, a national survey showed that the outreach programmes reached only 59% of MSM and 57% of waria individuals (Ministry of Health Republic of Indonesia, 2013). This low performance might explain the unsatisfactory results of HIV testing among these two populations. By mid-2014, the programme reported 115 077 MSM and 4868 transwomen tested for HIV or around 13 and 11% of the estimated number of each population, respectively (Ministry of Health Republic of Indonesia, 2015). This low performance indicates that some aspects of HIV prevention targeting these population in Indonesia should be improved, including the outreach approach.

This study aimed to answer the following questions: (i) How do MSM and waria in Indonesia perceive an outreach approach to promote HIV testing? and (ii) What can we learn from their experiences? To this end, we explored their experiences with HIV outreach services.

## MATERIALS

### Setting

We invited participants through local CSOs that implemented outreach targeting MSM and waria groups in every city. We recruited 6–8 MSM and waria individuals from each city, who (i) had received services from OWs; (ii) were aged 18 years or older and (iii) had never worked as an OW. We identified MSM participants based on their self-reported sexual practices in the past year (i.e. ever had sex with another man in the past 12 months), and identified waria participants based on their gender identity (i.e. self-identified waria).

### Data collection

We invited potential individuals to a location for an interview. We explained the purpose of the study to all participants and asked for their consent before participation. The participants were reassured that their identity and personal details would be kept anonymous and

confidential, and that they could withdraw their participation at any time. AN, PL, and one research assistant conducted semi-structured interviews in Bahasa Indonesia using an interview guide (see Table 1). At the end of the session, the participants received IDR 120 000 (~USD 10) as compensation for their time and transportation costs.

### Data management and analysis

All interviews were audio-recorded and transcribed verbatim by a trained transcriber. AN and PL read and checked all transcripts to ensure understanding and completeness. The study data were extracted from the transcripts.

Data analysis employed grounded theory approach (Corbin and Strauss, 1990). All completed transcripts were uploaded into NVivo 10 (QSR, Cambridge, MA) and textual data were coded into themes and categories inductively. At first, two coders (AN and KR) separately coded and analysed one set of transcripts to generate the initial codes independently and to identify the potential themes and categories. Results from both coders were then matched to find common themes and categories. In cases where the coders disagreed in their identification of the different themes, they examined and consolidated these differences through discussions. Subsequently, the coders continued a separate coding on another set of transcripts and discussed the new emerging themes and categories as well as the differences they found. The agreed themes and categories were then used for analysing the remaining set of transcripts.

Specific quotes were selected for use in this paper and translated into English by two bilingual translators. The two translators deliberated the differences between their translations and constructed a final English version. As an extra control, a third person translated the final English version back into Bahasa Indonesia.

**Table 1:** Guidance questions for focus group discussion

Topics	Guidance questions
Contact with OW	How do you usually get contacted by an OW? Do you feel comfortable to be contacted that way? If not, in which way do you think will make you feel more comfortable? How often did the OW contact you in the last 3 months? Do you feel satisfied with that number of contact?
Information/services received from OW	What information or services did you ever receive from the OW? Do you find the provided information or services beneficial for you? If not, what kind of information do you expect to be provided?

OW, outreach worker.

## RESULTS

### Study participants

A total of 32 individuals participated in the interviews; 21 were MSM and 11 were waria (see Table 2). The participants' age ranged from 19 to 55 years with almost a half (47%) were under 25 years; 34% were 25–34 years and 19% were 35 years or older. Without being prompted, three participants (2 MSM and 1 waria) independently disclosed their HIV-positive status during the interview.

### Themes and categories

We grouped the emerging themes into categories as follows: experience with outreach; HIV testing service and uptake and promotion strategy and media use.

### Experience with outreach

#### Contact with OWs

Most participants' first contact with an OW occurred at a common hangout and/or cruising place for MSM and

**Table 2:** Characteristics of study participants

Characteristics	N (32)	% (100.0)
Risk group		
MSM	21	65.6
Waria	11	34.4
Age group		
≤24 years	20	62.5
25+ years	12	37.5
City		
Medan	8	25.0
Jakarta	6	18.8
Bandung	6	18.8
Yogyakarta	6	18.8
Surabaya	6	18.8

MSM, men who have sex with men; Waria, Indonesian term for transwomen.

waria, such as city square, club or sex work area. Other first contacts happened through either Internet-based messaging or social media.

I went to a discotheque, and ran into A, R, R, and D [OWs]. They chatted with me and gave me information about HIV. I was interested, and then asked for A's contact details. I told him that I would like to learn more.  
(M, young MSM, Jakarta)

When we were hanging out, sister S approached us, and we've been friends since then. Together, we initiated 'arisan' which is now considered as a model for community-based circle for waria.  
(R, young waria, Surabaya)

Some participants first contacted an OW mainly after searching for gay-related information or after having experienced symptoms that appeared to be HIV-related. Follow-up contacts usually occurred weekly or monthly and most often took place in the participants' home or via the Internet. A common method of follow up among the waria participants was the so-called 'arisan', a periodic social gathering held by members of a group. The gatherings have an 'edutainment' character, where information is shared in an informal way, while at the same time there is also entertainment e.g. in the form of a raffle.

In our weekly 'arisan', health-related information is always shared, about edutainment events, condoms and other topics.  
(R, waria, Surabaya)

MSM participants regularly visiting cruising areas reported that they often received new insights, free condoms and lubricants from OWs every time they came to the area. The participants also reported that they usually received reminders about a gathering and/or mobile clinic through Blackberry messenger (BBM). This service was also often used whenever they needed to request a personal consultation from an OW.

I am usually in contact [with the OW] through SMS (short message service) or BBM, or if there is any gathering or something like that.  
(A, MSM, Bandung)

I often receive BBM texts, every month [reminder] for STI check or every three months for HIV testing.  
(R, waria, Medan)

Participants who worked as sex workers reported consistent condom use during sex with their clients. However, the consistent condom use became problematic during sex with their steady partners, both among MSM and waria participants. Participants who openly indicated their HIV status seemed to be aware about the

risk of reinfection if they had unprotected sex with individuals diagnosed with HIV.

They [waria sex worker] are good in negotiating condom use with their clients. They can be persistent but not so much when it comes to sex with their boyfriend, including me myself.  
(R, waria, Jakarta)

If we have sex without condom and our sex partner turns out HIV-positive, while I am also HIV-positive, once his semen get into my body, that would be a loss of my own ...  
(Y, MSM, Jakarta)

### Services received from outreach

Participants reported that OWs delivered basic HIV information, including information on HIV transmission, prevention options and HIV services available. Participants with HIV also reported that OWs helped them access the national health scheme they need for HIV care and treatment. Some participants indicated that OWs also offered services through their CBO that went beyond HIV, namely professional skills workshops and social activities.

I learned about STIs and about how to deal with the health system which was very useful because I now can have my CD4 count checked. I have access to anti-retroviral meds and I always get informed whenever there is a free CD4 check.  
(I, HIV-positive waria, Jakarta)

I was involved in many activities that HA [a local CBO] conducted, and have obtained a lot of information, not only about HIV, but also about STIs, and other knowledge such as management, public speaking, even cooking classes.  
(A, MSM, Bandung)

Participants commonly perceived the OW as a friend who provides HIV information in a safe environment in an understandable way. OWs were also considered as one of their motivators in accessing HIV care for testing. Once the relationship was established, the clients seemed to demand services ranging from obtaining information on HIV, testing reminders, to assistance for accessing care, support and treatment. The waria indicated that getting along with an OW and being attached to a CBO—where the OW worked or was affiliated with could make them feel more respected.

If there is no OW ... we will be puzzled every time we get ill, while to find information at hospital will be impossible. We might end up asking friends who are not necessarily capable of giving us the right information. It

[having an OW] isn't just about having a source of information, but they also provide us a safe space.

(Y, MSM, Yogyakarta)

One of the benefits is that they [OWs] can 'save' us ... moreover, we as waria have always been deemed as disgraceful persons. So, if people can see that we belong to a sort of organization, they may perceive us better.

(V, waria, Medan)

### Attitudes towards outreach

Most participants felt comfortable when approached by OWs who were peers—those who shared the same age, gender or sexuality—especially waria. This was especially the case when waria were approached by waria. The benefits of outreach activities mentioned by the participants include being able to learn about HIV and AIDS, sharing information with peers, and having a constant reminder regarding HIV testing schedules.

Their services are quite good, really .... We often get reminded to get tested [for HIV]. We also often receive broadcasted messages through BBM for VCT (voluntary counseling and testing) schedules.

(R, MSM, Surabaya)

Because I am a waria, I am more comfortable with fellow waria ... I love sharing with sister R [an OW waria], she seems knowledgeable. Besides, she isn't an arrogant person, so I can easily ask questions about anything.

(D, waria, Medan)

Some respondents indicated that there is an assumption that only those infected by HIV can be hired as OWs. To some, if an OW is known to be diagnosed with HIV, it could cause HIV prevention messages to be unconvincing. Scepticism about younger OWs was also raised by an older participant.

When one OW gets infected with HIV, the community won't trust him anymore. They [OWs] always advise us [for condom use] but they themselves got infected. [To be an OW] A person should be really committed and truly capable of doing the job. Although I believe there is a sort of pre-training, some people still will say like 'that kid gets hired as an OW, really? He is just a kid! ...' then he gets ignored.

(T, MSM, Yogyakarta)

Some participants expected to obtain up-to-date information and new skills regarding HIV when they attended an outreach gathering. However, this expectation was not always met, making some participants feel like they had wasted their time.

Some gatherings were held without any clear agenda or specific activity ... so sometimes I felt like I came for nothing.

(R, MSM, Bandung)

On the other hand, some participants felt bored with outreach meetings that solely focused on HIV. They expected various activities that were more appealing and offered deeper insights. They expected to get information not only related to HIV, condoms, and STIs, but also about sexual and reproductive health in general.

They are poor in approaching people because they just keep on talking about VCT, HIV gatherings, and whatever ... they should probably create more appealing events ... so people would be interested in joining.

(D, MSM, Surabaya)

I would suggest that they try and find new topics ... every time there is an HIV gathering coming up, my friends often said 'don't bother to come, they always talk about the same thing, it's just boring!'

(S, waria, Yogyakarta)

### HIV testing services and uptake

#### Personal testing barriers

MSM participants pointed out that many men only came to an HIV clinic when they got ill, regardless of the availability of information about the benefit of knowing their HIV status early and about the locations of friendly clinics. They also stated that shame and fear were the most common causes preventing them from accessing HIV services.

What I heard from most people is that they are afraid to find out if they are HIV-positive. Because they think their life will end soon. Many people said like that.

(M, MSM, Jakarta)

They actually want to [get HIV tested] but most of them are too shy for that. They are scared people will notice if they turn out to be HIV-positive or to have an STI.

(A, MSM, Bandung)

Most participants indicated coming to HIV clinic for their subsequent HIV test unaccompanied because they are aware about their risk of HIV. Some having steady partners reported that they also encouraged their partners to get HIV tested.

I did my second and third [HIV] test because of my self-awareness, VCT is supposed to be done voluntarily, right? ... I did my last test together with my boyfriend.

(R, waria, Surabaya)

#### HIV testing services

Some participants suggested that provision of an HIV clinic directory with a geo-location feature would be helpful. The directory should feature comprehensive information about HIV services in each clinic. This would enable them to select HIV care which is accessible and

suits their needs and comfort level. They also recommended conducting more mobile HIV testing at places where MSM and waria hang out. For STI checks, waria indicated a preference for being examined by male doctors or nurses.

We need information about procedures of VCT and where their locations are.  
(H, MSM, Medan)

I wish we could do the VCT here [at our place] so we don't have to go to the clinic. It would be better if the clinic staff come to our place so all waria can just gather around [for testing].  
(F, waria, Yogyakarta)

Waria should be examined by a male staff [doctor/medic], not female. I don't feel awkward with men, but I do with women. I think male staff know us better . . .  
(M, waria, Bandung)

The participants wished HIV service providers to be more stigma free. Some of them indicated negative experiences they had received from staff at public clinics (which were generally less gay friendly than community-based service clinics) because of their sexual orientation. One participant pointed out that some MSM would rather access HIV testing at public clinics than at community-based service clinics as it was more anonymous.

I forget which *puskesmas* [primary health center] exactly. I once took a friend to this *puskesmas*, and was asked [by a clinic-staff] why I came and why I wanted to get tested. As soon as I said I am gay, the staff preached to me that I should repent [from being gay] so I won't get infected.  
(M, young MSM, Jakarta)

My ex-boyfriend is a very discreet person. We are totally the opposite of each other. He didn't feel comfortable coming to GN [a local CBO] for HIV testing, so we went to *puskesmas* instead. But when I get HIV tested myself, I'd rather go to GN.  
(D, MSM, Surabaya)

They also mentioned the inconvenient opening hours of the services, particularly for waria and MSM who were office employees or students.

We wanted to bring our friend for [HIV] testing but no clinics were available because their office hours coincided with operational hours of the clinics. Clinics are open for service only until 12 or 1 PM.  
(A, MSM, Surabaya)

One thing that I find inconvenient is about the opening hours. It's too early for us, we are still in bed. (J, waria, Surabaya)

### Promotion strategy and media use

The participants expected the availability of a wider choice of media to keep themselves updated with the latest HIV information.

Maybe, a specific website which offers many kinds of information. Do more promo activities in many channels such as newspaper, radio, and so on, so more people would notice. Also, you might want to do a poll to check if the information provided is satisfactory. The name of the website should be easy to remember . . .  
(A, MSM, Bandung)

The other needs mentioned included edutainment activities, provision of information centres, and the need to present negative or fear-based messages in a more positive light.

Speaking as a young people, please do not scare us . . . I think it's better to show us some role models of persons who live with HIV and still can perform their daily life as well as other [non-HIV-infected] people. That would be more encouraging than only frightening us.  
(M, young MSM, Jakarta)

Respondents reflected that outreach via the Internet will not replace face-to-face contact. This type of contact was still considered useful for avoiding miscommunication. In addition, not all MSM and waria used the Internet and social media frequently.

I feel more comfortable asking questions in person, so we get a clear explanation about everything we ask. That's what is lacking from the Internet, we might misunderstand things . . .  
(A, MSM, Yogyakarta)

I feel more comfortable talking face-to-face [with an OW]. The internet has a very sophisticated language, and many of them are in English, while I don't understand English like at all. It is better to meet in person. Also, using the Internet costs mobile data. I bet most people would feel reluctant . . .  
(J, waria, Surabaya)

## DISCUSSION

Our results indicate that the outreach clients in this study generally have a positive attitude towards the outreach approach promoting HIV testing. They perceived OWs as their motivators in accessing HIV testing as well as HIV care and treatment. OWs also served as persons providing understandable HIV information. Clients expected OWs to be well-trained and more creative in performing outreach. They thought that the Internet and

social media have helped them stay in touch with OWs. Yet, they expressed that such virtual contacts could not simply replace the face-to-face contact, especially for waria. Furthermore, clients suggested that outreach should be delivered in a more appealing manner, for example through activities that may facilitate clients learning some professional or life skills. They emphasized that outreach needs to use more positive framing and go beyond HIV and health contents, especially for the youth.

Our findings support the studies that have revealed HIV message fatigue among high-risk populations (Stockman *et al.*, 2004; Rowniak, 2009; Thomas *et al.*, 2012). This emphasizes the importance of developing new outreach programme strategies. For example, it should go beyond condom promotion and address the diverse health and social needs of MSM and waria in Indonesia. Our data further indicate that incorporating HIV information into a non-HIV social-gathering activity is useful, primarily for waria individuals who form a socially cohesive group (Saggurti *et al.*, 2013; Shaikh *et al.*, 2016). Others have recommended organizing activities that can facilitate bonding and empowerment among the two groups (Parker, 1996; Beeker *et al.*, 1998; Evans *et al.*, 2010), which we call 'edutainment' activities. The MSM participants in our study also highlighted their keen interest in activities which allow them to gain professional skills.

In line with our findings, other studies indicated that MSM and transwomen preferred not to know their HIV status until they experienced illness (Safren *et al.*, 2006). Furthermore, studies have shown that predominant barriers to HIV testing include the fear of being diagnosed with HIV and fear of discrimination due to positive testing result (Risher *et al.*, 2013; Zhang *et al.*, 2013; Goncalves *et al.*, 2016). Therefore, it is crucial to formulate appropriate messages that address such barriers. Consistent with a study from India (Thomas *et al.*, 2012), our findings recommend promoting more positive images of healthy HIV-positive individuals to eliminate some fear-based negative messages about living with HIV. Such negative messages only discourage people from testing, primarily among younger MSM.

In line with the results of a study from South Africa (Tucker *et al.*, 2013), this study reveals that it is crucial for OWs to gain the trust of their clients. As soon as the trust is gained, it is more likely that the reached clients would embrace services offered by OWs and stay in contact for follow up. Moreover, although OWs' primary task in Indonesia is to get their clients tested for HIV, the participants reported that some OWs provided services beyond testing. We found that some clients who

were diagnosed with HIV were also helped by their OWs in accessing HIV care and treatment. Similarly, a previous study highlighted OWs' significant role in engaging and retaining clients living with HIV in care (Rajabiun *et al.*, 2007). This indicates the importance for OWs to be knowledgeable in HIV case management, and for future outreach programmes in Indonesia to revisit the job description assigned for OWs.

Some participants in this study indicated that they initiated contact with an OW because of information they had received via the Internet. Other studies showed that the Internet can help HIV programmes reach a larger number of individuals (Zou *et al.*, 2013; Brennan *et al.*, 2015). The Internet also seems to help clients stay connected with their OWs and keep them updated with information about HIV services.

### Strengths and limitations

This study provides information about the strengths and limitations of an outreach approach to improving HIV testing targeting MSM and transgender women in Indonesia. The study employs the clients' perspective to evaluate an outreach approach for HIV testing that targets the two vulnerable groups. Although the number of participants in our study is relatively small, it is sufficient for a qualitative analysis. Therefore, our findings offer evidence-based information of two hard-to-reach groups that can be used to improve approaches targeting these populations. This study has some limitations. First, all interviews were conducted in an urban setting. Thus, the findings might not be applicable to rural settings. Second, we applied non-probabilistic sampling. Therefore, generalizing the findings outside our study sample should be done with caution. Third, we recruited participants through an organization implementing outreach, which means that the participants generally have a good relationship with the organization. Such a method may lead to social desirability bias even though we emphasized that all information would be kept anonymous during all interviews. Nevertheless, most participants appeared to express openly their opinions and experiences. Fourth, the political and social situations change rapidly in the country and our findings may reflect a situation that has changed by now. Nonetheless, we believe that the data present worthwhile lessons learned for future programme improvement.

### Implications for practice

This study has revealed several elements that should be considered when improving future outreach as a public health programme. First, managing performance,

through human resources management, by recruiting OWs who share characteristics of their target populations, ensure the provision of tailored training programme for the OWs that build on their knowledge and skills in outreach, and rigorous monitoring and evaluation upon their performances. Having a peer-to-peer training package available for the outreach team may also be useful for maintaining OWs' capacity in performing outreach (Nugroho *et al.*, 2018). The second is innovation. As outreach is a form of communication strategy, it should always progress following the evolution of how people communicate, which may include a mix of physical and virtual communications. OWs should aim applying a client-centred approach by providing tailored services based on the level and type of support clients need (Grimsrud *et al.*, 2016). This could also be performed by applying different generations of outreach for different clients, i.e. make use online and virtual platforms in accordance to portion and levels of sophistication of clients' use (FHI 360-LINKAGES Project, 2019). The third is partnership with relevant stakeholders. Apart from comprehensively refining the outreach approach, structural barriers at HIV services and care should continuously be tackled by, for example, (i) providing accessible stigma-free HIV services and care and (ii) providing various types of HIV services that may suit diverse clients' characteristics (Differentiated Service Delivery, 2018).

## ETHICAL APPROVAL

We asked for consent from all participants in this study prior to their participation. Participants joined this study at their own will and could withdraw any time during the study process. The Research Ethics Board of Atma Jaya University Indonesia Jakarta approved this study with registration number: 504/III/LPPM-PM.10.05/06/2014.

## ACKNOWLEDGEMENTS

The authors would like to express a sincere gratitude to all the participants in this study for their time and openness to share their experience. We highly appreciate the support and assistance from Rosidin M. Al Harbi during the data collection process. We also gratefully thank Dr Alfi Yasmina for her assistance in co-translating the quotes for the manuscript and Dr Sandra O. Wesselink for her assistance in data analysis.

## FUNDING

Costs for conducting all interviews and transcribing the audio-recording were funded by the Global Fund to fight AIDS, TB

and Malaria (GF-ATM) Round 10 grant through a regional grant to the ISEAN Hivos Program. GWL-INA Network is a Sub Recipient of ISEAN Hivos Program in Indonesia. The funding had no role in study design, data analysis or preparation of the manuscript.

## REFERENCES

- Agüero, L. K. W.-D., Gorsky, R. D. and Seeman, G. M. (1996) Cost of outreach for HIV prevention among drug users and youth at risk. *Drugs & Society*, **9**, 185–197.
- Becker, C., Guenther-Grey, C. and Raj, A. (1998) Community empowerment paradigm drift and the primary prevention of HIV/AIDS. *Social Science & Medicine*, **46**, 831–842.
- Brennan, D. J., Lachowsky, N. J., Georgievski, G., Rosser, B. R., MacLachlan, D., Murray, J. *et al.* (2015) Online outreach services among men who use the Internet to seek sex with other men (MISM) in Ontario, Canada: an online survey. *Journal of Medical Internet Research*, **17**, e277.
- Coates, T. J., Richter, L. and Caceres, C. (2008) Behavioural strategies to reduce HIV transmission: how to make them work better. *The Lancet*, **372**, 669–684.
- Corbin, J. M. and Strauss, A. (1990) Grounded theory research: procedures, canons, and evaluative criteria. *Qualitative Sociology*, **13**, 3–21.
- Differentiated Service Delivery. (2018) *A Decision Framework for HIV Testing Services*. <http://www.differentiatedcare.org/> (last accessed 14 May 2019).
- Evans, C., Jana, S. and Lambert, H. (2010) What makes a structural intervention? Reducing vulnerability to HIV in community settings, with particular reference to sex work. *Global Public Health*, **5**, 449–461.
- FHI 360-LINKAGES Project. (2019) *A Vision for Going Online to Accelerate the Impact of HIV Programs*. FHI 360, Washington, DC.
- Goncalves, V. F., Kerr, L. R., Mota, R. S., Macena, R. H., Almeida, R. L., Freire, D. G. *et al.* (2016) Incentives and barriers to HIV testing in men who have sex with men in a metropolitan area in Brazil. *Cadernos de Saúde Pública*, **32**, doi: 10.1590/0102-311x00049015.
- Grimsrud, A., Bygrave, H., Doherty, M., Ehrenkranz, P., Ellman, T., Ferris, R. *et al.* (2016) Reimagining HIV service delivery: the role of differentiated care from prevention to suppression. *Journal of the International AIDS Society*, **19**, 21484.
- Hankins, C. A. and de Zalduondo, B. O. (2010) Combination prevention: a deeper understanding of effective HIV prevention. *AIDS*, **24**(Suppl. 4), S70–S80.
- Higa, D. H., Crepaz, N., Marshall, K. J., Kay, L., Vosburgh, H. W., Spikes, P. *et al.* (2013) A systematic review to identify challenges of demonstrating efficacy of HIV behavioral interventions for gay, bisexual, and other men who have sex with men (MSM). *AIDS and Behavior*, **17**, 1231–1244.
- Jain, V. and Deeks, S. G. (2010) When to start antiretroviral therapy. *Current HIV/AIDS Reports*, **7**, 60–68.
- Ministry of Health Republic of Indonesia. (2013) *Fact Sheet Integrated Biological and Behavioural Survey (IBBS): High*



- Risk Group in Indonesia, 2013*. <http://www.slideshare.net/ditkeswa/finalstbp-totaldraft13-januari1> (last accessed 10 December 2018).
- Ministry of Health Republic of Indonesia. (2015) *National Action Plan of HIV Response 2015–2019*. [http://www.aidsdatahub.org/dmdocuments/IBBS\\_2011\\_Report\\_Indonesia.pdf](http://www.aidsdatahub.org/dmdocuments/IBBS_2011_Report_Indonesia.pdf) (last accessed 10 December 2018).
- National AIDS Commission of Indonesia. (2012) *National Strategy and Action Plan of HIV and AIDS Response Year 2010–2014*. [https://www.ilo.org/wcmsp5/groups/public/—ed\\_protect/—protrav/—ilo\\_aids/documents/legaldocument/wcms\\_173075.pdf](https://www.ilo.org/wcmsp5/groups/public/—ed_protect/—protrav/—ilo_aids/documents/legaldocument/wcms_173075.pdf) (last accessed 10 December 2018).
- Nugroho, A., Coulter, R. W. S., Erasmus, V., Laksmono, P. A., Mihari, T. S. and Richardus, J. H. (2018) Evaluation of a training aimed at building capacity for outreaching to men who have sex with men and transgender women in Indonesia. *Health Education Research*, **34**, 223–233.
- Parker, R. G. (1996) Empowerment, community mobilization and social change in the face of HIV/AIDS. *AIDS*, **10**(Suppl. 3), S27–S31.
- Platteau, T., Wouters, K., Apers, L., Avonts, D., Nostlinger, C., Sergeant, M. *et al.* (2012) Voluntary outreach counselling and testing for HIV and STI among men who have sex with men in Antwerp. *Acta Clinica Belgica*, **67**, 172–176.
- Rajabiun, S., Mallinson, R. K., McCoy, K., Coleman, S., Drainoni, M. L., Rebholz, C. *et al.* (2007) “Getting me back on track”: the role of outreach interventions in engaging and retaining people living with HIV/AIDS in medical care. *AIDS Patient Care STDS*, **21**(Suppl. 1), S20–S29.
- Risher, K., Adams, D., Sithole, B., Ketende, S., Kennedy, C., Mnisi, Z. *et al.* (2013) Sexual stigma and discrimination as barriers to seeking appropriate healthcare among men who have sex with men in Swaziland. *Journal of the International AIDS Society*, **16**(3 Suppl. 2), 18715.
- Rowniak, S. (2009) Safe sex fatigue, treatment optimism, and serosorting: new challenges to HIV prevention among men who have sex with men. *Journal of the Association of Nurses in AIDS Care*, **20**, 31–38.
- Safren, S. A., Martin, C., Menon, S., Greer, J., Solomon, S., Mimiaga, M. J. *et al.* (2006) A survey of MSM HIV prevention outreach workers in Chennai, India. *AIDS Education and Prevention*, **18**, 323–332.
- Saggurti, N., Mishra, R. M., Proddutoor, L., Tucker, S., Kovvali, D., Parimi, P. *et al.* (2013) Community collectivization and its association with consistent condom use and STI treatment-seeking behaviors among female sex workers and high-risk men who have sex with men/transgenders in Andhra Pradesh, India. *AIDS Care*, **25**(Suppl. 1), S55–S66.
- Shaikh, S., Mburu, G., Arumugam, V., Mattipalli, N., Aher, A., Mehta, S. *et al.* (2016) Empowering communities and strengthening systems to improve transgender health: outcomes from the Pehchan programme in India. *Journal of the International AIDS Society*, **19**, 20809.
- Simoni, J. M., Nelson, K. M., Franks, J. C., Yard, S. S. and Lehavot, K. (2011) Are peer interventions for HIV efficacious? A systematic review. *AIDS and Behavior*, **15**, 1589–1595.
- Stockman, J. K., Schwarcz, S. K., Butler, L. M., de Jong, B., Chen, S. Y., Delgado, V. *et al.* (2004) HIV prevention fatigue among high-risk populations in San Francisco. *Journal of Acquired Immune Deficiency Syndromes*, **35**, 432–434.
- Thomas, B., Mimiaga, M. J., Mayer, K. H., Closson, E. F., Johnson, C. V., Menon, S. *et al.* (2012) Ensuring it works: a community-based approach to HIV prevention intervention development for men who have sex with men in Chennai, India. *AIDS Education and Prevention*, **24**, 483–499.
- Tucker, A., de Swardt, G., Struthers, H. and McIntyre, J. (2013) Understanding the needs of township men who have sex with men (MSM) health outreach workers: exploring the interplay between volunteer training, social capital and critical consciousness. *AIDS and Behavior*, **17**(Suppl. 1), S33–S42.
- UNAIDS. (2013) *HIV in Asia and the Pacific*. [http://www.unaids.org/sites/default/files/media\\_asset/2013\\_HIV-Asia-Pacific\\_en\\_0.pdf](http://www.unaids.org/sites/default/files/media_asset/2013_HIV-Asia-Pacific_en_0.pdf) (last accessed 10 December 2018).
- van Griensven, F. and de Lind van Wijngaarden, J. W. (2010) A review of the epidemiology of HIV infection and prevention responses among MSM in Asia. *AIDS (London, England)*, **24**(Suppl. 3), S30–S40.
- Walensky, R. P., Freedberg, K. A., Weinstein, M. C. and Paltiel, A. D. (2007) Cost-effectiveness of HIV testing and treatment in the United States. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*, **45**(Suppl. 4), S248–S254.
- WHO SEARO. (2010) *HIV/AIDS Among Men Who have Sex With Men and Transgender Populations in South-East Asia*. [http://apps.searo.who.int/PDS\\_DOCS/B4568.pdf?ua=1](http://apps.searo.who.int/PDS_DOCS/B4568.pdf?ua=1) (last accessed 10 December 2018).
- Zhang, L., Xiao, Y., Lu, R., Wu, G., Ding, X., Qian, H-Z. *et al.* (2013) Predictors of HIV testing among men who have sex with men in a large Chinese city. *Sexually Transmitted Diseases*, **40**, 235–240.
- Zou, H., Wu, Z., Yu, J., Li, M., Ablimit, M., Li, F. *et al.* (2013) Internet-facilitated, voluntary counseling and testing (VCT) clinic-based HIV testing among men who have sex with men in China. *PLoS One*, **8**, e51919.