

# Evaluation of the SF-36 questionnaire for assessment of the quality of life of endometriosis patients undergoing treatment: A systematic review and meta-analysis

ROMINA-MARINA SIMA<sup>1,2\*</sup>, LIANA PLEȘ<sup>1,2\*</sup>, BOGDAN SOCEA<sup>3,4</sup>, PANAGIOTIS SKLAVOUNOS<sup>5</sup>, IONUȚ NEGOI<sup>3,6</sup>, ANCA-DANIELA STĂNESCU<sup>1,2</sup>, IOAN-IULIAN IORDACHE<sup>5</sup>, BASHAR HAJ HAMOUD<sup>5</sup>, MARC PHILLIP RADOSA<sup>7</sup>, INGOLF JUHASZ-BOESS<sup>5</sup>, ERICH FRANZ SOLOMAYER<sup>5</sup>, MIHAI C.T. DIMITRIU<sup>1,8</sup>, CĂTĂLIN CÎRSTOVEANU<sup>9,10</sup>, DRAGOȘ ȘERBAN<sup>3,11</sup> and JULIA CAROLINE RADOSA<sup>5</sup>

<sup>1</sup>Department of Obstetrics and Gynecology, ‘Carol Davila’ University of Medicine and Pharmacy, 020021 Bucharest;

<sup>2</sup>Department of Obstetrics and Gynecology, ‘Sf. Ioan’ Hospital, Bucur Maternity, 040294 Bucharest;

<sup>3</sup>Department of Surgery, ‘Carol Davila’ University of Medicine and Pharmacy, 020021 Bucharest; <sup>4</sup>Department of Surgery,

‘Sf. Pantelimon’ Emergency Hospital, 021659 Bucharest, Romania; <sup>5</sup>Department of Gynecology, Obstetrics and Reproductive Medicine, Saarland University Hospital, D-66421 Homburg, Germany; <sup>6</sup>Department of Surgery,

Clinical Emergency Hospital of Bucharest, 014461 Bucharest, Romania; <sup>7</sup>Department of Gynecology, University Hospital Leipzig, D-04103 Leipzig, Germany; <sup>8</sup>Department of Obstetrics and Gynecology,

‘Sf. Pantelimon’ Emergency Hospital, 021659 Bucharest; <sup>9</sup>Department of Pediatrics,

‘Carol Davila’ University of Medicine and Pharmacy, 020021 Bucharest; <sup>10</sup>Pediatrics Department,

‘Maria Sklodowska Curie’ Emergency Children Clinical Hospital, 041451 Bucharest;

<sup>11</sup>Department of Surgery, University Emergency Hospital, 050098 Bucharest, Romania

Received July 16, 2021; Accepted August 16, 2021

DOI: 10.3892/etm.2021.10718

**Abstract.** Endometriosis has a negative influence on the physical, psychological, and social aspects of a patient's life; therefore, it affects the health-related quality of life (HRQoL). The current review aimed to investigate the efficiency of a 36-item generic questionnaire survey (SF-36) for patients with endometriosis who were undergoing medical or surgical treatment. A search strategy including the key words ‘endometriosis’, ‘quality of life’ (QOL), and ‘questionnaire SF-36’ was applied using the PubMed/MEDLINE, EMBASE, and Cochrane databases in order to include articles that evaluated the QOL among women with endometriosis using the SF-36. Only articles that included interviews of patients both before and after surgical or medical endometriosis treatment or those articles that compared study groups were considered. The qualitative analysis was based on 37 articles, whereas the quantitative analysis utilized 14 articles.

The research participants included 11,101 women, among whom 6,888 patients were diagnosed with endometriosis. The analysis recorded 17 studies dealing with all types of endometriosis, 9 studies dealing with deep infiltrative endometriosis (DIE), and 9 studies dealing with bowel endometriosis or DIE with bowel involvement. QOL was evaluated using only SF-36 in 12 studies that collectively included 1,912 women and using SF-36 in association with other questionnaires in 25 studies that collectively included 8,022 women. For patients with endometriosis, physical functioning [odds ratio (OR), 78.87; 95% confidence interval (CI), 68.97-88.77;  $I^2=98.77\%$ ;  $P\leq 0.001$ ] was the most affected life parameter. This parameter showed the highest improvement after surgical intervention (OR, 63.39; 95% CI, 48.71-78.07;  $I^2=97.65\%$ ;  $P\leq 0.001$ ) or hormonal treatment (OR, 38.65; 95% CI, 14.39-62.91;  $I^2=38.65\%$ ;  $P\leq 0.001$ ). The 36-item survey generic questionnaire seems to be an efficient tool for assessment of the QOL of life of women with endometriosis who are undergoing surgical or medical treatment. It can be applied before and after the procedure, and it can also be used for comparing study groups.

*Correspondence to:* Dr Bogdan Socea, Department of Surgery, ‘Carol Davila’ University of Medicine and Pharmacy, 37 Dionisie Lupu Street, 020021 Bucharest, Romania  
E-mail: bogdan.socea@umfcd.ro

\*Joint senior authorship

**Key words:** endometriosis, quality of life, SF-36, endometriosis therapy, laparoscopy

## Contents

1. Introduction
2. Research methods
3. Results of the study research
4. Discussion

## 1. Introduction

Endometriosis is a condition resulting from ectopic endometrial cell implants that grow progressively and determine an inflammatory response (1). The ovaries are the most common locations for endometriotic lesion formation, followed by the anterior and posterior cul-de-sac, the posterior surface of the broad ligaments, the uterosacral ligaments, the serosa of the uterus, the fallopian tubes, the sigmoid colon, the appendix, and the round ligament (2). The therapeutic options are based on clinical presentation, including infertility, pain, or mass; they are individualized according to disease severity, patient age, and reproductive desires while considering the medication side effects, costs and morbidity of the surgical procedures (3,4).

Endometriosis negatively affects the physical, psychological, and social aspects of women's lives (5), and therefore has a great impact on the health-related quality of life (HRQoL). It is difficult to characterize the quality of life (QOL) among specific patient groups because it is a broad concept that includes life satisfaction, good health, education, personal and family safety, adequate housing, employment, interrelationships, and leisure pursuits (6). Globally, in the scientific literature, QOL is evaluated using a multitude of validated and non-validated questionnaires, such as the 36-item survey generic questionnaire (SF-36). It includes eight conceptual health domains: General Health (GH), Physical Functioning (PF), Bodily Pain (BP), Role Physical (RP), Vitality (VT), Social Functioning (SF), Mental Health (MH), and Role Emotional (RE); these are summarized in the physical (PCS) and mental (MCS) component scores (7).

Recently, a number of original studies and systematic reviews have evaluated the impact of endometriosis on patient QOL (8-10). Studies have also reported the impact of different therapeutic approaches on the QOL of patients with endometriosis (11). A systematic review detailed the quality of sexual life after endometriosis-targeted laparoscopic surgery (12). Another recent meta-analysis showed that endometriosis-targeted surgical resection can improve the major domains of QOL (13). However, systematic reviews that evaluate the impact of surgical and non-surgical treatments on the QOL of patients with endometriosis are scarce.

The main objective of the present review was to investigate the efficiency of questionnaire SF-36 in assessing the QOL of patients with endometriosis who were undergoing medical or surgical treatment.

## 2. Research methods

We utilized the 'Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)' guideline to conduct this study' (14).

*Data sources and search strategies.* A combination of key word terms including 'endometriosis', 'quality of life' and 'questionnaire SF-36' were utilized for the literature research conducted in October 2018. The utilized databases included PubMed/Medline, Embase, and Cochrane; some other additional sources and references of the identified articles were also utilized. The search strategy that was utilized for PubMed/MEDLINE is outlined as follows.

*Study selection.* Articles that evaluated the quality of life (QOL) and used the SF-36 to assess women with endometriosis were evaluated. Articles that interviewed patients both before and after endometriosis-related surgical or medical treatment were included in our analysis. We also utilized the reports of women with endometriosis that were compared with a control group. Studies that focused on any type of endometriosis and were published in English, German, or French were considered in our analysis. There were no restrictions based on publishing year, journal, author, or institution.

*Data extraction.* All the selected studies were independently evaluated by two reviewers. Any disagreements regarding the articles were solved by the third more experienced author and the senior authors. The data were extracted in tables that consisted of first author's name, publication year, study type, number of initial participants, treatment type, presence of a control group or lack thereof, endometriosis type, treatment, follow-up, and the final number of participants.

*Quality assessment.* To evaluate the quality of the included studies, the modified Newcastle-Ottawa Scale (NOS) was applied (15). This scale helped to assess the risk of bias in the remaining studies. The studies were classified as 'good' when they fulfilled at least 70% of the 12 criteria, 'fair' when they fulfilled at least 50%, and 'poor' when they fulfilled less than 50%.

*Statistical analysis.* Since we observed between-study heterogeneity, we applied random-effects meta-analyses, using odds ratios (OR) and 95% confidence intervals (95% CI). The results were presented as pooled prevalence values. Studies that reported data by comparing various endometriosis types were included in the data for the most severe, whereas studies that dealt with different types of surgical treatment were considered the most difficult interventions. Statistical analysis was performed using the Open-Meta analyst [CEBM Brown, open-source, OpenMetaAnalyst for Windows 10 (64-bit)].

In our study, statistical results were calculated as follows. The first report was based on a comparison between patients with endometriosis and a control group (healthy women); the second considered surgical treatment, and the third group considered hormonal treatment.

## 3. Results of the study research

*Study research.* In accordance with our research plan, we initially identified the records of 657 articles that corresponded to our search strategy. Furthermore, 25 articles were sourced generated from additional sources (cross references). Any duplicates were removed, and the final database consisted of 345 articles. Excluding some articles due to language issues, we obtained reviews, case reports, or abstracts from 127 records for screening. Among these, 55 studies were excluded based on the inclusion criteria. Finally, full-text evaluation was conducted for 72 studies. The qualitative analysis was based on 37 articles, whereas the quantitative analysis utilized 14 articles.

Regarding the study types, our qualitative analyses included 21 prospective studies, 3 retrospective studies,

and 7 randomized clinical trials. A total of 25 studies that associated other questionnaires with the SF-36 to evaluate the QOL of patients with endometriosis were identified.

Based on our study protocol, we observed that 13 studies did not apply the SF-36 questionnaire before and after treatment. The results were specified only in the graphics and not in the text. Some studies reported the results as a total number; thus, it was impossible to include them in the statistics. Only studies that had all the reported numerical data from the questionnaires were selected for the meta-analyses.

*Quality and the risk of bias.* Based on the inclusion criteria, we obtained 37 studies; among these, 16 articles were considered as being of 'good' quality (between 6 and 8 points), and 11 articles were considered as being of 'fair' quality (between 3 and 5 points). Table I includes evaluations for each study.

*Study characteristics.* The characteristics for each study are recorded in Table II. The research included 1,1101 women, among whom 6,888 patients were diagnosed with endometriosis. Based on the endometriosis type, 17 studies dealt with all types of endometriosis, 9 studies focused on deep infiltrative endometriosis, and another 9 studies identified bowel endometriosis or deep infiltrative endometriosis with bowel involvement. QOL was evaluated by using only the SF-36 in 12 studies including 1,912 women and by using SF-36 associated with various other questionnaires in 25 studies that had a collective total of 8,022 women.

As part of the SF-36 questionnaire's general evaluation, it was applied for studies that presented the QOL of the patients with endometriosis in comparison to that of healthy women or patients with endometriosis that were interviewed using SF-36 just once (without any follow-up period). Based on this classification, the review included 15 studies with a general overview on endometriosis. Among them, 6 studies evaluated the QOL of patients with endometriosis compared with that of healthy women, and 9 studies compared different types of treatment among patients with endometriosis. On this topic, 6,810 women were evaluated, and 6,393 had endometriosis. The classification based on endometriosis subtypes confirmed that 5,410 patients had all the endometriosis types, 416 had deep infiltrative endometriosis (DIE), and 198 had minimal endometriosis.

For this group of the review, the quantitative analysis could be performed in 6 studies. Our study revealed that patients with endometriosis had an altered QOL compared to the control healthy women group. The pooled prevalence showed that the selected parameters were influenced as follows: Physical Functioning (PF): OR, 78.87 (95% CI, 68.97-88.77;  $I^2=98.77%$ ;  $P\leq 0.001$ ); Bodily Pain (BP): OR, 51.63 (95% CI, 45.23-58.03;  $I^2=92.62%$ ;  $P\leq 0.001$ ); Role Physical (RP): OR, 56.12 (95% CI, 45.08-67.12;  $I^2=95.40%$ ;  $P\leq 0.001$ ); Vitality (VT): OR, 43.09 (95% CI, 32.23-53.95;  $I^2=98.67%$ ;  $P\leq 0.001$ ); Social Functioning (SF): OR, 58.10 (95% CI, 47.28-68.91;  $I^2=97.67%$ ;  $P\leq 0.001$ ); Mental Health (MH): OR, 55.65 (95% CI, 45.58-65.72;  $I^2=98.41%$ ;  $P\leq 0.001$ ); and Role Emotional (RE): OR, 57.84 (95% CI, 47.21-68.48;  $I^2=95.19%$ ;  $P\leq 0.001$ ). Based on pooled prevalence, for patients with endometriosis, the most affected parameter was Physical Functioning (PF): OR, 78.87 (95% CI, 68.97-88.77;  $I^2=98.77%$ ;  $P\leq 0.001$ ).

Qualitative analysis regarding QOL for patients that underwent surgical treatment for endometriosis was evaluated with a SF-36 tool, and it included 21 studies with a collective population of 3,368 patients (Table III). In all the selected studies, the questionnaire was completed before and after surgery. The various studies' follow-up periods ranged between 7 weeks and 3 years. The types of endometriosis were all types of endometriosis (3 studies), DIE (14 studies), or intestinal endometriosis (3 studies). Statistical analyses were performed for 5 studies that included mean values for each result. The studies confirmed that the parameters evaluated with SF-36 improved after surgical treatment. Pooled prevalence revealed that each parameter improved as follows: Physical Functioning (PF): OR, 63.39 (95% CI, 48.71-78.07;  $I^2=97.65%$ ;  $P\leq 0.001$ ); General Health (GH): OR, -0.71 (95% CI, -1.20--0.23;  $I^2=90.04%$ ;  $P\leq 0.001$ ); Bodily Pain (BP): OR, 41.25 (95% CI, 38.95-43.55;  $I^2=97.33%$ ;  $P\leq 0.001$ ), Role Physical (RP): OR, 43.16 (95% CI, 32.74-53.58;  $I^2=91.83%$ ;  $P\leq 0.001$ ); Vitality (VT): OR, 37.99 (95% CI, 30.54-45.44;  $I^2=93.45%$ ;  $P\leq 0.001$ ); Social Functioning (SF): OR, 48.09 (95% CI, 30.67-63.50;  $I^2=98.69%$ ;  $P\leq 0.001$ ); Mental Health (MH): OR, 51.78 (95% CI, 44.70-58.87;  $I^2=92.73%$ ;  $P\leq 0.001$ ); and Role Emotional (RE): OR, 50.65 (95% CI, 41.37-59.93;  $I^2=88.39%$ ;  $P\leq 0.001$ ). In the case of patients who had their endometriosis surgically treated, among the parameters evaluated by SF-36, the highest rate of improvement was observed in Physical Functioning (PF): OR, 63.39 (95% CI, 48.71-78.07;  $I^2=97.65%$ ;  $P\leq 0.001$ ).

Quantitative analysis was conducted for 7 studies on patients who underwent hormone treatments, which were evaluated with SF-36, whereas the statistic forest plots were based on 3 studies. The patients with endometriosis (392 in number) were compared with a control group that did not receive the same treatment (placebo or control). The studies showed that parameters evaluated with SF-36 improved after hormonal treatment. Pooled prevalence revealed that each parameter improved as follows: Physical Functioning (PF): OR, 38.65 (95% CI, 14.39 -62.91;  $I^2=38.65%$ ;  $P\leq 0.001$ ); General Health (GH): OR, -0.60 (95% CI, -1.78-0.57;  $I^2=94.13%$ ;  $P\leq 0.001$ ); Bodily Pain (BP): OR, 28.51 (95% CI, 10.05-41.96;  $I^2=97.33%$ ;  $P\leq 0.001$ ); Role Physical (RP): OR, 32.42 (95% CI, 0.62-64.49;  $I^2=98.62%$ ;  $P\leq 0.001$ ); Vitality (VT): OR, 31.49 (95% CI, 6.58-56.39;  $I^2=98.58%$ ;  $P\leq 0.001$ ); Social Functioning (SF): OR, 35.86 (95% CI, 12.01-59.7;  $I^2=97.92%$ ;  $P\leq 0.001$ ); Mental Health (MH): OR, 35.11 (95% CI, 5.72-64.50;  $I^2=98.85%$ ;  $P\leq 0.001$ ); and Role Emotional (RE): OR, -0.39 (95% CI, -0.90-0.20;  $I^2=74.57%$ ;  $P\leq 0.001$ ). After the hormonal treatment, the parameter that showed the most improvement was Physical Functioning (PF): OR, 38.65 (95% CI, 14.39 -62.91;  $I^2=38.65%$ ;  $P\leq 0.001$ ).

#### 4. Discussion

Our meta-analyses revealed that the SF-36 questionnaire had a heterogenous applicability in all the studies. Recently, Chauvet *et al* performed a recent systematic review, which proved that the SF-36 was the most frequently used scale, followed by the EHP-30, for evaluating quality of life (QOL) of patients with endometriosis (16). The design of our study was different from previous meta-analyses as we included articles

Table I. Quality assessment of the studies.

| Authors, year                                 | Selection                            |                                 |                           |                                 | Outcome                               |                       |                               | Total Score (Refs.) |
|---|--------------------------------------|---------------------------------|---------------------------|---------------------------------|---------------------------------------|-----------------------|-------------------------------|---------------------|
|   | Representativeness of exposed cohort | Selection of non-exposed cohort | Ascertainment of exposure | Outcome not present at baseline | Compatibility of cohort (matched for) | Assessment of outcome | Sufficient follow-up duration |                     |
| Van Aken <i>et al.</i> , 2017                 | +                                    | +                               | +                         |                                 | +                                     | +                     | +                             | Fair (18)           |
| Angioni <i>et al.</i> , 2015                  | +                                    |                                 | +                         | +                               | +                                     | +                     |                               | Good (31)           |
| Silveira da Cunha Araujo <i>et al.</i> , 2014 | +                                    |                                 | +                         |                                 | +                                     |                       | +                             | Fair (26)           |
| Bassi <i>et al.</i> , 2011                    | +                                    |                                 | +                         | +                               |                                       | +                     |                               | Good (27)           |
| Caruso <i>et al.</i> , 2015                   | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (28)           |
| Chauvet <i>et al.</i> , 2018                  | +                                    |                                 | +                         | +                               | +                                     |                       |                               | Fair (16)           |
| Comptour <i>et al.</i> , 2020                 | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (13)           |
| Darai <i>et al.</i> , 2010                    | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (40)           |
| Darai <i>et al.</i> , 2017                    | +                                    |                                 | +                         | +                               | +                                     | +                     |                               | Fair (37)           |
| Dubernard <i>et al.</i> , 2008                | +                                    |                                 | +                         | +                               | +                                     | +                     |                               | Good (29)           |
| Friedl <i>et al.</i> , 2015                   | +                                    |                                 | +                         | +                               | +                                     |                       |                               | Good (24)           |
| Garavaglia <i>et al.</i> , 2018               | +                                    |                                 | +                         | +                               | +                                     | +                     |                               | Good (38)           |
| De Graaff <i>et al.</i> , 2013                | +                                    |                                 | +                         | +                               | +                                     | +                     |                               | Fair (39)           |
| Grandi <i>et al.</i> , 2015                   | +                                    | +                               | +                         | +                               | +                                     |                       | +                             | Good (54)           |
| Jones <i>et al.</i> , 2004                    | +                                    |                                 | +                         | +                               | +                                     | +                     |                               | Good (56)           |
| Laursen <i>et al.</i> , 2005                  | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (19)           |
| Ledu <i>et al.</i> , 2018                     | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (33)           |
| Lovkvist <i>et al.</i> , 2016                 | +                                    | +                               | +                         |                                 | +                                     | +                     |                               | Fair (55)           |
| Mabrouk <i>et al.</i> , 2011                  | +                                    | +                               | +                         | +                               | +                                     |                       |                               | Fair (43)           |
| Mabrouk <i>et al.</i> , 2012                  | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (36)           |
| Melis <i>et al.</i> , 2015                    | +                                    | +                               | +                         | +                               | +                                     |                       |                               | Fair (47)           |
| Miller, 2000                                  | +                                    |                                 | +                         | +                               | +                                     | +                     | +                             | Good (44)           |
| Marki <i>et al.</i> , 2017                    | +                                    |                                 | +                         | +                               | +                                     |                       |                               | Fair (45)           |
| Nnoaham <i>et al.</i> , 2011                  | +                                    | +                               | +                         | +                               | +                                     |                       | +                             | Good (51)           |
| Nunes <i>et al.</i> , 2014                    | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (48)           |
| Petrelluzzi <i>et al.</i> , 2008              | +                                    | +                               | +                         | +                               | +                                     | +                     |                               | Good (52)           |
| Ribeiro <i>et al.</i> , 2014                  | +                                    |                                 | +                         | +                               | +                                     | +                     | +                             | Good (30)           |
| Riiskjær <i>et al.</i> , 2018                 | +                                    |                                 | +                         | +                               | +                                     | +                     | +                             | Good (34)           |

Table I. Continued.

| Authors, year                | Selection                            |                                 |                           |                                 | Outcome                               |                       |                               | Total Score (Refs.) |                    |
|------------------------------|--------------------------------------|---------------------------------|---------------------------|---------------------------------|---------------------------------------|-----------------------|-------------------------------|---------------------|--------------------|
|                              | Representativeness of exposed cohort | Selection of non-exposed cohort | Ascertainment of exposure | Outcome not present at baseline | Compatibility of cohort (matched for) | Assessment of outcome | Sufficient follow-up duration |                     | Adequate follow-up |
| Roman <i>et al</i> , 2012    | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (21)          |
| Roman <i>et al</i> , 2018    | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (32)          |
| Roman <i>et al</i> , 2018    | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Fair (32)          |
| Teixeira <i>et al</i> , 2017 | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (49)          |
| Touboul <i>et al</i> , 2015  | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (41)          |
| Valentin <i>et al</i> , 2017 | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Fair (25)          |
| Verket <i>et al</i> , 2018   | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (20)          |
| Bi and Xie, 2018             | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (22)          |
| Zhao <i>et al</i> , 2012     | +                                    | +                               | +                         | +                               | +                                     | +                     | +                             | +                   | Good (46)          |

that evaluated patients with endometriosis using the SF-36 questionnaire before and after surgical or medical treatment or as a comparison tool between groups. The literature did not reveal a similar approach. For example, Arcoverde *et al* performed another meta-analysis utilizing generic questionnaires and revealed that surgery improved the quality of life for patients with all types of endometriosis (17). We excluded cases with hormonal approach and cases that were evaluated using different questionnaires (18). There was no meta-analyses that evaluated the efficiency of SF-36 for patients with endometriosis treated with hormonal therapy or as a comparison method between study groups.

Regarding the first issue of our classification, we identified 6,393 women who received treatment for endometriosis and answered the SF-36 questionnaire. Based on this approach, we observed that, for women with endometriosis, compared with the control groups, physical functioning was the main parameter that received the most alteration. Two peculiar studies were included within this group of studies. Laursen *et al* (19) compared patients with fibromyalgia/whiplash, endometriosis, lower back pain, or rheumatoid arthritis and healthy women. We included this study because of its data accuracy; furthermore, there was no other pathological influence on the women with endometriosis. The second study was the Verket *et al* study (20), which evaluated women from the general population, women with endometriosis, and women with rheumatoid arthritis. We incorporated it into the analyses of this study because it did not have a control group of healthy women for comparison to women with endometriosis, and the arthritis had no impact. For both of these studies, we considered only the data for the endometriosis patients.

Van Aken *et al* demonstrated that, for endometriosis patients, pain cognition and QOL are independently associated (18). Verket *et al* highlighted that patients with moderate to severe endometriosis had overall impaired QOL compared to women from the general population (20). Women with DIE reported constipation, defecation pain, appetite disorders, longer evacuation time, and increased stool consistency without laxatives (21). Neuromuscular electro-stimulation was found to be an efficient method for alleviating endometriosis-related pain (22).

The present analysis of QOL for patients with endometriosis who received surgical treatment included data from 3,368 patients. The pre- and post-surgery questionnaire in all the studies showed that physical functioning was the principal parameter that improved after the interventions. The studies proved that endometriosis symptoms such as pain showed improvement after surgical treatment (23), even though endometriosis also influenced biopsychosocial variables (24). Patients with minimal endometriosis seldom reported surgery as a good treatment for improving QOL (25). For patients with bowel endometriosis, QOL improved significantly after laparoscopic treatment (26,27), with long-term results (28-30). The study by Angioni *et al* (31), a randomized clinical trial, included a surgery group as well as a hormonal treatment group. For this group, we considered the arm A results, with a complete resection of DIE. The study by Roman *et al* (2018) (32) was analyzed with regard to the segmental resection of the rectum, and the study by Ledu *et al* (33) was utilized for the group consisting of patients with stoma; this was because of the

Table II. Characteristics of the studies.

| Authors, year                                 | Selection                                  |   |   |   |  | Outcome   |                         |                               |
|---|--|---|---|---|--|---|-------------------------|-------------------------------|
|   | Type of study                              | Number of included patients/lost to follow-up | Additional questionnaires used with SF-36 | Comparison and control groups   | Type of endometriosis  | Treatment   | Follow-up               | Final no. of patients (Refs.) |
| Van Aken <i>et al.</i> , 2017                 | Cross-sectional questionnaire-based survey | 92/9  | EHP-30, PCS, PVAQ, PASS and NRS           | Patients with endometriosis compared with healthy women                               | Women with variety in their disease severity. The majority of patients (74%) presented a moderate-to-severe stage of endometriosis | Therapeutic surgical interventions  | /                       | 83 (18)                       |
| Angioni <i>et al.</i> , 2015                  | Randomized clinical trial                  | 159/0   | /   | Complete laparoscopic excision of DIE (Arm A) and incomplete excision of DIE (Arm B). | DIE  | GnRHa after laparoscopic excision of DIE (Arm 2A) or incomplete excision of DIE (Arm 2B). | 3, 6 and 12 months      | 159 (31)                      |
| Silveira da Cunha Araújo <i>et al.</i> , 2014 | Observational prospective cohort           | 45/0  | /   | /   | DIE with colorectal involvement  | Laparoscopy with colorectal resection   | 12 and 48 months        | 45 (26)                       |
| Bassi <i>et al.</i> , 2011                    | Prospective                                | 151/0   | /   | /   | DIE with colorectal involvement  | Resection of a segment of the rectosigmoid  | 12 months               | 151 (27)                      |
| Caruso <i>et al.</i> , 2015                   | Prospective                                | 92/0  | VAS, FSFI, FSDS                           | Study group and control group   | All types  | Study group (DNG) and control group (AINS)  | 3 and 6 months          | 92 (28)                       |
| Chauvet <i>et al.</i> , 2018                  | Cross-sectional                            | 1,156/243                                     | EHP-30                                    | /   | All types  | Confirmed surgical diagnosis of endometriosis   | /                       | 913 (16)                      |
| Comptour <i>et al.</i> , 2020                 | Prospective, multicenter cohort            | 1237/256                                      | VAS                                       | /   | All types  | Laparoscopic treatment  | 3 years                 | 981 (13)                      |
| Darai <i>et al.</i> , 2010                    | Single-center, retrospective               | 29/0  | IPSS, BFLUTS                              | Laparoscopy group and laparotomy group  | Extensive pelvic endometriosis   | Radical en bloc hysterectomy and colorectal resection (REHCR)                             | 16 months and 11 months | 29 (40)                       |

Table II. Continued.

| Authors, year                  | Type of study                              | Selection                                     |   |   | Outcome                        |  |                             |                               |
|--------------------------------|--|---|---|---|--------------------------------|--|-----------------------------|-------------------------------|
|                                |  | Number of included patients/lost to follow-up | Additional questionnaires used with SF-36 | Comparison and control groups   | Type of endometriosis          | Treatment  | Follow-up                   | Final no. of patients (Refs.) |
| Darai <i>et al</i> , 2017      | Prospective                                | 46/26   | /   | /   | DIE and colorectal involvement | Osteopathic manipulative therapy   | /                           | 20 (37)                       |
| Dubernard <i>et al</i> , 2008  | Prospective                                | 93/35   | /   | /   | Colorectal endometriosis       | Laparoscopic segmental colorectal resection  | 22.5 months                 | 58 (29)                       |
| Friedl <i>et al</i> , 2015     | Prospective                                | 61/0  | HADS-D                                    | /   | All types                      | Previous operations  | /                           | 61 (24)                       |
| Garavaglia <i>et al</i> , 2018 | Prospective observational                  | 20/0  | /   | /   | Bowel endometriosis            | Laparoscopic colorectal resection  | 12 months                   | 20 (38)                       |
| De Graaff <i>et al</i> , 2013  | Cross-sectional questionnaire-based survey | 1450/519                                      | GSWH instrument                           | /   | All types                      | Laparoscopic and/or histological diagnosis of endometriosis, who had at least one contact related to endometriosis-associated symptoms | /                           | 931 (39)                      |
| Grandi <i>et al</i> , 2015     | Prospective observational                  | 40/6  | VAS                                       | DNG group and NSAID group   | All types                      | 19 in the E2V/DNG group and 15 in the NSAID group  | After 6 cycles of treatment | 34 (54)                       |
| Jones <i>et al</i> , 2004      | Prospective observational                  | 66/26   | EHP-30                                    | /   | All types                      | Conservative surgery   | 4 months                    | 40 (56)                       |
| Laursen <i>et al</i> , 2005    | Prospective observational                  | 40/0  | VAS                                       | Fibromyalgia/whiplash group, endometriosis group, low back pain (group, or rheumatoid arthritis group and healthy women | All types                      | Experimental pain stimulus [hyperalgesia to pressure pain threshold (PPT)]   | /                           | 40 (19)                       |
| Ledu <i>et al</i> , 2018       | Retrospective                              | 134/87  | /   | Stoma group vs. non-stoma group   | DIE with rectal involvement    | Discooid resection   | /                           | 47 (33)                       |
| Lovkvist <i>et al</i> , 2016   | Observational                              | 800/369                                       | /   | /   | All types                      | Treatment  | /                           | 431 (55)                      |
| Mabrouk <i>et al</i> , 2011    | Retrospective                              | 241/135                                       | VAS                                       | COC users, and COC nonusers.  | Uncomplicated posterior DIE    | Laparoscopic surgery   | /                           | 106 (43)                      |

Table II. Continued.

| Authors, year                    | Type of study                     | Number of included patients/lost to follow-up | Additional questionnaires used with SF-36       | Selection  |                               | Outcome                               |                 |                               |
|----------------------------------|-----------------------------------|---|---|--|-------------------------------|---------------------------------------|-----------------|-------------------------------|
|                                  |                                   |   |   | Comparison and control groups  | Type of endometriosis         | Treatment                             | Follow-up       | Final no. of patients (Refs.) |
| Mabrouk <i>et al.</i> , 2012     | Prospective                       | 47/0  | VAS   | /  | Colorectal. endometriosis     | Laparoscopic segmental resection      | 18 months       | 47 (36)                       |
| Melis <i>et al.</i> , 2015       | Prospective                       | 41/0  | FSFI, BAT                                       | Women with deep endometriosis compared with healthy women  | DIE                           | Surgery                               | /               | 41 (47)                       |
| Miller, 2000                     | Prospective, randomized           | 120/0   | VAS   | /  | All types                     | Therapy with depot leuprolide acetate | 4 weeks         | 120 (44)                      |
| Marki <i>et al.</i> , 2017       | Prospective, cross-sectional      | 210/17  | /   | /  | All types                     | Treatment at the participating clinic | /               | 193 (45)                      |
| Nnoaham <i>et al.</i> , 2011     | Prospective cross-sectional study | 1,669/924                                     | WPAI:GH questionnaire                           | i) Women with endometriosis; ii) symptomatic control women without endometriosis; and iii) sterilization control women without endometriosis | All types                     | Laparoscopy                           | /               | 745 (51)                      |
| Nunes <i>et al.</i> , 2014       | cross-sectional                   | 510/253                                       | ACR criteria were used to evaluate fibromyalgia | Women with endometriosis compared with women with no history of endometriosis  | All types                     | Laparoscopy                           | /               | 257 (48)                      |
| Petrelluzzi <i>et al.</i> , 2008 | Prospective                       | 175/82  | VAS, PSQ  | Women with endometriosis compared with healthy women   | All types                     | Laparoscopy, laparotomy               | /               | 93 (52)                       |
| Ribeiro <i>et al.</i> , 2014     | Prospective observational         | 45/5  | /   | /  | Intestinal deep endometriosis | Laparoscopic colorectal resections    | 6 and 12 months | 40 (30)                       |
| Riiskjar <i>et al.</i> , 2018    | Prospective, observational        | 185/10  | NRS   | /  | Rectosigmoid endometriosis.   | Laparoscopic bowel resection          | 12 months       | 175 (34)                      |



Table II. Continued.

| Authors, year                | Type of study                             | Number of included patients/lost to follow-up | Selection                                 |   | Outcome                            |  |                 |                               |
|------------------------------|---|---|---|---|------------------------------------|--|-----------------|-------------------------------|
|                              |   |   | Additional questionnaires used with SF-36 | Comparison and control groups   | Type of endometriosis              | Treatment                                      | Follow-up       | Final no. of patients (Refs.) |
| Roman <i>et al</i> , 2012    | Cohort prospective                        | 116/0   | GIQLI, KESS                               | Endometriosis involving the Douglas pouch, deep endometriosis without digestive infiltration vs. deep endometriosis infiltrating the rectum | DIE                                | Surgery  | 1 and 3 years   | 116 (21)                      |
| Roman <i>et al</i> , 2018    | Randomized trial                          | 60/0  | VAS, KESS<br>GIQLI, USP                   | Conservative surgery group vs. rectal segmental resection   | DIE infiltrating the rectum        | Conservative surgery or radical rectal surgery | 24 months       | 60 (32)                       |
| Sesti <i>et al</i> , 2007    | Randomized comparative trial              | 234/12  | VAS                                       | Placebo vs. GnRH-a vs. dietary therapy  | Endometriosis stage III-IV (r-AFS) | Conservative pelvic surgery                    | 12 months       | 222 (50)                      |
| Teixeira <i>et al</i> , 2017 | Randomized controlled trial               | 41/0  | VAS                                       | Potentized estrogen vs. placebo   | DIE                                | Potentized estrogen                            | 24 weeks        | 41 (49)                       |
| Touboul <i>et al</i> , 2015  | Randomized controlled trial               | 52/0  | /   | Open surgery vs. laparoscopically assisted colorectal resection   | Colorectal endometriosis           | Laparoscopically assisted to open surgery      | 13.8 months     | 52 (41)                       |
| Valentin <i>et al</i> , 2017 | Prospective and multicenter observational | 198/31  | /   | /   | Minimal endometriosis              | Laparoscopic procedure                         | /               | 167 (25)                      |
| Verket <i>et al</i> , 2018   | Cross-sectional questionnaire study       | 1,149/992                                     | /   | Women from the general population, vs. women with endometriosis compared with women with rheumatoid arthritis (RA).                         | All types                          | Surgically confirmed diagnosis.                | /               | 157 (20)                      |
| Xie and Bi, 2018             | Retrospective study                       | 154/0   | NRS and ESSS                              | Treatment group vs. control group   | All types                          | Neuromuscular electrical stimulation           | After treatment | 154 (22)                      |
| Zhao <i>et al</i> , 2012     | Controlled, randomized study              | 50/0  | STAI                                      | Patients with endometriosis compared with control group   | All types                          | Leuprolide                                     | After treatment | 50 (46)                       |

Table III. Patient characteristics.

| Study                         | Study group characteristics  | Control group characteristics  | Type of endometriosis   | Therapeutic approach   | (Refs.) |
|-------------------------------|--|--|---|--|---------|
| Van Aken <i>et al.</i> , 2017 | 50 women with laparoscopically and/or magnetic resonance imaging-confirmed endometriosis             | 42 healthy control women   | Two-thirds with DIE   | Half the patients (56%) had a history of one or more therapeutic surgical interventions; 10 patients (35%) had undergone major: surgery hysterectomy, ovariectomy, bowel or bladder procedures, extended adhesiolysis, or a combination of procedures. | (18)    |
| Angioni <i>et al.</i> , 2015  | 80 patients underwent complete laparoscopic excision of DIE (Arm A)                                  | 79 patients underwent incomplete surgery (Arm B)                     | DIE   | After surgery each surgical arm was randomized in two groups: no treatment groups 1A [40 pts.] and 1B [40 pts.] and GnRH $\alpha$ treatment for 6 months groups 2A [40 pts.] and 2B  | (31)    |
| Caruso <i>et al.</i> , 2015   | 54 women with endometriosis received 2 mg/daily DNG  | 48 with endometriosis received non-steroidal anti-inflammatory drugs | Women affected by chronic pelvic pain with a clinical diagnosis of endometriosis  | 2 mg/daily DNG for 6 months  | (28)    |
| Darai <i>et al.</i> , 2010    | 16 women who underwent laparoscopy   | 13 that underwent laparotomy   | Extensive pelvic endometriosis  | Radical en bloc hysterectomy and colorectal resection (REHCR)  | (40)    |
| Friedl <i>et al.</i> , 2015   | 62 patients with a histologically confirmed diagnosis of endometriosis                               | 61 healthy women recruited by the internet in a student community    | Confirmed endometriosis   | Without details about previous surgical interventions  | (24)    |
| Grandi <i>et al.</i> , 2015   | 19 patients with endometriosis that received E2V (quadruphasic association of E2 valerate)/DNG group | 15 patients with endometriosis that received NSAID                   | All types (diagnosed by previous laparoscopy or laparotomy or the presence of an ovarian endometrioma diagnosed by transvaginal ultrasound) | Patients with endometriosis that were not scheduled for first or repeated surgery  | (54)    |
| Laursen <i>et al.</i> , 2005  | 10 patients with endometriosis   | 41 healthy women for control group                                   | Chronic endometriosis   | Experimental pain stimulus (hyperalgesia to pressure pain threshold (PPT))   | (19)    |
| Ledu <i>et al.</i> , 2018     | 33 patients who received protective stoma  | 14 patients without protective stoma                                 | DIE with rectal involvement   | Segmental or discoid resection of the rectum   | (33)    |
| Mabrouk <i>et al.</i> , 2011  | 75 patients who had received COCs during preoperative period   | 31 nonusers, who had not received any hormone treatment              | Uncomplicated posterior DIE   | Laparoscopic surgery   | (43)    |
| Melis <i>et al.</i> , 2015    | 41 women with deep endometriosis   | 40 healthy women without endometriosis                               | DIE   | Most participants were after surgical diagnosis without any hormonal treatment, other patients entered the study before surgery  | (47)    |

Table III. Continued.

| Study                            | Study group characteristics   | Control group characteristics   | Type of endometriosis       | Therapeutic approach   | (Refs.) |
|----------------------------------|---|---|-----------------------------|--|---------|
| Nnoaham <i>et al.</i> , 2011     | Women with endometriosis  | Symptomatic control women without endometriosis; and sterilization control women without endometriosis                      | All types                   | Prospectively recruited consecutive premenopausal women, scheduled for a laparoscopy: i) to investigate endometriosis-associated pelvic pain; or ii) to be sterilized        | (51)    |
| Nunes <i>et al.</i> , 2014       | 257 women with a laparoscopic and histopathological diagnosis of endometriosis                                | 253 women with no signs or symptoms of endometriosis  | All grades of endometriosis | Previous laparoscopy   | (48)    |
| Petrelluzzi <i>et al.</i> , 2008 | 93 women with endometriosis   | 82 healthy women  | All types                   | Laparoscopy or laparotomy for diagnosis  | (52)    |
| Roman <i>et al.</i> , 2012       | 53 women with deep endometriosis infiltrating the rectum  | 21 women with endometriosis involving the Douglas pouch and 42 women with deep endometriosis without digestive infiltration | DIE                         | Surgically treated patients  | (21)    |
| Roman <i>et al.</i> , 2018       | 27 patients who received conservative surgery   | 33 patients who received segmental resection  | DIE infiltrating the rectum | Conservative surgery or radical rectal surgery   | (32)    |
| Roman <i>et al.</i> , 2018       | 119 patients with endometriosis who received postoperative adjunctive medical or dietary therapy for 6 months | 115 patients with endometriosis who received placebo for 6 months   | Endometriosis stage III-IV  | Laparoscopic or laparotomic diagnosis of severe endometriosis stage III-IV. Randomization was achieved at the time of postoperative control (7 days after surgery)           | (32)    |
| Teixeira <i>et al.</i> , 2017    | 23 patients with endometriosis who received potentized estrogen   | 27 patients with endometriosis who received placebo   | DIE                         | Potentized estrogen administered after diagnosis of deeply infiltrating endometriosis based on magnetic resonance imaging or transvaginal ultrasound after bowel preparation | (49)    |
| Touboul <i>et al.</i> , 2015     | 26 patients who underwent open surgery  | 26 patients who underwent laparoscopy   | Colorectal endometriosis    | Laparoscopically assisted or open colorectal resection for endometriosis   | (41)    |
| Verket <i>et al.</i> , 2018      | 156 women with endometriosis  | 156 women from the general population, and 837 women with rheumatoid arthritis (RA)   | All types                   | No surgical or medical treatment   | (20)    |
| Xie and Bi, 2018                 | 83 patients with endometriosis were in the treatment group  | 71 patients with endometriosis represented the control group  | All types                   | Neuromuscular electrical stimulation   | (22)    |
| Zhao <i>et al.</i> , 2012        | 50 patients with endometriosis  | 50 patients with endometriosis represented the control group  | All types                   | Patients received gonadotrophin-releasing hormone (GnRH) agonist therapy   | (46)    |

rectal resection for DIE with rectal involvement (the study results were relevant for this group).

Some authors have recommended surgery for rectosigmoid endometriosis in cases where the disease is unresponsive to conservative treatments (34-36). Daraï *et al* supported the finding that osteopathic manipulative therapy has a good impact on endometriosis symptoms in women with colorectal involvement (37). However, although the radical surgical approach can have a major positive impact on QOL, it does not improve all symptoms, as reported by Garavaglia *et al* (38). De Graaff *et al* concluded that many patients with endometriosis experienced better QOL and continued to have endometriosis-associated symptoms even when they were treated in tertiary care centers (39). Roman *et al* reported that conservative surgery should be reserved for patients with large deep rectal endometriosis (32). Daraï *et al* recommended laparoscopy for patients that required radical en bloc hysterectomy and colorectal resection (40). Laparoscopic colorectal resection was also indicated for cases of lower intraoperative and postoperative complications for patients with rectal endometriosis due to higher spontaneous post-surgery pregnancy rates (41). Araujo *et al* reported that gonadotropin-releasing hormone agonist (GnRHa) administration was followed by a temporary improvement in pain among patients with incomplete resection of deep infiltrative endometriosis (DIE), but it did not affect postsurgical pain when DIE implants were completely excised (42).

The study by Angioni *et al* (31) was considered for the medical treatment group because it applied postoperative randomization (of initial groups with complete or incomplete resection of endometriosis) to receivers or non-receivers of GnRHa. It selected a group with complete resection of endometriosis, which received GnRH, and compared it with complete resection of endometriosis without medical treatment. Regarding statistics, the Caruso group (28) was not considered because it compared GnRH receivers with AINS receivers.

Mabrouk *et al* confirmed the role of combined oral contraceptive therapy in curbing the progression of dysmenorrhea and dyspareunia and the growth of deep endometriotic nodules (43). Meanwhile, Miller concluded that the stimulatory phase of the GnRHa therapy could be correlated with an increase in pain and a decrease in the QOL (44). Márki *et al* emphasized that pain management and emotion regulation strategies can be utilized for improving the QOL of patients with endometriosis. The authors suggest that it could be obtained through simultaneous application with physical treatment and psychological care (45).

Zhao *et al* conducted the first study to explore only the effects of psychosomatic therapy on the QOL of endometriosis patients and demonstrated that progressive muscular relaxing (PMR) training could serve as a reference for future psychosomatic interventions regarding endometriosis. PRM works efficiently for anxiety and depression related to endometriosis (46). Melis *et al* further suggested that a patient-centered approach that is integrated with various medical, psychological, and sexual issues should be applied in the treatment of patients with endometriosis (47).

Nunes *et al* compared the QOL of patients with fibromyalgia and women with and without endometriosis. The QOL of women with endometriosis showed the highest impairment (48).

Teixeira *et al* demonstrated that potentated estrogens significantly reduced endometriosis-associated pain compared to a placebo (49). Sesti *et al* demonstrated that, for endometriosis stage III-IV, QOL can be improved through postoperative hormonal suppression treatment or dietary therapy compared to surgery as well as a placebo for relieving pain (50).

Regardless of the differences across country- and ethnicity-level contexts, endometriosis was found to impair QOL and work productivity for women who continued to experience diagnostic delays in primary care (51). Even if hypocortisolism could be sometimes considered a consequence of the aversive symptoms of endometriosis, the etiopathology remains unelucidated (52,53). The relationship between endometriosis and ovarian cancer was studied by Grandi *et al* and the preventive effect of inhibiting ovulation and menstruation was highlighted (54).

EHP-30 and SF-36 questionnaires are sensitive to changes related to endometriosis, especially correlations between increased pressure pain sensitivity and hyperalgesia and the impairment of QOL parameters, as reported by Laursen *et al* (19). Thus, healthcare resource allocators should focus on women with endometriosis as the disease may have a different impact for each woman (55). The SF-36 seems to be a valid instrument for evaluating QOL in women with endometriosis (56,57).

In conclusion, the 36-item survey generic questionnaire, SF-36, seems to represent an efficient/valid method for assessing the QOL of patients with endometriosis who are undergoing surgical or hormonal treatments. It can be applied before and after procedures or in order to compare study groups.

### Acknowledgements

Not applicable.

### Funding

No funding was received.

### Availability of data and materials

The information and data contained in this review and meta-analysis are documented by relevant references. Any other details can be obtained upon request.

### Authors' contributions

LP and JCR performed the conceptualization of the study and had major contributions in writing the manuscript. BHH contributed to the methodology. MCTD and CC analyzed and interpreted the patient data. PS, III, MPR, IJB and EFS provided the formal analysis and visualization. IN, ADS and BS conducted the literature investigation and data curation. RMS provided resources and contributed to writing the manuscript (review and editing). MCTD, BS, and DS searched the literature for similar work and articles and contributed to writing the manuscript. LP and JCR supervised the study and provided project administration. All authors read and approved the final manuscript for publication.

**Ethics approval and consent to participate**

Not applicable.

**Patient consent for publication**

Not applicable.

**Competing interests**

The authors declare no conflict or competing interests.

**References**

- Giudice LC: Clinical practice. Endometriosis. *N Engl J Med* 362: 2389-2398, 2010.
- Gustofson RL, Kim N, Liu S and Stratton P: Endometriosis and the appendix: A case series and comprehensive review of the literature. *Fertil Steril* 86: 298-303, 2006.
- Practice Committee of the American Society for Reproductive Medicine: Treatment of pelvic pain associated with endometriosis: A committee opinion. *Fertil Steril* 101: 927-935, 2014.
- Bodean O, Bratu O, Bohiltea R, Munteanu O, Marcu D, Spinu DA, Vacaroiu IA, Socea B, Diaconu CC, Gradinaru DF and Cirstoiu M: The efficacy of synthetic oral progestin pills in patients with severe endometriosis. *Rev Chim (Bucharest)* 69: 1411-1415, 2018.
- Moradi M, Parker M, Sneddon A, Lopez V and Ellwood D: Impact of endometriosis on women's lives: A qualitative study. *BMC Womens Health* 14: 123, 2014.
- Bergner M: Quality of life, health status, and clinical research. *Med Care* 27 (Suppl 3): S148-S156, 1989.
- Ware JE Jr and Sherbourne CD: The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 30: 473-483, 1992.
- Yoldemir T: Quality of life for women with endometriosis: Premenopausal and postmenopausal perspectives. *Climacteric* 21: 411-412, 2018.
- Kashi AM, Moradi Y, Chaichian S, Najmi Z, Mansori K, Salehin F, Rastgar A and Khateri S: Application of the World Health Organization quality of life instrument, short form (WHOQOL-BREF) to patients with endometriosis. *Obstet Gynecol Sci* 61: 598-604, 2018.
- Gallagher JS, DiVasta AD, Vitonis AF, Sarda V, Laufer MR and Missmer SA: The impact of endometriosis on quality of life in adolescents. *J Adolesc Health* 63: 766-772, 2018.
- Fritzer N, Tammaa A, Salzer H and Hudelist G: Dyspareunia and quality of sex life after surgical excision of endometriosis: A systematic review. *Eur J Obstet Gynecol Reprod Biol* 173: 1-6, 2014.
- Franck C, Poulsen MH, Karampas G, Giraldi A and Rudnicki M: Questionnaire-based evaluation of sexual life after laparoscopic surgery for endometriosis: A systematic review of prospective studies. *Acta Obstet Gynecol Scand* 97: 1091-1104, 2018.
- Comptour A, Lambert C, Chauvet P, Figuiet C, Gremeau AS, Canis M, Pereira B and Bourdel N: Long-term evolution of quality of life and symptoms following surgical treatment for endometriosis: Different trajectories for which patients? *J Clin Med* 9: 2461, 2020.
- Liberati A, Altman DG, Tetzlaff J, Mulrow C, Gøtzsche PC, Ioannidis JP, Clarke M, Devereaux PJ, Kleijnen J and Moher D: The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate health care interventions: Explanation and elaboration. *J Clin Epidemiol* 62: e1-e34, 2009.
- Lo CK, Mertz D and Loeb M: Newcastle-ottawa scale: Comparing reviewers' to authors' assessments. *BMC Med Res Methodol* 14: 45, 2014.
- Chauvet P, Guiguet-Auclair C, Comptour A, Denouël A, Gerbaud L, Canis M and Bourdel N: Feelings and expectations in endometriosis: Analysis of open comments from a cohort of endometriosis patients. *J Gynecol Obstet Hum Reprod* 47: 281-287, 2018.
- Arcoverde FV, Andres MP, Borrelli GM, Barbosa PA, Abrão MS and Kho RM: Surgery for endometriosis improves major domains of quality of life: A systematic review and meta-analysis. *J Minim Invasive Gynecol* 26: 266-278, 2019.
- van Aken MA, Oosterman JM, van Rijn CM, Ferdek MA, Ruijt GS, Peeters BW, Braat DD and Nap AW: Pain cognition versus pain intensity in patients with endometriosis: Toward personalized treatment. *Fertil Steril* 108: 679-686, 2017.
- Laursen BS, Bajaj P, Olesen AS, Delmar C and Arendt-Nielsen L: Health related quality of life and quantitative pain measurement in females with chronic non-malignant pain. *Eur J Pain* 9: 267-275, 2005.
- Verket NJ, Uhlig T, Sandvik L, Andersen MH, Tanbo TG and Qvigstad E: Health-related quality of life in women with endometriosis, compared with the general population and women with rheumatoid arthritis. *Acta Obstet Gynecol Scand* 97: 1339-1348, 2018.
- Roman H, Ness J, Suci N, Bridoux V, Gourcerol G, Leroi AM, Tuech JJ, Ducrotté P, Savoye-Collet C and Savoye G: Are digestive symptoms in women presenting with pelvic endometriosis specific to lesion localizations? A preliminary prospective study. *Hum Reprod* 27: 3440-3449, 2012.
- Bi XL and Xie CX: Effect of neuromuscular electrical stimulation for endometriosis-Associated pain: A retrospective study. *Medicine (Baltimore)* 97: e11266, 2018.
- Comptour A, Chauvet P, Canis M, Grémeau AS, Pouly JL, Rabischong B, Pereira B and Bourdel N: Patient quality of life and symptoms after surgical treatment for endometriosis. *J Minim Invasive Gynecol* 26: 717-726, 2019.
- Friedl F, Riedl D, Fessler S, Wildt L, Walter M, Richter R, Schüller G and Böttcher B: Impact of endometriosis on quality of life, anxiety, and depression: An Austrian perspective. *Arch Gynecol Obstet* 292: 1393-1399, 2015.
- Valentin L, Canis M, Pouly JL, Lasnier C, Jaffeux P, Aublet-Cuvelier B and Bourdel N: SF-36 preoperative interest of predicting improvement of quality of life after laparoscopic management of minimal endometriosis. *J Gynecol Obstet Hum Reprod* 46: 137-142, 2017.
- Silveira da Cunha Araújo R, Ribeiro HS, Sekula VG, da Costa Porto BT and Ribeiro PA: Long-term outcomes on quality of life in women submitted to laparoscopic treatment for bowel endometriosis. *J Minim Invasive Gynecol* 21: 682-688, 2014.
- Bassi MA, Podgaec S, Dias JA Jr, Filho ND, Petta CA and Abrao MS: Quality of life after segmental resection of the rectosigmoid by laparoscopy in patients with deep infiltrating endometriosis with bowel involvement. *J Minim Invasive Gynecol* 18: 730-733, 2011.
- Caruso S, Iraci M, Cianci S, Casella E, Fava V and Cianci A: Quality of life and sexual function of women affected by endometriosis-associated pelvic pain when treated with dienogest. *J Endocrinol Invest* 38: 1211-1218, 2015.
- Dubernard G, Rouzier R, David-Montefiore E, Bazot M and Daraï E: Use of the SF-36 questionnaire to predict quality-of-life improvement after laparoscopic colorectal resection for endometriosis. *Hum Reprod* 23: 846-851, 2008.
- Ribeiro PA, Sekula VG, Abdalla-Ribeiro HS, Rodrigues FC, Aoki T and Aldrighi JM: Impact of laparoscopic colorectal segment resection on quality of life in women with deep endometriosis: One year follow-up. *Qual Life Res* 23: 639-643, 2014.
- Angioni S, Pontis A, Dessole M, Surico D, De Cicco Nardone C and Melis I: Pain control and quality of life after laparoscopic en-block resection of deep infiltrating endometriosis (DIE) vs. incomplete surgical treatment with or without GnRHa administration after surgery. *Arch Gynecol Obstet* 291: 363-370, 2015.
- Roman H, Bubenheim M, Huet E, Bridoux V, Zacharopoulou C, Daraï E, Collinet P and Tuech JJ: Conservative surgery versus colorectal resection in deep endometriosis infiltrating the rectum: A randomized trial. *Hum Reprod* 33: 47-57, 2018.
- Ledu N, Rubod C, Piessen G, Roman H and Collinet P: Management of deep infiltrating endometriosis of the rectum: Is a systematic temporary stoma relevant? *J Gynecol Obstet Hum Reprod* 47: 1-7, 2018.
- Riiskjr M, Forman A, Kesmodel US, Andersen LM, Ljungmann K and Seyer-Hansen M: Pelvic pain and quality of life before and after laparoscopic bowel resection for rectosigmoid endometriosis: A prospective, observational study. *Dis Colon Rectum* 61: 221-229, 2018.
- Constantin V, Carăp A, Bobic S, Păun S, Brătîlă S, Socea B, Morosan AM and Mirancea N: Accurate diagnosis of sigmoid colon endometriosis by immunohistochemistry and transmission electron microscopy—a case report. *Chirurgia* 110: 482-485, 2015.
- Mabrouk M, Spagnolo E, Raimondo D, D'Errico A, Caprara G, Malvi D, Catena F, Ferrini G, Paradisi R and Seracchioli R: Segmental bowel resection for colorectal endometriosis: Is there a correlation between histological pattern and clinical outcomes? *Hum Reprod* 27: 1314-1319, 2012.

37. Daraï C, Bendifallah S, Foulot H, Ballester M, Chabbert-Buffet N and Daraï E: Impact of osteopathic manipulative therapy in patient with deep with colorectal endometriosis: A classification based on symptoms and quality of life. *Gynecol Obstet Fertil Senol* 45: 472-477, 2017 (In French).
38. Garavaglia E, Inversetti A, Ferrari S, De Nardi P and Candiani M: Are symptoms after a colorectal segmental resection in deep endometriosis really improved? The point of view of women before and after surgery. *J Psychosom Obstet Gynecol* 39: 248-251, 2018.
39. De Graaff AA, D'Hooghe TM, Dunselman GA, Dirksen CD and Hummelshoj L; WERF EndoCost Consortium; Simoens S: The significant effect of endometriosis on physical, mental and social wellbeing: Results from an international cross-sectional survey. *Hum Reprod* 28: 2677-2685, 2013.
40. Daraï E, Ballester M, Chereau E, Coutant C, Rouzier R and Wafo E: Laparoscopic versus laparotomic radical en bloc hysterectomy and colorectal resection for endometriosis. *Surg Endosc* 24: 3060-3067, 2010.
41. Touboul C, Ballester M, Dubernard G, Zilberman S, Thomin A and Daraï E: Long-term symptoms, quality of life, and fertility after colorectal resection for endometriosis: Extended analysis of a randomized controlled trial comparing laparoscopically assisted to open surgery. *Surg Endosc* 29: 1879-1887, 2015.
42. Araujo D, Passos RB, Souza CP, Silva AP and Marques M: Cost effectiveness analysis of goserelin empiric therapy for deep endometriosis treatment. *Value in health* 12: A164, 2009.
43. Mabrouk M, Frascà C, Geraci E, Montanari G, Ferrini G, Raimondo D, Alvisi S, Paradisi R, Villa G and Seracchioli R: Combined oral contraceptive therapy in women with posterior deep infiltrating endometriosis. *J Minim Invasive Gynecol* 18: 470-474, 2011.
44. Miller JD: Quantification of endometriosis-associated pain and quality of life during the stimulatory phase of gonadotropin-releasing hormone agonist therapy: A double-blind, randomized, placebo-controlled trial. *Am J Obstet Gynecol* 182: 1483-1488, 2000.
45. Márki G, Bokor A, Rigó J and Rigó A: Physical pain and emotion regulation as the main predictive factors of health-related quality of life in women living with endometriosis. *Hum Reprod* 32: 1432-1438, 2017.
46. Zhao L, Wu H, Zhou X, Wang Q, Zhu W and Chen J: Effects of progressive muscular relaxation training on anxiety, depression and quality of life of endometriosis patients under gonadotrophin-releasing hormone agonist therapy. *Eur J Obstet Gynecol Reprod Biol* 162: 211-215, 2012.
47. Melis I, Litta P, Nappi L, Agus M, Melis GB and Angioni S: Sexual function in women with deep endometriosis: Correlation with quality of life, intensity of pain, depression, anxiety, and body image. *Int J Sex Heal* 27: 175-185, 2015.
48. Nunes FR, Ferreira JM and Bahamondes L: Prevalence of fibromyalgia and quality of life in women with and without endometriosis. *Gynecol Endocrinol* 30: 307-310, 2014.
49. Teixeira MZ, Podgaec S and Baracat EC: Potentized estrogen in homeopathic treatment of endometriosis-associated pelvic pain: A 24-week, randomized, double-blind, placebo-controlled study. *Eur J Obstet Gynecol Reprod Biol* 211: 48-55, 2017.
50. Sesti F, Pietropolli A, Capozzolo T, Broccoli P, Pierangeli S, Bollea MR and Piccione E: Hormonal suppression treatment or dietary therapy versus placebo in the control of painful symptoms after conservative surgery for endometriosis stage III-IV. A randomized comparative trial. *Fertil Steril* 88: 1541-1547, 2007.
51. Nnoaham KE, Hummelshoj L, Webster P, d'Hooghe T, de Cicco Nardone F, de Cicco Nardone C, Jenkinson C, Kennedy SH and Zondervan KT; World Endometriosis Research Foundation Global Study of Women's Health consortium: Impact of endometriosis on quality of life and work productivity: A multi-center study across ten countries. *Fertil Steril* 96: 366-373, 2011.
52. Petrelluzzi KFS, Garcia MC, Petta CA, Grassi-Kassisse DM and Spadari-Bratfisch RC: Salivary cortisol concentrations, stress and quality of life in women with endometriosis and chronic pelvic pain. *Stress* 11: 390-397, 2008.
53. Matei AM, Draghici-Ionescu AM, Cioplea M, Zurac SA, Boda D, Serban I, Caruntu C, Ilie MA and Fekete LG: Skin endometriosis: A case report and review of the literature. *Exp Ther Med* 21: 532, 2021.
54. Grandi G, Toss A, Cortesi L, Botticelli L, Volpe A and Cagnacci A: The association between endometriomas and ovarian cancer: preventive effect of inhibiting ovulation and menstruation during reproductive life. *Biomed Res Int* 2015: 751571, 2015.
55. Lövkvist L, Boström P, Edlund M and Olovsson M: Age-related differences in quality of life in Swedish women with endometriosis. *J Womens Health (Larchmt)* 25: 646-653, 2016.
56. Jones G, Jenkinson C and Kennedy S: The impact of endometriosis upon quality of life: A qualitative analysis. *J Psychosom Obstet Gynaecol* 25: 123-33, 2004.
57. Bourdel N, Chauvet P, Billone V, Douridas G, Fauconnier A, Gerbaud L and Canis M: Systematic review of quality of life measures in patients with endometriosis. *PLoS One* 14: e0208464, 2019.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.