



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Corrigendum to “The Cumulative Rate of SARS-CoV-2 Infection in Chinese Hemodialysis Patients”  
[Kidney Int. Rep. 2020 1416–1421]

*Kidney Int Rep* (2021) 6, 2525; <https://doi.org/10.1016/j.ekir.2021.07.021>

Xin Xu<sup>1,13</sup>, Sheng Nie<sup>1,13</sup>, Jian Sun<sup>1,13</sup>, Yaozhong Kong<sup>2,13</sup>, Min Liang<sup>1</sup>, Dongfeng Li<sup>3</sup>,  
Tean Ma<sup>4</sup>, Jiaqing Peng<sup>5</sup>, Shikui Gao<sup>6</sup>, Yong Shao<sup>7</sup>, Yan Zha<sup>8</sup>, Gangyi Peng<sup>9</sup>, Chanjun Ren<sup>10</sup>,  
Ailong Huang<sup>11</sup> and Fan Fan Hou<sup>1,12</sup>

<sup>1</sup>State Key Laboratory of Organ Failure Research; National Clinical Research Center for Kidney Disease, Nanfang hospital, Southern Medical University, Guangzhou, China; <sup>2</sup>Department of Nephrology, The First People’s Hospital of Foshan, Foshan, China; <sup>3</sup>The people’s Hospital of Honghu, Honghu, China; <sup>4</sup>Department of Nephrology, The First People’s Hospital of Jingzhou, Jingzhou, China; <sup>5</sup>Department of Nephrology, Jingzhou Central Hospital, Jingzhou, China; <sup>6</sup>Department of Nephrology, Honghu Traditional Chinese Medicine Hospital, Honghu, China; <sup>7</sup>Department of Nephrology, The Second People’s Hospital of Honghu, Honghu, China; <sup>8</sup>Department of Nephrology, Guizhou Provincial People’s Hospital, Guizhou University, Guiyang, China; <sup>9</sup>Bureau of Medical Administration, Health Commission of Guangdong, Guangdong, China; <sup>10</sup>Wuhan Kingmed Medical Laboratory, Wuhan, China; <sup>11</sup>Key Laboratory of Molecular Biology on Infectious Diseases, Ministry of Education; Chongqing Medical University, Chongqing, China; and <sup>12</sup>Guangzhou Regenerative Medicine and Health Guangdong Laboratory, Guangzhou, China

<sup>13</sup>XX, SN, JS, and YK contributed equally to this work.

The authors regret that the printed version of the above article contained a number of errors. The correct and final version follows. The authors would like to apologise for any inconvenience caused.

DOI of original article: <https://doi.org/10.1016/j.ekir.2020.07.010>

DOI of corrected article: [10.1016/j.ekir.2021.07.022](https://doi.org/10.1016/j.ekir.2021.07.022)

**Correspondence:** Fan Fan Hou, Renal Division, Nanfang Hospital, Southern Medical University, 1838 North Guangzhou Avenue, Guangzhou 510515, China. E-mail: [ffhouguangzhou@163.com](mailto:ffhouguangzhou@163.com)

---