

CORRECTION

Open Access



# Correction to: Septic patients without obvious signs of infection at baseline are more likely to die in the ICU

Francesco Campanelli<sup>1</sup>, Agnès Soudry-Faure<sup>2</sup>, Aurélie Avondo<sup>1</sup>, Jean-Baptiste Roudaut<sup>3</sup>, Jean-Pierre Quenot<sup>3,4,5</sup>, Patrick Ray<sup>1</sup> and Pierre-Emmanuel Charles<sup>3,4\*</sup>

## Correction to: *BMC Infectious Diseases* (2022) 22:205

<https://doi.org/10.1186/s12879-022-07210-y>

Following publication of the original article [1], the authors identified that the given name and family name were erroneously transposed.

The incorrect authors' names list is:

Campanelli Francesco<sup>1</sup>, Soudry-Faure Agnès<sup>2</sup>, Avondo Aurélie<sup>1</sup>, Roudaut Jean-Baptiste<sup>3</sup>, Quenot Jean-Pierre<sup>3,4,5</sup>, Ray Patrick<sup>1</sup> and Charles Pierre-Emmanuel<sup>3,4\*</sup>

The correct authors' names list is:

Francesco Campanelli<sup>1</sup>, Agnès Soudry-Faure<sup>2</sup>, Aurélie Avondo<sup>1</sup>, Jean-Baptiste Roudaut<sup>3</sup>, Jean-Pierre Quenot<sup>3,4,5</sup>, Patrick Ray<sup>1</sup> and Pierre-Emmanuel Charles<sup>3,4\*</sup>

The author group has been updated above and the original article [1] has been corrected.

## Author details

<sup>1</sup>Centre Régional Universitaire Des Urgences, Hôpital F. Mitterrand, C.H.U. DIJON, Bd Mal de Lattre de Tassigny, Dijon, France. <sup>2</sup>Centre d'Investigation Clinique, Hôpital F. Mitterrand, C.H.U. Dijon, 14 rue Gafarel, Dijon, France. <sup>3</sup>Service

de Médecine Intensive Réanimation, Hôpital F. Mitterrand, C.H.U. DIJON, 14 rue Gafarel, B.P. 7790821079 Dijon Cedex, France. <sup>4</sup>Laboratoire Lipness, U.M.R. 1231, INSERM, Université de Bourgogne-Franche Comté, 7 Bd Jeanne d'Arc, Dijon, France. <sup>5</sup>INSERM, CIC 1432, Module Épidémiologie Clinique, Université de Bourgogne-Franche Comté, Dijon, France.

Published online: 16 March 2022

## Reference

1. Campanelli F, Soudry-Faure A, Avondo A, Roudaut J-B, Quenot J-P, Ray P, Charles P-E. Septic patients without obvious signs of infection at baseline are more likely to die in the ICU. *BMC Infect Dis.* 2022;22:205. <https://doi.org/10.1186/s12879-022-07210-y>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The original article can be found online at <https://doi.org/10.1186/s12879-022-07210-y>.

\*Correspondence: pierre-emmanuel.charles@chu-dijon.fr

<sup>3</sup> Service de Médecine Intensive Réanimation, Hôpital F. Mitterrand, C.H.U. DIJON, 14 rue Gafarel, B.P. 7790821079 Dijon Cedex, France

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.