

A tool for examining the role of the zinc finger myelin transcription factor 1 (Myt1) in neural development: Myt1 knock-in mice

Lynn D. Hudson · Elena Romm · Jo Ann Berndt ·
Joseph A. Nielsen

Received: 28 March 2010/Accepted: 23 November 2010/Published online: 26 January 2011
© The Author(s) 2011. This article is published with open access at Springerlink.com

Abstract The Myt1 family of transcription factors is unique among the many classes of zinc finger proteins in how the zinc-stabilized fingers contact the DNA helix. To examine the function of Myt1 in the developing nervous system, we generated mice in which Myt1 expression was replaced by an enhanced Green Fluorescent Protein fused to a Codon-improved Cre recombinase as a protein reporter. *Myt1* knock-in mice die at birth, apparently due to improper innervation of their lungs. Elimination of Myt1 did not significantly affect the number or distribution of neural precursor cells that normally express Myt1 in the embryonic spinal cord. Nor was the general pattern of differentiated neurons altered in the embryonic spinal cord. The *Myt1* knock-in mice should provide an important tool for identifying the *in vivo* targets of Myt1 action and unraveling the role of this structurally distinct zinc finger protein in neural development.

Keywords Oligodendrocytes · Transcriptional networks · Corepressors · Myelinogenesis · Zinc finger transcription factors

Introduction

During development, oligodendrocyte precursor cells arise in multiple locations and differentiate into the myelinating cells of the central nervous system. Much progress has been made in the identification of individual transcription factors required for orchestrating the multiple waves of oligodendrocyte development. (Kessaris et al. 2006; Nicolay et al. 2007; Wegner 2008). The myelin transcription factor 1 (*Myt1*) family represents zinc finger proteins of a structurally novel CCHC class that were originally cloned by their binding to the most abundantly expressed myelin gene in oligodendrocytes: proteolipid protein (*Plp1*; Kim and Hudson 1992). Myt1 zinc fingers recognize DNA in a distinct manner, with a whole zinc finger sitting snugly in the major groove of double-stranded DNA (Gamsjaeger et al. 2008). This manner of interaction contrasts with the classical GATA-type and steroid hormone zinc finger proteins, which all utilize an α -helix to position DNA-contacting residues in the major groove. Myt1 physically associates with the Sin3B protein, a corepressor that modifies chromatin structure by

L. D. Hudson (✉) · E. Romm · J. A. Berndt ·

J. A. Nielsen

Section of Developmental Genetics, National Institute of Neurologic Disorders and Stroke, National Institutes of Health, Building 1, Room 228, Bethesda, MD 20892-0166, USA
e-mail: hudson1@od.nih.gov

Present Address:

J. A. Nielsen

Center for Devices and Radiological Health, Food and Drug Administration, Silver Spring, MD, USA

binding class I histone deacetylases (HDACs; Romm et al. 2005). Of note, oligodendrocytes contain multiple forms of Sin3B, both a long form that includes a HDAC binding site and short forms that lack a HDAC binding site. Myt1 binding to target genes can recruit HDAC via Sin3B, with an accompanying localized hypo-acetylation of core histones and resultant transcriptional repression. But in the face of sufficient amounts of Sin3B short forms, the Sin3B-Myt1 complexes bound to target genes may preclude HDAC binding, leading to a state of transcriptional activation. Within the nucleus, Myt1 is located in domains that are distinct from the nucleolus, coiled bodies, and other nuclear subdomains (Armstrong et al. 1995). The discrete punctate pattern of Myt1 immunostaining coincided with the target PLP gene, which was associated with the nuclear periphery, in half of the cells observed in both progenitors and differentiated oligodendrocytes (Nielsen et al. 2002). Mature oligodendrocytes do not express appreciable amounts of Myt1 and the subcellular distribution of Myt1 shifts shortly before cells stop expressing Myt1; consequently the Myt1 protein in mature oligodendrocytes is transiently detected in the cytoplasmic compartment (Armstrong et al. 1995).

Myt1 modulates the proliferation and differentiation of oligodendrocyte lineage cells in rodents (Nielsen et al. 2004), as observed following the introduction of a dominant negative construct containing the four zinc-finger binding domain of Myt1 into oligodendrocyte primary cultures. Expression of the dominant negative construct inhibited the proliferation of progenitors as well as their differentiation into oligodendrocytes, as assessed by bromodeoxyuridine incorporation, morphology, migratory characteristics, and myelin gene expression. Myt1 alters neuronal differentiation in *Xenopus* (Bellefroid et al. 1996), and together with the basic-helix-loop-helix protein Neurogenin 3 forms a “feed-forward” expression loop to promote endocrine islet cell differentiation (Ahnfelt-Ronne et al. 2007; Wang et al. 2008). Selective knock-out of Myt1 in the pancreas compromises islet cell differentiation; of note is the induction of the paralogs Myt1l and Myt3 in these animals (Wang et al. 2007). In the nervous system, Myt1 is expressed in neuroepithelial germinal zones in rodent embryos (Kim et al. 1997) and continues to be expressed in germinal zones of adults (Armstrong et al. 1995). Myt1 cannot be detected in cells

immunostained with TuJ1, which recognizes neurons undergoing terminal mitosis and accumulates with further differentiation (Kim et al. 1997). Myt1 is upregulated in gliomas (Armstrong et al. 1997) as well as following injury: spinal cord contusion (Wrathall et al. 1998), murine hepatitis virus-induced demyelination (Vana et al. 2007) and in multiple sclerosis lesions (Vana et al. 2007).

An important tool for establishing the *in vivo* role of transcription factors whose expression patterns implicate them in oligodendrocyte development is loss-of-function transgenic mice. Loss of Myt1 is hypothesized to affect not only the commitment to the oligodendrocyte lineage but also the subsequent differentiation of oligodendrocyte progenitors and the normal myelinating functions of mature oligodendrocytes. As suggested by *in vitro* overexpression studies, the *Myt1*^{−/−} may have reduced proliferative capacity as well as altered differentiation abilities (Nielsen et al. 2004). The *Myt1* null mice may also be expected to suffer from neuronal dysfunction as suggested from the detection of Myt1 in neural precursors and selected populations of mature neurons, including motor neurons (Armstrong et al. 1995). In this paper we describe the construction and initial characterization of *Myt1* knock-in mice.

Materials and methods

Construction of the Myt1 targeting vector

The murine myelin transcription factor 1 (*Myt1*) gene is MGI:1100535 located on Chr2:181498037–181562502 bp. Synonyms include: mKIAA0835, NZF-2a, NZF-2b, Nzf2, Nztf2. A building vector containing FLP recombinase target binding sites (*frt*) flanking a neomycin gene as well as a Cre recombinase target binding sites (*loxP*; pOSfrtLoxP) was supplied by Dr. Randy Thresher, University of North Carolina. The following cassettes were subcloned into this building vector:

1. the 4.9 kb *Myt1* 5' region of homology, which corresponds to the portion of intron 1 that ends at exon 2. This segment was cloned from 129/SvEv mouse genomic DNA using the forward primer 5'TATATATAGCGGCCGCTTACCAGGTAGC

- CTAATTGG3' and the reverse primer 5'GTTA ATCGGCCGTTCTGAGCTCACCTGCAAAG3'. 2. the *eGFP* gene subcloned from peGFP-C2 (Clontech; Mountain View, CA) using *Hind*III and *Kpn*I. 3. the *iCre* recombinase subcloned from pBlue.iCre ((Shimshek et al. 2002); gift of R. Sprengel) using *Hind*III and *Kpn*I. The *iCre* was first inserted upstream of the SV40 polyadenylation site in peGFP-C2 and in frame with GFP in order to create an *eGFP-iCre* fusion protein. The coding region of the fusion protein is directly followed by the polyadenylation site. 4. the 1.4 kb *Myt1* 3' region of homology, which corresponds to intron 2 and exon 3. This segment was cloned from 129/SvEv mouse genomic DNA using the forward primer 5'GCATCGATTCA GTGCCAATGCAGTCTGGCC3' and the reverse primer 5'ATGCTAGCTGGAGTTTGGGTC ATTGTGG3'.

To ensure initiation of translation with the eGFP-iCre fusion protein, the *Myt1* initiation codon located at the beginning of exon 2 was changed to a GTG. The final 14.548 kb plasmid was sequenced *in toto*, which also confirmed that premature initiation of translation would not occur upstream of eGFP.

The NIH Institutional Biosafety Committee approved the Recombinant DNA Registration Document RD-95-I-06 for the use of this construct in mice.

Derivation and maintenance of transgenic mice

Electroporation of 129/SvEv embryonic stem cells with the *Myt1* targeting construct was carried out by Dr. Randy Thresher at the University of North Carolina Animal Models Core Facility. Clones were screened by PCR using the forward primer: 5'CACACCGC GTCACCTTAATATGCGAAG3' and the reverse primer: 5'CAGCAAGCCCTTGTAAGTCACAGT C3' in order to amplify a 3.4 Kb band in transgenic animals.

We carried out Southern analysis to identify the correctly targeted ES cells, which were subsequently implanted at UNC. Four founders were then transferred to NIH and two lines (*Myt1* A2A2 and *Myt1* C6H5) were selected for detailed analysis. To remove the neomycin cassette, mice were bred for 5 generations with B6;SJL-Tg(ACTFLPe)9205Dym/J mice

(Jackson Laboratories, Bar Harbor, Maine). This transgenic strain expresses a variant of the *Saccharomyces cerevisiae* *FLP1* recombinase gene under the direction of the human *ACTB* promoter. The *FLP1* recombinase excises the DNA sequence flanked by *frt* sites.

Southern and PCR analysis was carried out to assess whether recombination had occurred correctly at the *Myt1* locus and to document the removal of the neomycin cassette. For the PCR analysis of resulting mice, the forward primer 5'ATACGAAGTTATT CGAAAGTTCCTATTCTCTAGAAAG3' and reverse primer 5'ACCCTTAAATACTGCATCAAGGAACAA TCGAGGAGC3' detected a 1,435 bp band in mice containing the neomycin sequence and a 230 bp band in mice lacking the neomycin sequence.

Neo- animals were back-crossed for at least 10 generations against C57BL/6 mice. Each generation was genotyped by PCR, and periodically confirmed with Southern Blot experiments. The final back-crosses are maintained as two heterozygous lines lacking neomycin (*Myt1*+/*Myt1*-, *neo*-/*neo*-). All the breedings are documented in a computerized Filemaker-based animal breeding database. Animals were handled according to IACUC guidelines under animal protocol #1155-04. These "Myt1GFPicre-KnockIn" mice are freely available through the Mutant Mouse Regional Resource Centers (MMRRC) under the MMRRC ID#:031888.

Southern/Northern analysis and routine genotyping

Southern analysis was conducted on DNA purified from liver or whole embryos using a standard phenol extraction followed in some cases by a DNeasy treatment (Qiagen; Valencia, CA). Following restriction enzyme treatment, DNA was electrophoresed on 0.7% agarose gels. The marker lane consisted of the 1 kb DNA ladder Plus (Fermentas; Burlington, Ontario). Gels were pretreated with 0.2 N HCl for 10 min, denatured in 1.5 N NaCl/0.5 N NaOH for 45 min, and neutralized in 1.5 N NaCl/1 M Tris pH7.4 for 30 min prior to capillary transfer onto a positively charged nylon transfer membrane Hybond-N⁺ (Amersham; Piscataway, NJ). Probes were radiolabeled with $\alpha^{32}\text{P}$ dCTP (Roche; Branchburg, NJ), and included the following: a 605 bp probe from the *Myt1* 5'region of homology; a 791 bp probe from the

Myt1 3' region of homology; a 800 bp neomycin probe (from pPNT, LDN 33). The Myt1 5' and 3' homology probes were cloned from 129/SvEv genomic DNA using PfuUltra high fidelity DNA polymerase (Stratagene; Garden Grove, CA) with the following primers: For the 605 bp 5' homology probe, the forward primer was 5'GTTCCAAGA CAGCTATTACATGGTGGC3', and reverse primer was 5'GGAATTGTATCTCTCCAGGGTCAGTG3'. For the 791 bp 3' homology probe, the forward primer was 5'GACTGTGACTTACAAAGGGCTT GCTG3', and the reverse primer was 5'CTGACT GATGGGTACCTCTCAATAGG3'. Each PCR fragment was subcloned into the pCR-Blunt II TOPO vector (Invitrogen; Carlsbad, CA). The 605 bp 5' homology probe can be excised from the pCR-5' Myt1outSB605 plasmid with *Eco*RI, and the 791 bp 3' homology probe can be excised from the pCR-3' Myt1outSB791 plasmid with *Not*I and *Kpn*I. Hybridizations were carried out in Hybrisol I solution (Millipore; Billerica, MA) with 100 µg/ml salmon sperm DNA (Phoenix Bio Tech; Phoenix, AZ).

Northern analysis

RNA was prepared from E15 brains by homogenization in TRIzol reagent (Invitrogen) using an IKA Ultra-Turrax T8 power homogenizer, followed by precipitation with isopropyl alcohol. RNA was electrophoresed on 1% agarose gels containing ethidium bromide together with a RNA Millenium molecular weight marker (Ambion; Austin, TX), and blotted with NorthernMax-Gly reagents (Ambion). Capillary transferred onto positively charged nylon transfer membrane Hybond-N⁺(Amersham). Probes were radiolabeled with $\alpha^{32}\text{P}$ dCTP (Roche; Branchburg, NJ) and included the following: a 580 bp cDNA probe, cloned from pmMyt1 plasmid containing murine Myt1 cDNA using PfuUltra high fidelity DNA polymerase (Stratagene) with the forward primer: 5' GAATTCGGAGCTCTCAATGGTT-CATC 3', and the reverse primer: 5'GAATTCCAG CTTGCTTGATCGTTCC 3'. This 580 bp PCR fragment was then subcloned into the pCR-Blunt II TOPO vector (Invitrogen) to generate the pCR-Myt1NB580 plasmid. The 580 bp Myt1 cDNA fragment can be excised from the Myt1NB580

plasmid with *Eco*RI or directly labeled with $\alpha^{32}\text{P}$ dCTP (Roche). Hybridizations were carried out in Hybrisol I solution (Millipore) with 100 µg/ml salmon sperm DNA (Phoenix Bio Tech).

Genotyping by PCR

Tail or toe lysates were prepared by incubating samples from 3–16 h in 50 µl of 1 M Tris pH 8.0, 5 M NaCl, 0.5 M EDTA, 10% Tween 20, 10% NP-40, 0.4 µg/ml freshly added proteinase K. Following centrifugation to remove debris, 1 µl of DNA lysate was added to 0.1 µl of each primer (0.4 µl total), 0.2 µl Taq DNA polymerase (Promega), 5 µl 1% Triton X-100 and 18.4 µl H₂O in EasyStart Micro 50 tubes (M β P; San Diego, CA). Two primer pairs were used in each tube in order to discriminate between transgenic homozygous (*Myt1*^{−/−}), heterozygous (*Myt1*[±]), and wild-type mice (*Myt1*^{+/+}). To amplify a 325 bp transgenic fragment, the forward primer was 5'CAGCCATACCACTACATTGAGAGG3' and the reverse primer was 5'TGCATTGGCACTGAATC GATGG3'. To amplify a 188 bp wild type fragment, the forward primer was 5'CCAACCTGCTTCTTT TGGTCCC3' and the reverse primer was 5'GGCT CGCTTGTCACTCACTTC3'. The PCR cycling was carried out for a total of 30 cycles in a Hybaid, PCR Express HBPX110 (Thermo; Essex, UK) under the following conditions: melt at 94°C for 5 min, denature at 94°C for 30 s, anneal at 55°C for 30 s, extend 72°C for 1 min (30 cycles). PCR products were analyzed on a 2% agarose gel containing ethidium bromide. The marker lane consists of the 100 bp DNA ladder Plus (Fermentas). In this assay, *Myt1*^{−/−} mice display a single 325 bp band, *Myt1*[±] display both the 325 bp and 188 bp bands, and *Myt1*^{+/+} mice display a single 188 bp band.

Immunohistochemistry and statistical analysis

E18 spinal cords were immersion fixed in phosphate-buffered 4% paraformaldehyde with picric acid (FD Neurotechnologies; Baltimore, MD). Following cryo-protection treatment in 0.1 M phosphate buffer (pH 7.4) containing 20% sucrose for 72 h at 4°C, spinal cords were rapidly frozen and stored at −75°C. Serial cryostat sections (20 µm) were cut transversely

through the whole spinal cord. Every 1st, 2nd and 3rd section of each series of 10 sections were mounted on Superfrost Plus microscope slides (3 sections per slide). Upon drying, sections were stored at -20°C . Following washes in 0.01 M phosphate-buffered saline (PBS; pH 7.4), sections were incubated overnight at 4°C in PBS containing 1% normal donkey serum, 4% BSA (Jackson ImmunoResearch, West Grove, PA), 0.3% Triton X-100 (Sigma, St. Louis, MO), the polyclonal rabbit anti-eGFP IgG (1:200, Invitrogen, Carlsbad, CA) and the biotinylated monoclonal mouse anti-NeuN antibody (1:100, Millipore, Billerica, MA). This was followed by incubation of sections at room temperature in the PBS/Triton X solution containing Alexa Fluor[®] 488 donkey anti-rabbit IgG (Molecular Probe, Eugene, OR) for 1 h, and then in the PBS/Triton X containing streptavidin conjugated Alexa Fluor[®] 594 (Molecular Probe) for another hour. Each step was followed by washes in PBS. Sections were coverslipped using Vectashield[®] with DAPI (Vector Labs, Burlingame, CA) and stored at 4°C .

Photographs of fixed 18-day mouse embryo spinal cord sections were captured with the Olympus 1 \times 70 fluorescent microscope using the 4X UPLanF1 objective lens. The photographs were processed and digitized using the CoolSNAP fX camera (Photometrics, Roper Scientifics, Tucson, AZ) and the IP Lab version 3.9.5 rl program (Scientific Image Processing, BDBiosciences, San Jose, CA). The photographs were further processed with Adobe Photoshop CS3, Version 10.0.1 (Adobe Systems Inc., San Jose, CA) utilizing the Channel window to prepare separate fluorescent images for the eGFP (Alexa Fluor[®]488), NeuN (Alexa Fluor[®]594) and DAP.

Cells that fluoresced for eGFP, NeuN and DAPI were counted using the ImageJ program (Rasband, W.S., ImageJ, National Institutes of Health, Bethesda, MD, USA, <http://rsb.info.nih.gov/ij/> 1997–2009 (also see Abramoff et al. 2004). Two separate sets of 18 day embryo spinal cords from the wild type, heterozygote and homozygote animals were counted. The cells in fourteen sections representing 14 different areas of each 18 day mouse embryo spinal cord (wild type, heterozygote and homozygote) from the cervical to sacral were counted. The 2 sets were averaged and graphed with Microsoft Excel (Microsoft Corp., Redmond, WA).

Results

Generating *Myt1* knock-in mice

Myt1 can be transcribed from two alternative promoters to yield either a 6 zinc finger protein (6ZF*Myt1* or *Myt1a*) or a 7 zinc finger protein (7ZF*Myt1* or *Myt1b*; Matsushita et al. 2002; Nielsen et al. 2004). 7ZF*Myt1* is by far the predominant product of the *Myt1* locus in all tissues examined (Matsushita et al. 2002), (Nielsen et al. 2004); (Gu et al. 2004). To establish the *in vivo* role of *Myt1*, we designed a knock-in construct so that the promoter for the seven zinc finger *Myt1* isoform (7ZF*Myt1*) would drive expression of a green fluorescent protein (eGFP) reporter together with the *cre* recombinase (Fig. 1). Cassettes for these two reporter genes were fused in frame, so that translation of the eGFP-iCre fusion protein would occur from the *eGFP-iCre* initiation codon within exon 2. The initiation codon for 7ZF*Myt1* in exon 2 was altered. Also included was a *neo* selectable marker flanked with *frt* sites so that the *neo* cassette, which can impact the expression of the targeted gene, could be subsequently excised by crossing transgenic mice with mice expressing the flip recombinase.

Correctly recombined transgenes were documented in two lines of *Myt1* knock-in mice by Southern analysis (Fig. 2). Using a probe that is distal to the 5' region of homology (panel b), the expected 5.9 kb *KpnI* band (lanes 2 and 3) and 6.3 kb *HindIII* band (lanes 5 and 6) were observed in both *Myt1*[±], *Neo*[±] and *Myt1*^{−/−}, *Neo*^{−/−} mice. Using a probe that is distal to the 3' region of homology (panel a), the expected 8.4 kb *HindIII* band was observed in *Myt1*[±], *Neo*[±] (lane 2) and *Myt1*^{−/−}, *Neo*^{+/+} (lane 3) animals. This band was also detectable with a *neo* probe (lane 8). A smaller *HindIII* band of 7.2 kb was observed in animals that were derived from matings with mice expressing the flip recombinase (lanes 4 and 5), indicating that *neo* had been excised from the targeted *Myt1* locus in these animals. The 5' region of homology probe and the 3' region of homology probe together confirm the correct homologous recombination of the electroporated knock-in construct.

No message corresponding to *Myt1* was detectable by Northern analysis in homozygous *Myt1* knock-in mice at embryonic day 15, a time when *Myt1* is normally highly expressed (lanes 3 and 6, Fig. 3).




Fig. 1 Construction of Myt1 knock-in Mice. The targeting vector is designed to eliminate exon 2, which contains the initiation codon for 7ZFMyt1, (depicted as AUG with a red arrow) and drive expression of the eGFP/cre fusion gene directly from the 7ZFMyt1 promoter. The probes used on Southern blots to document the recombinant event are shown in pink and are distal to the 5' region of homology (in red) and

the 3' region of homology (in light blue). The frt recombination sites flanking the Neomycin (*Neo*) are marked with triangles. Note that following excision of the neomycin cassette by breeding with mice expressing the flip recombinase, a single frt site remains, together with a loxP site. The 22 exons are pictured as boxes (not drawn to scale). (Color figure online)




Fig. 2 Southern Analysis of Myt1 knock-in Mice. In panel a, Hind III-digested genomic DNA was hybridized with either the 3' Myt1 outside region of homology probe (lanes 1–5) or the neo probe (lanes 7–9). lane 1 wild type DNA: Myt1^{+/+}, Neo^{-/-}; lane 2 Myt1[±], Neo[±]; lane 3 Myt1^{-/-}, Neo^{++/+}; lane 4 Myt1[±], Neo^{-/-}; lane 5 Myt1^{-/-}, Neo^{-/-}; lane 6 no DNA; lane 7 Myt1^{++/+}, Neo^{-/-}; lane 8 Myt1^{-/-}, Neo^{++/+}; lane 9 Myt1^{-/-}, Neo^{-/-}. The 3' Myt1 probe identifies a 12.2 kb HindIII band in wild type mice and an 8.4 kb HindIII band in transgenics, which also displays hybridization to the neo probe. In transgenics lacking Neo (lanes 4 and 5), the 3' Myt1 probe

identifies a 7.2 kb HindIII band. In panel b, KpnI (lanes 1–3) or HindIII-digested (lanes 4–6) genomic DNA was hybridized with the 5' Myt1 outside region of homology probe. lane 1 Myt1^{++/+}, Neo^{-/-}; lane 2 Myt1[±], Neo[±]; lane 3 Myt1^{-/-}, Neo^{++/+}; lane 4 Myt1^{++/+}, Neo^{-/-}; lane 5 Myt1[±], Neo[±]; lane 6 Myt1^{-/-}, Neo^{++/+}. The 5' Myt1 probe identifies an 8.7 kb KpnI band in wild type mice and a 5.9 kb KpnI band in transgenics. It also reveals a 12.2 kb HindIII band in wild type mice and a 6.3 kb HindIII band in transgenics. Both panels include a 1 kb DNA ladder Plus as a molecular weight marker

Heterozygous embryos expressed approximately half as much of the 5.1 kb *Myt1* transcripts at E15 when either total RNA (lane 2) or polyA-selected RNA (lane 5) was analyzed.

Phenotype of *Myt1* knock-in mice

In our initial PCR genotyping and Southern blot analysis of transgenic animals, no homozygous animals were present in litters analyzed at postnatal day 7. To determine if there was a viability problem with homozygous animals, and to pinpoint the time at which survival may be affected, we set up a series of matings between heterozygous animals. After collecting embryos throughout prenatal development, we determined that homozygous animals survived through the prenatal period but died shortly after birth. Table 1 summarizes the results from an embryonic and a postnatal time point. At embryonic day 18, the observed numbers of wild type, homozygous and heterozygous closely approximate the expected Mendelian ratio, with 26.8% of the offspring representing homozygous animals. No living homozygous animals were found from the 15 matings analyzed at birth or the 18 matings that were analyzed at later postnatal ages. One mutant allele had no effect on viability.

To determine the cause of death, heterozygous animals were mated and a Caesarean section performed shortly before the scheduled birth. Pups were kept at 37°C during the hour observation period. Unlike wild type or heterozygous animals, the homozygous animals all died within 20 min. Homozygous animals never progressed from gasping to normal breathing. Their lungs did not appear white in color and were not inflated as evidenced by their lack of buoyancy when placed in a container of phosphate buffered saline.

Heterozygous animals appeared undistinguishable from wild type animals and reproduced normally.

Neural cells with an active *Myt1* promoter can be tracked with eGFP in knock-in mice

Myt1 expressing cells in null or heterozygous embryos can be identified through the eGFP marker (Fig. 4). At E18, eGFP cells are distributed throughout the spinal cord (panels c). These eGFP cells were





Fig. 3 Northern Analysis of the *Myt1* knock-in Mice. This analysis was carried out after the crossing with B6;SJL-Tg(ACFLPe)9205Dym/J mice that express the *Saccharomyces cerevisiae* *FLP1* recombinase gene. All mice were genotyped to verify that the neo gene was absent. In panel a, a *Myt1* cDNA probe identified a 5.1 kb band in either total RNA (lanes 1–3) or mRNA (lanes 4–6) in mice containing at least one wild-type *Myt1* allele. lane 1 *Myt1*^{+/+}, lane 2 *Myt1*^{+/−}, lane 3 *Myt1*^{−/−}, RNA Millenium molecular weight marker; lane 4 *Myt1*^{+/+}, lane 5 *Myt1*^{+/−}, lane 6 *Myt1*^{−/−}. No *Myt1* transcripts in any size range are detectable in the null mice. In panel b, the ethidium bromide stained gel shows the integrity of the RNA samples

not double labeled with Neuronal Nuclei protein (NeuN; panels a), suggesting that *Myt1* neural progenitors have not differentiated into the mature neurons identified by NeuN staining at this age. Moreover, the pattern of NeuN staining in transgenic embryos was similar to that in wild type embryos (panels b). The lack of overlap between eGFP-positive cells and NeuN cells is shown in a higher

Table 1 Genotyping the progeny from *Myt1* \pm Matings

Genotype	E18 embryos (from 48 matings)			Postnatal day 7 mice (from 18 matings)		
	<i>Myt1</i> +/+	<i>Myt1</i> \pm	<i>Myt1</i> -/-	<i>Myt1</i> +/+	<i>Myt1</i> \pm	<i>Myt1</i> -/-
Number of offspring	102	190	107	37	52	0
% of offspring	25.6	47.6	26.8	41.6	58.4	0

**Fig. 4** Immunohistochemistry of Spinal Cord Sections from *Myt1* knock-in Mice. Spinal cord sections from E18 wild type mice (+/+), heterozygotes (\pm) or null mice (-/-) were triple stained with: panel **a**, eGFP (green), NeuN (red) and DAPI.

panel **b** shows NeuN alone; panel **c** shows eGFP alone; and panel **d** shows DAPI alone. The size bar = 10 microns. (Color figure online)

magnification view of a heterozygous spinal cord (Fig. 5). Quantitation of the immunohistochemistry is shown in Fig. 6. No significant difference in the number of NeuN cells was apparent in null or heterozygous animals compared to wild type. Likewise, there was no significant difference in the number of eGFP cells in heterozygous or null animals.

Discussion

The *Myt1* knock-in mice should provide an invaluable resource for: (1) defining the role of the *Myt1* gene family in the specification of neural lineages and the development of oligodendrocytes, and (2) determining the set of gene targets that this structurally




Fig. 5 Immunohistochemistry of Spinal Cord Sections from Myt1 knock-in Mice. A spinal cord section from an E18 homozygote ($-/-$) mouse was triple stained with eGFP (green), NeuN (red) and DAPI. The size bar = 10 microns. (Color figure online)

unique class of zinc finger transcription factors binds to in vivo. Recently, attention to the Myt1 gene family has focused on the discovery that the Myt1 paralog, Myt1l, is one of a simple trio of genes necessary for the efficient conversion of mouse fibroblasts into functional neurons in vitro (Vierbuchen et al. 2010). While documenting that a Myt1 gene family member is instrumental in neural specification, this advance is short on the molecular details of what genes are activated or repressed and in what stages of the cells. In the Myt1 knock-in mice, lineage tracing is possible due to the incorporation of a *cre* coding region fused in frame to the GFP reporter. Heterozygous transgenic animals mated to the ROSA26 Cre indicator strain (Soriano 1999) will display β -galactosidase expression in all Myt1 lineage cells. Additional Cre-reporter strains that would be helpful include the Z/AP line that contains a human placental alkaline phosphatase (hPLAP) reporter (Lobe et al. 1999) and the R26R-EYFP line with an enhanced yellow fluorescent protein (EYFP) reporter (Srinivas et al. 2001).

Elimination of Myt1 expression disrupts neural development. Another Myt1 transgenic model constructed by Gu and colleagues displayed a breathing defect attributable to improper innervation of the




Fig. 6 Quantitation of Immunohistochemistry. The average of the total number of cells labeled with NeuN, eGFP or DAPI per E18 spinal cord section are shown for wild type mice (+/+; blue bars), heterozygotes (\pm ; red bars) or null mice ($-/-$; yellow bars). Black lines represent standard deviations. (Color figure online)

diaphragm in the transgenic mice (Wang et al. 2007). Likewise, the homozygous knock-in Myt1 mice in the present study presented with breathing difficulties shortly after birth. Given that the transgenic line constructed by Gu and colleagues is potentially a dominant negative model, since altered Myt1 transcripts are present (Wang et al. 2007), it is reassuring that this phenotype is seen in both transgenic models. The new data provided in the present study includes the observation that eliminating Myt1 does not affect the number or distribution of neurons. These data would not have been predicted based on earlier observations that Myt1 is expressed by rat cortical neuronal progenitors at E11 and is up-regulated 2.7-fold between E11 and E13 (Nielsen et al. 2009). In the Myt1 eGFP/cre knock-in mice, the distribution and number of differentiated neurons in the spinal cord appears normal, as does the pattern of neural

cells in which the Myt1 promoter actively drives the expression of the eGFP reporter (Figs. 4, 5 and 6). Of note, the eGFP reporter is not expressed by the neuronal population labeled by NeuN in the spinal cord. Oligodendrocytes and motor neurons were thought to derive from a common precursor expressing Olig 1 and Olig 2 (Zhou and Anderson 2002; Lu et al. 2000), but subsequent fate mapping studies identified multiple sources generating oligodendrocytes in the brain and spinal cord (Kessaris et al. 2006; Cai et al. 2005). Further lineage tracing studies with the cre reporter may help resolve the origin of some of these oligodendrocyte subpopulations.

Myt1 is also critical for islet cell differentiation, as shown in a series of elegant experiments by Gu and Hecksher-Sorensen and their colleagues (Ahnfelt-Ronne et al. 2007; Wang et al. 2007, 2008). The loss-of-function Myt1 model constructed by targeting exon 9 has a Myt1 mRNA level reduced by over fourfold; heterozygous mice displayed normal pancreatic development, making it unlikely that the Myt1- allele behaves in a dominant negative fashion in this transgenic model (Wang et al. 2007). Compensatory activation of the paralogs Myt1l and Myt3 occurs in the embryonic Myt1^{-/-} pancreas. Of particular interest is the observation that Myt1 and neurogenin 3 (Ngn3) positively regulate each other's expression in the developing pancreas, with Notch mediated signaling regulating the number of Ngn3+ cells. A Notch signaling pathway is likewise a feature of the switch from neurogenesis to gliogenesis in the spinal cord, which involves not only the induction of Olig2 expression, but also the down-regulation of neurogenin 1 (Ngn1) and neurogenin 2 (Ngn2) via Notch signaling (Zhou et al. 2001).

Genomic approaches to oligodendrocyte biology have greatly expanded our horizon by showing the multiple different temporal transcription factor expression patterns occurring during oligodendrocyte differentiation. (Dugas et al. 2006; Nielsen et al. 2006). While Myt1 is down-regulated during oligodendrocyte differentiation, Myt is in the company of sixty other transcription factors that are up-regulated over twofold in cortical neuronal progenitors between embryonic day 11 and 13 (Nielsen et al. 2009). Seventy percent of the transcription factors up-regulated in cultured oligodendrocytes during differentiation in the absence of neurons are also up-regulated in acutely isolated developing oligodendrocytes (Dugas

et al. 2006; Nielsen et al. 2006). Many of these transcription factors await functional studies before their role in the complex network regulating oligodendrocyte development can be assigned. In Myt1 knock-in mice, the absence of Myt1 is expected to lead to the inactivation of an array of target genes. Transgenic mice provide a handle for identifying target genes of transcription factors when combined with the powerful “omics” technologies to identify which transcripts are over or under-expressed in the presence or absence of a given transcription factor. Given the wealth of molecular detail available on how Myt1 burrows into the DNA groove and recruits histone deacetylase to regulate neural transcription, the Myt1 knock-in mice offer an invaluable tool for further exploration of the transcriptional circuitry in neural cells.

Acknowledgments The authors thank Dr. Eva Mezey for insights on quantitative immunohistochemistry, Wayne Rasband for assistance with the imageJ program, and Dr. Heinz Arnheiter for training and access to his computerized Filemaker-based animal breeding database program. Dr. Gabor Lovas' insights on the phenotype of the knock-in mice are gratefully acknowledged. This work was supported by National Institute of Neurological Disorders and Stroke (NINDS)/National Institutes of Health intramural funds to LDH for project# 1 Z01 NS002528.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

- Abramoff MD, Magelhaes PJ, Ram SJ (2004) Image processing with image. *J Biophotonics Int* 11(7):36–42
- Ahnfelt-Ronne J, Hald J, Bodker A, Yassin H, Serup P, Hecksher-Sorensen J (2007) Preservation of proliferating pancreatic progenitor cells by Delta-Notch signaling in the embryonic chicken pancreas. *BMC Dev Biol* 7:63
- Armstrong RC, Kim JG, Hudson LD (1995) Expression of myelin transcription factor I (MyTI), a “zinc-finger” DNA-binding protein, in developing oligodendrocytes. *Glia* 14:303–321
- Armstrong RC, Migneault A, Shegog ML, Kim JG, Hudson LD, Hessler RB (1997) High-grade human brain tumors exhibit increased expression of myelin transcription factor I (MYTI), a zinc finger DNA-binding protein. *J Neuropathol Exp Neurol* 56:772–781
- Bellefroid EJ, Bourguignon C, Hollemann T, Ma Q, Anderson DJ, Kintner C, Pieler T (1996) X-MyT1, a Xenopus

- C2HC-type zinc finger protein with a regulatory function in neuronal differentiation. *Cell* 87:1191–1202
- Cai J, Qi Y, Hu X, Tan M, Liu Z, Zhang J, Li Q, Sander M, Qiu M (2005) Generation of oligodendrocyte precursor cells from mouse dorsal spinal cord independent of Nkx6 regulation and Shh signaling. *Neuron* 45:41–53
- Dugas JC, Tai YC, Speed TP, Ngai J, Barres BA (2006) Functional genomic analysis of oligodendrocyte differentiation. *J Neurosci* 26:10967–10983
- Gamsjaeger R, Swanton MK, Kobus FJ, Lehtomaki E, Lowry JA, Kwan AH, Matthews JM, Mackay JP (2008) Structural and biophysical analysis of the DNA binding properties of myelin transcription factor 1. *J Biol Chem* 283:5158–5167
- Gu G, Wells JM, Dombkowski D, Preffer F, Aronow B, Melton DA (2004) Global expression analysis of gene regulatory pathways during endocrine pancreatic development. *Development* 131:165–179
- Kessaris N, Fogarty M, Iannarelli P, Grist M, Wegner M, Richardson WD (2006) Competing waves of oligodendrocytes in the forebrain and postnatal elimination of an embryonic lineage. *Nat Neurosci* 9:173–179
- Kim JG, Hudson LD (1992) Novel member of the zinc finger superfamily: a C2-HC finger that recognizes a glia-specific gene. *Mol Cell Biol* 12:5632–5639
- Kim JG, Armstrong RC, Agoston D, Robinsky A, Wiese C, Nagle J, Hudson LD (1997) Myelin transcription factor 1 (Myt1) of the oligodendrocyte lineage, along with a closely related CCHC zinc finger, is expressed in developing neurons in the mammalian central nervous system. *J Neurosci Res* 50:272–290
- Lobe CG, Koop KE, Kreppner W, Lomeli H, Gertsenstein M, Nagy A (1999) Z/AP, a double reporter for cre-mediated recombination. *Dev Biol* 208:281–292
- Lu QR, Yuk D, Alberta JA, Zhu Z, Pawlitzky I, Chan J, McMahon AP, Stiles CD, Rowitch DH (2000) Sonic hedgehog-regulated oligodendrocyte lineage genes encoding bHLH proteins in the mammalian central nervous system. *Neuron* 25:317–329
- Matsushita F, Kameyama T, Marunouchi T (2002) NZF-2b is a novel predominant form of mouse NZF-2/Myt1, expressed in differentiated neurons especially at higher levels in newly generated ones. *Mech Dev* 118:209–213
- Nicolay DJ, Doucette JR, Nazarali AJ (2007) Transcriptional control of oligodendrogenesis. *Glia* 55:1287–1299
- Nielsen JA, Hudson LD, Armstrong RC (2002) Nuclear organization in differentiating oligodendrocytes. *J Cell Sci* 115:4071–4079
- Nielsen JA, Berndt JA, Hudson LD, Armstrong RC (2004) Myelin transcription factor 1 (Myt1) modulates the proliferation and differentiation of oligodendrocyte lineage cells. *Mol Cell Neurosci* 25:111–123
- Nielsen JA, Maric D, Lau P, Barker JL, Hudson LD (2006) Identification of a novel oligodendrocyte cell adhesion protein using gene expression profiling. *J Neurosci* 26:9881–9891
- Nielsen JA, Lau P, Maric D, Barker JL, Hudson LD (2009) Integrating microRNA and mRNA expression profiles of neuronal progenitors to identify regulatory networks underlying the onset of cortical neurogenesis. *BMC Neurosci* 10:98
- Romm E, Nielsen JA, Kim JG, Hudson LD (2005) Myt1 family recruits histone deacetylase to regulate neural transcription. *J Neurochem* 93:1444–1453
- Shimshek DR, Kim J, Hubner MR, Spergel DJ, Buchholz F, Casanova E, Stewart AF, Seeburg PH, Sprengel R (2002) Codon-improved Cre recombinase (iCre) expression in the mouse. *Genesis* 32:19–26
- Soriano P (1999) Generalized lacZ expression with the ROSA26 Cre reporter strain. *Nat Genet* 21:70–71
- Srinivas S, Watanabe T, Lin CS, William CM, Tanabe Y, Jessell TM, Costantini F (2001) Cre reporter strains produced by targeted insertion of EYFP and ECFP into the ROSA26 locus. *BMC Dev Biol* 1:4
- Vana AC, Lucchinetti CF, Le TQ, Armstrong RC (2007) Myelin transcription factor 1 (Myt1) expression in demyelinated lesions of rodent and human CNS. *Glia* 55:687–697
- Vierbuchen T, Ostermeier A, Pang ZP, Kokubu Y, Sudhof TC, Wernig M (2010) Direct conversion of fibroblasts to functional neurons by defined factors. *Nature* 463:1035–1041
- Wang S, Zhang J, Zhao A, Hipkens S, Magnuson MA, Gu G (2007) Loss of Myt1 function partially compromises endocrine islet cell differentiation and pancreatic physiological function in the mouse. *Mech Dev* 124:898–910
- Wang S, Hecksher-Sorensen J, Xu Y, Zhao A, Dor Y, Rosenberg L, Serup P, Gu G, Ahnfelt-Ronne J, Hald J, Bodker A, Yassin H (2008) Myt1 and Ngn3 form a feed-forward expression loop to promote endocrine islet cell differentiation. Preservation of proliferating pancreatic progenitor cells by Delta-Notch signaling in the embryonic chicken pancreas. *Dev Biol* 317:531–540
- Wegner M (2008) A matter of identity: transcriptional control in oligodendrocytes. *J Mol Neurosci* 35:3–12
- Wrathall JR, Li W, Hudson LD (1998) Myelin gene expression after experimental contusive spinal cord injury. *J Neurosci* 18:8780–8793
- Zhou Q, Anderson DJ (2002) The bHLH transcription factors OLIG2 and OLIG1 couple neuronal and glial subtype specification. *Cell* 109:61–73
- Zhou Q, Choi G, Anderson DJ (2001) The bHLH transcription factor Olig2 promotes oligodendrocyte differentiation in collaboration with Nkx2.2. *Neuron* 31:791–807