



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review

Review of Non-bacterial Infections in Respiratory Medicine: Viral Pneumonia[☆]



José María Galván,^a Olga Rajas,^b Javier Aspa^{b,*}

^a Servicio de Medicina Interna, Hospital Universitario de la Princesa, IIS-IP, Madrid, Spain

^b Servicio de Neumología, Hospital Universitario de la Princesa, IIS-IP, Madrid, Spain

ARTICLE INFO

Article history:

Received 6 October 2014

Accepted 23 February 2015

Available online 3 October 2015

Keywords:

Viral pneumonia

Influenza virus

Respiratory syncytial virus

Coronavirus

Co-infection

Procalcitonin

Polymerase chain reaction

Antivirals

Vaccination

ABSTRACT

Although bacteria are the main pathogens involved in community-acquired pneumonia, a significant number of community-acquired pneumonias are caused by viruses, either directly or as part of a co-infection. The clinical picture of these different pneumonias can be very similar, but viral infection is more common in the pediatric and geriatric populations, leukocytes are not generally elevated, fever is variable, and upper respiratory tract symptoms often occur; procalcitonin levels are not generally affected. For years, the diagnosis of viral pneumonia was based on cell culture and antigen detection, but since the introduction of polymerase chain reaction techniques in the clinical setting, identification of these pathogens has increased and new microorganisms such as human bocavirus have been discovered. In general, influenza virus type A and syncytial respiratory virus are still the main pathogens involved in this entity. However, in recent years, outbreaks of deadly coronavirus and zoonotic influenza virus have demonstrated the need for constant alert in the face of new emerging pathogens. Neuraminidase inhibitors for viral pneumonia have been shown to reduce transmission in cases of exposure and to improve the clinical progress of patients in intensive care; their use in common infections is not recommended. Ribavirin has been used in children with syncytial respiratory virus, and in immunosuppressed subjects. Apart from these drugs, no antiviral has been shown to be effective. Prevention with anti-influenza virus vaccination and with monoclonal antibodies, in the case of syncytial respiratory virus, may reduce the incidence of pneumonia.

© 2014 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Revisión sobre las infecciones no bacterianas del aparato respiratorio: neumonías víricas

RESUMEN

Aunque las bacterias son los principales patógenos involucrados en la neumonía adquirida en la comunidad, algunos virus son responsables directos o en coinfección de un importante número de neumonías adquiridas en la comunidad. La clínica de estas neumonías puede ser muy similar, en el caso de los virus afectan más frecuentemente a la población infantil y geriátrica, con frecuencia no elevan la cifra de leucocitos, la fiebre es inconstante y frecuentemente se acompañan de síntomas de vías respiratorias altas. Característicamente no elevan la procalcitonina. Durante años el diagnóstico ha recaído en cultivos celulares y en detección de antígenos; desde la incorporación en la clínica de la PCR, la identificación de estos patógenos ha aumentado, descubriéndose nuevos microorganismos como el bocavirus. En general, el virus influenza A y el virus respiratorio sincitial siguen siendo los principales virus implicados. Sin

Palabras clave:

Neumonía vírica

Virus influenza

Virus respiratorio sincitial

Coronavirus

Coinfección

Procalcitonina

Reacción en cadena de la polimerasa

Antivirales

Vacunación

[☆] Please cite this article as: Galván JM, Rajas O, Aspa J. Revisión sobre las infecciones no bacterianas del aparato respiratorio: neumonías víricas. Arch Bronconeumol. 2015;51:590–597.

* Corresponding author.

E-mail address: jasper@separ.es (J. Aspa).

embargo, la irrupción en los últimos años de epidemias con alta letalidad de coronavirus y de zoonosis de virus influenza hace que sea necesario mostrarse alerta ante estos nuevos patógenos emergentes. Los inhibidores de la neuraminidasa para neumonías víricas han demostrado disminuir la transmisión en casos expuestos y mejorar la evolución clínica en pacientes en Cuidados Intensivos; su uso en infecciones banales no está recomendado. La ribavirina ha sido utilizada en niños con infecciones por virus respiratorio sincitial, así como en inmunodeprimidos. Fuera de estos fármacos, ningún otro antiviral ha probado su eficacia. Las medidas de prevención con vacunación para virus influenza y con anticuerpos monoclonales para virus respiratorio sincitial podrían disminuir la incidencia de neumonía.

© 2014 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

According to WHO estimates for 2012, around 450 million cases of pneumonia occur worldwide every year, causing 3 million deaths, and accounting for 5.5% of overall mortality worldwide.^{1,2} It is the fourth cause of death worldwide, and is a particularly a serious threat to children and the elderly.^{1,3}

Bacterial infections, being more common, have been more extensively studied. In contrast, research into viral community-acquired pneumonia (CAP), despite its growing epidemiological significance in developing countries and in the pediatric population, has been limited.⁴ Assuming the rate of diagnosis to still be lower than real incidence, around 200 million cases of viral pneumonia occur annually throughout the world, half of which are in children.⁵ Viral pneumonia is of great interest due to its impact on infant mortality, its role as a facilitator of bacterial infections (co-infections), and its ease of transmission, a factor which has transformed it into a worldwide threat.

In this review, we will focus on CAP caused by respiratory viruses in immunocompetent patients.

Differentiating Between Viral and Bacterial Pneumonia

It is important to distinguish between CAP of viral and bacterial origin. Clinical, radiological and laboratory variables that are commonly used to distinguish between these entities are listed in Table 1.

Epidemiological studies have been published by many authors. Ruiz-González et al.⁶ included patients with viral and intracellular bacterial pneumonia in the same group. They concluded that pneumonia caused by intracellular pathogens affected older patients, had a more insidious disease course, and often did not produce leukocytosis. Johnstone et al.⁷ found that patients with viral pneumonia had more cardiac comorbidities and were older; Ma et al.⁸ found that institutionalization led to a greater risk of viral CAP. Liu et al.⁹ reported that viral pneumonia caused more cough and less pleuritic pain, while Jennings et al.¹⁰ found myalgia to be the symptom most commonly associated with the viral disease. Despite the many publications, predicting the viral etiology from clinical

parameters is difficult and often inaccurate, as affirmed recently by Viasus et al.¹¹

With regard to radiological changes, focal alveolar infiltrates have been traditionally associated with the bacterial entity, and bilateral, interstitial infiltrates with the viral form.¹² However, recent studies using chest computed tomography (CT) have shown that a viral etiology cannot be ruled out by the appearance of localized alveolar infiltrates, and these may even signal the onset of many viral pneumonias.¹³

Viral infections do not usually affect the number of leukocytes, so the use of acute phase reactants, such as procalcitonin, as biomarkers may be of great help in reaching a diagnosis.¹⁴ Procalcitonin production depends on the presence of circulating tumor necrosis factor (TNF- α); in viral infections, macrophages produce interferon- α that can inhibit TNF- α , suppressing the elevation of procalcitonin, thus suggesting a viral origin.¹⁵

Despite these assertions, there is no gold standard for differentiating the etiology of pneumonia.¹⁶ Moreover, we must not forget that CAP, whether viral or bacterial, is a dynamic entity: differences in biomarker values or the appearance of infiltrates on radiology are only snapshots of an active process that can vary widely from day to day.

Types of Viruses

The list of viruses that can cause respiratory infection is long (Table 2). In this review, we will focus particularly on seasonal respiratory viruses.

Other viruses that have recently received considerable media attention, such as H5N1 influenza virus or coronaviruses (responsible for severe acute respiratory syndrome or the Middle East respiratory syndrome, MERS-CoV) will be examined in less depth.

The relative incidence and clinical and epidemiological characteristics of the different respiratory viruses are listed in Table 3.

Overall, syncytial respiratory virus (SRV) remains the primary causative agent of CAP in children, and the main cause of severe pneumonia in this population.^{17,18} Since the widespread introduction of PCR techniques, rhinoviruses are more often detected as the causative agent of viral pneumonia in children.^{19–22} However, because it is so frequently detected in asymptomatic individuals

Table 1
Differential Factors Between Viral Pneumonia and Bacterial Pneumonia.

	Suggestive of viral origin	Suggestive of bacterial origin
Age	Younger than 5 and older than 65 years	Adults
Epidemic status	Seasonal or epidemic outbreaks	Throughout the year
Disease course	Slow onset	Rapid onset
Clinical profile	Most frequently rhinitis and wheezing	Most frequently high fever and tachypnea
Total leukocyte count on admission	<10 \times 10 ⁶ c/L	>15 \times 10 ⁶ c/L and <4 \times 10 ⁶ c/L
C-reactive protein on admission	<20 mg/L	>60 mg/L
Serum procalcitonin on admission	<0.1 μ g/L	>0.5 μ g/L (>1 μ g/L with greater specificity)
Chest X-ray	Bilateral, interstitial infiltrates	Lobar alveolar infiltrates
Response to antibiotic treatment	Slow response or no response	Rapid

Adapted from Ruuskanen et al.³

Table 2
Viruses Related with Community-Acquired viral Pneumonia in Children and Adults.

Syncytial respiratory virus
Rhinovirus
Influenza A, B and C virus
Human metapneumovirus
Parainfluenza virus type 1, 2, 3 and 4
Human bocavirus
Coronavirus type 229E, OC43, NL63, HKU1, SARS and MERS-CoV
Adenovirus
Enterovirus
Varicella zoster virus, Epstein-Barr virus, human herpesvirus 6 and 7, cytomegalovirus
Hantavirus
Parechovirus
Mimivirus
Measles virus

Adapted from Ruuskanen et al.³

(15%), the etiological role of rhinovirus continues to be questioned, although it could be an indication of real, subclinical infections.²³ Recently identified pathogens, such as metapneumovirus and human bocavirus,^{24–27} are less commonly encountered. Although the prevalence of viral pneumonia caused by adenovirus is low (2%–12%), it must be identified, as it can lead to necrotizing pneumonia.²⁸ In this case, PCR techniques are much more sensitive than antigen detection techniques.²⁹

In the case of adults, the most commonly detected viruses are influenza virus (IV), SRV and parainfluenza,^{5,10,30,31} although incidence varies depending on the diagnostic techniques used. These viruses are also the most important in Spain.³² Pneumonias caused by other viruses are more rarely reported and include outbreaks of rhinovirus,^{33,34} adenovirus³⁵ (particularly serotype 14 in military institutions³⁶), coronavirus,³⁷ metapneumovirus,³⁸ and even bocavirus (in immunocompromised patients).³⁹

Co-infection

Infections involving both respiratory bacteria and viruses or 2 different viruses are common. The most widely accepted hypothesis is that the viral infection occurs first, followed by the bacterial form. Viral-mediated activation of proinflammatory molecules, such as interleukin-10, is thought to attract large numbers of neutrophils and macrophages to the lung. The arrival of these cytokines amplifies the immune response, causing inflammatory damage and preventing the proper clearance of bacteria.⁴⁰ Bacterial superinfection worsens the prognosis of the original viral infection. Indeed, research into the influenza pandemics of 1918, 1957, and 1968 shows that most deaths were caused by a secondary bacterial infection.⁴¹ During the 2009 H1N1 pandemic, 4%–24% of cases presented secondary bacterial infection.^{42–44} In infections caused by other viruses, however, especially H5N1 avian influenza, the associated pneumonia appears to be caused (more frequently) by direct viral action.⁴⁵

In clinical practice, this type of co-infection is particularly common in children (up to 45% of cases with CAP), and mainly involves pneumococcus,^{46–49} thus increasing clinical severity.⁵⁰ *Mycoplasma pneumoniae* and several species of *Chlamydia*⁵¹ are also common,⁵¹ while simultaneous co-infection with 2–3 viruses is not unusual.⁵²

CAP of mixed etiology has been characterized less in adults than in children, and prevalence is estimated at less than 5%.^{7,13} The most common combinations reported are rhinovirus+pneumococcus and influenza A virus+pneumococcus. More serious infections have been identified with the combination of viruses with *Legionella pneumophila*.⁵³

Data on morbidity and mortality in bacterial/viral co-infection are contradictory. Hong et al.⁵⁴ consider that these co-infections are no more severe than purely bacterial infections, and affect older patients and those with chronic lung diseases. In contrast, Johansson et al.⁵⁵ and Seki et al.⁵⁶ found that mixed etiology pneumonia was associated with higher severity scale scores and poorer progress.

Respiratory Virus Outbreaks: Experience with H1N1 Influenza Virus in 2009, Avian Influenza (H5N1 and H7N9), SARS and Middle East Coronavirus (MERS-CoV)

Some very well-known families of respiratory viruses have produced new species and some very virulent serotypes, which in recent years have caused epidemics with significant associated morbidity and mortality.⁵⁷

The great H1N1 IV-A epidemic of 2009 was unusual: transmission was enhanced by the lack of prior immunity, causing a serious medical situation worldwide.⁴² Studies performed subsequently showed that the younger population and certain risk groups were more affected, resulting in the loss of many more life-years.⁵⁸ However, from an inter-generational perspective, the burden of death and major complications was no higher than that caused by normal seasonal influenza.^{59,60}

In 2003, an outbreak of coronavirus in the Far East caused a severe acute respiratory syndrome. In the Middle East in 2012, another outbreak of a new coronavirus occurred, called Middle East respiratory syndrome coronavirus (MERS-CoV).⁶¹ In both epidemics, the incidence was reduced to a few hundred cases, partly due to the protective measures implemented,⁶² but mortality was high due to high virulence.⁶³ Clinical symptoms were not only limited to rapidly developing bilateral pneumonia, but also included acute renal failure and severe hematological disturbances.^{63,64}

Recombinations in animals of different IV subtypes are a global concern, as they are potentially capable of passing the species barrier, and occasionally propagating among humans.⁶⁵ The serious cases of H5N1 avian influenza recorded in 2005 in Southeast Asia,⁶⁶ or more recently the H3N2-porcine variant in the United States (2012),⁶⁷ and H7N9 avian influenza in China (2013),⁶⁸ are examples of this new threat. Fortunately, surveillance systems for monitoring animal reserves and activating a rapid medical response are improving.⁶⁹

Microbiological Diagnosis

The detection of virus or viral antigens in upper (nasopharyngeal aspirates) and lower (bronchoalveolar lavage and induced sputum) respiratory tract samples is usually based on culture and immunofluorescence microscopy. The detection of antibodies generated during the process of viral infection has also been used for years: seroconversion in 2 samples obtained over a period of time is suggestive of a new exposure to the pathogen. The introduction of polymerase chain reaction (PCR) techniques has improved the detection of respiratory viruses, particularly with the use of real-time techniques.³

The preferred specimen for upper respiratory samples, particularly in children, is nasopharyngeal aspirate, which combines a mixture of nasal and retropharyngeal secretions.⁷⁰ The use of sterile cotton wool swabs yields a similar sensitivity for most viruses, with the exception of SRV.^{71,72} Devices that combine a nylon fiber swab and a container of universal transport medium provide a high diagnostic yield at all ages, comparable to nasopharyngeal lavage.^{30,73}

Samples from the lower respiratory tract, being obtained at the site of the infection, have obvious advantages when determining the cause of pneumonia. However, contamination may occur during

Table 3
Characteristics of the Main Viruses Involved in Viral Pneumonia.

Virus	Family	Subtype	Incidence of CAP		Risk factors		Seasonality	Differential clinical factors	Treatment
			Children	Adults	Infection	Poor evolution			
Rhinovirus	<i>Picornaviridae</i>	–	≈18%	≈6%	All ages, but more in children	Asthma. Cellular immunosuppression	All year (more in autumn)	Upper airway symptoms: rhinorrhea, cough and nasal congestion	Pleconaril (compassionate use)
Syncytial respiratory virus (SRV)	<i>Paramyxoviridae</i>	1 and 2	≈11%	≈3%	Newborn and premature babies. Immunosuppression	COPD. Asthma. Stem cell transplant. Immunosuppression	End of autumn, beginning of Winter	Marked bronchial reactivity	Inhaled ribavirin (children), IV ribavirin (immunosuppression)
Influenza virus (IV)	<i>Orthomyxoviridae</i>	A and B seasonal	≈10%	≈8%	Children and geriatrics	>65 years. Comorbidities. Gestation. BMI ↑	End of autumn and winter	General asthenia. Influenza-like syndrome	NAI (OSE±resistant) Amantadines (not in B)
		H1N1 09 pandemic	–	–	<65 years	Gestation. Homeless. Obesity	Specific outbreaks in waves	More pneumonias, ICU and mortality	NAI (ZAN and PER in critical patients)
		H5N1	–	–	Contact with birds	Neutropenia and delayed diagnosis	Outbreaks throughout the year	Thrombocytopenia and kidney failure	High does NAI. Amantadines not beneficial
Parainfluenza virus (PIV)	<i>Paramyxoviridae</i>	1, 2, 3 and 4	≈8%	≈2%	Geriatric care homes	Lung and stem cell transplant. Fragile elderly	Autumn (PIV1-2) Spring (PIV-3)	Laryngeal croup (children with PIV-1)	Ribavirin iv (immunosuppression)
Metapneumovirus	<i>Paramyxoviridae</i>	–	≈8%	≈1%	Children <5 years	SRV coinfection. Immunosuppression	End of Winter and Spring	Wheezing. Asthma exacerbations	Ribavirin iv (immunosuppression)
Coronavirus	<i>Coronaviridae</i>	229E, NL63 OC43, KU1	≈7%	≈5%	Geriatric care homes	Asthma. Immunosuppression	Winter	Diarrhea (OC43, and intermittent)	No proven treatment. Chloroquine
		SARS	–	–	Bats and civets in Asia. Healthcare personnel	Elderly. DM. Hepatitis B. (Pediatric population protective factor)	Outbreaks throughout the year	Prodrome with fever and myalgia followed by a respiratory distress	No specific treatment. Corticosteroids used
Adenovirus	<i>Adenoviridae</i>	7, 14, 16	≈3%	≈2%	Prisons (outbreaks)	Pneumococcus	All year	Conjunctivitis, diarrhea, encephalitis	Cidofovir (proven in immunosuppression)
Bocavirus	<i>Parvoviridae</i>	–	≈5%	<1%	Children <2 years	Poorly defined	End of autumn, beginning of winter	Otitis media and pneumonia (few studies)	No specific treatment

BMI: body mass index; COPD: chronic obstructive pulmonary disease; IV: intravenous; NAI: neuraminidase inhibitors (OSE: oseltamivir; PER: peramivir; ZAN: zanamivir).

sampling, as the specimen passes through the upper airways. Induced sputum is often obtained from children, but bronchoalveolar lavage is preferred in adults.³ In contrast, the invasive character of other forms of diagnosis, such as transthoracic puncture, has led to their disuse.⁷⁰

In sample processing, PCR-based systems are 2–5 times more sensitive than conventional methods for the detection of respiratory viruses.³⁰ This technique is particularly useful in the elderly, who often have a lower nasopharyngeal viral burden than children.^{74,75} Moreover, some viruses (such as bocavirus) are hard to culture, and antigen testing may be equivocal (e.g. parainfluenza or adenovirus), so they can only be satisfactorily identified using PCR.^{76–78} A standard identification test has been developed that uses a PCR platform combined with microarrays to simultaneously determine more than 15 different respiratory viruses.⁷⁹ These platforms are the techniques most commonly used in our hospitals at the moment.⁸⁰

Etiological Treatment

Since the end of the last century, empirical antibiotic treatment of CAP has been initiated as soon as possible, as this has been shown to reduce morbidity and mortality.⁸¹

However, no prospective studies have examined the benefits of antibiotics in pneumonias with a high suspicion of viral origin. The possibility of bacterial superinfection or a mixed viral-bacterial etiology, progressive improvements in antibiotic tolerability, and the medical-legal requirement to “use of all available measures”, mean that physicians are unlikely to resist prescribing antibiotics in a patient presenting with pneumonia. This dilemma has only been prospectively studied in children, in the context of SRV bronchiolitis epidemics; the results, while not definitive, do not support the generalized administration of antibiotics.⁸²

Experience in the use of antiviral drugs in respiratory viruses can be divided into 3 areas: (a) treatment for influenza virus, which is relatively effective and well documented, thanks mainly to studies performed during the 2009 pandemic⁸³; (b) treatment for SRV, that has been mostly explored in children and immunosuppressed populations, although results remain unclear⁸⁴; and (c) treatment of other respiratory viruses, which has not been studied in depth and remains largely unevaluated.⁸⁵

Treatment of Influenza Virus

Neuraminidase, an enzyme of the IV capsid, is essential for intercellular viral propagation. Selective inhibitors of this enzyme (oseltamivir, zanamivir and peramivir) have been seen to contain infection during the peak of viral replication – the first 24–72 hours – improving clinical symptoms and reducing morbidity and mortality.^{86,87} Retrospective studies of patients with a diagnosis of influenza have shown reductions in the incidence of pneumonia in patients who received early treatment.⁸⁸ Although these drugs are active against both subtypes, seasonal epidemics of oseltamivir-resistant H1N1 IV-A have been described, so some guidelines recommend zanamivir as a first-line approach.⁸³ The pandemic H1N1 strain of 2009 did not generally show this resistance profile, and was widely treated with oseltamivir, with no detriment to outcomes.⁸⁹ In children, the clinical effect is less clear, but administration appears to be safe.⁹⁰ Laninamivir octanoate, a new neuraminidase inhibitor for inhaled administration only, has been seen to be effective in the treatment of IV infection, including oseltamivir-resistant strains.⁹¹

Adamantanes (amantadine and rimantadine) are drugs that are conventionally used in IV infection. They are exclusively specific to influenza virus type A. This, along with their potential for side

effects and rapid development of resistances, has meant that they have fallen into disuse in clinical practice.⁹² Drugs that can block RNA production and others that limit viral integration are under investigation, but no clinical trials have been started.⁹³ Studies are also being conducted with immunomodulators, with the aim of reducing viral-mediated inflammation and alleviating its effect on the host.⁹⁴

Treatment of Syncytial Respiratory Virus

None of the treatments tested has been significantly useful in the management of acute episodes of pneumonia or for improving respiratory parameters in follow-up. According to the largest meta-analysis performed in children to date, inhaled ribavirin during pneumonia can reduce hospital stay and time on mechanical ventilation, but does not significantly improve overall mortality.⁹⁵ Intravenous or oral administration of ribavirin has been used almost exclusively in severely immunocompromised patients (bone marrow or lung transplant recipients). Outcomes have been positive, but the results were not applicable to other patients.⁹⁶ The use of immunoglobulins in children has not shown any benefits over supportive treatment alone.⁹⁷ Generalized use of bronchodilators, corticosteroids or antibiotics is not recommended in the American pediatric guidelines for SRV bronchiolitis.⁹⁸ Trials are currently underway on new products such as catelicidin LL-37 (a vitamin D derivative)⁹⁹ and dingchuan decoction, a Chinese medicinal product,¹⁰⁰ and results are promising.

Treatment of Other Respiratory Viruses

Until recently, only supportive treatment has been available. However, some antiviral medications are currently under investigation. The cytidine analog, cidofovir, originally developed as treatment for CMV, has shown success *in vitro* against adenoviruses, although the response in immunocompromised patients with several forms of severe pneumonia was poor.¹⁰¹ Intravenous ribavirin has been used with success in lung transplant recipients with respiratory infections caused by metapneumovirus.¹⁰² Pleconaril, which incorporates itself into the capsid of *Rhinovirus* and *Enterovirus*, has been successfully used in limited case series; it is not yet on the market and is limited to compassionate use.¹⁰³ There is little doubt that intravenous acyclovir is beneficial in the rare cases of varicella zoster pneumonia in immunocompromised patients.^{85,104} Lastly, treatment with high-dose corticosteroids can improve the clinical course of viral CAP,⁸⁵ although this application is still controversial.

Prevention Measures

In infectious and contagious diseases, particularly of the respiratory tract, barrier methods are essential for preventing infection. Use of masks and gloves and hand-washing have been shown to be effective in reducing transmission rates in the healthcare setting.¹⁰⁵ Social isolation of patients during the clinical phase of the disease is also strongly recommended and reduces overall incidence.¹⁰⁶ However, it is difficult to implement these measures properly.¹⁰⁶

Immunization plays a very important role in prevention, but is only available for a few viruses. Anti-influenza A and B vaccines have been shown to reduce transmission during seasonal influenza epidemics in the general population,¹⁰⁷ but their effect on the course of pneumonia or on mortality is not so clear¹⁰⁸; nor is their efficiency in children younger than 2 years, although they continue to be administered in many countries.¹⁰⁹ In contrast, they appear to be very effective in elderly institutionalized subjects.¹¹⁰ They are currently recommended in Spain for patients with respiratory comorbidities or immunosuppression, individuals over 65 years of age, and healthcare workers.¹¹¹

In addition to vaccines, chemoprophylaxis with neuroaminidase inhibitors has been successfully tested during seasonal influenza epidemics.¹¹² As yet, no effective vaccine is available for SRV, but palivizumab has been used as chemoprophylaxis. This is a humanized monoclonal antibody that has shown a reduction of up to 50% in the incidence of pneumonia and associated hospital admissions in neonates with a high risk of infection.¹¹³

Conflict of Interests

The authors declare that they have no conflict of interests.

References

- WHO. Disease and injury regional mortality estimates, 2000–2012 – global summary estimates; 2012. Available at: <http://www.who.int/entity/healthinfo/global.burden.disease/GHE.DthGlobal.2000.2012.xls> [accessed 02.09.14].
- Rudan I, Boschi-Pinto C, Biloglav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bull World Health Organ.* 2008;86:408–16.
- Ruuskanen O, Lahti E, Jennings LC, Murdoch DR. Viral pneumonia. *Lancet.* 2011;377:1264–75.
- McIntosh K. Community-acquired pneumonia in children. *N Engl J Med.* 2002;346:429–37.
- Rohde GGU. The role of viruses in CAP. Monograph 63: community-acquired pneumonia. *Eur Respir Monogr.* 2014;63:74–87.
- Ruiz-González A, Falguera M, Vives M, Nogués A, Porcel JM, Rubio-Caballero M. Community-acquired pneumonia: development of a bedside predictive model and scoring system to identify the aetiology. *Respir Med.* 2000;94:505–10.
- Johnstone J, Majumdar SR, Fox JD, Marrie TJ. Viral infection in adults hospitalized with community-acquired pneumonia: prevalence, pathogens, and presentation. *Chest.* 2008;134:1141–8.
- Ma HM, Lee KP, Woo J. Predictors of viral pneumonia: the need for viral testing in all patients hospitalized for nursing home-acquired pneumonia. *Geriatr Gerontol Int.* 2013;13:949–57.
- Liu YF, Gao Y, Chen MF, Cao B, Yang XH, Wei L. Etiological analysis and predictive diagnostic model building of community-acquired pneumonia in adult outpatients in Beijing, China. *BMC Infect Dis.* 2013;13:309.
- Jennings LC, Anderson TP, Beynon KA, Chua A, Laing RT, Werno AM, et al. Incidence and characteristics of viral community-acquired pneumonia in adults. *Thorax.* 2008;63:42–8.
- Viasus D, Marinescu C, Villoslada A, Cordero E, Gálvez-Acebal J, Fariñas MC, et al. Community-acquired pneumonia during the first post-pandemic influenza season: a prospective, multicentre cohort study. *J Infect.* 2013;67:185–93.
- Tarver RD, Teague SD, Heitkamp DE, Conces DJ. Radiology of community-acquired pneumonia. *Radiol Clin North Am.* 2005;43:497–512, viii.
- Shiley KT, van Deerlin VM, Miller WT. Chest CT features of community-acquired respiratory viral infections in adult inpatients with lower respiratory tract infections. *J Thorac Imaging.* 2010;25:68–75.
- Gilbert DN. Use of plasma procalcitonin levels as an adjunct to clinical microbiology. *J Clin Microbiol.* 2010;48:2325–9.
- Johansson N, Kalin M, Backman-Johansson C, Larsson A, Nilsson K, Hedlund J. Procalcitonin levels in community-acquired pneumonia – correlation with aetiology and severity. *Scand J Infect Dis.* 2014;1–5.
- Musher DM, Roig IL, Cazares G, Stager CE, Logan N, Safar H. Can an etiologic agent be identified in adults who are hospitalized for community-acquired pneumonia: results of a one-year study. *J Infect.* 2013;67:11–8.
- Berkley JA, Munywoki P, Ngama M, Kazungu S, Abwao J, Bett A, et al. Viral etiology of severe pneumonia among Kenyan infants and children. *JAMA.* 2010;303:2051–7.
- Nair H, Nokes DJ, Gessner BD, Dherani M, Madhi SA, Singleton RJ, et al. Global burden of acute lower respiratory infections due to respiratory syncytial virus in young children: a systematic review and meta-analysis. *Lancet.* 2010;375:1545–55.
- Hayden FG. Rhinovirus and the lower respiratory tract. *Rev Med Virol.* 2004;14:17–31.
- Louie JK, Roy-Burman A, Guardia-Labar L, Boston EJ, Kiang D, Padilla T, et al. Rhinovirus associated with severe lower respiratory tract infections in children. *Pediatr Infect Dis J.* 2009;28:337–9.
- Calvo C, García-García ML, Blanco C, Pozo F, Flecha IC, Pérez-Breña P. Role of rhinovirus in hospitalized infants with respiratory tract infections in Spain. *Pediatr Infect Dis J.* 2007;26:904–8.
- Miller EK, Lu X, Erdman DD, Poehling KA, Zhu Y, Griffin MR, et al., for the New Vaccine Surveillance Network. Rhinovirus-associated hospitalizations in young children. *J Infect Dis.* 2007;195:773–81.
- Jartti T, Jartti L, Peltola V, Waris M, Ruuskanen O. Identification of respiratory viruses in asymptomatic subjects: asymptomatic respiratory viral infections. *Pediatr Infect Dis J.* 2008;27:1103–7.
- Williams JV, Harris PA, Tollefson SJ, Halburnt-Rush LL, Pingsterhaus JM, Edwards KM, et al. Human metapneumovirus and lower respiratory tract disease in otherwise healthy infants and children. *N Engl J Med.* 2004;350:443–50.
- Fry AM, Lu X, Chittaganpitch M, Peret T, Fischer J, Dowell SF, et al. Human bocavirus: a novel parvovirus epidemiologically associated with pneumonia requiring hospitalization in Thailand. *J Infect Dis.* 2007;195:1038–45.
- Milder E, Arnold JC. Human metapneumovirus and human bocavirus in children. *Pediatr Res.* 2009;65:78R–83R.
- Heugel J, Martin ET, Kuypers J, Englund JA. Coronavirus-associated pneumonia in previously healthy children. *Pediatr Infect Dis J.* 2007;26:753–5.
- Ou ZY, Zeng QY, Wang FH, Xia HM, Lu JP, Xia JQ, et al. Retrospective study of adenovirus in autopsied pulmonary tissue of pediatric fatal pneumonia in South China. *BMC Infect Dis.* 2008;8:122.
- Freythum F, Vabret A, Galateau-Salle F, Ferey J, Eugene G, Petitjean J, et al. Detection of respiratory syncytial virus, parainfluenzavirus 3, adenovirus and rhinovirus sequences in respiratory tract of infants by polymerase chain reaction and hybridization. *Clin Diagn Virol.* 1997;8:31–40.
- Johansson N, Kalin M, Tiveljung-Lindell A, Giske CG, Hedlund J. Etiology of community-acquired pneumonia: increased microbiological yield with new diagnostic methods. *Clin Infect Dis.* 2010;50:202–9.
- Lieberman D, Shimoni A, Shemer-Avni Y, Keren-Naos A, Shtainberg R, Lieberman D. Respiratory viruses in adults with community-acquired pneumonia. *Chest.* 2010;138:811–6.
- Angeles Marcos M, Camps M, Pumarola T, Antonio Martinez J, Martinez E, Mensa J, et al. The role of viruses in the aetiology of community-acquired pneumonia in adults. *Antivir Ther.* 2006;11:351–9.
- Hicks LA, Shepard CW, Britz PH, Erdman DD, Fischer M, Flannery BL, et al. Two outbreaks of severe respiratory disease in nursing homes associated with rhinovirus. *J Am Geriatr Soc.* 2006;54:284–9.
- Louie JK, Yagi S, Nelson FA, Kiang D, Glaser CA, Rosenberg J, et al. Rhinovirus outbreak in a long term care facility for elderly persons associated with unusually high mortality. *Clin Infect Dis.* 2005;41:262–5.
- Esposito DH, Gardner TJ, Schneider E, Stockman LJ, Tate JE, Panozzo CA, et al. Outbreak of pneumonia associated with emergent human adenovirus serotype 14–Southeast Alaska, 2008. *J Infect Dis.* 2010;202:214–22.
- Tate JE, Bunning ML, Lott L, Lu X, Su J, Metzgar D, et al. Outbreak of severe respiratory disease associated with emergent human adenovirus serotype 14 at a US air force training facility in 2007. *J Infect Dis.* 2009;199:1419–26.
- Reina J, López-Causapé C, Rojo-Molinero E, Rubio R. Clinico-epidemiological characteristics of acute respiratory infections caused by coronavirus OC43, NL63 and 229E. *Rev Clin Esp.* 2014;214:499–504.
- Walsh EE, Peterson DR, Falsely AR. Human metapneumovirus infections in adults: another piece of the puzzle. *Ann Intern Med.* 2008;168:2489–96.
- Kupfer B, Vehreschild J, Cornely O, Kaiser R, Plum G, Viazov S, et al. Severe pneumonia and human bocavirus in adult. *Emerg Infect Dis.* 2006;12:1614–6.
- McCullers JA. Insights into the interaction between influenza virus and pneumococcus. *Clin Microbiol Rev.* 2006;19:571–82.
- Morens DM, Taubenberger JK, Fauci AS. Predominant role of bacterial pneumonia as a cause of death in pandemic influenza: implications for pandemic influenza preparedness. *J Infect Dis.* 2008;198:962–70.
- Bautista E, Chotpitayasunondh T, Gao Z, Harper SA, Shaw M, Uyeki TM, et al., Writing Committee of the WHO Consultation on Clinical Aspects of Pandemic (H1N1) 2009 Influenza. Clinical aspects of pandemic 2009 influenza A (H1N1) virus infection. *N Engl J Med.* 2010;362:1708–19.
- Louie JK, Acosta M, Winter K, Jean C, Gavali S, Schechter R, et al., for the California Pandemic (H1N1) Working Group. Factors associated with death or hospitalization due to pandemic 2009 influenza A(H1N1) infection in California. *JAMA.* 2009;302:1896–902.
- Kumar A, Zarychanski R, Pinto R, Cook DJ, Marshall J, Lacroix J, et al. Critically ill patients with 2009 influenza A(H1N1) infection in Canada. *JAMA.* 2009;302:1872–9.
- Beigel JH, Farrar J, Han AM, Hayden FG, Hyer R, de Jong MD, et al., Writing Committee of the WHO Consultation on human influenza A/H5N1. Avian influenza A (H5N1) infection in humans. *N Engl J Med.* 2005;353:1374–85.
- Lehtinen P, Jartti T, Virkki R, Vuorinen T, Leinonen M, Peltola V, et al. Bacterial coinfections in children with viral wheezing. *Eur J Clin Microbiol Infect Dis.* 2006;25:463–9.
- Xiang Z, Gonzalez R, Xie Z, Xiao Y, Liu J, Chen L, et al. Human rhinovirus C infections mirror those of human rhinovirus A in children with community-acquired pneumonia. *J Clin Virol.* 2010;49:94–9.
- Korppi M. Mixed microbial aetiology of community-acquired pneumonia in children. *APMIS.* 2002;110:515–22.
- Levine OS, O'Brien KL, Deloria-Knoll M, Murdoch DR, Feikin DR, deLuca AN, et al. The Pneumonia etiology research for child health project: a 21st century childhood pneumonia etiology study. *Clin Infect Dis.* 2012;54 Suppl. 2: S93–101.
- Harada Y, Kinoshita F, Yoshida LM, Minh le N, Suzuki M, Morimoto K, et al. Does respiratory virus coinfection increase the clinical severity of acute respiratory infection among children infected with respiratory syncytial virus. *Pediatr Infect Dis J.* 2013;32:441–5.
- Chen CJ, Lin PY, Tsai MH, Huang CG, Tsao KC, Wong KS, et al. Etiology of community-acquired pneumonia in hospitalized children in northern Taiwan. *Pediatr Infect Dis J.* 2012;31:e196–201.
- Honkinen M, Lahti E, Österback R, Ruuskanen O, Waris M. Viruses and bacteria in sputum samples of children with community-acquired pneumonia. *Clin Microbiol Infect.* 2012;18:300–7.
- Diederer BM, van der Eerden MM, Vlasopolder F, Boersma WG, Kluytmans JA, Peeters MF. Detection of respiratory viruses and *Legionella* spp. by real-time

- polymerase chain reaction in patients with community acquired pneumonia. *Scand J Infect Dis.* 2009;41:45–50.
54. Hong HL, Hong SB, Ko GB, Huh JW, Sung H, Do KH, et al. Viral infection is not uncommon in adult patients with severe hospital-acquired pneumonia. *PLOS ONE.* 2014;9:e95865.
 55. Johansson N, Kalin M, Hedlund J. Clinical impact of combined viral and bacterial infection in patients with community-acquired pneumonia. *Scand J Infect Dis.* 2011;43:609–15.
 56. Seki M, Kosai K, Yanagihara K, Higashiyama Y, Kurihara S, Izumikawa K, et al. Disease severity in patients with simultaneous influenza and bacterial pneumonia. *Intern Med.* 2007;46:953–8.
 57. Gautret P, Gray GC, Charrel RN, Odezulu NG, Al-Tawfiq JA, Zumla A, et al. Emerging viral respiratory tract infections–environmental risk factors and transmission. *Lancet Infect Dis.* 2014;14:1113–22.
 58. Simonsen L, Spreuwenberg P, Lustig R, Taylor RJ, Fleming DM, Kroneman M, et al. Global mortality estimates for the 2009 Influenza Pandemic from the GLaMOR project: a modeling study. *PLoS Med.* 2013;10:e1001558.
 59. Lemaître M, Carrat F, Rey G, Miller M, Simonsen L, Viboud C. Mortality burden of the 2009 A/H1N1 influenza pandemic in France: comparison to seasonal influenza and the A/H3N2 pandemic. *PLoS ONE.* 2012;7:e45051.
 60. Hayward AC, Fragaszy EB, Bermingham A, Wang L, Copas A, Edmunds WJ, et al. Comparative community burden and severity of seasonal and pandemic influenza: results of the Flu Watch cohort study. *Lancet Respir Med.* 2014;2:445–54.
 61. Coleman CM, Frieman MB. Coronaviruses: important emerging human pathogens. *J Virol.* 2014;88:5209–12.
 62. Graham RL, Donaldson EF, Baric RS. A decade after SARS: strategies for controlling emerging coronaviruses. *Nat Rev Microbiol.* 2013;11:836–48.
 63. Arabi YM, Arifi AA, Balkhy HH, Najm H, Aldawood AS, Ghabashi A, et al. Clinical course and outcomes of critically ill patients with Middle East respiratory syndrome coronavirus infection. *Ann Intern Med.* 2014;160:389–97.
 64. Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA. Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med.* 2012;367:1814–20.
 65. Medina RA, García-Sastre A. Influenza A viruses: new research developments. *Nat Rev Microbiol.* 2011;9:590–603.
 66. Beigel JH, Farrar J, Han AM, Hayden FG, Hyer R, de Jong MD, et al. Avian influenza A (H5N1) infection in humans. *N Engl J Med.* 2005;353:1374–85.
 67. Bowman AS, Nelson SW, Page SL, Nolting JM, Killian ML, Sreevatsan S, et al. Swine-to-human transmission of Influenza A(H3N2) virus at agricultural fairs, Ohio, USA, 2012. *Emerg Infect Dis.* 2014;20:1472–80.
 68. Gao R, Cao B, Hu Y, Feng Z, Wang D, Hu W, et al. Human infection with a novel avian-origin influenza A (H7N9) virus. *N Engl J Med.* 2013;368:1888–97.
 69. Al-Tawfiq JA, Zumla A, Gautret P, Gray GC, Hui DS, Al-Rabeeah AA, et al. Surveillance for emerging respiratory viruses. *Lancet Infect Dis.* 2014;14:992–1000.
 70. Loens K, van Heirstraeten L, Malhotra-Kumar S, Goossens H, Ieven M. Optimal sampling sites and methods for detection of pathogens possibly causing community-acquired lower respiratory tract infections. *J Clin Microbiol.* 2009;47:21–31.
 71. Heikkinen T, Marttila J, Salmi AA, Ruuskanen O. Nasal swab versus nasopharyngeal aspirate for isolation of respiratory viruses. *J Clin Microbiol.* 2002;40:4337–9.
 72. Ruohola A, Waris M, Allander T, Ziegler T, Heikkinen T, Ruuskanen O. Viral etiology of common cold in children, Finland. *Emerg Infect Dis.* 2009;15:344–6.
 73. Lieberman D, Lieberman D, Shimoni A, Keren-Naus A, Steinberg R, Shemer-Avni Y. Identification of respiratory viruses in adults: nasopharyngeal versus oropharyngeal sampling. *J Clin Microbiol.* 2009;47:3439–43.
 74. Talbot HK, Faisey AR. The diagnosis of viral respiratory disease in older adults. *Clin Infect Dis.* 2010;50:747–51.
 75. She RC, Polage CR, Caram LB, Taggart EW, Hymas WC, Woods CW, et al. Performance of diagnostic tests to detect respiratory viruses in older adults. *Diagn Microbiol Infect Dis.* 2010;67:246–50.
 76. Arens MQ, Buller RS, Rankin A, Mason S, Whetsell A, Agapov E, et al. Comparison of the Eragen Multi-Code Respiratory Virus Panel with conventional viral testing and real-time multiplex PCR assays for detection of respiratory viruses. *J Clin Microbiol.* 2010;48:2387–95.
 77. Kuypers J, Wright N, Ferrenberg J, Huang ML, Cent A, Corey L, et al. Comparison of real-time PCR assays with fluorescent-antibody assays for diagnosis of respiratory virus infections in children. *J Clin Microbiol.* 2006;44:2382–8.
 78. Arnold JC, Singh KK, Spector SA, Sawyer MH. Undiagnosed respiratory viruses in children. *Pediatrics.* 2008;121:e631–7.
 79. Tiveljung-Lindell A, Rotzén-Ostlund M, Gupta S, Ullstrand R, Grillner L, Zweyberg-Wirgart B, et al. Development and implementation of a molecular diagnostic platform for daily rapid detection of 15 respiratory viruses. *J Med Virol.* 2009;81:167–75.
 80. Culebras E, Betriu C, Vázquez-Cid E, López-Varela E, Rueda S, Picazo JJ. Detection and genotyping of human respiratory viruses in clinical specimens from children with acute respiratory tract infections. *Rev Esp Quimioter.* 2013;26:47–50.
 81. Bartlett JG, Breiman RF, Mandell LA, File TM. Community-acquired pneumonia in adults: guidelines for management. *The Infectious Diseases Society of America. Clin Infect Dis.* 1998;26:811–38.
 82. Friis B, Andersen P, Brenøe E, Hornsleth A, Jensen A, Knudsen FU, et al. Antibiotic treatment of pneumonia and bronchiolitis. A prospective randomised study. *Arch Dis Child.* 1984;59:1038–45.
 83. Harper SA, Bradley JS, Englund JA, File TM, Gravenstein S, Hayden FG, et al. Seasonal influenza in adults and children – diagnosis, treatment, chemoprophylaxis, and institutional outbreak management: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis.* 2009;48:1003–32.
 84. Turner TL, Kopp BT, Paul G, Landgrave LC, Hayes D, Thompson R. Respiratory syncytial virus: current and emerging treatment options. *Clinicoecon Outcomes Res.* 2014;6:217–25.
 85. Cheng VC, Tang BS, Wu AK, Chu CM, Yuen KY. Medical treatment of viral pneumonia including SARS in immunocompetent adult. *J Infect.* 2004;49:262–73.
 86. Moscona A. Neuraminidase inhibitors for influenza. *N Engl J Med.* 2005;353:1363–73.
 87. Kaiser L, Wat C, Mills T, Mahoney P, Ward P, Hayden F. Impact of oseltamivir treatment on influenza-related lower respiratory tract complications and hospitalizations. *Ann Intern Med.* 2003;139:1667–72.
 88. Yu H, Liao Q, Yuan Y, Zhou L, Xiang N, Huai Y, et al. Effectiveness of oseltamivir on disease progression and viral RNA shedding in patients with mild pandemic 2009 influenza A H1N1: opportunistic retrospective study of medical charts in China. *BMJ.* 2010;341:c4779.
 89. Shobugawa Y, Saito R, Sato I, Kawashima T, Daput C, Daput IC, et al. Clinical effectiveness of neuraminidase inhibitors – oseltamivir, zanamivir, laninamivir, and peramivir – for treatment of influenza A(H3N2) and A(H1N1)pdm09 infection: an observational study in the 2010–2011 influenza season in Japan. *J Infect Chemother.* 2012;18:858–64.
 90. Wang K, Shun-Shin M, Gill P, Perera R, Harnden A. Neuraminidase inhibitors for preventing and treating influenza in children (published trials only). *Cochrane Database Syst Rev.* 2012;4:Cd002744.
 91. Watanabe A, Chang SC, Kim MJ, Chu DW, Ohashi Y, MARVEL Study Group. Long-acting neuraminidase inhibitor laninamivir octanoate versus oseltamivir for treatment of influenza: a double-blind, randomized, noninferiority clinical trial. *Clin Infect Dis.* 2010;51:1167–75.
 92. Jefferson T, Demicheli V, di Pietrantonj C, Rivetti D. Amantadine and rimantadine for influenza A in adults. *Cochrane Database Syst Rev.* 2006;2:Cd001169.
 93. Król E, Rychłowska M, Szewczyk B. Antivirals – current trends in fighting influenza. *Acta Biochim Pol.* 2014;61:495–504.
 94. Simmons C, Farrar J. Insights into inflammation and influenza. *N Engl J Med.* 2008;359:1621–3.
 95. Ventre K, Randolph A. WITHDRAWN. Ribavirin for respiratory syncytial virus infection of the lower respiratory tract in infants and young children. *Cochrane Database Syst Rev.* 2010;5:Cd000181.
 96. Marcelin JR, Wilson JW, Razonable RR. Mayo Clinic Hematology/Oncology and Transplant Infectious Diseases Services. Oral ribavirin therapy for respiratory syncytial virus infections in moderately to severely immunocompromised patients. *Transpl Infect Dis.* 2014;16:242–50.
 97. Fuller HL, del Mar CB. WITHDRAWN. Immunoglobulin treatment for respiratory syncytial virus infection. *Cochrane Database Syst Rev.* 2010;9:Cd004883.
 98. American Academy of Pediatrics Subcommittee on Diagnosis and Management of Bronchiolitis. Diagnosis and management of bronchiolitis. *Pediatrics.* 2006;118:1774–93.
 99. Currie SM, Findlay EG, McHugh BJ, Mackellar A, Man T, Macmillan D, et al. The human cathelicidin LL-37 has antiviral activity against respiratory syncytial virus. *PLOS ONE.* 2013;8:e73659.
 100. Li L, Yu CH, Ying HZ, Yu JM. Antiviral effects of modified dingchuan decoction against respiratory syncytial virus infection in vitro and in an immunosuppressive mouse model. *J Ethnopharmacol.* 2013;147:238–44.
 101. Louie JK, Kajon AE, Holodny M, Guardia-LaBar L, Lee B, Petru AM, et al. Severe pneumonia due to adenovirus serotype 14: a new respiratory threat. *Clin Infect Dis.* 2008;46:421–5.
 102. Hopkins P, McNeil K, Kermeen F, Musk M, McQueen E, Mackay I, et al. Human metapneumovirus in lung transplant recipients and comparison to respiratory syncytial virus. *Am J Respir Crit Care Med.* 2008;178:876–81.
 103. Rotbart HA, Webster AD, Pleconaril Treatment Registry Group. Treatment of potentially life-threatening enterovirus infections with pleconaril. *Clin Infect Dis.* 2001;32:228–35.
 104. Frangides CY, Pneumatikos I. Varicella-zoster virus pneumonia in adults: report of 14 cases and review of the literature. *Eur J Intern Med.* 2004;15:364–70.
 105. Jefferson T, Foxlee R, del Mar C, Dooley L, Ferroni E, Hewak B, et al. Physical interventions to interrupt or reduce the spread of respiratory viruses: systematic review. *BMJ.* 2008;336:77–80.
 106. Bell D, Nicoll A, Fukuda K, Horby P, Monto A, Hayden F, et al., World Health Organization Writing Group. Non-pharmaceutical interventions for pandemic influenza, national and community measures. *Emerg Infect Dis.* 2006;12:88–94.
 107. Osterholm MT, Kelley NS, Sommer A, Belongia EA. Efficacy and effectiveness of influenza vaccines: a systematic review and meta-analysis. *Lancet Infect Dis.* 2012;12:36–44.
 108. Michiels B, Govaerts F, Remmen R, Vermeire E, Coenen S. A systematic review of the evidence on the effectiveness and risks of inactivated influenza vaccines in different target groups. *Vaccine.* 2011;29:9159–70.
 109. Jefferson T, Rivetti A, di Pietrantonj C, Demicheli V, Ferroni E. Vaccines for preventing influenza in healthy children. *Cochrane Database Syst Rev.* 2012;8:Cd004879.
 110. Jefferson T, Rivetti D, Rivetti A, Rudin M, di Pietrantonj C, Demicheli V. Efficacy and effectiveness of influenza vaccines in elderly people: a systematic review. *Lancet.* 2005;366:1165–74.
 111. Salleras L, Saenz MC, Castrodeza J, de Juanes JR, Arrazola MP. Consenso sobre vacunaciones del adulto de la Sociedad Española de Medicina Preventiva Salud

- Pública e Higiene: Vacunación antigripal. Gripe epidémica. *Vacunas*. 2010;11 Suppl. 2:46–55.
112. Welliver R, Monto AS, Carewicz O, Schatteman E, Hassman M, Hedrick J, et al. Effectiveness of oseltamivir in preventing influenza in household contacts: a randomized controlled trial. *JAMA*. 2001;285:748–54.
 113. Andabaka T, Nickerson JW, Rojas-Reyes MX, Rueda JD, Bacic Vrca V, Barsic B. Monoclonal antibody for reducing the risk of respiratory syncytial virus infection in children. *Cochrane Database Syst Rev*. 2013;4: Cd006602.