

Review: brain neurobiology of gambling disorder based on rodent models

This article was published in the following Dove Press journal:
Neuropsychiatric Disease and Treatment

Gabriel C Quintero Garzola 

Florida State University – Republic of
Panama, Panama City, Republic of Panama

Abstract: Different literature reviews of gambling disorder (GD) neurobiology have been focused on human studies, others have focused on rodents, and others combined human and rodent studies. The main question of this review was: which are the main neurotransmitters systems and brain structures relevant for GD based on recent rodent studies? This work aims to review the experimental findings regarding the rodent's neurobiology of GD. A search in the Pub Med database was set (October 2012–October 2017) and 162 references were obtained. After screening, 121 references were excluded, and only 41 references remained from the initial output. More, other 25 references were added to complement (introduction section, neuroanatomical descriptions) the principal part of the work. At the end, a total of 66 references remained for the review. The main conclusions are: 1) according to studies that used noninvasive methods for drug administration, some of the neurotransmitters and receptors involved in behaviors related to GD are: muscarinic, N-methyl-D-aspartate (NMDA), cannabinoid receptor 1 (CB₁), cannabinoid receptor 2 (CB₂), dopamine 2 receptor (D₂), dopamine 3 receptor (D₃), and dopamine 4 receptor (D₄); 2) moreover, there are other neurotransmitters and receptors involved in GD based on studies that use invasive methods of drug administration (eg, brain microinjection); example of these are: serotonin 1A receptor (5-HT_{1A}), noradrenaline receptors, gamma-aminobutyric acid receptor A (GABA_A), and gamma-aminobutyric acid receptor B (GABA_B); 3) different brain structures are relevant to behaviors linked to GD, like: amygdala (including basolateral amygdala (BLA)), anterior cingulate cortex (ACC), hippocampus, infralimbic area, insular cortex (anterior and rostral agranular), nucleus accumbens (NAc), olfactory tubercle (island of Calleja), orbitofrontal cortex (OFC), medial prefrontal cortex (mPFC), prefrontal cortex (PFC) – subcortical network, striatum (ventral) and the subthalamic nucleus (STN); and 4) the search for GD treatments should consider this diversity of receptor/neurotransmitter systems and brain areas.

Keywords: gambling disorder, review, nervous system, murine paradigm

Introduction

Literature reviews regarding gambling disorder (GD) neurobiology have been specialized on human,^{1,2} rodents,^{3,4} or combination of both.^{5,6} The main question of this review was: which are the main neurotransmitters systems and brain structures relevant for GD based on recent rodent studies? Let me define first gambling and its epidemiological traits, before going in its neurobiological aspects.

Gambling conduct can be described as to put in peril anything significant, and to confide on the assumption of obtaining a gain in return.⁷ GD is characterized by gaming behaviors that seriously disrupt the finances, social relations, and professional advancement of a fellow.⁸ The lifetime prevalence of GD has been estimated

Correspondence: Gabriel C Quintero Garzola
Florida State University – Republic of
Panama, Jacinto Palacios Cobos Street,
#227, City of Knowledge, Clayton,
Panama City, Republic of Panama
Tel +1 507 317 0367
Fax +1 507 317 0366
Email gquintero@fsu.edu

at 0.4% to 4.2%.⁹ Moreover, GD is presently included in the Diagnostic and Statistical Manual of Mental Disorders-5 (DSM)-5, in a novel category, within the division of addictions (behavioral addictions).¹⁰

There have been some literature reviews of the neurobiology of GD focused on human clinical studies.^{1,2,11-14} Specifically, the Lemieux and al'Absi's review proposed that psychological and neurobiological aspects of the stress play a significant role in the starting, prolongation, and relapse of the addictions (including GD). Moreover, the mechanisms include interactions between biological mediators of the stress and the reward system; also, interactions between mediators of the stress and other systems related to addiction (endogenous opioids, the sympathetic-adrenal-medullary system, and endocannabinoids).¹

Another review work by Grant et al posed that GD is linked with alteration across different cognitive domains related to impulsivity and compulsivity;² moreover, it pointed that, based on imaging reports, GD relates to anatomical and functional anomalies of nexus involved in the reward processing and top-down monitoring.² In addition, it pointed that probably, diverse neural systems are involved in the pathophysiology (related to serotonin (5-HT), glutamate, dopamine (DA), opioids, and norepinephrine).²

Then again, the Goulet-Kennedy et al's review points that prefrontal cortex (PFC) and the striatum are the main conductors of decision processes, based on clinical studies.¹¹ Furthermore, that literature review states that the traits of decision making's neural networks can be characterized by means of imaging technology; also, they consider that non-invasive neural stimulation in the PFC, and its network (striatum and others) have elucidated the neurobiological basis of decision-making processes;¹¹ Decision making is involved in different aspects of our daily life,¹¹ including all the spectrums of gambling behavior. Hence, a better understanding of the decision process' neurobiology could be useful for a better quality of life of patients.¹¹

The publications review of the Banz et al group emphasizes in the capacity of neurobiological data to help in the promotion of improved norms and strategies for treatment and prevention.¹² Furthermore, another review by Levy and Glimcher concludes that imaging investigations in humans suggest the existence of a brain network that codifies the values of rewards by means of a standard neural scale.¹³ Based on the authors, the brain

area linked with this standard neural scale is a zone of the ventromedial prefrontal cortex (vmPFC)/orbitofrontal cortex (OFC). The authors propose that a better comprehension of brain mechanism for estimating and deciding might provide basic discernments of abnormal choice conducts like those of gambling.¹³

Also, an imaging meta-analyses review by Meng et al,¹⁴ reports that GD fellows display a significantly higher activation (compared to healthy controls) in brain areas like right lentiform nucleus and left middle occipital gyrus. Moreover, the South Oaks Gambling Screen scores were linked with overactivity in the right lentiform nucleus and in the bilateral parahippocampus; but the scores were negatively linked to right middle frontal gyrus. Altogether, this suggests dysfunction within the frontostriatal cortical pathway in fellows with GD.¹⁴

In addition, other reviews focus on both human and rodent.^{5,6,15,16} The Norbury and Husain publications review points that a high level of sensation seeking is a factor related to gambling and substance addiction.⁵ Moreover, these authors support the existence of a relationship between sensation seeking and dopaminergic transmission, especially in the D₂ receptors. Specifically, fellows with marked sensation seeking display also elevated DA tonic levels and an over-responsive midbrain dopaminergic responses to signals of future reward.⁵ Moreover, Norbury and Husain propose that even for stimuli of similar strength, reactive responses could vary in terms of approach-avoidance displayed by the subject; the authors propose that these variations stem from differences in the efficiency of DA transmission at the level of the striatum.⁵

Additionally, another review by Quintero concludes that pathological and nonpathological gamblers can differ in terms of brain's anatomy, brain's physiology, electroencephalography (EEG) profile, executive and cognitive efficiency.⁶ For instance, fellows with GD can denote alterations in the insula, OFC, and frontal lobe;⁶ moreover, fellows with GD compared to nonpathological gambler show differences in frontoparietal activation pattern (if winning or losing a game) and insular activity (altered cognitive interpretation of near-miss results and trial success) related to gaming.⁶ With respect to anatomical differences between gamblers and non-gamblers, the first ones show more gray-matter volume compared to normal subjects, based on magnetic resonance imaging technology; furthermore, gamblers have a smaller size of right thalamus, right hippocampus, and left putamen compared to normal subjects.⁶ Regarding research on

rodent, this review states that the correctness of gambling decision is affected by the action of DA receptors and brain areas like insular cortex (rostral agranular zone), infralimbic, and prelimbic.⁶

Another review by Potenza stresses that diverse neurotransmitters like glutamate, noradrenaline, DA, 5-HT, opioid, and brain structures like insula, ventral striatum, and vmPFC (among other areas) are linked to gambling and GD.¹⁵ Furthermore, a literature review by van den Bos et al did focus on three almost ignored aspects of GD: developmental sex differences in GD, adolescence as a sensitive period for developing GD, and paths for upgrading ecological validity of investigative tools.¹⁶

Finally, another set of reviews has been specialized on rodents.^{3,4,17,18} Particularly, a literature review by Alguacil and González-Martin stressed on the “umbrella category” of reward deficiency syndrome; this syndrome includes diverse neuropsychiatric and addiction disorders (including gambling).³ More, these disorders share dysfunctional reward sensitivity, inadequate impulsivity, and/or compulsive conduct. That review considers that further investigation about the reward deficiency syndrome could ease the design of new drugs that are efficient for that cluster of disorders.³

A publications review by Winstanley and Clark emphasizes that the adequate laboratory models for GD should screen fundamental cognitive procedures and have adequate translatability to different species;⁴ moreover, models with these characteristics have a potential capacity to contribute to decision/neuroscience and the investigation of addictive behaviors.⁴ Another review by Anselme stresses that deprivation and randomness, either psychological or physiological, increase the motivation for searching valuable stimuli;¹⁷ moreover, this increase in motivation relates to the organism’s hardness for forecasting relevant environment’s stimulus and incidents.¹⁷

Finally, a review by Cocker and Winstanley states that cognitive biases are important in the evolution of GD;¹⁸ furthermore, these biases can be recreated in rodent models. In effect, some evidences suggest that biases can be linked to dopaminergic activity, especially in the D₄ receptor; the authors suggest the exploration of the D₄ receptor as an alternative for treating GD.¹⁸

The present review aims to integrate recent research findings about the rodent neurobiology of GD and related behaviors. It is expected that this integration could ease the elaboration of most complete pharmacological and/or

behavioral approaches for treating GD in human populations.

Materials and methods

Inclusion and exclusion criteria

The publications were selected based on the next inclusion criteria: a) rodent studies (mice or rat), b) experimental or quasi-experimental design, c) publications that include description about the relationship between the nervous system (brain and/or neurotransmitter) and gambling behavior or GD, d) publications that detail the number of animals, e) the sex of the animals could be male, female or not specified in the publication, f) publications written in English (at least title and abstract), and g) publications released within a recent five years temporal range: October 2012–October 2017. As a reference, some reviews were added, but principally for the introduction and discussion parts.

With respect to the exclusion criteria of the publications, these included the next: a) the non-compliance of the inclusion criteria, and b) it should not be an abstract, nor a publication of a scientific meeting, nor a publication included in non-scientific literature.

Inquiry strategy

A screening of publications in the Pub Med database was carried out based on the five recent years (October/01/2012–October/20/2017). The search terms included: “Gambling” AND “Brain”, “Gambling” AND “Neurobiology”, “Gambling Disorder” AND “Brain”, and “Gambling Disorder” AND “Neurobiology”. The next filters were added for the searching process: text availability (Abstract), species (other animals), Languages (English). Initially, 162 references were obtained in the Pub med search. A total of 121 references was eliminated by different factors (literature review or meta-analysis type, non-English language, human specie, duplicates, and others), resulting in a total of 41 references for subsequent analysis. In addition, another 25 complementary references were detected through references scanning or web searching, and added to the manuscript. As a reference, these 25 references related mainly to the background in the field (included in the introduction section) and to neuroanatomical references. The complete (full) forms of these 66 references were obtained on the web or solicited directly to the authors; later these references were evaluated for the preparation of the review.

More information is described in [Figure 1](#) (Flow diagram of publication selection process). As a guide for the

reader, the first part of the manuscript (results section) includes experimental works that manipulate neurotransmitters and receptors in a non-invasive way (for instance: subcutaneous administration (sc) or intraperitoneal administration (ip)). The second part describes brain structures, and also neurotransmitters/receptors that were evaluated in an invasive way (for instance, brain microinjection).

For the elaboration of this literature review, it was followed by the ethical principles and guidelines of the Helsinki Declaration.

Results

There are two tables summarizing the results: [Table 1](#) entitled “Summary of main studies included in the review – Neurotransmitters”, and [Table 2](#) entitled “Summary of main studies included in the review – Brain structures”. These tables detail different aspects of the studies revised like: neurotransmitters, messengers and receptors systems studied, the brain area, drug name, drug effects, drug route of administration, mental process and/or conduct analyzed, behavioral test (paradigm) used,




Figure 1 Flow diagram of publication selection process. The diagram presents the plan used for publications choice, starting from initial Pub Med database search, up to the final articles incorporated in the publication.

Table 1 Summary of main studies included in the review – Neurotransmitters

Author & year	Silveira et al, 2015 ¹⁹	Yates et al, 2015 ²⁰	Yates et al, 2015 ²⁰	Gueye et al, 2016 ²¹	Di Ciano et al, 2015 ³⁰	Pes et al, 2017 ⁴³
Messenger & receptor	Acetylcholine – muscarinic	Glutamate –NMDA	Glutamate –NMDA	Cannabinoids – CB ₁ /CB ₂ receptors	Dopamine –D ₄ receptor	Dopamine –D ₂ /D ₃ receptor
Brain area	na	na	na	na	na	na
Drug name, effect, route	Scopolamine, antagonist, sc	MK-801, antagonist, sc	Ketamine, antagonist, ip	WIN 55, 212–2, agonist, ip	PD 168077, agonist, ip, L-745,870, antagonist, ip	Pramipexole, agonist, sc
Mental process and/or conduct analyzed	Decision making	Decreased sensitivity to delayed and uncertain reinforcement	Decreased sensitivity to reinforcer amount without altering delay/probability discounting	Improved choice strategy and increased choice latency (SO group); but increased perseverations punished (O group)	- D ₄ agonist small decrease latencies - D ₄ antagonist small increase latencies	Increase unfavorable decisions, impaired the discounting of probabilistic losses, and augment risk-taking behaviors
Behavioral test	Rodent gambling task	Delayed/probabilistic reinforcement, sensitivity to reinforcer amount	Delayed/probabilistic reinforcement, sensitivity to reinforcer amount	Rodent gambling task	Rodent gambling task	Probability discounting task
Species	Rats	Rats	Rats	Rats	Rats	Rats
Relevance of study	Receptor affect decision making under conditions of risk and uncertainty	- NMDA differ mediating impulsivity - MK-801 increased risky decision - MK-801 may be ineffective for GD	- NMDA differ mediating impulsivity	Agonism of CB ₁ /CB ₂ affect gambling choice differentially in healthy and dysfunctional subjects	Modest effect on latency in decision making	- Pramipexole distorted representation of rewards and impaired ability to discern contingencies (unfavorable and favorable) - Pramipexole conduct effects dissociates from changes in mesolimbic dopamine

Abbreviations: CB₁, cannabinoid receptor 1; CB₂, cannabinoid receptor 2; D₂, dopamine 2 receptor; D₃, dopamine 3 receptor; D₄, dopamine 4 receptor; GD, gambling disorder; ip, intraperitoneal administration; NMDA, N-methyl-D-aspartate; na, not applicable (because non-specific brain area was studied); O group, optimal group; sc, subcutaneous administration; SO group, suboptimal group.

Table 2 Summary of main studies included in the review – Brain structures

Author & year Brain area	Fitoussi et al, 2015 ⁴⁴ Amygdala	Tremblay et al, 2014 ⁴⁵ Basolateral amygdala	Zeeb & Winstanley, 2013 ⁴⁶ Orbitofrontal cortex – basolateral amygdala pathway	Cocker et al, 2016 ⁴⁸ Anterior cingulate cortex	Cocker et al, 2016 ⁴⁸ Cingulate cortex (anterior)	Pittaras et al, 2016 ⁵⁰ Hippocampus
Messenger & receptor	Serotonin, Serotonergic receptor	—	—	GABA, GABA _A and GABA _B	Dopamine, D ₄	Dopamine, serotonin and noradrenalin
Drug name, effect, route	—	—	—	Muscimol, agonist, infusion and baclofen, agonist, infusion	PD168077, agonist, infusion	—
Behavioral test	Rat gambling task	Loss – chasing task, the betting task	Rat gambling task	Rodent slot machine task	Rodent slot machine task	Mice gambling task
Other technique/ treatment used	—	Lesion of basolateral amygdala	Disconnection of pathway	—	—	—
Mental process and/or conduct analyzed	Decision making	Risk seeking	—	—	—	—
Species	Rat	Rat	Rat	Rat	Rat	Mice
Relevance of study	Amygdala: low serotonergic metabolism and constant action related to poor decision	- Amygdala plays a role in choice biases related to losses. - Basolateral amygdala lesions reduced loss aversion. - Communication is vital for adequate assessment of reward value to influence choice.	- Disconnection retarded acquisition of the task. - Disconnection prevented changes in the value of a specific reward from contributing appropriately to cost-benefit decision making	Impaired differentiation of wins and loses outcomes in the task. Anterior cingulate cortex is involved in analyzing the best response when competing stimulus and outcome associations are activated	Qualitatively equal effect (compared to left column – GABA), but augmented reward expectancy was only evident on archetypal “near-miss” trials	High levels of dopamine, serotonin, and noradrenalin predicted the emergence of more exploratory and risky behaviors in healthy inbred mice

(Continued)

Table 2 (Continued)

Author & year	Koot et al, 2013 ⁵¹	Ishii et al, 2015 ⁵⁴	Ishii et al, 2015 ⁵⁴	Ishii et al, 2015 ⁵⁴	Pushparaj et al, 2015 ⁵⁵
Brain area	Insular cortex	Insular cortex	Anterior insular cortex	Anterior insular cortex	Caudal agranular Insular cortex
Messenger & receptor	Corticosteroid hormones	GABA, GABA _A , and GABA _B	Dopamine, D ₂	Serotonin, 5-HT _{1A}	GABA, GABA _A , and GABA _B
Drug name, effect, route	Corticosteroid hormones, Agonist, sc	Muscimol, agonist, infusion and Baclofen, agonist, infusion	Eticlopride hydrochloride, antagonist, infusion	WAY100635, antagonist, infusion	Muscimol, agonist, infusion and baclofen, agonist, infusion
Behavioral test	Rodent gambling task	The amount gambling task, and the delay gambling task	Rodent gambling task (modified)	Rodent gambling task (modified)	Rodent gambling task
Other technique/ treatment used	c-fos immuno - histochemistry	—	—	—	Lesion by infusion of ibotenic acid
Mental process and/or conduct analyzed	Decision making	Risk taking behaviors	Risk preference	Risk preference	Decision making: does not favor options linked to high reward frequency and low punishment
Species	Rat	Rat	Rat	Rat	Rat
Relevance of study	Stress (corticosteroid rapid action) disrupts decision making (reward-based)	Insular cortex inactivation induces risk-taking behaviors linked to altered decision-making during tests	Anterior insular cortex inactivation decreased risk preference	Anterior insular cortex blockade increased risk preference, after losing in a previous risky choice	Caudal agranular Insular cortex is not involved in decision-making behavior under risk

(Continued)

Table 2 (Continued)

Author & year	Pushparaj et al, 2015 ⁵⁵	Fitoussi et al, 2015 ⁴⁴	Koot et al, 2013 ⁵¹	Zeep et al, 2015 ⁴⁹	Zeep et al, 2015 ⁴⁹
Brain area	Rostral agranular insular cortex	Lateral ventricle	Infralimbic area	Infralimbic area	Infralimbic area
Messenger & receptor	GABA, GABA _A , and GABA _B	—	Serotonin	GABA, GABA _A , and GABA _B	Dopamine, D ₂
Drug name, effect, route	Muscimol, agonist, infusion and baclofen, agonist, infusion	—	—	Muscimol, agonist, infusion and baclofen, agonist, infusion	Eticlopride hydrochloride, antagonist, infusion
Behavioral test	Rodent gambling task	Rodent gambling task	Rodent gambling task	Rodent gambling task	Rodent gambling task
Other technique/ treatment used	Lesion by means of ibotenic acid infusion	Maternal vitamin D depletion (before and during pregnancy)	c-fos immunohistochemistry	—	—
Mental process and/or conduct analyzed	Decision making: favor options with high reward frequency and low punishment	Decision making	Decision making	Decision making (disrupted by inactivation)	Choice preference (not altered)
Species	Rat	Rat	Rat	Rat	Rat
Relevance of study	Rostral agranular insular cortex is involved in decision-making behavior under risk	Lateral ventricles enlargement did not alter decision making of the offspring	Infralimbic higher serotonergic metabolism relates to poor decision making	Inactivation of infralimbic area has effects on option choice: increase for bad and decrease for good choices	Infralimbic disruption does not alter choice preference nor optimal performance

(Continued)

Table 2 (Continued)

Author & year	Zeeb et al, 2015 ⁴⁹	Peak et al, 2015 ⁵⁶	Roitman & Loriaux, 2014 ⁵⁷	Lobo et al, 2015 ⁵⁸	Ishii et al, 2012 ⁵³	Ishii, et al 2015 ⁵⁴
Brain area	Prelimbic cortex	Neocortex	Nucleus accumbens	Olfactory Tubercle (Islands of Calleja, and islands of Calleja Major)	Orbitofrontal cortex	Orbitofrontal cortex
Messenger & receptor	GABA, GABA _A , and GABA _B	—	—	D ₃	GABA, GABA _A and GABA _B	Serotonin, 5-HT _{1A}
Drug name, effect, route	Muscimol, agonist, infusion and baclofen, agonist, infusion	—	—	—	Muscimol, agonist, infusion and baclofen, agonist, infusion	WAY100635, antagonist, infusion
Behavioral test	Rodent gambling task	Rodent gambling task	Go - no/go task	Rodent gambling task	The amount of gambling task. The delay gambling task	Rodent gambling task (modified)
Other technique/ treatment used	—	Maternal vitamin D depletion (before and during pregnancy) Decision making	Electrophysiological recordings of neurons Impulsivity/Restrain (Nucleus accumbens cue responses correlated with action, regardless of cue type or accuracy)	In situ hybridization (messenger ribonucleic acid levels of D ₃)	—	—
Mental process and/or conduct analyzed	Choice selection was unaffected by inactivation	Thinner neocortex did not alter decision making of the offspring	Individual cue-responsive neurons showed either increases or decreases in activity at cue onset	—	Risk preference	Risk preference
Species	Rat	Rat	Rat	Rat	Rat	Rat
Relevance of study	Prelimbic inactivation: does not alter choice preference	Thinner neocortex did not alter decision making of the offspring	Individual cue-responsive neurons showed either increases or decreases in activity at cue onset	messenger ribonucleic acid levels of D ₃ in Islands of Calleja (r = -0.91), and islands of Calleja Major (r = 0.62) are correlated with performance in task	Orbitofrontal cortex inactivation increase risk preference	Orbitofrontal cortex antagonism decreased risk preference

(Continued)

Table 2 (Continued)

Author & year	Ishii et al, 2015 ⁵⁴	Zeeb et al, 2015 ⁴⁹	Paine et al, 2013 ⁶⁰	Paine et al, 2015 ⁶¹	Baarendse et al, 2013 ⁶²
Brain area	Orbitofrontal cortex	Orbitofrontal cortex	Medial prefrontal cortex	Medial prefrontal cortex	Medial prefrontal cortex
Messenger & receptor	Dopamine, D ₁ and D ₂	GABA, GABA _A , and GABA _B	—	GABA and GABA _A	—
Drug name, effect, route	SCH 23390 hydrochloride, antagonist, infusion Eticlopride, hydrochloride, antagonist, infusion	Muscimol, agonist, infusion and baclofen, agonist, infusion	—	Bicuculline methiodide, antagonist and infusion	—
Behavioral test	Rodent gambling task (modified)	Rodent gambling task	Rodent gambling task	Rodent gambling task	Rodent gambling task
Other technique/treatment used	—	—	- Excitotoxic lesion by ibotenic acid - But, SCH 23390 hydrochloride lessened the lesion induced deficit	—	- Social Isolation: P ₂₁ - P ₄₂ - Cellular Electrophysiology - Five-Choice Serial Reaction Time task
Mental process and/or conduct analyzed	Risk preference	Decision making	Decision making impaired: fewer good choices, and chose the best choice less frequently	Decision making disrupted	Impulsivity/restrain and decision making
Species	Rats	Rats	Rats	Rats	Rats
Relevance of study	Orbitofrontal cortex antagonist did not alter risk preference	Orbitofrontal cortex inactivation did not affect decision making	Medial prefrontal cortex lesion impaired decision making	Disruptions in GABA dynamics contribute to decision-making deficits	- Isolation: impaired impulsive and decision processes under novel or demanding context. - Isolation induced lasting changes in pyramidal neurons and synapses of the medial prefrontal cortex

(Continued)

Table 2 (Continued)

Author & year	Fitoussi et al, 2015 ⁴⁴	Cocker et al, 2012 ⁶³	Gadziola and Wesson, 2016 ⁶⁴	Phillips et al, 2016 ⁶⁵	Aleksandrova et al, 2013 ⁶⁶
Brain area	Prefrontal – subcortical network	Striatum	Ventral striatum, olfactory tubercle	Ventral striatum	Subthalamic nucleus
Messenger & receptor	Dopamine and serotonin	Dopamine, D ₂ and D ₃	—	—	—
Drug name, effect, route	—	—	—	—	—
Behavioral test	Rodent gambling task	Novel task for decision making	Novel water - motivated instrumental task	Stereotypic behavior paradigm	Rodent gambling task
Other technique/ treatment used	- Histology and chromatographic analyses of post-mortem sample, - c-fos immuno - histochemistry	Micro – positron emission tomography, and autoradiography using ¹¹ C raclopride	In vivo electrophysiological recording	Immunohistochemical staining of FosB and ΔFosB technique	- Deep brain stimulation - Histology
Mental process and/or conduct analyzed	Decision making	Bet sensitivity	- Organization of goal directed behavior. - Process of reward information	Stereotypic behavior	Impulsivity/restrain
Species	Rats	Rats	Mice	Mice	Rats
Relevance of study	- Good decision: wider network, a disengagement of key prefrontal areas (insular and infralimbic areas) and the amygdala. - Poor decision: making: low network action and sustained amygdala activity	Low striatal D ₂ and D ₃ density correlates to high wager sensitivity	Encode the onset and progression of motivated behaviors, and discriminate the type and magnitude of a reward	- Elevated activity of the ventral striatum relates to stereotypic behavior - Related to the compulsivity seen in ludopathy	Sustained deep brain stimulation in the subthalamic nucleus, induced increase of premature responding in the task, and this persisted after finishing deep brain stimulation

Abbreviations: [¹¹C] carbon 11; D₁, dopamine 1 receptor; D₂, dopamine 2 receptor; D₃, dopamine 3 receptor; D₄, dopamine 4 receptor; GABA, gamma-aminobutyric acid; GABA_A, gamma-aminobutyric acid receptor A; GABA_B, gamma-aminobutyric acid receptor B; P₂₁, post-natal day 21; P₄₂, post-natal day 42; 5-HT_{1A}, serotonin 1A receptor; 5-HT_{2A}, serotonin 2A receptor; sc, subcutaneous administration.

specie (rat or mice), author and year of publication, and relevance of the study.

Neurotransmitter

Acetylcholine receptor (cholinergic system)

The antagonism of muscarinic receptors (scopolamine, sc) but not of nicotinic receptors (mecamylamine hydrochloride, sc) impaired decision making in rat gambling tasks. Hence, muscarinic receptors can specifically disrupt decision making under conditions of risk and uncertainty (like those found in gambling).¹⁹

NMDA antagonists

The blockade of NMDA receptors (but not AMPA (alpha-amino-3-hydroxy-5-methylisoxazolepropionate) receptors) with the antagonist MK-801 hydrogen maleate (non-competitive antagonist; sc) decreased sensitivity to delayed and uncertain reinforcement in rats, based on the delayed/probabilistic reinforcement, and on the sensitivity to reinforcer amount tests (operant conditioning chambers).²⁰ Moreover, the antagonism with ketamine hydrochloride (uncompetitive antagonist, ip) decreased sensitivity to reinforcer amount without altering delay/probability discounting in the same tests.²⁰ These findings suggest that NMDA receptors differentially mediate impulsivity, with MK-801 hydrogen maleate reducing impulsive choice, but augmenting risky decisions. It is relevant to consider this contrast for treating individuals displaying different psychiatric disorders characterized by impulsivity or risky decisions. In this case, a subject displaying marked impulsive choices would be treated better by means of a drug similar to MK-801 hydrogen maleate; nevertheless, the same medication might be inadequate for someone with GD.²⁰

CB₁ and CB₂

The blockade of CB₁ (antagonist AM 4113, ip) or CB₂ receptors (antagonist/inverse agonist AM 630, ip), or the inhibition of fatty-acid amide hydrolase (URB 597, ip) did not influence the rat gambling task performance.²¹ However, the agonism of CB₁ and CB₂ receptors (WIN 55, 212-2, ip) improved choice strategy, and increased choice latency in the suboptimal group; but only increased perseverative behavior when punished, in the optimal group.²¹ This could be interpreted as the stimulation of cannabinoid receptors could induce different gambling choice conducts based on the type of subjects; specifically, in healthy subjects (optimal group) induce inadequate

conducts, but in dysfunctional subjects (suboptimal group) induced adequate conducts.²¹

As a reference, it has been pointed out that the endocannabinoid system is associated with the reinforcing effects of drugs of abuse.²² CB₂ receptors have been linked to central functions, including a role in addictive processes.^{23,24} Moreover, CB₁ receptors are located pre-synaptically, inhibit synaptic transmission, and allow synaptic modulation.²⁵ Furthermore, CB₂ receptors are located in different zones of the nervous system: periphery,^{26,27} striatum, hippocampus, thalamus,²⁸ and ventral tegmental area (VTA).²⁹

DA receptors

An investigation did report that either D₄ receptors agonism (PD 168077, ip) or D₄ receptors antagonism (L-745,870, ip) had a minimal effect on latency measures and decision making, during the rodent gambling task.³⁰ Additionally, neither the D₃ receptor agonism (PD 128907, ip) nor the D₃ receptor antagonism (SB 277011-A, ip) influenced decision making.³⁰ Also, the antagonism of D₂ receptor (L-741,626, sc) did not affect decision making.³⁰ In general, D₂, D₃, and D₄ ligands did not influence significantly the choice behaviors of the rodent gambling task.

As reference, the D₄ receptors can be found in the next areas within the nervous system: cerebral cortex, amygdala, hypothalamus, pituitary gland, visual system (retina),³¹ and the basal ganglia.^{32–36} Furthermore, D₃ receptors are localized in the islands of Calleja, mammillary bodies, nucleus accumbens (NAc) shell, frontoparietal cortex, the substantia nigra/VTA, basolateral amygdala (BLA), and lateral habenula.^{37–40} In general, some authors agree that the specific localizations of D₃ and D₄ dopamine receptors in the nervous system support their roles in cognition and emotion.^{41,42}

Another investigation reported that pramipexole (an agonist of D₂ and D₃ receptors, sc) induced GD tendencies based on a probability discounting task in rats.⁴³ Specifically, pramipexole augmented unfavorable decisions, disrupted the discounting of probabilistic losses, augmented risk-taking behaviors, distorted the representation of rewards, and impaired the ability to discern favorable from unfavorable contingencies.⁴³ Moreover, the results of complementary studies (voltammetry recordings and High Performance Liquid Chromatography (HPLC)) focused in the NAc suggested that pramipexole behavioral effects were separated from the dynamic changes related to mesolimbic DA release.⁴³

As a reference, the HPLC also measured besides dopamine level, the level of serotonin and norepinephrine. Moreover, the pump speed (Shimadzu LC-6A liquid chromatograph, Columbia, Maryland, United States of America) was 1.5 mL/min. The reverse-phase column utilized was a Rexchrom (Regis Technologies, Morton Grove, Illinois, United States of America) S50100-ODS C18 column with a length of 25 cm and an internal diameter of 4.6 mm. The compounds were measured at +0.7 V using a Shimadzu L-ECD-6A electrochemical detector.

Brain structures

Amygdala

Rat studies did show that amygdala low serotonergic metabolism, or its sustained activity related to poor decision making in the rat gambling task.⁴⁴ Moreover, the lesions of the rat BLA related to reduced risk seeking for losses, but intact risk aversion for gains, based on the loss – chasing task, and the betting task;⁴⁵ these data supported the hypothesis that the amygdala plays a more prominent role in choice biases related to losses. The Tremblay's data suggested that risk seeking for losses being explained by changes in the amygdalar activity, because the amygdala roles in representing the negative affect, and the aversive emotional reaction to loss. Also, these findings discouraged the explanation of risk seeking for losses, because the aberrant estimations of probability or loss magnitude.⁴⁵ This result suggested that communication between these areas is vital for the appropriate assessment of reward value to influence choice.⁴⁵

In addition, another rat investigation explored the disconnection between BLA pathways and the OFC, and found a retarded acquisition in the gambling task.⁴⁶ Based on Zeeb opinion, this disconnection prevented modifications in the value of a specific reward for contributing appropriately to cost-benefit decision making.⁴⁶ Also, it seems that pathways from the OFC to BLA are important in the decision process, and for the adequate assessment of reward value to influence choice.⁴⁶

As a reference, a rat study reported that this specie has decision-making processes that are influenced by a previous reference point,⁴⁷ this study used a modified T maze paradigm. Specifically, the modification did consist of adding “pockets” at both sides (right and left) of the T's stem; because, these pockets in the stem stored pellets, rats could set reference values for each arm of the maze, before selecting.

The “previous expectation” had been previously reported in human research.⁴⁷ It is known that decision-making processes can be disrupted in GD. If the decision making can be influenced by the “previous expectation”, then this expectation should be studied, and its neurobiology for easing the treatment of GD. It is still necessary to find out which brain structure(s) is(are) involved in a “previous expectation”.

Cingulate cortex

Researches regarding the inactivation of the anterior cingulate cortex (ACC) by means of GABA_A and GABA_B agonism reported opposed results. Specifically, one study stated that inactivation of the ACC by means of a mixture of the GABA_A agonist (muscimol; infusion in the brain) and the GABA_B agonist (baclofen; infusion in the brain) impaired rodent's ability to differentiate winning from losing outcomes in a rat slot machine task.⁴⁸ However, the other study that inactivated ACC by means of GABA_B (baclofen hydrochloride, brain microinfusion) and GABA_A (muscimol hydrobromide, brain microinfusion) receptors agonism reported no effect on decision making based on a rat gambling task.⁴⁹

Moreover, another study explored the effect of D₄ agonism (PD168077; infusion in the brain) in the ACC, and it found a disruption of the rat's ability to differentiate winning from losing outcomes in the slot machine task.⁴⁸ Also, it found an augmentation of the reward expectancy, but only on archetypal “near-miss” trials (ie, when the first two of three stimuli in the array were concordant with a rewarding outcome, and only the last stimulus critically signaled a non-win);⁴⁸ Cocker considered that the ACC is fundamental for analyzing the adequate response when competing stimulus and outcome associations are activated; also, this author considered that the D₄ receptor antagonists might be an effective treatment for GD.⁴⁸

Hippocampus

High levels of DA, 5-HT, and noradrenaline in the hippocampus predicted the emergence of more exploratory and risky behaviors in a strain of healthy inbred mice, based on a gambling task.⁵⁰ In this study, they focused on postmortem brain analysis, rather than administration of drugs, or experimental treatments on living brains.

Insular cortex

Some studies evaluated different areas of the insular cortex like: overall insular cortex, anterior insular cortex, or agranular insular cortex (rostral or caudal). The insular cortex

did seem relevant to the rapid (30 mins) disruptive action of corticosteroid hormones (C174 Corticosterone HBC-complex, sc) on decisions (reward based) of the rat's Iowa gambling task.⁵¹ This corticosteroid action was related to stress experience. As a reference, the disruption on decision was accompanied by significant changes in the insular cortex (based on c-fos immuno-histochemistry).

Another study showed that inactivation of insular cortex by a mixture of GABA_A (muscimol; brain microinjection) and GABA_B (baclofen hydrochloride; brain microinjection) receptors agonists induced risky behaviors linked to altered decisions, based on a rat gambling test (a radial arm maze).⁵²

Other studies investigated the anterior insular cortex relevance in gambling-related behaviors, and found mixed results. Specifically, the first investigation found a decreased in risk preference, based on two rats gambling tasks (the amount gambling task and the delay gambling task); the treatment was a mixture of GABA_A (muscimol; brain microinjection) and GABA_B (baclofen; brain microinjection) receptor agonists.⁵³ Moreover, the second investigation blocked the D₂ receptors of the anterior insular cortex (eticlopride hydrochloride, brain microinjection) and found augmentation of risk preference, after winning in a previous risky choice; also, the blockade of the 5-HT_{1A} receptors (WAY100635; brain microinjection) of the anterior insular cortex increased risk preference, after losing in a previous risky choice of a gambling task.⁵⁴ As reference, the antagonism of dopamine 1 receptor (D₁) (SCH23390 hydrochloride) or the antagonism of serotonin 2A receptor (5-HT_{2A}) (M100907, brain microinjection) in the anterior insular cortex did not alter risk preference in the rat gambling task.⁵⁴

Regarding the agranular insular cortex, its inactivation by a mixture of GABA_A (muscimol, brain microinjection) and GABA_B receptors (baclofen, brain microinjection) agonism decrease risk preference based on two different rat gambling tasks; besides, in risk-free control situations, the agranular insular cortex inactivation did not impair decision making.⁵³ Furthermore, the inactivation of the caudal agranular insular cortex by either lesion (ibotenic acid; brain microinjection) or by a mixture of GABA_A (muscimol; brain microinjection) and GABA_B (baclofen; brain microinjection) receptors agonists did not disrupt decision-making behavior under risk in a rat gambling task.⁵⁵

Lateral ventricles

Brain anatomical abnormalities like the enlargement of the lateral ventricles did not alter decision making in the rat

gambling task.⁵⁶ As a reference, the enlargement of ventricles was induced by a maternal (before and during pregnancy) diet deficient in vitamin D; subsequently, the whole litters were placed in a standard diet and evaluated.

Limbic area

Different experimental manipulations of the infralimbic area generated poor decision making in the rat gambling task; for instance: a higher serotonergic metabolism, a rapid action of corticosteroids hormones (30 mins), and inactivation by gamma-aminobutyric acid (GABA) receptors agonism. Specifically, the relationships between a higher serotonergic metabolism in the infralimbic area, and poor decision making in the rat gambling task was inferred based on the postmortem brain analysis after gambling behavioral tests.⁴⁴ Moreover, the rapid and disruptive action of corticosteroid hormones of stress (C174 Corticosterone HBC-complex; sc) on decision making of rats was performed by means of non-invasive brain manipulations.⁵¹ Furthermore, the disruption of decision making after agonism of GABA_A (muscimol hydrobromide; brain microinjection) and GABA_B (baclofen hydrochloride; brain microinjection) receptors was performed by means of direct brain injections; specifically, this inactivation allowed an augmented preference for disadvantageous options and reduced choice for optimal options.⁴⁹

However, other conditions like manipulations of the D₂ receptors did not alter neither choice preference nor optimal performance in the rat gambling task; specifically, the infralimbic cortex was treated by means of administering the D₂ receptor antagonist (eticlopride hydrochloride; brain microinjection).⁴⁹

On the other hand, another set of studies targeted the prelimbic cortex by the agonism of GABAergic receptors and the antagonism of D₂ receptors obtaining opposed results.⁴⁹ Specifically, the inactivation of prelimbic cortex by means of the agonism of GABA_A (muscimol hydrobromide; brain microinjection) and GABA_B (baclofen hydrochloride; brain microinjection) receptors disrupted decision making in the rat gambling task.⁴⁹ Moreover, this inactivation allowed an augmented choice for disadvantageous options, and a reduced choice for optimal options.⁴⁹ Under other conditions, the disruption of the prelimbic cortex by means of treatment with the D₂ receptor antagonist (eticlopride hydrochloride; brain microinjection) did not alter choice preference neither optimal performance in the rat gambling task.⁴⁹

Neocortex

Nervous system anatomical abnormalities like a tiny cerebral cortex did not modify decision making in the rat gambling task.⁵⁶ As a reference, the reduction of the cerebral cortex was produced by a maternal diet (before and during pregnancy) deficient in vitamin D; afterward, the offsprings were placed in a standard diet and evaluated.

NAc

A study found a relationship between the activity of specific cue responsive NAc neurons, and the cue onset during a go/no go task. Despite a gambling task was not used, the go/no go task has relevance to impulsivity, that is a core trait of GD.⁵⁷ Specifically, electrophysiological recordings of neurons in the NAc during the go/no go tasks showed that individual cue-responsive neurons displayed either increases or decreases in activity at the cue onset; NAc cue responses correlated with action, regardless of cue type or accuracy.⁵⁷

Olfactory tubercle

An investigation found a correlation between the messenger ribonucleic acid (mRNA) levels of D₃ receptors in the island of Calleja ($r=-0.91$), in the islands of Calleja major ($r=0.62$) and the performance of male rats in the rodent gambling task (this study only used males).⁵⁸ This finding was consistent with a human imaging study (positron emission tomography) that reported a link between D₃ receptor binding index and the severity of disordered gambling.⁵⁹

OFC

A group of studies with different manipulations (neurotransmitters and receptors systems) reported diverse results: eg, increase in risk preference (inactivation by GABAergic agonism, or rapid corticosteroid action),^{51,53} decrease in risk preference (antagonism of 5-HT_{1A}),⁵⁴ no effects on risk preference (antagonism of D₁, D₂, or 5-HT_{2A} receptors),⁵⁴ and no effect on decision making (inactivation by GABAergic agonism or D₂ receptor antagonism).⁴⁹ However, it is important to note that under risk-free control situations, the GABAergic agonism of OFC did not affect decision making; hence, the degree of risk of the task should be considered.⁵³ Specific details of all the previous reports are explained in the next paragraphs.

The OFC was inactivated by a mixture of GABA_A (muscimol; brain microinjection) and GABA_B (baclofen;

brain microinjection) receptors agonists; this treatment augmented risk preference in the rats, based on two gambling tasks (the amount gambling task, and the delay gambling task).⁵³ However, under risk-free control situations, the inhibition of the OFC did not disrupt decision making. According to the authors, the OFC denoted relevance at the time of accepting or declining a risk.⁵³ Moreover, the lateral OFC did show relevance for the rapid (30 mins) disruptive action of corticosteroid hormone (C174 Corticosterone HBC-complex; sc) on decision making (reward based) in a rat Iowa gambling task.⁵¹ This was inferred because the disruption on decision process was accompanied by significant changes in gene expression in the lateral OFC (increase in c-fos expression, based on c-fos immuno-histochemistry).⁵¹

Nevertheless, the antagonism of 5-HT_{1A} receptors (WAY100635; brain microinjection) in the rat's OFC decreased risk preference on a modified gambling task.⁵⁴ Finally, other studies reported absence of effects of OFC manipulation; specifically, the antagonism of either D₁ receptors (SCH 23390 hydrochloride; brain microinjection), D₂ receptors (eticlopride hydrochloride; brain microinjection), or 5-HT_{2A} receptors (M100907; brain microinjection) did not alter risk preference on a modified gambling task.⁵⁴ Furthermore, the inactivation of the OFC by means of GABA_A (muscimol hydrobromide; brain microinjection) and GABA_B (baclofen hydrochloride; brain microinjection) receptors did not affect decision making in rats, based on a gambling task.⁴⁹ Moreover, D₂ receptor antagonism (eticlopride hydrochloride; brain microinjection) did not affect decision making in the same paradigm.⁴⁹

mPFC

Some studies showed that direct manipulation (ibotenic acid lesion, GABAergic antagonism) or developmental manipulation (adolescence/juvenile social isolation) of the mPFC disrupted decision making.⁶⁰⁻⁶² More details about the previous reports are explained in the next paragraphs.

First, the excitotoxic lesion of mPFC induced by ibotenic acid (brain microinjection) worsened decision making (fewer selection of advantageous or optimal choices) based on a rat gambling task;⁶⁰ however, this deficit in decision making was attenuated after treatment with D₁ receptors antagonism (SCH23390; ip). However, the D₂ receptors antagonism (haloperidol; sc) did not attenuate the deficit.⁶⁰

Furthermore, the antagonism of GABA_A receptors (bicuculline methiodide; brain microinjection) in the mPFC disrupted decision making in the rat gambling task. Despite this study described this application for schizophrenia treatment, this finding is also useful for GD treatment, because it studied decision process during the rat gambling task.⁶¹ Finally, social isolation from early adolescent to juvenile period (post-natal day 21 (P₂₁) to post-natal day 42 (P₄₂)) induced lasting cellular and synaptic changes in the pyramidal neurons of the adult mPFC.⁶² Besides, isolation consequences counteract the DA enhancement induced by a DA agonism bolsterer (amphetamine sulfate; ip) or by a DA reuptake inhibitor (GBR12909 dihydrochloride; ip) in the five-choice serial reaction time task (challenging conditions).⁶² Also, the social isolation decreased sensitivity to DA in the pyramidal neurons of the mPFC. Impulsivity was measured in the rat gambling task and other tests. Also, social isolation impaired impulsive action and decision making under novel or challenging circumstances based on the rat gambling task and other tests. However, impulsive choices were not affected by social isolation.⁶²

PFC – subcortical network and related structures

A rats' study combined gambling tasks, post-mortem analysis (DA and 5-HT turnovers), and c-fos immuno-detection in the brain prefrontal – subcortical network.⁴⁴ Differences between good and bad decision making was found. Good decision making was characterized by a wider network (but once good choices had been made), and a disengagement of the key prefrontal areas (insular and infralimbic cortices) and the amygdala. On the other hand, poor decision making was related to a lower network recruitment and to a sustained amygdala activity.⁴⁴ Besides, poor decision making was linked to an imbalance of monoaminergic metabolism (ie: a higher infralimbic vs a lower amygdalar serotonergic metabolism), and to an aberrant low recruitment of brain areas linked to executive functions and affective valence during decision processes.⁴⁴

Striatum

Studies have looked at the relevance of the striatum in general (rats), and its specific zones like olfactory tubercle and ventral striatum (mice). In general, it was found that striatum activity was linked to wager sensitivity, motivated behavior, discrimination of rewards, stereotypical behavior, and compulsivity.^{63–65} Additional technical details are described in the next paragraphs.

Specifically, lower striatal D₂ and D₃ receptors densities correlated to high wager sensitivity, based on a novel task for decision making in rats, micro-positron emission tomography, and autoradiography using [11C] raclopride.⁶³

In addition, a mice study reported that the olfactory tubercle (a sub-region of the ventral striatum) robustly encoded the onset and progression of motivated behaviors (organization of goal-directed behaviors), and discriminated the type and magnitude of a reward (process of reward information).⁶⁴ As reference, this mice investigation did use a novel water-motivated instrumental task, and “in vivo” electrophysiological recordings; despite this investigation did not perform explicit behavioral tests about gambling, it was proposed by the authors, that the findings were conceptually/theoretically related to GD.⁶⁴ Finally, another investigation performed on mice found that an augmented activity of the ventral striatum related to stereotypical behavior;⁶⁵ the mice were evaluated by means of a stereotypical behavior paradigm, and the brains analyzed by immunohistochemical staining of FosB and delta FosB.⁶⁵ The authors (Phillips et al) proposed that the stereotypy observed could be relevant to the compulsivity described in GD and other disorders (eating and drug seeking).⁶⁵

STN

There was scarce research regarding the involvement of the STN in gambling behavior. Specifically, a study reported that several sessions of bilateral deep brain stimulation (DBS) of the STN induced a subsequent increment of premature responding in the gambling task; this increment even persisted after finishing the stimulation.⁶⁶ As a reference, DBS of the STN had been also associated with impulsivity in the absence of Parkinsonism (under specific conditions).⁶⁶

Discussion

The main question of this review was: which are the main neurotransmitters systems and brain structures relevant for GD based on recent rodent studies? This question was answered in this section by contrasting the present review main points and those from previous literature reviews in the field (reviews cited in the Introduction). The present review found that NMDA receptor antagonism influence reinforcement sensitivity and impulsivity; also, that D₂ and D₃ receptors' agonism induces GD tendencies. These points agree with the publications review by Grant et al,² it concluded that probably diverse neural systems

participate in the pathophysiology of GD like those related to glutamate and DA among others messengers.²

The present work considers that the BLA–OFC pathway is relevant for the assessment of reward among other functions in the rat gambling task. This partially agrees with the Levy and Glimcher's review;¹³ precisely, those authors proposed that the vmPFC/OFC is part of a brain network that codify the values of rewards by means of a standard neural scale (based on human neuroimaging).¹³ It seems that the OFC is related to the assessment and codification of rewards in gambling activities.

Based on the present review, a higher level of DA, norepinephrine, and 5-HT in the hippocampus predicted exploratory and risky behaviors in gambling. Related to this, a review by Meng et al pointed out that bilateral overactivity of the parahippocampus among other structures, positively correlated with South Oaks Gambling Screen scores.¹⁴ Taking these together, it seems that a higher metabolism and activity of the zone of the hippocampus and its surroundings (parahippocampus) relates to more risky gambling tendencies.

Furthermore, this work found that insular cortex activity relates to decision making in the rat gambling task; in the same sense, a previous literature review stated that the insular cortex (including rostral agranular zone) among difference structures influence the correctness of gambling decision.⁶ Taking these together, it seems that insular cortex relates to decision making in gambling tasks.

Moreover, the present review found that infralimbic area relates to decision making in the rat gambling task; this agrees with another publications review that states that infralimbic area among other structures is involved in the correctness of gambling decision on rodent tasks.⁶

The present review found that OFC activity is related to risk preference; relevant to this, another review states that alterations in the OFC among other structures are found in fellows with GD.⁶ It seems that alteration of the OFC activity is relevant for GD. Additionally, this review found that mPFC is involved in decision making, and its disruption impairs decision making. Other reviews have proposed similar ideas; Goulet-Kennedy et al's review pointed that PFC among other structures is fundamental for decision processes based on clinical studies.¹¹ Moreover, a review by Potenza states that vmPFC among other areas is relevant for GD.¹⁵ In general, it seems that PFC (including medial and ventromedial area) is relevant in the dynamics of GD.

Besides, the present review points that PFC–subcortex network activity is linked to poor decision making if lower

network action and sustained activity of the amygdala are present; moreover, PFC–subcortex network is associated with good decision if a wider network and disengagement of key prefrontal areas and the amygdala are present. These points agree with Grant et al's review;² Grant et al work concludes that based on imaging reports, GD relates to anatomical and functional anomalies of nexus involved in reward processing and top-down monitoring.² Hence, both reviews agree that disruption of top-down circuits is a common element in problems linked to gambling.

Regarding the striatum, the present literature review states that the striatum's density of dopamine receptors relates to wager sensitivity; also, the ventral striatum activity relates to stereotypy (like the GD compulsivity). Moreover, the olfactory tubercle relates to the onset and progression of motivated behaviors and reward's discrimination. Similarly, other publication reviews like the one by Goulet-Kennedy et al pointed that the striatum is a conductor of decision processes (which are relevant to gambling behaviors) based on clinical studies.¹¹

Furthermore, the review by Norbury and Husain states that marked sensation seeking relates to GD, and to dopaminergic transmission;⁵ specifically, fellows with marked sensation seeking display high tonic DA levels and over-responsive midbrain dopaminergic responses to signals of future reward.⁵ Also, differences in subject reactions (variability in approach – avoidance reactions) to stimuli stems from differences in the efficiency of DA transmission at the level of striatum.⁵ Another review by Potenza, stresses that the ventral striatum among other structures is linked to gambling and GD.¹⁵ Integrating, the reviews state that striatum relates to wager sensitivity (based on DA receptor density), stereotypy (ventral zone), conduction of decision processes (including those of gambling behaviors), and variability in approach – avoidance to stimuli.

The different points contrasted above in the Discussion section, between the present and other reviews published, have been integrated for elaborating clinical indications. These indications are the next: a) glutamate and DA seem relevant in the pathophysiology of GD; however, other neurotransmitters should also be considered, b) the OFC is relevant for the assessment and codification of rewards in gambling activities, c) a higher metabolism and activity of the hippocampus and its surroundings (parahippocampus) relates to risky gambling tendencies, d) the insular cortex and the infralimbic area are relevant for gambling-

related decisions, e) the alteration of OFC activity is relevant for GD, f) the PFC (including mPFC and vmPFC) is relevant for the dynamic of GD, g) the disruption of top (cortical)–down (subcortical) circuits can be relevant to gambling problems, and h) the striatum relates to wager sensitivity, stereotypy, decision processes, and approach/avoidance to stimuli related to gambling.

Conclusion

Based on the studies revised that used noninvasive methods for drug administration, some of the receptors involved in behaviors related to GD are: muscarinic, NMDA, CB₁, CB₂, D₂, D₃, and D₄ receptors. Moreover, based on the studies revised that used invasive methods for drug administration, some of the neurotransmitters and receptors involved in GD are: 5-HT_{1A}, noradrenaline receptors, GABA_A, and GABA_B. According to this work, the next brain structures are involved in behaviors related to GD: amygdala (including BLA), BLA–OFC pathways, ACC, hippocampus, infralimbic area, prelimbic cortex, insular cortex (including anterior and rostral agranular zones), NAc, olfactory tubercle (island of Calleja and the island of Calleja major), OFC, mPFC, PFC–subcortical network, striatum (including ventral zone and olfactory tubercle), and STN. The present review and others described in the field agree that DA and glutamate, among other neurotransmitters, are relevant to GD. The present review and others described in the field agree that the next brain areas are relevant for GD: OFC, hippocampus/parahippocampus, insular cortex, infralimbic area, PFC, PFC–subcortical network, and striatum. The search for GD treatments should consider and integrate this diversity of neurotransmitters, receptors, and brain areas.

Abbreviation list

AMPA, alpha-amino-3-hydroxy-5-methylisoxazolepropionate; ACC, anterior cingulate cortex; BLA, basolateral amygdala; CB₁, cannabinoid receptor 1; CB₂, cannabinoid receptor 2; DBS, deep brain stimulation; DSM-5, Diagnostic and Statistical Manual of Mental Disorders-5; DA, dopamine; D₁, dopamine 1 receptor; D₂, dopamine 2 receptor; D₃, dopamine 3 receptor; D₄, dopamine 4 receptor; EEG, electroencephalography; GD, gambling disorder; GABA, gamma-aminobutyric acid; GABA_A, gamma-aminobutyric acid receptor A; GABA_B, gamma-aminobutyric acid receptor B; HPLC, High Performance Liquid Chromatography; ip, intraperitoneal administration; mPFC, medial prefrontal cortex; mRNA, messenger ribonucleic acid; mins, minutes;

NMDA, N-methyl-D-aspartate; NAc, nucleus accumbens; OFC, orbitofrontal cortex; P₂₁, post-natal day 21; P₄₂, post-natal day 42; PFC, prefrontal cortex; SENACYT, Secretaria Nacional de Ciencia, Tecnologia e Innovacion (English: National Secretariat of Science, Technology and Innovation); 5-HT, serotonin; 5-HT_{1A}, serotonin 1A receptor; 5-HT_{2A}, serotonin 2A receptor; sc, subcutaneous administration; SNI, Sistema Nacional de Investigacion (English: National System of Investigation); STN, subthalamic nucleus; VTA, ventral tegmental area; vmPFC, ventromedial prefrontal cortex.

Acknowledgments

We would like to thank Open Access Publishing Fund from Florida State University Libraries for its funding. Dr. G. Quintero Garzola belongs and is backed by the SNI (Sistema Nacional de Investigacion) from SENACYT. SENACYT is a public organization located in Panama.

Disclosure

The author reports no conflicts of interest in this work.

References

- Lemieux A, al'Absi M. Stress psychobiology in the context of addiction medicine: from drugs of abuse to behavioral addictions. *Prog Brain Res*. 2016;223:43–62. doi:10.1016/bs.pbr.2015.08.001
- Grant JE, Odlaug BL, Chamberlain SR. Neural and psychological underpinnings of gambling disorder: a review. *Prog Neuropsychopharmacol Biol Psychiatry*. 2016;65:188–193. doi:10.1016/j.pnpbp.2015.10.007
- Alguacil LF, González-Martin C. Target identification and validation in brain reward dysfunction. *Drug Discov Today*. 2015;20(3):347–352. doi:10.1016/j.drudis.2014.10.014
- Winstanley CA, Clark L. Translational models of gambling-related decision-making. *Curr Top Behav Neurosci*. 2016;28:93–120. doi:10.1007/7854_2015_5014
- Norbury A, Husain M. Sensation-seeking: dopaminergic modulation and risk for psychopathology. *Behav Brain Res*. 2015;288:79–93. doi:10.1016/j.bbr.2015.04.015
- Quintero GC. A biopsychological review of gambling disorder. *Neuropsychiatr Dis Treat*. 2017;13:51–60. doi:10.2147/NDT.S118818
- Potenza MN, Kosten TR, Rounsaville BJ. Pathological gambling. *Jama*. 2001;286(2):141–144.
- National Opinion Research Center. *Gambling Impact and Behavior Study*. Chicago: NORC Chicago; 1999. Available from: <http://www.norc.org/PDFs/publications/GIBSFinalReportApril1999.pdf>. Accessed November 20, 2018.
- Lorains FK, Cowlishaw S, Thomas SA. Prevalence of comorbid disorders in problem and pathological gambling: systematic review and meta-analysis of population surveys. *Addiction*. 2011;106(3):490–498. doi:10.1111/j.1360-0443.2010.03300.x
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Arlington, VA: APA; 2013.
- Goulet-Kennedy J, Labbe S, Fecteau S. The involvement of the striatum in decision making. *Dialogues Clin Neurosci*. 2016;18(1):55–63.

12. Banz BC, Yip SW, Yau YH, Potenza MN. Behavioral addictions in addiction medicine: from mechanisms to practical considerations. *Prog Brain Res.* 2016;223:311–328. doi:10.1016/bs.pbr.2015.08.003
13. Levy DJ, Glimcher PW. The root of all value: a neural common currency for choice. *Curr Opin Neurobiol.* 2012;22(6):1027–1038. doi:10.1016/j.conb.2012.06.001
14. Meng YJ, Deng W, Wang HY, et al. Reward pathway dysfunction in gambling disorder: a meta-analysis of functional magnetic resonance imaging studies. *Behav Brain Res.* 2014;275:243–251. doi:10.1016/j.bbr.2014.08.057
15. Potenza MN. Neurobiology of gambling behaviors. *Curr Opin Neurobiol.* 2013;23(4):660–667. doi:10.1016/j.conb.2013.03.004
16. van Den Bos R, Davies W, Dellu-Hagedorn F, et al. Cross-species approaches to pathological gambling: a review targeting sex differences, adolescent vulnerability and ecological validity of research tools. *Neurosci Biobehav Rev.* 2013;37(10 Pt 2):2454–2471. doi:10.1016/j.neubiorev.2013.07.005
17. Anselme P. Dopamine, motivation, and the evolutionary significance of gambling-like behaviour. *Behav Brain Res.* 2013;256:1–4. doi:10.1016/j.bbr.2013.07.039
18. Cocker PJ, Winstanley CA. Irrational beliefs, biases and gambling: exploring the role of animal models in elucidating vulnerabilities for the development of pathological gambling. *Behav Brain Res.* 2015;279:259–273. doi:10.1016/j.bbr.2014.10.043
19. Silveira MM, Malcolme E, Shoaib M, Winstanley CA. Scopolamine and amphetamine produce similar decision-making deficits on a rat gambling task via independent pathways. *Behav Brain Res.* 2015;281:86–95. doi:10.1016/j.bbr.2014.12.029
20. Yates JR, Batten SR, Bardo MT, Beckmann JS. Role of ionotropic glutamate receptors in delay and probability discounting in the rat. *Psychopharmacology.* 2015;232(7):1187–1196. doi:10.1007/s00213-014-3747-3
21. Gueye AB, Trigo JM, Vemuri KV, Makriyannis A, Le Foll B. Effects of various cannabinoid ligands on choice behaviour in a rat model of gambling. *Behav Pharmacol.* 2016;27(2–3Spec Issue):258–269. doi:10.1097/FBP.0000000000000222
22. Le Foll B, Goldberg SR. Cannabinoid CB1 receptor antagonists as promising new medications for drug dependence. *J Pharmacol Exp Ther.* 2005;312(3):875–883. doi:10.1124/jpet.104.077974
23. Gamaledin I, Zvonok A, Makriyannis A, Goldberg SR, Le Foll B. Effects of a selective cannabinoid CB2 agonist and antagonist on intravenous nicotine self-administration and reinstatement of nicotine seeking. *PLoS One.* 2012;7(1):e29900. doi:10.1371/journal.pone.0029900
24. Navarrete F, Rodriguez-Arias M, Martin-Garcia E, et al. Role of CB2 cannabinoid receptors in the rewarding, reinforcing, and physical effects of nicotine. *Neuropsychopharmacol.* 2013;38(12):2515–2524. doi:10.1038/npp.2013.157
25. Melis M, Pistis M. Endocannabinoid signaling in midbrain dopamine neurons: more than physiology? *Curr Neuropharmacol.* 2007;5(4):268–277. doi:10.2174/157015907782793612
26. Maldonado R, Valverde O, Berrendero F. Involvement of the endocannabinoid system in drug addiction. *Trends Neurosci.* 2006;29(4):225–232. doi:10.1016/j.tins.2006.01.008
27. Maldonado R, Berrendero F, Ozaita A, Robledo P. Neurochemical basis of cannabis addiction. *Neuroscience.* 2011;181:1–17. doi:10.1016/j.neuroscience.2011.02.035
28. Gong JP, Onaivi ES, Ishiguro H, et al. Cannabinoid CB2 receptors: immunohistochemical localization in rat brain. *Brain Res.* 2006;1071(1):10–23. doi:10.1016/j.brainres.2005.11.035
29. Zhang HY, Gao M, Liu QR, et al. Cannabinoid CB2 receptors modulate midbrain dopamine neuronal activity and dopamine-related behavior in mice. *Proc Natl Acad Sci U S A.* 2014;111(46):E5007–E5015. doi:10.1073/pnas.1413210111
30. Di Ciano P, Pushparaj A, Kim A, et al. The impact of selective dopamine D2, D3 and D4 ligands on the rat gambling task. *PLoS One.* 2015;10(9):e0136267. doi:10.1371/journal.pone.0136267
31. Cohen AI, Todd RD, Harmon S, O'Malley KL. Photoreceptors of mouse retinas possess D4 receptors coupled to adenylate cyclase. *Proc Natl Acad Sci USA.* 1992;89(24):12093–12097. doi:10.1073/pnas.89.24.12093
32. Valerio A, Belloni M, Gorno ML, Tinti C, Memo M, Spano P. Dopamine D2, D3, and D4 receptor mRNA levels in rat brain and pituitary during aging. *Neurobiol Aging.* 1994;15(6):713–719.
33. O'Malley KL, Harmon S, Tang L, Todd RD. The rat dopamine D4 receptor: sequence, gene structure, and demonstration of expression in the cardiovascular system. *New Biol.* 1992;4(2):137–146.
34. Meador-Woodruff JH, Grandy DK, Van Tol HH, et al. Dopamine receptor gene expression in the human medial temporal lobe. *Neuropsychopharmacol.* 1994;10(4):239–248. doi:10.1038/npp.1994.27
35. Meador-Woodruff JH, Haroutunian V, Powchik P, Davidson M, Davis KL, Watson SJ. Dopamine receptor transcript expression in striatum and prefrontal and occipital cortex. Focal abnormalities in orbitofrontal cortex in schizophrenia. *Arch Gen Psychiatry.* 1997;54(12):1089–1095.
36. Mrzljak L, Bergson C, Pappy M, Huff R, Levenson R, Goldman-Rakic PS. Localization of dopamine D4 receptors in GABAergic neurons of the primate brain. *Nature.* 1996;381(6579):245–248. doi:10.1038/381245a0
37. Khaled MA, Pushparaj A, Di Ciano P, Diaz J, Le Foll B. Dopamine D3 receptors in the basolateral amygdala and the lateral habenula modulate cue-induced reinstatement of nicotine seeking. *Neuropsychopharmacol.* 2014;39(13):3049–3058. doi:10.1038/npp.2014.158
38. Heidbreder C. Novel pharmacotherapeutic targets for the management of drug addiction. *Eur J Pharmacol.* 2005;526(1–3):101–112. doi:10.1016/j.ejphar.2005.09.038
39. Bouthenet ML, Souil E, Martres MP, Sokoloff P, Giros B, Schwartz JC. Localization of dopamine D3 receptor mRNA in the rat brain using in situ hybridization histochemistry: comparison with dopamine D2 receptor mRNA. *Brain Res.* 1991;564(2):203–219.
40. Diaz J, Pilon C, Le Foll B, et al. Dopamine D3 receptors expressed by all mesencephalic dopamine neurons. *J Neurosci.* 2000;20(23):8677–8684.
41. Nakajima S, Gerretsen P, Takeuchi H, et al. The potential role of dopamine D(3) receptor neurotransmission in cognition. *Eur Neuropsychopharmacol.* 2013;23(8):799–813. doi:10.1016/j.euroneuro.2013.05.006
42. Le Foll B, Goldberg SR, Sokoloff P. The dopamine D3 receptor and drug dependence: effects on reward or beyond? *Neuropharmacology.* Sep. 2005;49(4):525–541.
43. Pes R, Godar SC, Fox AT, et al. Pramipexole enhances disadvantageous decision-making: lack of relation to changes in phasic dopamine release. *Neuropharmacology.* 2017;114:77–87. doi:10.1016/j.neuropharm.2016.11.014
44. Fitoussi A, Le Moine C, De Deurwaerdere P, et al. Prefrontal-subcortical imbalance characterizes poor decision-making: neurochemical and neural functional evidences in rats. *Brain Struct Funct.* 2015;220(6):3485–3496. doi:10.1007/s00429-014-0868-8
45. Tremblay M, Cocker PJ, Hosking JG, Zeeb FD, Rogers RD, Winstanley CA. Dissociable effects of basolateral amygdala lesions on decision making biases in rats when loss or gain is emphasized. *Cogn Affect Behav Neurosci.* 2014;14(4):1184–1195. doi:10.3758/s13415-014-0271-1
46. Zeeb FD, Winstanley CA. Functional disconnection of the orbitofrontal cortex and basolateral amygdala impairs acquisition of a rat gambling task and disrupts animals' ability to alter decision-making behavior after reinforcer devaluation. *J Neurosci.* 2013;33(15):6434–6443. doi:10.1523/JNEUROSCI.3971-12.2013
47. Bhatti M, Jang H, Kralik JD, Jeong J. Rats exhibit reference-dependent choice behavior. *Behav Brain Res.* 2014;267:26–32. doi:10.1016/j.bbr.2014.03.012

48. Cocker PJ, Hosking JG, Murch WS, Clark L, Winstanley CA. Activation of dopamine D4 receptors within the anterior cingulate cortex enhances the erroneous expectation of reward on a rat slot machine task. *Neuropharmacology*. 2016;105:186–195. doi:10.1016/j.neuropharm.2016.01.019
49. Zeeb FD, Baarendse PJ, Vanderschuren LJ, Winstanley CA. Inactivation of the prelimbic or infralimbic cortex impairs decision-making in the rat gambling task. *Psychopharmacology*. 2015;232(24):4481–4491. doi:10.1007/s00213-015-4075-y
50. Pittaras E, Callebert J, Chennaoui M, Rabat A, Granon S. Individual behavioral and neurochemical markers of unadapted decision-making processes in healthy inbred mice. *Brain Struct Funct*. 2016;221(9):4615–4629. doi:10.1007/s00429-016-1192-2
51. Koot S, Baars A, Hesselting P, van Den Bos R, Joels M. Time-dependent effects of corticosterone on reward-based decision-making in a rodent model of the Iowa gambling task. *Neuropharmacology*. 2013;70:306–315. doi:10.1016/j.neuropharm.2013.02.008
52. Mizoguchi H, Katahira K, Inutsuka A, et al. Insular neural system controls decision-making in healthy and methamphetamine-treated rats. *Proc Natl Acad Sci U S A*. 2015;112(29):E3930–E3939. doi:10.1073/pnas.1418014112
53. Ishii H, Ohara S, Tobler PN, Tsutsui K, Iijima T. Inactivating anterior insular cortex reduces risk taking. *J Neurosci*. 2012;32(45):16031–16039. doi:10.1523/JNEUROSCI.2278-12.2012
54. Ishii H, Ohara S, Tobler PN, Tsutsui K, Iijima T. Dopaminergic and serotonergic modulation of anterior insular and orbitofrontal cortex function in risky decision making. *Neurosci Res*. 2015;92:53–61. doi:10.1016/j.neures.2014.11.009
55. Pushparaj A, Kim AS, Musiol M, et al. Differential involvement of the agranular vs granular insular cortex in the acquisition and performance of choice behavior in a rodent gambling task. *Neuropsychopharmacol*. 2015;40(12):2832–2842. doi:10.1038/npp.2015.133
56. Peak JN, Turner KM, Burne TH. The effect of developmental vitamin D deficiency in male and female Sprague-Dawley rats on decision-making using a rodent gambling task. *Physiol Behav*. 2015;138:319–324. doi:10.1016/j.physbeh.2014.09.007
57. Roitman JD, Loriaux AL. Nucleus accumbens responses differentiate execution and restraint in reward-directed behavior. *J Neurophysiol*. 2014;111(2):350–360. doi:10.1152/jn.00350.2013
58. Lobo DS, Aleksandrova L, Knight J, et al. Addiction-related genes in gambling disorders: new insights from parallel human and pre-clinical models. *Mol Psychiatry*. 2015;20(8):1002–1010. doi:10.1038/mp.2014.113
59. Boileau I, Payer D, Chugani B, et al. The D2/3 dopamine receptor in pathological gambling: a positron emission tomography study with [11C](+)-propyl-hexahydro-naphtho-oxazin and [11C]raclopride. *Addiction*. 2013;108(5):953–963. doi:10.1111/add.12066
60. Paine TA, Asinof SK, Diehl GW, Frackman A, Leffler J. Medial prefrontal cortex lesions impair decision-making on a rodent gambling task: reversal by D1 receptor antagonist administration. *Behav Brain Res*. 2013;243:247–254. doi:10.1016/j.bbr.2013.01.018
61. Paine TA, O'Hara A, Plaut B, Lowes DC. Effects of disrupting medial prefrontal cortex GABA transmission on decision-making in a rodent gambling task. *Psychopharmacology*. 2015;232(10):1755–1765. doi:10.1007/s00213-014-3816-7
62. Baarendse PJ, Counotte DS, O'Donnell P, Vanderschuren LJ. Early social experience is critical for the development of cognitive control and dopamine modulation of prefrontal cortex function. *Neuropsychopharmacol*. 2013;38(8):1485–1494. doi:10.1038/npp.2013.47
63. Cocker PJ, Dinelle K, Kornelson R, Sossi V, Winstanley CA. Irrational choice under uncertainty correlates with lower striatal D (2/3) receptor binding in rats. *J Neurosci*. 2012;32(44):15450–15457. doi:10.1523/JNEUROSCI.0626-12.2012
64. Gadziola MA, Wesson DW. The neural representation of goal-directed actions and outcomes in the Ventral Striatum's olfactory tubercle. *J Neurosci*. 2016;36(2):548–560. doi:10.1523/JNEUROSCI.3328-15.2016
65. Phillips D, Choleris E, Ervin KS, et al. Cage-induced stereotypic behaviour in laboratory mice covaries with nucleus accumbens FosB/DeltaFosB expression. *J Neurosci*. 2016;301:238–242.
66. Aleksandrova LR, Creed MC, Fletcher PJ, Lobo DS, Hamani C, Nobrega JN. Deep brain stimulation of the subthalamic nucleus increases premature responding in a rat gambling task. *Behav Brain Res*. 2013;245:76–82. doi:10.1016/j.bbr.2013.02.011

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and

is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>