

ORAL PRESENTATION

Open Access

Off-pump coronary revascularization performed concomitant with non-cardiovascular surgery

JW Han*, KH Lee, HY Hwang, KB Kim

From 23rd World Congress of the World Society of Cardio-Thoracic Surgeons
Split, Croatia. 12-15 September 2013

Background

A concern about complications secondary to the cardiopulmonary bypass caused reluctance to perform combined cardiac and non-cardiovascular surgery. We studied the patients who underwent off-pump CABG concomitant with non-cardiovascular surgery.

Methods

Of 2083 patients who underwent isolated off-pump CABG between 1999 and 2012, 91 patients who underwent off-pump CABG concomitant with non-cardiovascular surgery (group 1) were compared with 1991 patients who underwent isolated off-pump CABG (group 2). In group 1, there were 49 malignancies and 42 benigns. Non-cardiovascular surgery included 34 thoracic surgeries (11 bullectomy, 11 pulmonary wedge resection, 5 thymectomy, 4 lobectomy, 1 tracheoesophageal fistulectomy, 1 radical mastectomy, and 1 diaphragm plication), 56 abdominal surgery (24 gastrectomy, 14 cholecystectomy, 7 colectomy, 5 herniorrhaphy, 2 mass excision, 2 nephrectomy, 1 feeding jejunostomy, and 1 perigastric LN biopsy), and 1 below-knee amputation.

Results

Mean age at operation were 67.8 ± 9.6 years (group 1) and $63.9 \text{ years} \pm 9.5$ (group 2), respectively. There were no significant differences in preoperative risk factors, including smoking, hypertension, diabetes mellitus, dyslipidemia, obesity, stroke, left ventricular dysfunction, and chronic renal failure between the 2 groups. Mean number of distal anastomosis was 2.93 ± 1.0 in group 1 and 3.09 ± 0.97 in group 2 ($p=0.133$). Operative mortality was 1.1% (1/91) in group 1 and 1.3% (26/1991) in group 2, respectively ($p=1.000$). There were no significant differences in postoperative morbidities, including mediastinitis,

re-exploration for bleeding, perioperative myocardial infarction, low cardiac output syndrome, atrial fibrillation, and stroke between the 2 groups.

Conclusions

Off-pump CABG concomitant with non-cardiovascular surgery was not associated with increased operative mortality and postoperative morbidities.

Published: 11 September 2013

doi:10.1186/1749-8090-8-S1-O179

Cite this article as: Han et al.: Off-pump coronary revascularization performed concomitant with non-cardiovascular surgery. *Journal of Cardiothoracic Surgery* 2013 **8**(Suppl 1):O179.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



* Correspondence: irene962063@naver.com

Adult Cardiac Surgery, Seoul National University Hospital, Seoul, Korea