

LncRNA SAMMSON Overexpression Suppresses Vascular Smooth Muscle Cell Proliferation via Inhibiting miR-130a Maturation to Participate in Intracranial Aneurysm

Wen Pan¹
Yuan Gao²
Weifeng Wan³
Wenfeng Xiao⁴
Chao You⁵

¹Brain Center, Shougang Shuigang General Hospital, Liupanshui City, Guizhou Province, 553000, People's Republic of China; ²Department of Neurosurgery, Liupanshui People's Hospital, Liupanshui City, Guizhou Province, 553001, People's Republic of China; ³Department of Neurosurgery, Affiliated Hospital of Southwest Medical University, Luzhou City, Sichuan Province, 646000, People's Republic of China; ⁴Department of Neurosurgery, Sichuan Mianyang 404 Hospital, Mianyang City, Sichuan Province, 510700, People's Republic of China; ⁵Department of Neurosurgery West China Hospital, Sichuan University, Chengdu Province, 610041, People's Republic of China

Background: MiR-130a is a recently identified critical player in vascular smooth muscle cell (VSMC) proliferation, which participates in intracranial aneurysm (IA). However, the involvement of miR-130a in IA and its upstream regulator are unknown. Our preliminary sequencing analysis revealed a close correlation between miR-130a and lncRNA SAMMSON across IA samples. Therefore, we further studied the crosstalk between SAMMSON and miR-130a in IA.

Methods: SAMMSON and miR-130a expression were measured using RT-qPCR. SAMMSON subcellular location was analyzed with nuclear fractionation assay. Their direct interaction was explored with RNA pull-down assay. The role of SAMMSON in miR-130a maturation was studied with overexpression analysis. VSMC cell proliferation was analyzed with BrdU assay.

Results: SAMMSON and premature miR-130a were deregulated in IA, while mature miR-130a was upregulated in IA. SAMMSON is localized in both the nucleus and cytoplasm, and direct interaction between SAMMSON and miR-130a was observed. SAMMSON overexpression suppressed miR-130a maturation in VSMCs and reduced the enhancing effects of miR-130a on VSMC cell proliferation.

Conclusion: SAMMSON is overexpressed in IA and suppresses VSMC proliferation via inhibiting miR-130a maturation.

Keywords: intracranial aneurysm, SAMMSON, miR-130a, maturation

Introduction

Intracranial aneurysm (IA), also widely known as brain aneurysm, is a severe cerebrovascular disorder caused by the cerebral vein or artery wall weakness.^{1,2} IA is commonly treated with ballooning or dilation of local blood vessels. IA may rupture or leak, and the ruptured IA could lead to bleeding in brain tissues, causing hemorrhagic stroke.^{3,4} It is estimated that more than 25% of IA patients will die after IA rupture, and IA complications cause deaths in another 25% of patients within 6 months after rupture.⁴⁻⁶ The ruptured IA is usually treated with surgical clipping or less invasive endovascular coiling.⁷ However, these two treatments might reduce blood flow to the brain and bleeding in the brain.⁷ At present, the prevention of IA rupture is still critical for IA treatment.

Correspondence: Chao You
Department of Neurosurgery West China Hospital, Sichuan University, 37 Guoxue Lane, Wuhou District, Chengdu Province, 610041, People's Republic of China
Email chaoyousichuan@163.com

Previous studies have elucidated the correlation between IA development and many risk factors, such as alcoholism, tobacco consumption, hypertension, obesity, head trauma, and infections.^{2,8,9} Besides, alterations in the expression of some molecular factors may also contribute to IA.¹⁰ For instance, lncRNAs and miRNAs affect protein production and regulate other noncoding RNAs to regulate IA growth and rupture.^{11,12} Although lacking protein-coding capacity, lncRNAs and miRNAs participate in human diseases, including IA, mainly by affecting the expression of protein-coding genes. Therefore, regulating the expression of certain miRNAs or lncRNAs with critical functions in IA may provide potentials for IA treatment.^{11,12}

MiR-130a enhances the proliferation of vascular smooth muscle cells (VSMCs),¹³ which participate in IA.¹⁴ It is still unknown whether miR-130a participates in IA. Our preliminary sequencing analysis revealed a close correlation between miR-130a and lncRNA SAMMSON, a well-studied cancer-related lncRNA,¹⁵ across IA samples. SAMMSON plays a critical role in cancer biology by interacting with critical signal pathways involved in cancer development, such as PI3K/Akt Pathway.¹⁵ However, the involvement of SAMMSON in IA is unknown. This study analyzed the potential interaction between SAMMSON and miR-130a in IA.

Materials and Methods

Patients and Blood Extraction

Aneurysmal tissues were collected from 30 patients with ruptured IAs at West China Hospital, Sichuan University during microsurgical clipping. In addition, superficial temporal arteries (STAs) were collected from other 30 patients who underwent lateral frontal craniotomies and pterional craniotomies due to injuries at the same hospital. All tissue samples were stored at -80°C prior to subsequent RNA isolations. The study was approved by the Ethics Committee of West China Hospital, Sichuan University. All IAs and controls signed informed consent. [Table 1](#) shows the clinical data of the two groups of participants.

VSMCs and Electric Transfection

Human VSMCs were purchased from Clonetics (San Diego, USA) and cultured in medium 231 containing smooth muscle growth supplements at 37°C in an incubator with 5% CO_2 . A total of 10^7 VSMCs were transfected with either 10 μg pcDNA3.1-SAMMSON expression vector, 50 nM miR-

Table 1 Clinical Data of Two Groups of Participants

	IAs (n=30)	STAs (n=30)
Gender (Female %)	11 (36.67%)	12 (40.00%)
IA Size (mm, mean \pm SD)	12.37 \pm 8.19	NA
Age (Years, mean \pm SD)	55.83 \pm 6.16	54.98 \pm 11.39
Hypertension %	19 (63.33%)	15 (50%)
Smokers %	8 (26.67%)	8 (26.67%)
Drinkers %	7 (23.33%)	7 (23.33%)

130a mimic, or their corresponding negative controls using Neon Electroporation Transfection system (Thermo Fisher Scientific). In addition, untreated cells were used as the control (C) cells. The subsequent analyses were performed at 48 h of post-transfection.

RNA Isolation and Analysis

Total RNAs were extracted from cells of each transfection group and tissue samples using the Direct-zol RNA Kit (ZYMO RESEARCH) and treated with DNase I to remove DNA. RNA quality was analyzed by Agilent 2100 Bioanalyzer (Agilent Technologies, USA). Only RNA samples with high quality were used for subsequent experiments. Otherwise, RNA isolations were repeated.

RT-qPCR

A total of 1000 ng total RNA were reverse transcribed into cDNA, followed by qPCR to determine the expression of SAMMSON and premature miR-130a using 18S rRNA as the endogenous control. Mature miR-130a expression was analyzed with All-in-OneTM miRNA qRT-PCR Reagent Kit (Genecopoeia) by qPCR with U6 as the endogenous control. The method of $2^{-\Delta\Delta\text{Ct}}$ was applied to normalize the expression of each gene to the corresponding controls.

Nuclear Fractionation Assay

Nuclear/Cytosol Fractionation Kit (BioVision, # K266) was applied to prepare nuclear and cytosolic samples from VSMCs. Briefly, after cell lysis, the homogenate was centrifuged for 10 min at 2500 g (4°C). The supernatant was used as the cytosolic fraction, and the nuclear pellet was collected to serve as the nuclear fraction. After that, RNA isolation from these two fractions was performed, followed by RTs and PCR to determine the expression of SAMMSON with GAPDH as the endogenous control. Electrophoresis was performed with 1% agarose gel to separate PCR products. After EB staining, images were taken with MyECL Imager (Bio-Rad).

RNA Pull-Down Assay

Vectors with T7 promoter expressing premature miR-130a or NC miRNA were used to prepare transcripts of both premature miR-130a or NC miRNA through in vitro transcription using the T7 RNA polymerase MEGAscript™ T7 Transcription Kit (Cat # AMB13345, Thermo Fisher Scientific). Both transcripts were labeled with biotin using the Pierce™ Biotin 3' End DNA Labeling Kit (Cat # PI89818, VWR) and named Bio-miR-130a and Bio-NC, respectively. The two miRNAs were then transfected into VSMCs as described above. After the preparation of cell lysis at 48h of post-transfection, biotin-ligated miRNAs were pulled down using streptavidin agarose magnetic beads (Life Technologies). The pull-down samples were subjected to RNA isolations and RT-qPCR to determine the expression of SAMMSON.

BrdU Incorporation Assay

Cell proliferation after transfections was analyzed by BrdU incorporation. In brief, transfected cells were seeded onto a 96-well plate to a density of 6000 cells per well. After cultured for 48 h, BrdU was added into each well to a final concentration of 10 μ M. After cultured for another 24 h, cells were fixed for 30 min and then incubated with peroxidase-coupled anti-BrdU-antibody (Sigma–Aldrich) for one hour. After washed with PBS three times, cells were further incubated with peroxidase substrate for 1 h. Finally, OD values at 450 nm were measured.

Statistical Analysis

All in vitro experiments were performed in three biological replicates. Two independent groups were compared by unpaired *t* test. Multiple independent groups were compared by ANOVA Tukey's test. A $P < 0.05$ was statistically significant.

Results

Changed SAMMSON and miR-130a Levels in IA

SAMMSON and miR-130a levels in IAs and STAs were analyzed by RT-qPCR. The results illustrated that the levels of SAMMSON (Figure 1A, $p < 0.01$) and premature miR-130a (Figure 1B, $p < 0.01$) were lower while mature miR-130a was higher in IA than in STA (Figure 1C, $p < 0.01$). Therefore, SAMMSON downregulation and increased miR-130a maturation are likely involved in IA. Correlation analysis revealed that SAMMSON was positively correlated with premature miR-130a (Figure 1D) but inversely correlated with mature miR-130a (Figure 1E).

Subcellular Location of SAMMSON and Its Direct Interaction with Premature miR-130a

Nuclear fractionation assay was performed to analyze the subcellular localization of SAMMSON in VSMCs. With




Figure 1 SAMMSON and miR-130a expression in IAs and STAs. IA and STA samples were subjected to RNA isolation, followed by RT-qPCR to analyze the differential expression of SAMMSON (A), premature miR-130a (B), and mature miR-130a (C). Pearson's correlation coefficient was performed to analyze the correlations of SAMMSON with premature miR-130a (D) and mature miR-130a (E). ** $p < 0.01$.

GAPDH as the endogenous control, our data clearly showed that SAMMSON was detected in both nuclear (N) and cytosolic (C) fractions. Therefore, SAMMSON might traffic between the nucleus and cytoplasm (Figure 2A). RNA pull-down assay was conducted with Bio-miR-130a to examine the direct interaction between premature miR-130a and SAMMSON. Compared to Bio-NC pull-down samples, a significantly higher SAMMSON level was detected in Bio-miR-130a pull-down samples (Figure 2B, $p < 0.01$). As premature miRNAs are localized in the nucleus, we concluded that SAMMSON in the nucleus might directly interact with premature miR-130a.

SAMMSON Suppresses miR-130a Maturation in VSMCs

SAMMSON or mature miR-130a was overexpressed in VSMCs. The follow-up RT-qPCR analyses illustrated the increased expression of both SAMMSON and miR-130a in VSMCs from 24 h to 72h (Figure 3A, $p < 0.05$). In addition, SAMMSON overexpression in VSMCs significantly increased premature miR-130a level (Figure 3B, $p < 0.05$) but significantly decreased mature miR-130a level (Figure 3C, $p < 0.05$) in VSMCs. Therefore, SAMMSON might suppress miR-130a maturation in VSMCs.

Roles of SAMMSON and miR-130a in VSMC Proliferation

BrdU incorporation assay was utilized to analyze VSMC proliferation after overexpression of SAMMSON and/or

miR-130a. The results clearly demonstrated an inhibitory effect of SAMMSON on cell proliferation and enhanced effects of miR-130a on cell proliferation ($p < 0.05$). Moreover, the co-transfection assay illustrated that SAMMSON suppressed the enhancing effects of miR-130a on VSMC proliferation (Figure 4, $p < 0.05$).

Discussion

Our study first reported the interaction between SAMMSON and miR-130a in IA. We observed an altered expression of SAMMSON and miR-130a (both mature and premature form) in IA. Remarkably, miR-130a maturation in VSMCs is regulated by SAMMSON, suggesting the potential application of SAMMSON and miR-130a in the treatment of IA.

A recent study showed that miR-30a is upregulated in a rat spontaneous hypertension model, and miR-30a overexpression targets GAX to promote VSMC proliferation.¹³ Hypertension is a major risk factor for IA.^{2,8,9} Therefore, miR-130a might also participate in IA. This study first reported upregulation of mature miR-130a and decreased premature miR-130a expression in IA. Therefore, the increased maturation of miR-130a but not the increased transcription of miR-130a may participate in IA. The study also confirmed the enhancing effects of miR-130a on VSMC proliferation. The altered VSMC proliferation plays different roles at different stages of IAs.¹⁶ The increased VSMC proliferation rate promotes the growth of IA at early stages and suppresses IA rupture at late




Figure 2 Subcellular location of SAMMSON and its direct interaction with premature miR-130a. Nuclear fractionation assay was conducted to analyze the subcellular localization of SAMMSON in VSMCs (A). RNA pull-down assay was performed with Bio-miR-130a to examine the direct interaction between premature miR-130a and SAMMSON (B). ** $p < 0.01$.




Figure 3 Role of SAMMSON in miR-130a maturation in VSMCs. SAMMSON or mature miR-130a was transfected in VSMCs. Their overexpression in VSMCs was confirmed from 24h to 72h (A). The role of SAMMSON in regulating premature miR-130a (B) and mature miR-130a (C) levels in VSMCs was also analyzed by RT-qPCR. *p<0.05.




Figure 4 Effects of SAMMSON and miR-130a in VSMC proliferation. BrdU incorporation assay was performed to analyze VSMC proliferation after overexpression of SAMMSON and/or miR-130a. *p<0.05.

stages. Therefore, accurate regulation of miR-130a expression might suppress IA growth and rupture.

At present, the upstream regulation of miR-130a in VSMC proliferation is unknown. This study showed that SAMMSON is downregulated in IA, and SAMMSON overexpression in VSMCs suppresses miR-130a maturation, suggesting that SAMMSON is a novel molecular inhibitor of miR-130a maturation. Furthermore, SAMMSON could be detected in both the nucleus and cytoplasm. It has been well established that premature miRNAs are mainly localized in the nucleus while mature miRNAs are mainly localized in the cytoplasm.¹⁷ Therefore, we hypothesized that SAMMSON might sponge premature miR-130a in the nucleus to suppress its maturation. This hypothesis is supported by the observation of the direct interaction between SAMMSON and miR-130a.

The study is limited by the relatively small size of patients and lack of in vivo functional analyses of SAMMSON and miR-130a. Therefore, our conclusions remain to be further validated.

Conclusion

Overall, our data illustrated that SAMMSON overexpression might suppress miR-130a maturation in VSMCs to participate in IAs. Therefore, regulating the expression of SAMMSON and miR-130a may improve the treatment of IA.

Availability of Supporting Data

The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to their containing information that could compromise the privacy of research participants.

Ethical Approval and Consent to Participate

All patients signed the written informed consent. All experiments were approved by the Ethics Committee of West China Hospital, Sichuan University. Procedures operated in this research completely complied with the standards set out in the Announcement of Helsinki and Laboratory Guidelines of Research in China.

Author Contributions

Wen Pan, Chao You: study concepts, literature research, clinical studies, data analysis, experimental studies, manuscript writing and review; Yuan Gao: study design, literature research, experimental studies and manuscript editing; Weifeng Wan: definition of intellectual content, clinical studies, data acquisition and statistical analysis; Wenfeng Xiao: data acquisition, manuscript preparation and data analysis.

All authors have read and approved the submission of the manuscript. All authors contributed to data analysis, drafting or revising the article, have agreed on the journal to which the article will be submitted, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Funding

There is no funding to report.

Disclosure

All authors have no conflicts of interest. We declare that we do not have any commercial or associative interest that represents a conflict of interest connected with the work submitted.

References

1. Wang X, Zhu C, Leng Y, Degnan AJ, Lu J. Intracranial aneurysm wall enhancement associated with aneurysm rupture: a systematic review and meta-analysis. *Acad Radiol.* 2019;26(5):664–673. doi:10.1016/j.acra.2018.05.005
2. Jin D, Song C, Leng X, Han P. A systematic review and meta-analysis of risk factors for unruptured intracranial aneurysm growth. *Int J Surg.* 2019;69:68–76. doi:10.1016/j.ijso.2019.07.023
3. Juvela S, Korja M. Intracranial aneurysm parameters for predicting a future subarachnoid hemorrhage: a Long-Term Follow-up Study. *Neurosurgery.* 2017;81(3):432–440. doi:10.1093/neuros/nyw049
4. de Souza MR, Fagundes CF, Rabelo NN, Teixeira MJ, Figueiredo EG. Association between intracranial aneurysm and meningiomas: an integrative survival Analysis with identification of prognostic factors. *Clin Neurol Neurosurg.* 2020;198:106128. doi:10.1016/j.clineuro.2020.106128
5. Niu X, Wang T, Li J, et al. An integrative survival analysis with identification of prognostic factors in the patients with coexisting glioma and intracranial aneurysm. *World Neurosurg.* 2018;111:e592–e600. doi:10.1016/j.wneu.2017.12.126
6. Ruiz-Sandoval JL, Cantú C, Chiquete E, et al. Aneurysmal subarachnoid hemorrhage in a Mexican multicenter registry of cerebrovascular disease: the RENAMEVASC study. *J Stroke Cerebrovasc Dis.* 2009;18(1):48–55. doi:10.1016/j.jstrokecerebrovasdis.2008.09.019
7. Darsaut TE, Findlay JM, Magro E, et al. Surgical clipping or endovascular coiling for unruptured intracranial aneurysms: a pragmatic randomised trial. *J Neurol Neurosurg Psychiatry.* 2017;88(8):663–668. doi:10.1136/jnnp-2016-315433
8. Zholdybayeva EV, Medetov YZ, Aitkulova AM, et al. Genetic risk factors for intracranial aneurysm in the Kazakh population. *J Mol Neurosci.* 2018;66(1):135–145. doi:10.1007/s12031-018-1134-y
9. Björkman J, Frösen J, Tähtinen O, et al. Aneurysm size is the strongest risk factor for intracranial aneurysm growth in the Eastern Finnish population. *Neurosurgery.* 2019;84(5):1098–1103. doi:10.1093/neuros/nyy161
10. Xu Z, Rui YN, Hagan JP, Kim DH. Intracranial aneurysms: pathology, genetics, and molecular mechanisms. *Neuromolecular Med.* 2019;21(4):325–343. doi:10.1007/s12017-019-08537-7
11. Li H, Wang W, Zhang L, et al. Identification of a long noncoding RNA-associated competing endogenous RNA network in intracranial aneurysm. *World Neurosurg.* 2017;97:684–692.e684. doi:10.1016/j.wneu.2016.10.016
12. Gao G, Zhang Y, Yu J, et al. Long non-coding RNA MALAT1/microRNA-143/VEGFA signal axis modulates vascular endothelial injury-induced intracranial aneurysm. *Nanoscale Res Lett.* 2020;15(1):139. doi:10.1186/s11671-020-03357-2
13. Wu WH, Hu CP, Chen XP, et al. MicroRNA-130a mediates proliferation of vascular smooth muscle cells in hypertension. *Am J Hypertens.* 2011;24(10):1087–1093. doi:10.1038/ajh.2011.116
14. Miyata T, Minami M, Kataoka H, et al. Osteoprotegerin prevents intracranial aneurysm progression by promoting collagen biosynthesis and vascular smooth muscle cell proliferation. *J Am Heart Assoc.* 2020;9(17):e015731. doi:10.1161/JAHA.119.015731
15. Ni H, Wang K, Xie P, Zuo J, Liu W, Liu C. LncRNA SAMMSON knockdown inhibits the malignancy of glioblastoma cells by inactivation of the PI3K/Akt pathway. *Cell Mol Neurobiol.* 2021;41(1):79–90. doi:10.1007/s10571-020-00833-2
16. Frösen J, Tulamo R, Paetau A, et al. Saccular intracranial aneurysm: pathology and mechanisms. *Acta Neuropathol.* 2012;123(6):773–786. doi:10.1007/s00401-011-0939-3
17. Li XG, Wang YB. SRPK1 gene silencing promotes vascular smooth muscle cell proliferation and vascular remodeling via inhibition of the PI3K/Akt signaling pathway in a rat model of intracranial aneurysms. *CNS Neurosci Ther.* 2019;25(2):233–244. doi:10.1111/cns.13043

Neuropsychiatric Disease and Treatment

Dovepress

Publish your work in this journal

Neuropsychiatric Disease and Treatment is an international, peer-reviewed journal of clinical therapeutics and pharmacology focusing on concise rapid reporting of clinical or pre-clinical studies on a range of neuropsychiatric and neurological disorders. This journal is indexed on PubMed Central, the 'PsycINFO' database and CAS, and

is the official journal of The International Neuropsychiatric Association (INA). The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/neuropsychiatric-disease-and-treatment-journal>