

LINE-1 methylation status of endogenous DNA double-strand breaks

Wichai Pornthanakasem¹, Narisorn Kongruttanachok², Chutipha Phuangphairoj¹, Chotika Suyarnsestakorn^{2,3}, Taweap Sanghangthum⁴, Sornjarod Oonsiri⁴, Wanpen Ponyeam¹, Thatchawan Thanasupawat¹, Oranart Matangkasombut⁵ and Apiwat Mutirangura^{1,*}

¹Center of Excellence in Molecular Genetics of Cancer and Human Diseases, Department of Anatomy, Faculty of Medicine, ²Inter-Department Program of Biomedical Sciences, Faculty of Graduate School, Chulalongkorn University, Bangkok 10330, ³The National Center for Genetic Engineering and Biotechnology, Pathumthani 12120, ⁴Department of Radiology, Division of Radiation Oncology, Faculty of Medicine and ⁵Department of Microbiology, Faculty of Dentistry, Chulalongkorn University, Bangkok 10330, Thailand

Received March 18, 2008; Revised and Accepted April 17, 2008

ABSTRACT

DNA methylation and the repair of DNA double-strand breaks (DSBs) are important processes for maintaining genomic integrity. Although DSBs can be produced by numerous agents, they also occur spontaneously as endogenous DSBs (EDSBs). In this study, we evaluated the methylation status of EDSBs to determine if there is a connection between DNA methylation and EDSBs. We utilized interspersed repetitive sequence polymerase chain reaction (PCR), ligation-mediated PCR and combined bisulfite restriction analysis to examine the extent of EDSBs and methylation at long interspersed nuclear element-1 (LINE-1) sequences nearby EDSBs. We tested normal white blood cells and several cell lines derived from epithelial cancers and leukemias. Significant levels of EDSBs were detectable in all cell types. EDSBs were also found in both replicating and non-replicating cells. We found that EDSBs contain higher levels of methylation than the cellular genome. This hypermethylation is replication independent and the methylation was present in the genome at the location prior to the DNA DSB. The differences in methylation levels between EDSBs and the rest of the genome suggests that EDSBs are differentially processed, by production, end-modification, or repair, depending on the DNA methylation status.

INTRODUCTION

Vilenchik and Knudson (1) estimated the existence of endogenous double-strand breaks (EDSBs) and suggested

that EDSBs could account for a substantial fraction of oncogenic events in human carcinomas. If EDSBs do not arise uniformly or are not processed at equal rates across the genome, mutation hot spots should be present (1). Our study helps to elucidate if DNA methylation influences EDSBs processing.

Several pieces of evidence suggest that DNA methylation may play an important role in maintaining genomic integrity. Genome-wide decreases in DNA methylation levels commonly occur in cancer (2–5), which leads to higher rates of mutations and genomic instability (6–8). In addition to alterations in the number of chromosomes, hypomethylation can result in chromosomal rearrangements and deletion of DNA, suggesting that DSBs are the intermediate products (6–8). Moreover, because the mutations occur spontaneously, the DSBs should occur endogenously. Studies in ICF syndrome (immunodeficiency, chromosomal instability and facial anomalies) (9), which is characterized by loss-of-function mutations in the cytosine DNA methyltransferase *DNMT3B*, and Wilm's tumor (10) demonstrated a direct association between loss of DNA methylation and rearrangements in the pericentromeric heterochromatin. Therefore, hypomethylation could lead to spontaneous mutations in *cis*, which are the epigenetic and genetic events occurring on the same chromosome. Consequently, evaluating methylation status of EDSBs may provide clues to better understanding how DNA methylation helps maintaining genomic integrity.

To determine whether DNA methylation affects EDSBs, we first developed a set of novel techniques to analyze the extent of methylation in genomic EDSBs. These techniques were devised from interspersed repetitive sequence (IRS) polymerase chain reaction (PCR) (11), ligation-mediated (LM) PCR-based assays (12–14) and combined bisulfite restriction analysis (COBRA) for

*To whom correspondence should be addressed. Tel: +662 256 4532; Fax: +662 254 1931; Email: apiwat.mutirangura@gmail.com

genome-wide methylation level analysis (4,5,15). LMPCR is a commonly used PCR technique designed to analyze locus-specific EDSBs during lymphoid development, such as V(D)J recombination (14,16,17) and hypermutation (18). LMPCR has also been used to detect DNA-associated proteins or chromatin accessibility to such proteins (19,20), and this technique has shown previous utility in genome mapping research (21,22). With these new assays, we evaluated the relationship between DNA methylation and EDSBs.

MATERIALS AND METHODS

Cells, cell lines and culture

Cell lines used were HeLa (cervical cancer), SW480 (colorectal adenocarcinoma), K562 (erythroleukemia), Daudi (B lymphoblast), Jurkat (T-cell leukemia) and Molt4 (T lymphoblast) (ATCC, Manassas, VA, USA). HeLa cells were synchronized at G0 phase by culture in serum deprivation medium, Dulbecco's modified Eagle's medium plus 0.2% fetal bovine serum, for 48 h. HeLa cells in G1/S and S phase cells were synchronized by the thymidine block method, and were cultured with 2 mM thymidine (Sigma-Aldrich, St Louis, MO, USA) to obtain cells at G1/S phase (23). Flow cytometry was used to determine stages of the cell cycle as well as fragmented and apoptotic cells. One millimolar of H₂O₂ was added for 24 h to induce apoptosis. For radiation treatment, the medium of G0 phase HeLa cells was replaced with 15 ml of ice-cold medium, and cells were exposed to γ - with a ⁶⁰Co source (Eldorado78).

High-molecular weight DNA preparation

To prepare high-molecular weight (HMW) DNA, 5×10^5 cells were embedded in 1% low-melting point agarose, lysed and digested in 400 μ l of 1 mg/ml proteinase K, 50 mM Tris, pH 8.0, 20 mM EDTA, 1% sodium lauryl sarcosine. The plugs were rinsed four times in Tris-EDTA (TE) buffer for 20 min. To polish overhang or cohesive-end EDSBs, T4 DNA polymerase (New England Biolabs, Beverly, MA, USA) and dNTPs were added and later inactivated by adding EDTA to a concentration of 20 mM for 5 min followed by rinsing four times in TE buffer for 20 min. To analyze blunt-end EDSBs, LMPCR was performed without T4 DNA polymerase. The modified LMPCR linkers (24) were prepared from the oligonucleotides

5'-AGGTAACGAGTCAGACCACCGATCGCTCGGAAGCTTACCTCGTGGACGT-3' and 5'-ACGTCCACGAG-3'. The linkers (50 pmol) were ligated to HMW DNA using T4 DNA ligase (New England Biolabs) at 25°C overnight. DNA was extracted from agarose plugs using a QIAquick gel extraction kit (Qiagen, Basel, Switzerland). For liquid DNA preparation, cells or HMW DNA were incubated in 1% sodium dodecyl sulfate/proteinase K (0.5 mg/ml), at 48°C overnight and subjected to phenol-chloroform extraction and ethanol precipitation. The precipitated DNA was resuspended in 20 μ l of TE buffer.

IRS-EDSB-LMPCR

The quantity of IRS-EDSB was measured by real-time PCR using a LightcyclerTM instrument (Roche Applied Science, Indianapolis, IN, USA) with the IRS primers, including long interspersed nuclear elements-1 (LINE-1s or L1s) primers 5'-CTCCCAGCGTGAGCGAC-3' (outward), 5'-AAGCCGGTCTGAAAAGCGCAA-3' (inward), Alu, Alu-CL2 5'-ACTGCACTCCAGCCTGGGC-3' or Tigger1 5'-CTCGCTGAAGGCTCAGATGATC-3'), the linker primer 5'-AGGTAACGAGTCAGACCACCGA-3' (24), and the Taqman probe homologous to the 3' linker sequence (6-fam) ACGTCCACGAGGT AAGCT TCCGAGCGA (tamra) (phosphate). Amplification was performed with 0.5 μ M of each primer, 0.4 μ M Taqman probe, 2 U of HotStarTaq (Qiagen, Valencia, CA, USA), 1 \times PCR buffer and 10 ng of ligated DNA for up to 40 cycles, with quantification after the extension steps. Two types of control DNA were used. The first was a 100-bp oligonucleotide sequence with the 5' linker sequence and 3' homology to L1 oligonucleotide sequences. The second was DNA digested with EcoRV and AluI and ligated to the LMPCR linkers. The amounts of EDSBs were compared with the ligated control digested DNA and reported as L1-EDSB-LMPCR templates per nanogram of DNA.

COBRA-IRS and COBRA-IRS-EDSB

Ligated HMW DNA was modified with bisulfite using a standard protocol (25). Bisulfite-modified DNA was recovered using a Wizard DNA clean-up kit (Promega, Madison, WI, USA) and desulfonated before PCR amplification. For PCR COBRA (15) of L1s (COBRA-L1) (4,5), bisulfite-treated DNA was subjected to 35 cycles of PCR with two primers, B-L1-inward 5'-CGTAAGGGTTAGGGAGTTTTT-3' and B-L1-outward 5'-RTAA AACCCTCCRAACCAATATAAAA-3' (4). Applying a hot-stop technique to prevent heteroduplex amplicons, α^{32} P-labeled-bisulfite-L1-outward oligo was added in the last PCR cycle. The amplicons were doubly digested in a 10 μ l reaction volume with 2 U of TaqI and 8 U of TasI in 1 \times TaqI buffer (MBI Fermentas, Vilnius, Lithuania) at 65°C for 4 h. The PCR is designed to detect unmethylated and methylated L1 sequences of 98 and 80 bp, respectively. The intensity of DNA fragments was measured with a PhosphorImager using Image Quant software (Molecular Dynamics, GE Healthcare, Slough, UK). The LINE-1 methylation level was calculated as the percentage of TaqI intensity divided by the sum of TaqI- and TasI-positive amplicons. For COBRA-L1-EDSB, the B-L1-inward oligo was replaced with B-LMPCR oligo, 5'-GTTTGGAAAGTTTATTTTGTGGAT-3', and 40 PCR cycles were carried out according to the same protocol. Bisulfite-treated Daudi, Jurkat and HeLa DNAs digested with EcoRV and AluI and ligated LMPCR linker were used as positive controls to normalize the inter-assay variation of all COBRA experiments. HeLa DNA without ligation was used as a negative control.

Southern blot hybridization

Southern blot was performed to compare between 5 μ g of HpaII- and MspI-digested HeLa DNA. L1-LMPCR amplicons from HMW and AluI-EcoRV-digested HeLa DNA were used as probes. L1-most-outward primer sequence was 5'-TATTCGGCCATCTTGGCTCCT-3'. Competitor DNA, COT-1 DNA, was used to prevent non-specific DNA hybridization, including sequence from L1s. Intensities in the >4 kb regions were measured with a PhosphorImager using Image Quant software (Molecular Dynamics). Semiquantitative methylation percentage was reported as the proportion of probes bound to HpaII-digested DNA to HpaII plus MspI-digested DNA.

Statistical analyses

Statistical significance was determined according to an independent sample *t*-test, a paired sample *t*-test or ANOVA using the SPSS program version 11.5 as specified.

RESULTS

Detection of genomic EDSB and methylation

First, we developed a new assay for the detection of EDSBs. This assay is based on a LMPCR, originally designed for the analysis of locus and cell-specific EDSBs (14,16–18). General EDSBs are believed to occur rarely and arbitrarily throughout the genome (1). Using repetitive sequences that randomly scatter throughout the human genome, we can detect genome-wide EDSBs in proximity to these repetitive sequences. We, therefore, combined LMPCR with IRSPCR or inter-Alu PCR (11) using the widely distributed L1s human retrotransposons (26) into a new assay called 'L1-EDSB-LMPCR'. In this assay, linker oligonucleotides are ligated to EDSBs in HMW DNA preparation and quantitatively analyzed by real-time PCR using an L1 primer and a Taqman probe complementary to the linker (Figure 1A). L1-PCR using a primer in the outward direction will amplify DNA sequences located outside the repetitive sequences.




Figure 1. L1-EDSB-LMPCR and COBRA-L1-EDSB. (A) Schematic representation of L1-EDSB-LMPCR showing L1 sequence and the ligated linker at a nearby EDSB. The white rectangle is a Taqman probe complementary to the LMPCR linker. Arrows are PCR primers. (B) The quantity of EDSBs detected with this method increased directly with the amount of experimentally induced DSBs, and significant amounts of EDSBs were detected in all tested cells but not in cells without ligation. An example of results of real-time L1-EDSB-LMPCR with tests and controls was demonstrated. The values 2, 10, 50, 250 and 1250 are quantities of restriction enzyme-digested (EcoRV and AluI) HeLa genomes ligated with LMPCR linker. dH₂O is water. Tested templates are HMW DNA from (1) HeLa without ligation, (2) Daudi blunt-end-EDSBs ligation, (3) Daudi polished-end-EDSBs ligation, (4) Jurkat blunt-end-EDSBs ligation and (5) Jurkat polished-end-EDSBs ligation. (C) Schematic representation of COBRA-L1 and COBRA-L1-EDSB, showing L1 sequence ligated by linker at an EDSB. Arrows are PCR primers, with asterisk indicating α -³²P-labeled primer for COBRA. AACCG and CCGA are L1 sequences; when treated with bisulfite and after undergoing PCR, unmethylated AACCG will be converted to AATTG (TasI site) and methylated CCGA to TCGA (TaqI site). (D) A typical example of results from COBRA-L1 and COBRA-L1-EDSB experiments indicating that the intensity between methylated, TaqI and unmethylated, TasI, bands of EDSBs were higher than the matched pair genomes. The arrow at 98 bp indicates TasI-digested unmethylated L1 sequences and the arrow at 80 bp indicates TaqI-digested methylated L1 sequences. -ve is dH₂O for COBRA-L1 and non-ligated HMW DNA for COBRA-L1-EDSB. TasI and TaqI are restriction enzymes added in each experiment.

Therefore, the sequences yielded from IRSPCR will represent human genome-wide sequences, including both unique and repetitive sequences (11,21). Increases in the L1-EDSB-LMPCR products corresponded with the amount of control DSBs generated by restriction enzyme digesting DNA. Without ligation, no L1-EDSB-LMPCR could be detected. Finally, significant amounts of EDSBs were detected in all tested cells (Figure 1B).

Next, we developed an assay to analyze the methylation level of EDSBs. Previously, we had extensively studied methylation status of L1s in several cancers and normal tissues by COBRA-L1. Treatment with bisulfite converts unmethylated cytosines, but not methylated cytosines, to uracils and then thymines after PCR. Therefore, this bisulfite treatment generates detectable methylation-dependent changes in the restriction pattern of PCR-amplified L1 sequences. Methylation level is then calculated and presented as a percentage of total DNA. We thus combined L1-EDSB-LMPCR with COBRA-L1 by treating linker-ligated DNA with bisulfite before PCR with L1/linker primers and restriction analysis (Figure 1C). With this new 'COBRA-L1-EDSB' assay, we can measure the methylation level of L1s near EDSBs, which reflects the methylation level of EDSBs in a genome-wide fashion. The degree of methylation between genomic L1 and L1-EDSB sequences was examined by COBRA-L1 and COBRA-L1-EDSB, respectively (Figure 1D).

With these new assays, we first evaluated what types of cells possess EDSBs and if the quantity of EDSBs reflects carcinogenic potentials. Using L1-EDSB-LMPCR, significant amounts of EDSBs were detected in all samples from several cancer cell lines, including Daudi, Jurkat, Molt4, K562, SW480 and HeLa cells, as well as in normal cells, including sperm and white blood cells (WBCs) from several individuals (Figure 2A). These data suggest that EDSBs can commonly be found in all cells both normal and cancer.

This assay prepared HMW DNA by in-gel preparation. This technique has been reported to decrease the number of DSBs generated during DNA preparation (27) and did not generate false positives in the LMPCR assay for the analysis of locus-specific EDSBs (14,18,27). To evaluate if in-gel HMW DNA preparation breaks genomic DNA, we compared the yield of the LMPCR from several sources of DNA, including, in-gel HMW DNA, liquid DNA, liquid DNA extracted from in-gel HMW DNA (gel-liquid) and in-gel DNA prepared from liquid DNA (liquid-gel). The quantities of L1-EDSB-LMPCR from liquid DNA were higher than in-gel HMW DNA (Figure 2B). Extracting liquid DNA from a gel increased the amount of LMPCR template over HMW DNA (Figure 2B). This result indicates that more DNA breaks were generated during the liquid DNA preparation process. Embedded liquid DNA into gel following by HMW DNA preparation protocol did not increase the amount of LMPCR template over liquid DNA alone (Figure 2B). Therefore, in-gel preparation did not generate significant additional DNA breaks. Moreover, the quantities of L1-EDSB-LMPCR from gel-liquid and liquid-gel were clearly lower than when L1-EDSB-LMPCR




Figure 2. Discovery and specificity of EDSB-LMPCR. (A) Using L1-EDSB-LMPCR, significant amounts of EDSBs were detected in all samples, including, cancer cell lines, sperm cells and WBCs from several individuals. (B) L1-EDSB-LMPCR of DNA from HeLa cells using different DNA-extraction methods. Gel, liquid, gel-liquid, liquid-gel and combined are L1-EDSB-LMPCR from DNA prepared in gel, liquid, extracted liquid DNA from gel-embedded HMW DNA, embedded liquid DNA into gel following by HMW DNA preparation protocol and 2 × admixture between 1:1 of HMW DNA and liquid DNA, respectively. Data represent means \pm SEM.

products from HMW DNA and liquid DNA were combined (Combined) (Figure 2B). Therefore, the significant amounts of LMPCR products from HMW DNA were not derived from DNA preparation. This experiment supported the presence of EDSBs.

General characteristics of EDSB-LMPCR

Repetitive sequences have been commonly recognized as locations of general recombination that destabilize human cancer genomes (28,29). It is interesting to investigate if the quantities of EDSBs differ between repetitive and unique sequences. Full length L1 is ~6 kb (26). L1-EDSB-LMPCR using the L1-inward primer will amplify L1 sequences, while the L1-outward primer should amplify unique sequences near L1 (11). We observed L1-EDSBs at similar levels regardless of the directions of L1 primers used (Figure 3A). The data indicate that the amount of EDSBs does not differ based on the nature of DNA sequences, between L1 repetitive and unique sequences. We found that IRS-LMPCR using primers for other types of repetitive sequences, including Alu and Tigger1, also yielded significant EDSBs in direct proportions to their copy numbers in the human genomes (30) (Figure 3B).




Figure 3. General characteristics and distribution of EDSB-LMPCR in the human genome. (A) L1-EDSB-LMPCR, using L1-inward and L1-outward primers, was performed in several cell types, and products were observed at similar levels regardless of the nature of linked EDSB sequences, L1, or unique sequences. Inward L1-EDSB-LMPCR was normalized by the proportion of L1 sequence copies in the human genome (www.ncbi.nlm.nih.gov). (B) Comparison of the amount of EDSB-LMPCRs using L1, Alu and Tigger1 primers in direct proportions to their copy numbers in the human genomes. (C) EDSBs were measured from HMW DNA with and without T4 polymerase treatment, respectively. Significant amount of L1-EDSB-LMPCRs of blunt and polished-end EDSBs from cancer cell lines were identified. Therefore, L1-EDSB ends are heterogeneous. (D) The quantity of EDSBs was not related to the proportion of fragmented cells. L1-EDSB-LMPCR quantity related to the percent of fragmented cells, documented by flow cytometry. Daudi DNase represents Daudi cells treated with DNase I before HMW DNA preparation.

L1s are AT rich, whereas Alu sequences are frequently located in CG-rich regions (31). Therefore, EDSBs are widely distributed in the human genome. In addition, because EDSBs are thought to occur rarely and randomly throughout the genome, LMPCR with primers specific to unique sequences yielded no positive locus-specific amplicons (data not shown). EDSB ends are heterogeneous, as significant quantities of L1-EDSB-LMPCR products were obtained, including two EDSB types: blunt- and polished-, blunt plus cohesive, end EDSBs (Figure 3C). Blunt-end DSBs are DNA that both strands terminate in a base pair. A cohesive or overhang is a stretch of unpaired nucleotides in the DNA end (14,16,17).

The quantity of EDSBs was not related to the proportion of fragmented cells (Figure 3D). Our EDSB-LMPCR protocol, particularly that for epithelial cells, minimizes contamination with apoptotic cells because these dying cells with fragmented DNA (32) usually lose attachment (33) and are thus washed off before cell collection for EDSB analysis. While there was positive L1-EDSB-LMPCR amplification, we were unable to detect any

fragmented DNA or apoptotic cells, as determined by LMPCR ladder (34) and flow cytometry (35), respectively (data not shown). Moreover, the apoptotic genome possesses normal levels of methylation and may not be detectable by COBRA-L1-EDSB. Fragmented DNA, as documented by electrophoresis, was collected from floating apoptotic HeLa cells after treatment with 1 mM H₂O₂ for 24 h (36). COBRA-L1 shows that apoptotic HMW DNA has similar L1 methylation to living cells, but COBRA-L1-EDSB yielded amplicons with multiple sizes that interfere with interpretation (data not shown).

L1-EDSB-LMPCR under different conditions

We utilized L1-EDSB-LMPCR on cells under different conditions known to associate with DSBs. Specifically, we used radiation and cell cycle synchronization. Radiation, which directly causes DNA damage, increased L1-EDSB-LMPCR levels directly correlating with the dosages of radiation used to treat the cells (Figure 4A). Nevertheless, there were wide ranges of the amount of




Figure 4. L1-EDSB-LMPCR under different conditions. L1-EDSB-LMPCR of (A) irradiated HeLa. (B) HeLa cells at G0, G1/S and 0, 3 and 5 h after the release into S phase from thymidine block. Control is without cell synchronization. G0 bore the least amount of EDSBs when compared with control, * $P < 0.05$ (independent one-tailed t -test). Data represent means \pm SEM.

EDSBs (Figure 4A). This result may be the consequence of several factors, including DSB end modifications and DSB repair rate. Therefore, even though L1-EDSB-LMPCR can detect radiation-induced DSBs, the technique lacks efficiency in evaluating the precise number of DSBs generated by radiation.

Because EDSBs were hypothesized to be preferentially produced in S phase from the conversion of single-strand lesions (1), we used HeLa cells to assess the frequency of EDSBs and their methylation status during various cell cycle stages: G0, G1/S and S. We found the impact of cell cycle effect is on the borderline of significance and EDSBs can be found in G0 phase (Figure 4B). Interestingly, there was a minor decrease in S phase cells (Figure 4B). Therefore, cell cycle stage is one of the conditions that may alter the amount of EDSBs. Nevertheless, similar to radiation the amount of EDSBs during cell phases may be influenced by several factors in addition to the production rate.

Specificity of COBRA-L1-EDSB

Figure 5A demonstrates a reconstitution experiment of EDSB-LMPCR. Because Daudi cells shows significantly higher level of genomic methylation than HeLa cells, we used DNA prepared from Daudi and HeLa cells as representative DNA with high and low methylation, respectively. We added varying ratios of HeLa and Daudi DNA digested with EcoRV and AluI (ligated to LMPCR linker) into HeLa genomic DNA. As expected, a COBRA-L1-EDSB analysis of the digested DNA, regardless of the presence of the HeLa genome, yielded equivalent levels of methylation to COBRA-L1. Furthermore, the methylation levels in these samples measured by both COBRA-L1 and COBRA-L1-EDSB are increased, which correlates with the proportion of highly methylated Daudi DNA contained in that sample (Figure 5A). The higher levels of methylation, compared to the rest of the genome, are a characteristic of EDSBs but not DSBs randomly generated by DNA shearing during sample preparation. Lower DSB methylation levels of liquid DNA, gel-liquid DNA and liquid-gel DNA were demonstrated when compared to HMW DNA (Figure 5B). In contrast to gDNA




Figure 5. Specificity of COBRA-L1-EDSB. (A) A reconstitution experiment of COBRA-L1-EDSB. COBRA-L1 and COBRA-L1-EDSB of the mixture between HeLa and Daudi DNA digested with AluI and EcoRV and ligated to EDSB-LMPCR linker. The 4:0, 3:1, 2:2, 1:3 and 0:4 are proportions of recombinant HeLa:Daudi DNA. COBRA-L1-EDSB 1/10 was the mixture between 2 ng of linker-ligated DNA and 18 ng of HeLa genomic DNA. Because Daudi cells shows significantly higher levels of genomic methylation compared to HeLa cells, we used DNA prepared from Daudi and HeLa cells as representative DNA with high methylation and low methylation, respectively. We added varying ratio of HeLa and Daudi DNA digested with EcoRV and AluI ligated to the LMPCR linker into HeLa genomic DNA. As expected, the methylation levels of the digested DNA, regardless of the presence of the HeLa genome, were measured by COBRA-L1-EDSB and COBRA-L1, and increases in final product correlated with the proportion of highly methylated Daudi DNA. (B) COBRA-L1 and COBRA-L1-EDSB methylation levels in a comparison of DNA preparation methods. Gel, gel-liquid, liquid and liquid-gel refer to DNA analyzed via L1-EDSB-LMPCR from DNA prepared in-gel, extracted liquid DNA from gel embedded HMW DNA, liquid and embedded liquid DNA into gel following by HMW DNA preparation protocol, respectively. COBRA-L1-EDSB methylation levels of HMW DNA were higher than liquid DNA. COBRA-L1-EDSB methylation levels of in-gel DNA prepared from liquid DNA, and liquid DNA prepared from HMW DNA, were lower than HMW but not different from liquid. Therefore, HMW DNA preparation does not cause hypermethylated DSBs. Data represent means \pm SEM.

prepared from cells cast into low melting point agarose plugs, the preparation protocol for liquid DNA generates DSBs (Figure 2B). Therefore, COBRA-L1-EDSB levels of liquid DNA should be derived from combination between EDSBs and DNA preparation producing DSBs. The lower DSB methylation levels of liquid DNA confirmed that the DNA preparation producing DSBs possess lower methylation levels than EDSBs. Moreover, DSB methylation levels of gel-liquid DNA and liquid-gel DNA are not different from levels of liquid DNA alone (Figure 5B). Therefore, higher methylation levels of COBRA-L1-EDSB, detected from HMW DNA, were unlikely to be derived from HMW DNA preparation.

EDSBs are ubiquitously hypermethylated

To compare the degree of methylation between L1 and L1-EDSB sequences, we examined by COBRA-L1 and

demonstrates that EDSBs are not only detectable but also widely distributed. Positive results were obtained from all types of selected IRS sequences. Nevertheless, LMPCR may not be able to detect some subset of EDSBs. LMPCR preparation requires ligation of a blunt-end DNA linker to a blunt and phosphorylated DSB end. In this study, we polished cohesive-end EDSBs by T4 polymerase. We found that both blunt and overhang ends are present and that the majority of EDSB ends are blunt. LMPCR fails, however, to detect some complex ends that cannot be polished such as a hairpin loop. Unlike studying V(D)J recombination (17,27), in this study, we did not apply Mung Bean nuclease to screen for these end-types because the enzyme can convert single-strand lesion into DSBs.

IRS-EDSB-LMPCR is the first method characterizing EDSBs. Consequently, there is no report of cells with different levels of EDSBs. Therefore, even though it is commonly accepted that in-gel HMW DNA preparation does not significantly affect DNA breaks (27), we performed additional experiments to disregard the possibility of error. The comparison among liquid DNA preparation, HMW DNA and combined experiments suggested that there are no detectable DSBs generated by the HMW DNA preparation process. Finally, cells treated under different conditions, including, serum deprivation and temperature (data not shown), altered EDSB levels. Because the differentially treated cells were processed for HMW DNA simultaneously, the possibility that the same DNA preparation method led to bias shearing of the DNA depending on different prior cellular conditions is remote.

LMPCR detects DSBs directly, while γ -H2AX foci staining aims to detect a cellular response to DSBs. Therefore, L1-EDSB-LMPCR analysis and staining for γ -H2AX foci may not yield the same information. For example, a recent report demonstrated that some genomic regions, particularly heterochromatin, are devoid of radiation-induced γ -H2AX foci (39). Consequently, L1-EDSB-LMPCR and γ -H2AX foci should assess radiation-induced DSBs differently depending on chromatin structures. In this study, we have demonstrated that, even though L1-EDSB-LMPCR products increased after radiation, the correlation to radiation dosage was not as precise as using γ -H2AX foci for detection (40). One explanation for the broad range of results when using L1-EDSB-LMPCR to detect radiation-induced DSBs could be that there is a wide variety of biological processes involved in radiation-induced DSB repairs (41,42).

This study demonstrates that EDSBs are present in all cell types. Moreover, EDSBs normally possess higher levels of methylation compared to the cellular genome, and this methylation pre-exists at the break sites. These findings were not only unprecedented but also not generally expected. Vilenchik and Knudson suggested that the causes of EDSBs are oncogenic events in human carcinomas (1). Moreover, genomic instability can be observed while cancer genomes are hypomethylated (6–8). Therefore, it is tempting to hypothesize that EDSBs should occur more frequently at hypomethylated sequences and, consequently, that COBRA-L1-EDSB should have been hypomethylated. Nonetheless, IRS-EDSB-LMPCR and COBRA-L1-EDSB are methods to measure the extent of

methylation at EDSBs during a given time. Therefore, the LMPCR levels reflect not only the rate of EDSB synthesis but also EDSB end modification and repair. Therefore, discovery of DSBs does not exclusively indicate DSB formation, but, the EDSB hypermethylation finding leads us to conclude that methylated and unmethylated DNA possess distinctive EDSB processing (i.e. repair) pathways. If these pathways have different precision, methylated and unmethylated DNA should have unequal rate of spontaneous mutations. In the future, it would be interesting to explore if and how cells process or repair EDSBs depending on DNA methylation. This finding may yield an important clue to prevent global hypomethylation-induced chromosomal rearrangements.

There may be several other mechanisms by which DNA methylation prevents IRS from inducing DNA rearrangements. Repetitive sequences have been commonly recognized as locations of general recombination that destabilizes human cancer genomes, particularly when genetic recombination occurred between different loci (28,29,43). A deletion at *Xist* alters chromatin conformation, usually associated with DNA methylation, of the inactive X chromosome and this change destabilizes both X chromosomes (44). In the future, it would be intriguing to explore if EDSBs play a role in these mechanisms and whether or not DNA methylation is important in preventing these chromosomal rearrangements.

ACKNOWLEDGEMENTS

We thank Profs Alfred G Knudson, Fox Chase Cancer Center and Stephen Buratowski, Harvard Medical School and Drs Vitrote Sriuranpong, Kanya Suphapeetiporn, Chulalongkorn University and Man Liu, Yale University for critical review and advice. This study was supported by the Thailand Research Fund and Faculty of Graduate School. N.K. and W.P. were supported by the Royal Golden Jubilee Ph.D. grant. Funding to pay the Open Access publication charges for this article was provided by the Thailand Research Fund.

Conflict of interest statement. None declared.

REFERENCES

- Vilenchik, M.M. and Knudson, A.G. (2003) Endogenous DNA double-strand breaks: production, fidelity of repair, and induction of cancer. *Proc. Natl Acad. Sci. USA*, **100**, 12871–12876.
- Feinberg, A.P. and Vogelstein, B. (1983) Hypomethylation distinguishes genes of some human cancers from their normal counterparts. *Nature*, **301**, 89–92.
- Feinberg, A.P. and Tycko, B. (2004) The history of cancer epigenetics. *Nat. Rev. Cancer*, **4**, 143–153.
- Chalitchagorn, K., Shuangshoti, S., Hourpai, N., Kongruttanachok, N., Tangkijvanich, P., Thong-ngam, D., Voravud, N., Sriuranpong, V. and Mutirangura, A. (2004) Distinctive pattern of LINE-1 methylation level in normal tissues and the association with carcinogenesis. *Oncogene*, **23**, 8841–8846.
- Mutirangura, A. (2007) Quantitative PCR analysis for methylation level of genome: clinical implications in cancer. *Asian Biomedicine*, **1**, 121–128.
- Lengauer, C., Kinzler, K.W. and Vogelstein, B. (1997) DNA methylation and genetic instability in colorectal cancer cells. *Proc. Natl Acad. Sci. USA*, **94**, 2545–2550.

7. Eden, A., Gaudet, F., Waghmare, A. and Jaenisch, R. (2003) Chromosomal instability and tumors promoted by DNA hypomethylation. *Science*, **300**, 455.
8. Chen, R.Z., Pettersson, U., Beard, C., Jackson-Grusby, L. and Jaenisch, R. (1998) DNA hypomethylation leads to elevated mutation rates. *Nature*, **395**, 89–93.
9. Xu, G.L., Bestor, T.H., Bourc'his, D., Hsieh, C.L., Tommerup, N., Bugge, M., Hulten, M., Qu, X., Russo, J.J. and Viegas-Pequignot, E. (1999) Chromosome instability and immunodeficiency syndrome caused by mutations in a DNA methyltransferase gene. *Nature*, **402**, 187–191.
10. Qu, G.Z., Grundy, P.E., Narayan, A. and Ehrlich, M. (1999) Frequent hypomethylation in Wilms tumors of pericentromeric DNA in chromosomes 1 and 16. *Cancer Genet. Cytogenet.*, **109**, 34–39.
11. Nelson, D.L., Ledbetter, S.A., Corbo, L., Victoria, M.F., Ramirez-Solis, R., Webster, T.D., Ledbetter, D.H. and Caskey, C.T. (1989) Alu polymerase chain reaction: a method for rapid isolation of human-specific sequences from complex DNA sources. *Proc. Natl Acad. Sci. USA*, **86**, 6686–6690.
12. Mueller, P.R. and Wold, B. (1989) In vivo footprinting of a muscle specific enhancer by ligation mediated PCR. *Science*, **246**, 780–786.
13. Pfeifer, G.P., Steigerwald, S.D., Mueller, P.R., Wold, B. and Riggs, A.D. (1989) Genomic sequencing and methylation analysis by ligation mediated PCR. *Science*, **246**, 810–813.
14. Schlissel, M., Constantinescu, A., Morrow, T., Baxter, M. and Peng, A. (1993) Double-strand signal sequence breaks in V(D)J recombination are blunt, 5'-phosphorylated, RAG-dependent, and cell cycle regulated. *Genes Dev.*, **7**, 2520–2532.
15. Xiong, Z. and Laird, P.W. (1997) COBRA: a sensitive and quantitative DNA methylation assay. *Nucleic Acids Res.*, **25**, 2532–2534.
16. Zhu, C. and Roth, D.B. (1995) Characterization of coding ends in thymocytes of scid mice: implications for the mechanism of V(D)J recombination. *Immunity*, **2**, 101–112.
17. Livak, F. and Schatz, D.G. (1997) Identification of V(D)J recombination coding end intermediates in normal thymocytes. *J. Mol. Biol.*, **267**, 1–9.
18. Papavasiliou, F.N. and Schatz, D.G. (2000) Cell-cycle-regulated DNA double-stranded breaks in somatic hypermutation of immunoglobulin genes. *Nature*, **408**, 216–221.
19. Pfeifer, G.P. and Tommasi, S. (2000) In vivo footprinting using UV light and ligation-mediated PCR. *Methods Mol. Biol.*, **130**, 13–27.
20. Espinoza, C.R. and Feeney, A.J. (2006) Quantifying chromatin accessibility of individual gene family members by combining ligation-mediated PCR with real-time PCR. *Biotechniques*, **41**, 404, 406, 408.
21. Mutirangura, A., Jayakumar, A., Sutcliffe, J.S., Nakao, M., McKinney, M.J., Buiting, K., Horsthemke, B., Beaudet, A.L., Chinault, A.C. and Ledbetter, D.H. (1993) A complete YAC contig of the Prader-Willi/Angelman chromosome region (15q11-q13) and refined localization of the SNRPN gene. *Genomics*, **18**, 546–552.
22. Mutirangura, A., Greenberg, F., Butler, M.G., Malcolm, S., Nicholls, R.D., Chakravarti, A. and Ledbetter, D.H. (1993) Multiplex PCR of three dinucleotide repeats in the Prader-Willi/Angelman critical region (15q11-q13): molecular diagnosis and mechanism of uniparental disomy. *Hum. Mol. Genet.*, **2**, 143–151.
23. Bostock, C.J., Prescott, D.M. and Kirkpatrick, J.B. (1971) An evaluation of the double thymidine block for synchronizing mammalian cells at the G1-S border. *Exp. Cell Res.*, **68**, 163–168.
24. Pornthanakasem, W. and Mutirangura, A. (2004) LINE-1 insertion dimorphisms identification by PCR. *Biotechniques*, **37**, 750, 752.
25. Cameron, E.E., Bachman, K.E., Myohanen, S., Herman, J.G. and Baylin, S.B. (1999) Synergy of demethylation and histone deacetylase inhibition in the re-expression of genes silenced in cancer. *Nat. Genet.*, **21**, 103–107.
26. Kazazian, H.H. Jr and Moran, J.V. (1998) The impact of L1 retrotransposons on the human genome. *Nat. Genet.*, **19**, 19–24.
27. Schlissel, M.S. (1998) Structure of nonhairpin coding-end DNA breaks in cells undergoing V(D)J recombination. *Mol. Cell Biol.*, **18**, 2029–2037.
28. Kidwell, M.G. and Holyoake, A.J. (2001) Transposon-induced hotspots for genomic instability. *Genome Res.*, **11**, 1321–1322.
29. Surtees, J.A., Argueso, J.L. and Alani, E. (2004) Mismatch repair proteins: key regulators of genetic recombination. *Cytogenet. Genome Res.*, **107**, 146–159.
30. Smit, A.F. (1999) Interspersed repeats and other mementos of transposable elements in mammalian genomes. *Curr. Opin. Genet. Dev.*, **9**, 657–663.
31. Lander, E.S., Linton, L.M., Birren, B., Nusbaum, C., Zody, M.C., Baldwin, J., Devon, K., Dewar, K., Doyle, M., FitzHugh, W. et al. (2001) Initial sequencing and analysis of the human genome. *Nature*, **409**, 860–921.
32. Inohara, N., Koseki, T., Chen, S., Benedict, M.A. and Nunez, G. (1999) Identification of regulatory and catalytic domains in the apoptosis nuclease DFF40/CAD. *J. Biol. Chem.*, **274**, 270–274.
33. Hengartner, M.O. (2000) The biochemistry of apoptosis. *Nature*, **407**, 770–776.
34. Staley, K., Blaschke, A.J. and Chun, J. (1997) Apoptotic DNA fragmentation is detected by a semi-quantitative ligation-mediated PCR of blunt DNA ends. *Cell Death Differ.*, **4**, 66–75.
35. Sherwood, S.W. and Schimke, R.T. (1995) Cell cycle analysis of apoptosis using flow cytometry. *Methods Cell Biol.*, **46**, 77–97.
36. Ren, J.G., Xia, H.L., Just, T. and Dai, Y.R. (2001) Hydroxyl radical-induced apoptosis in human tumor cells is associated with telomere shortening but not telomerase inhibition and caspase activation. *FEBS Lett.*, **488**, 123–132.
37. Olive, P.L., Banath, J.P. and Durand, R.E. (1990) Heterogeneity in radiation-induced DNA damage and repair in tumor and normal cells measured using the "comet" assay. *Radiat. Res.*, **122**, 86–94.
38. Stamato, T.D. and Denko, N. (1990) Asymmetric field inversion gel electrophoresis: a new method for detecting DNA double-strand breaks in mammalian cells. *Radiat. Res.*, **121**, 196–205.
39. Cowell, I.G., Sunter, N.J., Singh, P.B., Austin, C.A., Durkacz, B.W. and Tilby, M.J. (2007) gammaH2AX foci form preferentially in euchromatin after ionising-radiation. *PLoS ONE*, **2**, e1057.
40. Yoshida, K., Yoshida, S.H., Shimoda, C. and Morita, T. (2003) Expression and radiation-induced phosphorylation of histone H2AX in mammalian cells. *J. Radiat. Res.*, **44**, 47–51.
41. Hefferin, M.L. and Tomkinson, A.E. (2005) Mechanism of DNA double-strand break repair by non-homologous end joining. *DNA Repair*, **4**, 639–648.
42. Wyman, C., Ristic, D. and Kanaar, R. (2004) Homologous recombination-mediated double-strand break repair. *DNA Repair*, **3**, 827–833.
43. Tremblay, A., Jasin, M. and Chartrand, P. (2000) A double-strand break in a chromosomal LINE element can be repaired by gene conversion with various endogenous LINE elements in mouse cells. *Mol. Cell Biol.*, **20**, 54–60.
44. Diaz-Perez, S.V., Ferguson, D.O., Wang, C., Csankovszki, G., Wang, C., Tsai, S.C., Dutta, D., Perez, V., Kim, S., Eller, C.D. et al. (2006) A deletion at the mouse Xist gene exposes trans-effects that alter the heterochromatin of the inactive X chromosome and the replication time and DNA stability of both X chromosomes. *Genetics*, **174**, 1115–1133.