

Research



Cite this article: Roberts C, Allen R, Bird KE, Cunliffe M. 2020 Chytrid fungi shape bacterial communities on model particulate organic matter. *Biol. Lett.* **16**: 20200368. <http://dx.doi.org/10.1098/rsbl.2020.0368>

Received: 19 May 2020

Accepted: 8 September 2020

Subject Areas:
ecology

Keywords:
bacteria, fungi, particulate organic matter

Author for correspondence:

Michael Cunliffe

e-mail: micnli@mba.ac.uk

Electronic supplementary material is available online at <https://doi.org/10.6084/m9.figshare.c.5120417>.

Chytrid fungi shape bacterial communities on model particulate organic matter

Cordelia Roberts^{1,2}, Ro Allen¹, Kimberley E. Bird¹ and Michael Cunliffe^{1,2}

¹Marine Biological Association of the UK, The Laboratory, Citadel Hill, Plymouth, UK

²School of Biological and Marine Sciences, University of Plymouth, Plymouth, UK

CR, 0000-0003-0265-8714; RA, 0000-0002-7109-3721; KEB, 0000-0002-7244-5960; MC, 0000-0002-6716-3555

Microbial colonization and degradation of particulate organic matter (POM) are important processes that influence the structure and function of aquatic ecosystems. Although POM is readily used by aquatic fungi and bacteria, there is a limited understanding of POM-associated interactions between these taxa, particularly for early-diverging fungal lineages. Using a model ecological system with the chitin-degrading freshwater chytrid fungus *Rhizoclostridium globosum* and chitin microbeads, we assessed the impacts of chytrid fungi on POM-associated bacteria. We show that the presence of chytrids on POM alters concomitant bacterial community diversity and structure, including differing responses between chytrid life stages. We propose that chytrids can act as ecosystem facilitators through saprotrophic feeding by producing ‘public goods’ from POM degradation that modify bacterial POM communities. This study suggests that chytrid fungi have complex ecological roles in aquatic POM degradation not previously considered, including the regulation of bacterial colonization, community succession and subsequent biogeochemical potential.

1. Introduction

Particulate organic matter (POM) in aquatic ecosystems acts as ‘hotspots’ for bacteria [1–3] and fungi [4,5]. Microbial processing of POM has impacts on ecosystem functioning, including the biological carbon pump in the open ocean [6] and carbon transfer through freshwater food webs [7].

Bacteria–POM studies have characterized the microscale interactions between bacteria and particles, including the composition of colonizing communities [8–10], interactions between attached bacteria [11] and the dynamics of POM degradation [9–12]. Laboratory-based incubations with chitin microbeads as model POM have identified bacteria that colonize and degrade POM using extracellular enzymes, producing a pool of more freely available substrates, including dissolved organic matter (DOM), considered ‘public goods’ for other bacteria in the community to utilize [8].

Dikaryan fungi (Ascomycota and Basidiomycota) also attach to and degrade POM [13–18]. Given that POM-degrading fungi also use extracellular degradation mechanisms [19,20], it is likely that they produce ‘public goods’ for the wider community to exploit. Studies of freshwater leaf-degrading dikaryan fungi show that as bacteria lack key enzymes associated with plant polymer degradation [21], the production of low and intermediate weight DOM by fungi [22] may support enhanced bacterial growth on allochthonous leaf litter [21].

The roles of early-diverging saprotrophic fungal lineages, such as the Chytridiomycota (chytrids), in POM-associated processes are poorly understood.




Figure 1. (a) Schematic summary of experimental design. Experimental treatments: Control = yellow, Zoospores = blue, Established = red. (b) Bacterial diversity measured as the Shannon's H index of treatments over time. Bars represent standard error and asterisks denote level of significance as follows: ** $p < 0.01$, *** $p < 0.001$. (c) NMDS plot of bacterial community structure based on weighted UniFrac dissimilarity between bacterial communities found with treatment over time.

Chytrids are widespread fungi that produce motile zoospores to search for substrates to colonize, including allochthonous and autochthonous POM such as pollen [4], chitin-rich exuviae [23] and zooplankton carcasses [24], as well as living substrates, such as amphibian epidermises [25] and phytoplankton [26]. Once attached, a zoospore loses its flagellum before developing a walled sporangium with a rhizoid network, which attaches to and penetrates the substrate [27]. Chytrids subsequently feed saprotrophically via the rhizoids, which secrete extracellular enzymes to degrade POM to low molecular weight substrates for uptake and assimilation [28].

Even though chytrids and bacteria coexist in aquatic ecosystems, knowledge of chytrid–bacteria POM interactions is limited to niche overlap [4] and infection-associated dynamics on amphibian epidermises [29,30]. To our understanding, there is no current research on the direct influence of chytrids on POM-attached bacterial diversity and community structure. To address these knowledge gaps, we used the chitinophilic *Rhizoclosmatium globosum* in an experimental study with chitin microbeads to assess the interactions between chytrids, bacteria and POM. Using chitin microbeads as a POM experimental system removes the complex heterogeneity of natural particles (e.g. age and composition) while retaining ecological relevance since chitin is an important POM component in aquatic ecosystems [8]. We aimed to explore how the different chytrid life history stages (i.e. attaching zoospores versus established sporangia with rhizoid networks) impact concomitant attaching bacterial diversity and community structure.

2. Material and methods

(a) Experimental set-up

R. globosum JEL800 was maintained on PmTG agar [31] as described previously [27]. To harvest zoospores, established plates were flooded with 4 ml distilled H_2O and incubated at room temperature under laminar flow for 90 min. The zoospore suspension was passed through a 10 μm cell strainer and the concentration determined using a coulter Counter (Beckman Coulter, US).

Magnetic chitin microbeads (New England Bio) were used for the experiments using protocols adapted from [8]. Pond water containing a natural bacterial assemblage was collected from Efford Marsh pond (Plymouth, UK) and passed through a 40 μm mesh to remove detritus and large eukaryotes. Three

experimental treatments were set up as follows: 'Control', 40 μm filtered pond water and chitin microbeads; 'Zoospores', *R. globosum* zoospores, 40 μm filtered pond water and chitin microbeads; and 'Established', *R. globosum* grown initially on chitin microbeads for 24 h in 0.2 μm filtered pond water before addition of experimental 40 μm filtered pond water (figure 1a).

For each treatment, 23.5 ml pond water was added to a 25 ml vented culture flask. Each treatment was conducted with three replicate flasks per timepoint that were sampled destructively. Chitin microbeads were added to the flask and inverted several times to ensure even distribution. For chytrid treatments, zoospores were added to a concentration of approximately 3×10^4 cells ml^{-1} . Flasks were incubated in the dark at 22°C with mixing at 50 rpm. After 2 h, a washing step took place in the chytrid treatments, with the water replaced to ensure that the experiments proceeded with microbead-attached chytrids only ('Established' 40 ± 14.9 chytrids $microbead^{-1}$ and 'Zoospores' 37 ± 15.8 chytrids $microbead^{-1}$). After 2 (i.e. the wash step), 9, 24 and 34 h, all chitin microbeads from each flask were harvested using a magnet. The residual water was discarded, retaining the chitin microbeads, which were frozen in liquid nitrogen and stored at $-80^\circ C$.

(b) DNA extraction, 16S rRNA gene sequencing and bioinformatics

DNA was extracted from the chitin microbeads using the Zymo Research *Quick-DNA*TM Fecal/Soil Microbe Microprep Kit (Zymo Research, USA) following the manufacturer's instructions. The V4–V5 region of the 16S rRNA gene was amplified using primers 515FB and 926R [32], and sequenced using the Illumina MiSeq platform. Sequences were processed in R [33] using the DADA2 pipeline [34]. Demultiplexed reads were filtered and trimmed to remove primers and low-quality sequences. The DADA2 algorithm was used to infer amplicon sequence variants (ASVs) [34]. Paired-end reads were merged to obtain full denoised sequences. Chimeric sequences were removed, before taxonomy was assigned using the SILVA database (release 128) [35]. ASVs assigned as chloroplasts and mitochondria were removed. A maximum likelihood phylogenetic tree was estimated using the *phangorn* package (v.2.5.5) [36] and combined with the ASV table, taxonomic assignment and experimental metadata into a phyloseq object using the *phyloseq* package [37]. Sequences were rarefied to 4955 reads before further analysis.

(c) Data processing and statistical analyses

Shannon's index (H) was used to calculate diversity, and the effect of treatment and time on diversity was assessed using a two-way




Figure 2. Bacterial community composition. Bacteria are grouped by the top 10 most abundant orders; orders outside this are grouped as ‘other’.

ANOVA with Tukey’s HSD. Differences in community structure between samples (beta diversity) were calculated using a weighted UniFrac [38] distance matrix and visualized through non-metric multidimensional scaling (NMDS) ordination. Permutational multivariate analysis of variance (PERMANOVA) [39] was used to test the effect of treatment and time on community structure using the ‘adonis’ function in the R package *vegan* [40].

3. Results

The diversity of the bacterial communities attached to chitin microbeads in all treatments followed the same pattern of decline in the first 9 h of the experiment (figure 1*b*). After 9 h, bacterial diversity in the control treatment continued to decrease for the remainder of the study. Conversely, bacterial diversity in the presence of chytrids increased after 9 h compared with the control treatment (Tukey’s HSD $p < 0.05$). Attached bacterial diversity was greater in the zoospore treatment compared with the established chytrid treatment, although this trend was not statistically significant (Tukey’s HSD $p = 0.08$).

Bacterial community composition varied significantly between treatments and timepoints, with a significant interaction between these variables (PERMANOVA, all $p < 0.001$). After 2 h, the bacterial communities attached to the chitin microbeads in the established treatment were distinct from the other treatments (figure 1*c*), dominated by Burkholderiales, Chromatiales and to a lesser extent Neisseriales and Pseudomonadales (figure 2). Conversely, bacterial communities in the zoospore and control treatments were similar (figure 1*c*) and dominated by Burkholderiales (figure 2).

As the experiment progressed, the structure of the bacterial communities in the zoospore and established chytrid treatments converged, while bacterial communities in the control treatment remained distinct and with relatively limited variation through time (figure 1*c*). At 24 h, Cellvibrionales were also dominant in both the zoospore and established treatments, while Burkholderiales remained dominant in the

control treatment (figure 2). Together, these results suggest that the presence of chytrids on chitin particles impacts both initial bacterial colonization and community succession, with the effects specific to the chytrid life stage.

4. Discussion

Understanding microbial–POM interactions has been largely dominated by bacteria-focused studies. Limited work has been conducted on fungi–bacteria–POM interactions, and the roles of early-diverging fungal lineages in these interactions remains unresolved. Here we show that chytrid fungi impact POM colonizing bacteria and community succession, suggesting that chytrids play a role in shaping POM microbial communities, with implications for carbon cycling in aquatic ecosystems.

Community diversity across all treatments initially declined, following previously reported patterns of bacterial colonizing communities on chitin microbeads [8], suggesting that early colonization here was also governed by known mechanisms (e.g. attachment ability). However, as the presence of the established chytrids with developed rhizoids showed an initially distinct bacterial community structure from the zoospore treatment, this suggests that life-stage dependent ecological interactions occur between chytrids and bacteria that govern particle colonization.

Chitin degradation can result in the release of DOM in aquatic systems [41,42]. Prior to the addition of the bacteria, when the chytrids attach to and degrade the chitin microbeads, they likely produce extracellular *N*-acetylglucosamine (NAG) as a potential ‘public good’. Bacteria, including NAG-using bacteria that are unable to degrade chitin (i.e. ‘cheaters’), may be supported by this pool of ‘public goods’. Community-wide bacterial growth on chitin does not necessarily require chitin degradation by all community members and can be stimulated by the utilization of secondary degradation products including NAG [43,44]. Previous

experimental work has suggested that POM-derived DOM utilization involves a diverse assemblage of bacteria, with no single group dominating the consumption of NAG [45]. The initial structure of the colonizing bacterial community in the established treatment suggests that the bacteria present may be using a source of DOM, such as the proposed NAG pool, and that chytrids may play a role in DOM production from POM degradation in aquatic systems.

The temporal change in bacterial community composition may indicate the decline of the potential NAG pool, and a switch towards a chitin-degrading community at 24 h. As there is a similar switch in community composition seen in the zoospore treatment, this could have been stimulated, in part, by chitin degradation products from the chytrids acting as chemotactic signals for degraders [46,47]. Furthermore, *R. globosum* JEL800 rhizoids form grooves on the outer surface and penetrate chitin microbeads [27], modifying the POM structure and providing increased surface area for bacterial attachment and promoting chitin degradation. It is also possible that, because the chytrid cell wall also contains chitin, the differences in diversity between chytrid and control treatments could be due to an increased relative abundance of chitin-degrading bacteria using chitin from living chytrids or necromass. The predicted function of bacterial communities in this study approximated using Piphillin [48] provides support to this suggestion, showing a convergence in the established and zoospore treatments and divergence from the control treatment, which were distinct over time (electronic supplementary material, figure S1). Piphillin analysis predicted an initially elevated abundance of the NAG transporter gene in the established treatment only that declined over time, presumably as the NAG pool was depleted (electronic supplementary material, figure S2E). Subsequently, there was a predicted increase in chitinase and chitin-oligosaccharide transport genes [49] at 24 h in the established and zoospore treatments (electronic supplementary material, figure S2). Future studies should attempt to unravel the exact nature of the chytrid–bacteria interactions reported here, including the impacts of bacteria on saprotrophic chytrids and the direct assessment of bacterial function, such as through enzyme assays or metatranscriptomics.

POM degradation by microbes may also result in the indirect generation of diverse carbon substrates such as cell debris and metabolic by-products (e.g. organic acids) [8].

Enhanced chemical heterogeneity of these alternative carbon sources produced by both chytrids and bacteria could support the colonization of bacteria that use these products and drive the community dynamics reported here. As the colonization of these bacteria on the particle is likely to invoke competitive interactions, such as for space, the collective community function may diverge from a directly chitin-degrading community towards one that relies on secondary production, as shown in bacteria-only studies [8].

Chytrids have established roles in aquatic ecosystems, including parasitizing hosts and transferring resources via lipid-rich zoospores to higher trophic levels through the mycoloop [50]. Overall, our data indicate that independent of life stage, chytrids also influence the diversity and community structure of POM-colonizing bacterial communities. Increased diversity of bacteria associated with chytrids suggests that chytrids may produce DOM as a pool of ‘public goods’ supporting the growth of ‘cheaters’ and/or encouraging the chemotaxis of chitinolytic bacteria. The potential stimulation of bacterial chitin degradation by the presence of established chytrids, coupled with their own inherent degrading capability, implies that saprotrophic chytrids may have complex roles in regulating POM and DOM processing in aquatic ecosystems that are not yet considered.

Data accessibility. Sequence data have been deposited at the European Nucleotide Archive under project number PRJEB37940.

Authors' contributions. C.R. conceived and designed the study, carried out the study, conducted data analysis and interpretation of data and drafted the manuscript; R.A. supported data analysis, including sequence analysis, and contributed to interpretation of data and editing the manuscript; K.E.B. contributed to carrying out the study, including sample processing, interpretation of data and editing the manuscript; M.C. conceived and designed the study, interpreted data, and drafted and edited the manuscript. All authors gave final approval for publication and agree to be held accountable for the work performed herein.

Competing interests. We declare we have no competing interests.

Funding. C.R. is supported by an ARIES DTP PhD studentship funded from the UK Natural Environment Research Council (NERC). C.R., K.E.B., R.A. and M.C. are supported by the European Research Council (ERC) (MYCO-CARB project; grant no. 772584).

Acknowledgements. We thank Joyce Longcore (University of Maine) for providing *R. globosum* JEL800 from her culture collection (now curated by the Collection of Zoosporic Eufungi at the University of Michigan).

References

1. Azam F, Fenchel T, Field J, Gray J, Meyer-Reil L, Thingstad F. 1983 The ecological role of water-column microbes in the sea. *Mar. Ecol. Prog. Ser.* **10**, 257–263. (doi:10.3354/meps010257)
2. Azam F, Long RA. 2001 Sea snow microcosms. *Nature* **414**, 495–498. (doi:10.1038/35107174)
3. Simon M, Grossart HP, Schweitzer B, Ploug H. 2002 Microbial ecology of organic aggregates in aquatic ecosystems. *Aquat. Microb. Ecol.* **28**, 175–211. (doi:10.3354/ame028175)
4. Wurzbacher C, Rösel S, Rychła A, Grossart HP. 2014 Importance of saprotrophic freshwater fungi for pollen degradation. *PLoS ONE* **9**, e94643. (doi:10.1371/journal.pone.0094643)
5. Bochdansky AB, Clouse MA, Herndl GJ. 2017 Eukaryotic microbes, principally fungi and labyrinthulomycetes, dominate biomass on bathypelagic marine snow. *ISME J.* **11**, 362–373. (doi:10.1038/ismej.2016.113)
6. Boyd PW, Claustre H, Levy M, Siegel DA, Weber T. 2019 Multi-faceted particle pumps drive carbon sequestration in the ocean. *Nature* **568**, 327–335. (doi:10.1038/s41586-019-1098-2)
7. Tanentzap AJ, Szokan-Emilson EJ, Kielstra BW, Arts MT, Yan ND, Gunn JM. 2014 Forests fuel fish growth in freshwater deltas. *Nat. Commun.* **5**, 4077. (doi:10.1038/ncomms5077)
8. Datta MS, Sliwerska E, Gore J, Polz MF, Cordero OX. 2016 Microbial interactions lead to rapid micro-scale successions on model marine particles. *Nat. Commun.* **7**, 11965. (doi:10.1038/ncomms11965)
9. Enke TN, Datta MS, Schwartzman J, Cermak N, Schmitz D, Barrere J, Pascual-García A, Cordero OX. 2019 Modular assembly of polysaccharide-degrading marine microbial communities. *Curr. Biol.* **29**, 1528–1535. (doi:10.1016/j.cub.2019.03.047)
10. Wright RJ, Gibson MI, Christie-Oleza JA. 2019 Understanding microbial community dynamics to improve optimal microbiome selection. *Microbiome* **7**, 85. (doi:10.1186/s40168-019-0702-x)
11. Ebrahimi A, Schwartzman J, Cordero OX. 2019 Cooperation and spatial self-organization determine rate and efficiency of particulate organic matter degradation in marine bacteria. *Proc. Natl Acad. Sci.*

- USA **116**, 23 309–23 316. (doi:10.1073/pnas.1908512116)
12. Schmidt ML, Biddanda BA, Weinke AD, Chiang E, Januska F, Props R, Deneff VJ. 2020 Microhabitats are associated with diversity–productivity relationships in freshwater bacterial communities. *FEMS Microbiol. Ecol.* **96**, 29. (doi:10.1093/femsec/fiaa029)
 13. Claus H, Filip Z. 1998 Degradation and transformation of aquatic humic substances by laccase-producing fungi *Cladosporium cladosporioides* and *Polyporus versicolor*. *Acta Hydrochim. Hydrobiol.* **26**, 180–185. (doi:10.1002/(SICI)1521-401X(199805)26:3<180::AID-AHEH180>3.0.CO;2-9)
 14. Rojas-Jimenez K, Fonvielle JA, Ma H, Grossart HP. 2017 Transformation of humic substances by the freshwater ascomycete *Cladosporium* sp. *Limnol. Oceanogr.* **62**, 1955–1962. (doi:10.1002/lno.10545)
 15. Collado S, Oulego P, Suárez-Iglesias O, Díaz M. 2018 Biodegradation of dissolved humic substances by fungi. *Appl. Microbiol. Biotechnol.* **102**, 3497–3511. (doi:10.1007/s00253-018-8851-6)
 16. Perkins AK, Ganzert L, Rojas-Jimenez K, Fonvielle J, Hose GC, Grossart HP. 2019 Highly diverse fungal communities in carbon-rich aquifers of two contrasting lakes in northeast Germany. *Fungal Ecol.* **41**, 116–125. (doi:10.1016/j.funeco.2019.04.004)
 17. Cunliffe M, Hollingsworth A, Bain C, Sharma V, Taylor JD. 2017 Algal polysaccharide utilisation by saprotrophic planktonic marine fungi. *Fungal Ecol.* **30**, 135–138. (doi:10.1016/j.funeco.2017.08.009)
 18. Pilgaard B, Wilkens C, Herbst F-A, Vuillemin M, Rhein-Knudsen N, Meyer AS, Lange L. 2019 Proteomic enzyme analysis of the marine fungus *Paradendryphiella salina* reveals alginate lyase as a minimal adaptation strategy for brown algae degradation. *Scient. Rep.* **9**, 12338. (doi:10.1038/s41598-019-48823-9)
 19. Abdullah SK, Taj-Aldeen SJ. 1989 Extracellular enzymatic activity of aquatic and aero-aquatic conidial fungi. *Hydrobiologia* **174**, 217–223. (doi:10.1007/BF00008161)
 20. Abdel-Raheem A, Shearer CA. 2002 Extracellular enzyme production by freshwater ascomycetes. *Fungal Divers.* **11**, 1–19.
 21. Romani AM, Romani R, Fischer H, Mille-Lindblom C, Tranvik LJ. 2006 Interactions of bacteria and fungi on decomposing litter: differential extracellular enzyme activities. *Ecology* **87**, 2559–2569. (doi:10.1890/0012-9658(2006)87[2559:IOBAFO]2.0.CO;2)
 22. Fischer H, Mille-Lindblom C, Zwirnmann E, Tranvik LJ. 2006 Contribution of fungi and bacteria to the formation of dissolved organic carbon from decaying common reed (*Phragmites australis*). *Arch. Hydrobiol.* **166**, 79–97. (doi:10.1127/0003-9136/2006/0166-0079)
 23. Sparrow FK. 1960 *Aquatic phycocomycetes*. Ann Arbor, MI: University of Michigan Press.
 24. Tang KW, Bickel SL, Dzialis C, Grossart HP. 2009 Microbial activities accompanying decomposition of cladoceran and copepod carcasses under different environmental conditions. *Aquat. Microb. Ecol.* **57**, 89–100. (doi:10.3354/ame01331)
 25. Longcore JE, Pessier AP, Nichols DK. 1999 *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid pathogenic to amphibians. *Mycologia* **91**, 219–227. (doi:10.1080/00275514.1999.12061011)
 26. Canter HM, Lund JWG. 1948 Studies on plankton parasites. I. Fluctuations in the numbers of *Asterionella formosa* Hass. in relation to fungal epidemics. *New Phytol.* **47**, 238–261. (doi:10.1111/j.1469-8137.1948.tb05102.x)
 27. Laundon D, Christmas N, Wheeler G, Cunliffe M. 2019 Chytrid rhizoid morphogenesis resembles hyphal development in multicellular fungi and is adaptive to resource availability. *Proc. R. Soc. B* **287**, 20200433. (doi: 10.1098/rspb.2020.0433)
 28. Lange L, Barrett K, Pilgaard B, Gleason F, Tsang A. 2019 Enzymes of early-diverging, zoospore fungi. *Appl. Microbiol. Biotechnol.* **103**, 6885–6902. (doi:10.1007/s00253-019-09983-w)
 29. Walke JB, Becker MH, Loftus SC, House LL, Teotonio TL, Minbiole KPC, Belden LK. 2015 Community structure and function of amphibian skin microbes: an experiment with bullfrogs exposed to a chytrid fungus. *PLoS ONE* **10**, e0139848. (doi:10.1371/journal.pone.0139848)
 30. Bates KA *et al.* 2018 Amphibian chytridiomycosis outbreak dynamics are linked with host skin bacterial community structure. *Nat. Commun.* **9**, 693. (doi:10.1038/s41467-018-02967-w)
 31. Barr DJ. 1986 *Allochytridium expandens* rediscovered: morphology, physiology and zoospore ultrastructure. *Mycologia* **78**, 439–448. (doi: 10.1080/00275514.1986.12025267)
 32. Comeau AM, Douglas GM, Langille MGI. 2017 Microbiome Helper: a custom and streamlined workflow for microbiome research. *mSystems* **2**, e00127-16. (doi:10.1128/msystems.00127-16)
 33. R Core Team. 2019 *R: a language and environment for statistical computing, version 3.6.1*. Vienna, Austria: R Foundation for Statistical Computing. See <http://www.R-project.org/>.
 34. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016 DADA2: high-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583. (doi:10.1038/nmeth.3869)
 35. Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013 The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Res.* **41**, 590–596. (doi:10.1093/nar/gks1219)
 36. Schliep KP. 2010 phangorn: Phylogenetic analysis in R. *Bioinformatics* **27**, 592–593. (doi:10.1093/bioinformatics/btq706)
 37. McMurdie PJ, Holmes S. 2013 Phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE* **8**, e0061217. (doi:10.1371/journal.pone.0061217)
 38. Lozupone CA, Hamady M, Kelley ST, Knight R. 2007 Quantitative and qualitative β diversity measures lead to different insights into factors that structure microbial communities. *Appl. Environ. Microbiol.* **73**, 1576–1585. (doi:10.1128/AEM.01996-06)
 39. Anderson MJ. 2001 A new method for non-parametric multivariate analysis of variance. *Austral Ecol.* **26**, 32–46. (doi:10.1046/j.1442-9993.2001.01070.x)
 40. Oksanen J *et al.* 2015 *vegan: Community Ecology Package. R package version 2.5-6*. See <https://cran.r-project.org/web/packages/vegan/index.html>.
 41. Boyer JN. 1994 Aerobic and anaerobic degradation and mineralization of ^{14}C -chitin by water column and sediment inocula of the York River estuary, Virginia. *Appl. Environ. Microbiol.* **60**, 174–179. (doi:10.1128/AEM.60.1.174-179.1994)
 42. Kirchman DL, White J. 1999 Hydrolysis and mineralization of chitin in the Delaware estuary. *Aquat. Microb. Ecol.* **18**, 187–196. (doi:10.3354/ame018187)
 43. Riemann L, Azam F. 2002 Widespread *N*-acetyl- D -glucosamine uptake among pelagic marine bacteria and its ecological implications. *Appl. Environ. Microbiol.* **68**, 5554–5562. (doi:10.1128/AEM.68.11.5554-5562.2002)
 44. Beier S, Bertilsson S. 2013 Bacterial chitin degradation—mechanisms and ecophysiological strategies. *Front. Microbiol.* **4**, 149. (doi:10.3389/fmicb.2013.00149)
 45. Cottrell MT, Kirchman DL. 2000 Natural assemblages of marine proteobacteria and members of the *Cytophaga-Flavobacter* cluster consuming low- and high-molecular-weight dissolved organic matter. *Appl. Environ. Microbiol.* **66**, 1692–1697. (doi:10.1128/AEM.66.4.1692-1697.2000)
 46. Grossart H-P, Kjørboe T, Tang K, Ploug H. 2003 Bacterial colonization of particles: growth and interactions. *Appl. Environ. Microbiol.* **69**, 3500–3509. (doi:10.1128/AEM.69.6.3500-3509.2003)
 47. Mandel MJ, Schaefer AL, Brennan CA, Heath-Heckman EAC, DeLoney-Marino CR, McFall-Ngai MJ, Ruby EG. 2012 Squid-derived chitin oligosaccharides are a chemotactic signal during colonization by *Vibrio fischeri*. *Appl. Environ. Microbiol.* **78**, 4620–4626. (doi:10.1128/AEM.00377-12)
 48. Iwai S, Weinmaier T, Schmidt BL, Albertson DG, Poloso NJ, Dabbagh K, DeSantis TZ. 2016 Piphillin: improved prediction of metagenomic content by direct inference from human microbiomes. *PLoS ONE* **11**, e0166104. (doi:10.1371/journal.pone.0166104)
 49. Larsbrink J, Zhu Y, Kharade SS, Kwiatkowski KJ, Eijsink VGH, Koropatkin NM, McBride MJ, Pope PB. 2016 A polysaccharide utilization locus from *Flavobacterium johnsoniae* enables conversion of recalcitrant chitin. *Biotechnol. Biofuels* **9**, 260. (doi:10.1186/s13068-016-0674-z)
 50. Frenken T *et al.* 2017 Integrating chytrid fungal parasites into plankton ecology: research gaps and needs. *Environ. Microbiol.* **19**, 3802–3822. (doi:10.1111/1462-2920.13827)