



# An improved modified early warning score that incorporates the abdomen score for identifying multiple traumatic injury severity

Xiaobin Jiang

Emergency Department 1, Shanghai Ninth People's Hospital, Shanghai Jiaotong University School of Medicine, Shanghai, China

## ABSTRACT

**Background.** Rapid identification of trauma severity is essential for the timely triage of multiple trauma patients. Tools such as the modified early warning score (MEWS) are used for determining injury severity. Although the conventional MEWS is a good predictor of mortality, its performance assessing injury severity is moderate. This study hypothesized that adding an injury site severity-related score (e.g., abdomen score) may enhance the capability of the MEWS for identifying severe trauma.

**Method.** To validate the hypothesis, we propose an improved modified early warning score called MEWS-A, which incorporates an injury site-specific severity-related abdomen score to MEWS. The utility of MEWS and MEWS-A were retrospectively evaluated and compared for identifying trauma severity in adult multiple trauma patients admitted to the emergency department.

**Results.** We included 1,230 eligible multiple trauma patients and divided them into minor and severe trauma groups based on the injury severity score. Results of logistic regression and receiver operating characteristic (ROC) curve analyses showed that the MEWS-A had a higher area under the ROC curve (AUC: 0.81 95% CI [0.78–0.83]) than did the MEWS (AUC: 0.77 95% CI [0.74–0.79]), indicating that the MEWS-A is superior to the MEWS in identifying severe trauma. The optimal MEWS-A cut-off score is 4, with a specificity of 0.93 and a sensitivity of 0.54. MEWS-A  $\geq$  4 can be used as a protocol for decision-making in the emergency department.

**Conclusions.** Our study suggests that while the conventional MEWS is sufficient for predicting mortality risk, adding an injury site-specific score (e.g., abdomen score) can enhance its performance in determining injury severity in multiple trauma patients.

Submitted 30 April 2020  
Accepted 5 October 2020  
Published 27 October 2020

Corresponding author  
Xiaobin Jiang,  
xiaobin.jiang@hotmail.com

Academic editor  
Eric Bauman

Additional Information and  
Declarations can be found on  
page 10

DOI 10.7717/peerj.10242

© Copyright  
2020 Jiang

Distributed under  
Creative Commons CC-BY 4.0

OPEN ACCESS

**Subjects** Anesthesiology and Pain Management, Emergency and Critical Care, Gastroenterology and Hepatology, Internal Medicine, Public Health

**Keywords** Multiple trauma, Injury severity, Modified early warning score

## INTRODUCTION

In China, multiple trauma is one of the leading causes of death (Wang, Pan & Pan, 2017; Yin, Liang & Liu, 2015). Trauma resulting from motor vehicle and work-related accidents is responsible for more than 400,000 deaths in China annually; of these cases, approximately 1.0–1.8% involve multiple trauma (Zhang, Hong & Gregory, 2017; Yingcheng et al., 2014).

The management of multiple trauma presents important challenges to the emergency department (ED) in China owing to the high mortality risk associated with such injuries.

Rapid identification of trauma severity is essential for the timely triage of multiple trauma patients. However, the assessment of multiple trauma severity might require a sophisticated whole-body medical examination by a specialist, which is time-consuming. To overcome this challenge, scoring systems that can predict mortality risk and assess injury severity for trauma patients in the emergency department have been developed.

Trauma scoring systems can be broadly categorized into anatomical and physiological scoring systems. Anatomical trauma scores are mainly injury site-dependent scores such as the abbreviated injury scale (AIS) score (*Loftis, Price & Gillich, 2018*) and the AIS-derived injury severity score (ISS) (*Baker et al., 1974*), which evaluates the injury severity for each site and provides a weighted score for the sites with the most serious injuries. Although such tools are effective, they also require trauma specialists to examine each injury site, which is, again, time-consuming and impractical in an ED.

Physiological trauma scoring systems include the revised trauma score (RTS) (*Alvarez et al., 2016*), APACHE II (*Godinjak et al., 2016*), sepsis-related organ failure assessment (SOFA) score (*Raith et al., 2017*), and emergency trauma score (ETS) (*Raum et al., 2009*). These scores incorporate age, physiological variables (e.g., heart rate), and neurological variables (e.g., the Glasgow coma scale (GCS)) to identify injury severity. RTS may help in decision-making for hospital discharge and intensive care unit (ICU) admission (*Mansour, Eisha & Asaad, 2019*), but its accuracy in predicting mortality decreases with increasing age (*Kojima et al., 2019*) and exhibits poor correlation with anatomical injury severity (*Galvagno Jr et al., 2019*). APACHE II has proven to be a good predictor of mortality among brain injury patients (*Nik et al., 2018*), but it is inefficient for trauma triage in EDs because it has too many variables; its calculation requires 14 variables. The SOFA score has been used to quantify organ function and may predict mortality in ICU patients (*Bingold et al., 2015*). However, it is not well known among emergency doctors and its use is inconvenient because several days may be required for calculations aimed at verifying whether patients fulfill the strict criteria and require laboratory tests (*Machado et al., 2016*). The ETS is a simpler predictor of in-hospital mortality in trauma patients (*Park et al., 2017*), but a duration of 30 min may be required to measure the parameters (*Raum et al., 2009*). Overall, these scoring systems are reliable tools for identifying trauma severity, but they are also inconvenient to calculate and measure. The inherent complexity of sophisticated scoring tools makes their application as a rapid risk stratification tool that can be applied in a regular ED setting challenging.

Given the simplicity and speed required for tools to be effective in triaging multiple trauma patients in EDs, tools such as the modified early warning score (MEWS) (*Bozkurt et al., 2015*) have now been widely applied in EDs, particularly in China (*Xie et al., 2018*). The MEWS has been used for many purposes: to predict in-hospital mortality (*Xie et al., 2018*), determine hospital admission (*Wei et al., 2019*), identify critically ill patients (*Kruisselbrink et al., 2016*), and predict injury severity and need for ICU admission (*Salottolo et al., 2017*).

However, our previous study (*Jiang, Jiang & Mao, 2019*) suggests that although the MEWS is effective for predicting mortality, it is not an excellent predictor of trauma

severity, as its area under the receiver operating characteristic (ROC) curve (AUROC) for trauma severity prediction is less than 0.8. [Wong et al. \(2016\)](#) reported that adding an evaluation variable related to the injury site severity to an injury severity scoring system may improve the performance of the scoring system; thus, we hypothesized that adding an abdomen score, a variable for evaluating the severity of injury to the thorax and abdomen, to the MEWS may enhance its ability to predict injury severity of multiple trauma patients.

To validate the hypothesis, we proposed a scoring system called the MEWS-A, which combines the conventional MEWS with an abdomen score. We retrospectively evaluated and compared the utility of the MEWS and MEWS-A for identifying trauma severity of multiple trauma patients admitted to the ED.

## MATERIALS AND METHODS

### Ethical statement

The study protocol was approved by the Ethics Committee of Shanghai Ninth People's Hospital (approval no.: 2018-146-T132 and 2020-2018-146-T132-1), which waived the requirement to obtain informed consent.

### Settings and subjects

The study cohort included adult multiple trauma patients who were consecutively admitted to the author's institution between January 2014 and June 2018. The institution is an urban emergency medical center, admitting approximately 20,000 trauma patients each year, of which approximately 3,000 admissions present with multiple trauma. As in [Frink et al. \(2017\)](#), we defined multiple trauma as two or more body region injuries where one or a combination of multiple injuries may endanger the patient's life. We determined multiple trauma by reviewing medical records and computed tomography (CT).

Among all adult multiple trauma patients, those who had a pre-injury medical history or dysfunction were excluded because it would be challenging to identify outcomes directly related to injury severity ([Jiang, Jiang & Mao, 2019](#)). The study cohort was divided into minor and severe trauma groups according to ISS scores (ISS score  $<16$  and  $\geq 16$ , respectively). ISS  $\geq 16$  indicates that patients have a greater need for trauma care ([Galvagno Jr et al., 2015](#)), and it is a conventional threshold for defining severe trauma ([Kehoe et al., 2015](#)).

### MEWS and MEWS-A

[Table 1](#) shows the MEWS and MEWS-A scoring systems, which are used to assess the trauma severity of the included patients and compared. The conventional MEWS consists of four vital sign variables: systolic blood pressure (SBP), heart rate (HR), respiratory rate (RR), temperature (T), and one neurological variable, AVPU. AVPU measures the level of consciousness and has four possible outcomes: alert (A), verbal (V), pain (P), and unresponsive (U). As reported previously ([Mitsunaga et al., 2019](#)), the AVPU score is derived from the GCS as follows:  $A = 14-15$ ,  $V = 9-13$ ,  $P = 4-8$ , and  $U = 3$ . The MEWS is reported as an effective tool in identifying the risk of mortality ([Kruisselbrink et al., 2016](#)), but it is less sensitive in identifying trauma severity ([Jiang, Jiang & Mao, 2019](#)).

**Table 1** MEWS and MEWS-A score.

Variable	Score			
	0	1	2	3
Systolic blood pressure (mmHg)	101–199	81–100	70–80	<70
Heart rate (beats/min)	51–100	40–50	≥200	≥130
Respiratory rate (cycles/min)	9–14	101–110	<40	≥30
Temperature (°C)	35–38.4	15–20	111–129	≥30
AVPU score	Alert	15–20	<9	≥30
Abdomen score	35–38.4	21–29	<35	≥38.5
	Alert	Reacts to voice	Reacts to pain	Unresponsive
	Abdomen and Thorax non-tender	Abdomen or Thorax tender	Abdomen rigid or Flail chest	

The MEWS-A is generated by adding an abdomen score variable to the MEWS in order to improve its ability to assess injury severity. Designing new criteria (e.g., score range and calculation method) for the abdomen score is challenging; hence, we used an abdomen score that has been proposed in another scoring system, the Circulation, Respiration, Abdomen, Motor, Speech (CRAMS) scale ([Gormican, 1982](#)), which has been proven valid and reliable for assessing mortality risks of multiple trauma patients ([Chen et al., 2017](#)). Further, the abdomen score is preferable in EDs because it is convenient to measure and easy to remember. Abdomen score can be assessed by checking for tenderness (e.g., pain or discomfort) in the abdomen and thorax by palpation with no medical devices. Specifically, as shown in [Table 2](#), the abdomen score ranges from 0 to 2: 0, 1, and 2 points represent conditions of non-tender abdomen and thorax, tender abdomen or thorax, and rigid abdomen or flail chest, respectively.

### Statistical analysis

Statistical analysis was conducted on both scoring systems to evaluate their performance in assessing trauma severity in multiple trauma patients using R, version 3.5.2 (R Foundation for Statistical Computing, Vienna, Austria). Multiple imputation was applied for the included patients who had missing values for one to three vital signs. The MEWS and MEWS-A score, shown in [Table 1](#), were then computed for the minor and severe trauma groups.

Numerical and categorical outcomes are presented as medians (interquartile ranges [IQRs]) and frequencies (percentages), respectively. Categorical outcomes were analyzed by Pearson's chi-square test. Normality tests were performed on numerical outcomes. Normally distributed numerical outcomes were analyzed by the *t*-test while non-normally distributed outcomes were analyzed by the Wilcoxon-Mann-Whitney test.


**Table 2** Characteristics of the study cohort.

	Total	Minor trauma	Severe trauma	P-value
No.	1230	475	755	
Male, n (%)	904 (73.5)	323 (68.0)	581 (77.0)	0.001
Age (years)	48 (38, 59)	50 (39, 60)	48 (38, 58)	0.099
Temperature (°C)	37.0 (36.7, 37.0)	37.0 (36.8, 37.0)	36.8 (36.5, 37.0)	<0.001
Heart rate (beats/min)	85 (75, 96)	83 (76, 90)	85 (75, 100)	0.012
Respiratory rate (cycles/min)	20 (18, 21)	20 (19, 20)	20 (18, 21)	0.012
Systolic blood pressure (mmHg)	130 (115, 147)	136 (121, 150)	127 (107, 145)	<0.001
AVPU	0 (0, 1)	0 (0, 0)	1 (0, 2)	<0.001
LOS	12.0 (5.0, 21.0)	9.0 (6.0, 15.0)	15.0 (4.0, 26.0)	<0.001
Time of transport (hour)	2.0 (1.0, 3.0)	2.0 (1.0, 3.0)	2.0 (1.0, 3.0)	<0.001
Cause of injury, n (%)				<0.001
Motor vehicle crashes	683 (55.5)	217 (45.7)	466 (61.7)	
High fall	222 (18.0)	69 (14.5)	153 (20.3)	
Crushing injury	41 ( 3.3)	11 ( 2.3)	30 ( 4.0)	
Cut/piercing injury	45 ( 3.7)	29 ( 6.1)	16 ( 2.1)	
Burn	4 ( 0.3)	2 ( 0.4)	2 ( 0.3)	
Tumble injury	169 (13.7)	112 (23.6)	57 ( 7.5)	
Striking injury	66 ( 5.4)	35 ( 7.4)	31 ( 4.1)	
Primary injury site–n (%)				<0.001
Face	19 ( 1.5)	17 ( 3.6)	2 ( 0.3)	
Head and neck	537 (43.7)	102 (21.5)	435 (57.6)	
Thorax	318 (25.9)	129 (27.2)	189 (25.0)	
Abdomen and visceral pelvis	111 ( 9.0)	42 ( 8.8)	69 ( 9.1)	
Bony pelvis and extremities	243 (19.8)	184 (38.7)	59 ( 7.8)	
External structures	2 ( 0.2)	1 ( 0.2)	1 ( 0.1)	
Discharge status, n (%)				<0.001
Died in the hospital	185 (15.0)	2 ( 0.4)	183 (24.2)	
Discharged home	583 (47.4)	303 (63.8)	280 (37.1)	
Discharged against medical advice	73 ( 5.9)	29 ( 6.1)	44 ( 5.8)	
Discharged home with recommended self-care	344 (28.0)	112 (23.6)	232 (30.7)	
Transferred to another hospital	45 ( 3.7)	29 ( 6.1)	16 ( 2.1)	
Score				
MEWS	2 (1, 3)	1 (1, 2)	2 (1, 4)	<0.001
MEWS-A	3 (1, 4)	1 (1, 2)	4 (2, 5)	<0.001

To evaluate the scoring systems' performance in identifying trauma severity, the AUROC for the following logistic regression model was compared for the two scoring systems.

$$y = \frac{1}{1 + e^{-(b+wx)}} \quad (1)$$

where  $b$ ,  $w$ , and  $x$  indicate the bias, the weight, and the score, respectively.



**Figure 1** Study cohort.

Full-size  DOI: [10.7717/peerj.10242/fig-1](https://doi.org/10.7717/peerj.10242/fig-1)

To find the optimal decision threshold for identifying severe trauma, optimal cut-off points of both scoring systems were calculated. Further, the corresponding accuracy rate, sensitivity, and specificity at the point was analyzed.

## RESULTS

### Setting and subjects

**Figure 1** summarizes the study cohort. In total, 1,230 adult patients presenting with multiple trauma met the inclusion criteria. The minor injury and severe injury groups included 475 (38.6%) and 755 (61.4%) cases, respectively. Of 1,230 patients, 52 (4.2%) had missing vital signs data, among whom 30 had missing data of HR, RR, and SBP, 9 had missing data of SBP, 8 had missing data of RR and SBP, 2 had missing values of RR, 1 had missing data of HR and SBP, 1 had missing data of T and SBP, and 1 had missing data of T.

Normality testing results showed that distributions of all numerical variables could not be approximated as normal distributions. As shown in [Table 2](#), male patients, whose median (IQR) age was 48 (38–59) years, made up 73.5% of the sample. Most patients (90.4%) were transported to the ED by an ambulance. Comparing the length of stay (LOS) of the two groups, as expected, severe trauma patients had longer stays (15 [4–26] vs. 9 [6–15] days) in the hospital. Patients' injuries were primarily the result of motor vehicle accidents (55.5%). Most minor trauma patients suffered from injuries to the bony pelvis and extremities (38.7%) while most severe trauma patients suffered from injuries to the head and neck (57.6%).

The median MEWS and MEWS-A score (IQR) of the two groups were 1 (1-2) vs. 2 (1-4) and 1 (1-2) vs. 4 (2-5), respectively ( $p < 0.001$ ).

### Performances of the MEWS and MEWS-A score

As shown in [Fig. 2](#), the MEWS-A has a greater AUROC value (AUROC = 0.81, 95% CI [0.78–0.83]) than the MEWS (AUROC = 0.77, 95% CI [0.74–0.79]) for multiple trauma severity assessment, showing that incorporating the abdomen variable improves the trauma severity assessment performance. DeLong's test was used to compare the two ROCs and the  $p$ -value was  $p < 0.001$ .


The optimal cut-off values of  $\geq 3$  for the MEWS and  $\geq 4$  for the MEWS-A score were established. The corresponding accuracy, sensitivity, and specificity rates of the MEWS and MEWS-A score were 0.67 vs. 0.69, 0.52 vs. 0.54, and 0.92 vs. 0.93, respectively.

## DISCUSSION

This study aimed to validate the hypothesis that adding an injury site severity score (abdomen score) to the conventional MEWS may improve its ability to identify trauma severity. Our results demonstrate that the MEWS-A score had a higher AUROC value (AUROC = 0.81) for trauma severity assessment than the conventional MEWS (AUROC = 0.77), indicating that incorporating the abdomen score improves the performance of the conventional MEWS.

The conventional MEWS has been applied for identifying mortality risk and determining trauma severity. The MEWS was previously reported to be an excellent predictor of in-hospital mortality because the MEWS achieves high area under ROC (AUROC ranges from 0.83 to 0.90 in [Xie et al. \(2018\)](#), [Jiang, Jiang & Mao \(2019\)](#), and [Smith et al., 2013](#)), although the performance may decrease for elderly patients ([Mitsunaga et al., 2019](#)). However, the performance of the MEWS in identifying injury severity is moderate (AUROC < 0.8). Statistical analyses ([Salottolo et al., 2017](#)) showed that the MEWS is less associated with severe injury than in-hospital mortality. ([Jiang, Jiang & Mao, 2019](#)) also showed a lower AUROC value (AUROC = 0.77) of identifying trauma severity. Similarly, our results showed that the MEWS is a moderate predictor (AUROC = 0.77) for assessing injury severity.

By adding the abdomen score to MEWS, its ability to identify injury severity was improved. The conventional MEWS includes vital signs and AVPU. The vital signs may correlate well with mortality, but they show a poor correlation with injury severity. Even



**Figure 2** Receiver operating characteristic (ROC) curves for MEWS and MEWS-A.

Full-size  DOI: [10.7717/peerj.10242/fig-2](https://doi.org/10.7717/peerj.10242/fig-2)

severe trauma patients may have a low MEWS and stable physiological measures (*Gottschalk et al., 2006*). As reported previously, no physiological measure has a sensitivity high enough for a negative result (e.g., normal physiological value) to be confidently used for concluding that a patient is not seriously injured (*Annette et al., 2018*). In addition, injury sites can be anatomically categorized into 6 body regions: head, face, chest, abdomen, extremities (including pelvis), and external (*Quah, 2016*). The AVPU score is derived from the GCS, a triage tool that assess traumatic brain injury (*Gill et al., 2006*). Hence, the AVPU score could play a major role in identifying trauma severity at the head and neck. With the additional incorporation of an abdomen score, the MEWS-A score may help in evaluating injury severity in the chest and abdomen, leading to better performance in identifying overall injury severity.

Although the MEWS-A has a higher AUROC value, the specificity (correctly identified as minor trauma) and sensitivity (correctly identified as severe trauma) are marginally higher than those of the conventional MEWS. Several further modifications may improve the function of MEWS-A and will be addressed in future research. First, replacing AVPU with GCS may improve the function because AVPU is estimated from GCS and the association



of AVPU with GCS may vary with age (Nuttall, Paton & Kemp, 2018). Second, the proposed MEWS-A has the variables to assess trauma severity of the head, face, chest, and abdomen, but has no variables to evaluate severity of injuries to the extremities and external region. One solution to this is to add an AIS score for extremities and external region (Wong et al., 2016), but evaluation of the AIS score may require a CT scan and specialist knowledge of injuries, which will present difficulties for the nurses in EDs. Development of an extremity and external injury score variable, which can be easily used by nurses, will be included in future research. As shown in Table 2, the MEWS-A's ability to predict injury severity is marginally better than that of the MEWS because the majority of the cohort's injury sites are the head and neck. Additional data on patients presenting with chest and abdominal injuries will also be collected in future research.

The present study also showed the optimal cut-off threshold of the MEWS-A score  $\geq 4$  for identifying injury severity in multiple trauma patients with high specificity (0.93) and good sensitivity (0.54). The MEWS-A  $\geq 4$  can be used in decision-making protocol for ICU admission, surgery, and immediate medical intervention.

The MEWS-A score can be rapidly calculated even though incorporating an abdomen score increases the complexity of the scoring system. The abdomen score can be directly measured through palpation, avoiding the challenges associated with the requirement of special medical devices and transportation of patients between rooms and floors for examination. Empirically, nurses can accomplish the assessment of the MEWS-A within 5 min of the patient's arrival at the ED, thus requiring much less time than the 30 min needed to determine the ETS (Raum et al., 2009). Hence, the MEWS-A is convenient and can be used for multiple trauma triage in the ED.

Our study has several limitations. First, it was retrospective and was limited to a relatively small sample of patients from a single center. Owing to the retrospective design, it is possible that the calculated scores may have been different had they been obtained in real time because of the time pressure experienced at EDs and the variability due to scoring by multiple individuals. A prospective study based on a larger cohort from multiple emergency centers is, therefore, necessary to validate the findings. Most multiple injury patients in the cohort presented with head and neck injuries, which may cause bias in the results. In future research, more medical records of multiple trauma patients should be analyzed for further evaluation of injury severity assessment using the MEWS and MEWS-A score.

## CONCLUSIONS

The conventional MEWS is suitable for predicting mortality but is a relatively poor predictor of injury severity. However, the MEWS-A score, which incorporates an abdomen score into the conventional MEWS, is superior in identifying injury severity in multiple trauma patients, indicating that incorporating a site-specific injury severity score (e.g., abdomen score) can improve the performance of the MEWS. The MEWS-A is a simple and rapid tool for application and can be used for the timely triage of multiple trauma patients in the ED.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

The author received no funding for this work.

### Competing Interests

The author declares there are no competing interests.

### Author Contributions

- Xiaobin Jiang conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

### Ethics

The following information was supplied relating to ethical approvals (i.e., approving body and any reference numbers):

This retrospective cohort study was approved by the ethics committee of Shanghai Ninth People's Hospital, which is affiliated with Shanghai JiaoTong University School of Medicine (approval no.: 2018-146-T132 and 2020-2018-146-T132-1). The need for informed consent from the patients was waived.

### Data Availability

The following information was supplied regarding data availability:

Raw data and a code book are available in the [Supplementary Files](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10242#supplemental-information>.

## REFERENCES

- Alvarez BD, Razente D, Lacerda DA, Lothar NS, Von-Bahten IC, Stahlschmidt C, Menini M. 2016.** Analysis of the Revised Trauma Score (RTS) in 200 victims of different trauma mechanisms. *Revista do Colégio Brasileiro de Cirurgiões* **43(5)**:334–340 DOI 10.1590/0100-69912016005010.
- Annette MT, Tamara PC, Maya EO, Craig DN, Mohamud D, Rongwei F, Ngoc W, Erica LH, Roger C. 2018.** *Physiologic predictors of severe injury: systematic review*. Rockville: Agency for Healthcare Research and Quality (US).
- Baker SP, o'Neill B, Haddon Jr W, Long WB. 1974.** The injury severity score: a method for describing patients with multiple injuries and evaluating emergency care. *Journal of Trauma and Acute Care Surgery* **14(3)**:187–196 DOI 10.1097/00005373-197403000-00001.
- Bingold TM, Lefering R, Zacharowski K, Meybohm P, Waydhas C, Rosenberger P, Scheller B, Group DICR. 2015.** Individual organ failure and concomitant risk of

mortality differs according to the type of admission to ICU—a retrospective study of SOFA score of 23,795 patients. *PLOS ONE* **10(8)**:e0134329

DOI [10.1371/journal.pone.0134329](https://doi.org/10.1371/journal.pone.0134329).

- Bozkurt S, Köse A, Arslan ED, Erdoğan S, Üçbilek E, Çevik İ, Ayrık C, Sezgin O. 2015.** Validity of modified early warning, Glasgow Blatchford, and pre-endoscopic Rockall scores in predicting prognosis of patients presenting to emergency department with upper gastrointestinal bleeding. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* **23(1)**:109 DOI [10.1186/s13049-015-0194-z](https://doi.org/10.1186/s13049-015-0194-z).
- Chen X, Yang W, Sun S, Wang G. 2017.** Application of ISS and CRAMS score for the assessment of the prognosis of patients with multiple trauma. *Chinese Journal of Emergency Medicine* **26(6)**:664–668 DOI [10.3760/cma.j.issn.1671-0282.2017.06.013](https://doi.org/10.3760/cma.j.issn.1671-0282.2017.06.013).
- Frink M, Lechler P, Debus F, Ruchholtz S. 2017.** Multiple trauma and emergency room management. *Deutsches Ärzteblatt International* **114(29–30)**:497–503 DOI [10.3238/arztebl.2017.0497](https://doi.org/10.3238/arztebl.2017.0497).
- Galvagno Jr SM, Massey M, Bouzat P, Vesselinov R, Levy MJ, Millin MG, Stein DM, Scalea TM, Hirshon JM. 2019.** Correlation between the revised trauma score and injury severity score: implications for prehospital trauma triage. *Prehospital Emergency Care* **23(2)**:263–270 DOI [10.1080/10903127.2018.1489019](https://doi.org/10.1080/10903127.2018.1489019).
- Galvagno Jr SM, Sikorski R, Hirshon JM, Floccare D, Stephens C, Beecher D, Thomas S. 2015.** Helicopter emergency medical services for adults with major trauma. *Cochrane Database of Systematic Reviews* **12(12)**:CD009228 DOI [10.1002/14651858.CD009228.pub3](https://doi.org/10.1002/14651858.CD009228.pub3).
- Gill M, Steele R, Windemuth R, Green SM. 2006.** A comparison of five simplified scales to the out-of-hospital Glasgow Coma Scale for the prediction of traumatic brain injury outcomes. *Academic Emergency Medicine* **13(9)**:968–973 DOI [10.1197/j.aem.2006.05.019](https://doi.org/10.1197/j.aem.2006.05.019).
- Godinjak A, Igllica A, Rama A, Tančica I, Jusufović S, Ajanović A, Kukuljac A. 2016.** Predictive value of SAPS II and APACHE II scoring systems for patient outcome in a medical intensive care unit. *Acta Medica Academica* **45(2)**:97–103 DOI [10.5644/ama2006-124.165](https://doi.org/10.5644/ama2006-124.165).
- Gormican SP. 1982.** CRAMS scale: field triage of trauma victims. *Annals of Emergency Medicine* **11(3)**:132–135 DOI [10.1016/S0196-0644\(82\)80237-0](https://doi.org/10.1016/S0196-0644(82)80237-0).
- Gottschalk S, Wood D, DeVries S, Wallis LA, Bruijns S. 2006.** The cape triage score: a new triage system South Africa Proposal from the cape triage group. *Emergency Medicine Journal* **23(2)**:149–153 DOI [10.1136/emj.2005.028332](https://doi.org/10.1136/emj.2005.028332).
- Jiang X, Jiang P, Mao Y. 2019.** Performance of Modified Early Warning Score (MEWS) and Circulation, Respiration, Abdomen, Motor, and Speech (CRAMS) score in trauma severity and in-hospital mortality prediction in multiple trauma patients: a comparison study. *PeerJ* **7**:e7227 DOI [10.7717/peerj.7227](https://doi.org/10.7717/peerj.7227).
- Kehoe A, Smith J, Edwards A, Yates D, Lecky F. 2015.** The changing face of major trauma in the UK. *Emergency Medicine Journal* **32(12)**:911–915 DOI [10.1136/emermed-2015-205265](https://doi.org/10.1136/emermed-2015-205265).

- Kojima M, Endo A, Shiraishi A, Otomo Y. 2019.** Age-related characteristics and outcomes for patients with severe trauma: analysis of Japan's nationwide trauma registry. *Annals of Emergency Medicine* **73**(3):281–290  
[DOI 10.1016/j.annemergmed.2018.09.034](https://doi.org/10.1016/j.annemergmed.2018.09.034).
- Kruisselbrink R, Kwizera A, Crowther M, Fox-Robichaud A, O'Shea T, Nakibuuka J, Ssinabulya I, Nalyazi J, Bonner A, Devji T, Wong J, Cook D. 2016.** Modified Early Warning Score (MEWS) identifies critical illness among ward patients in a resource restricted setting in Kampala, Uganda: a prospective observational study. *PLOS ONE* **11**(3):e0151408 [DOI 10.1371/journal.pone.0151408](https://doi.org/10.1371/journal.pone.0151408).
- Loftis KL, Price J, Gillich PJ. 2018.** Evolution of the abbreviated injury scale: 1990–2015. *Traffic Injury Prevention* **19**(sup2):S109–S113 [DOI 10.1080/15389588.2018.1512747](https://doi.org/10.1080/15389588.2018.1512747).
- Machado FR, De Assunção MSC, Cavalcanti AB, Japiassú AM, De Azevedo LCP, Oliveira MC. 2016.** Getting a consensus: advantages and disadvantages of Sepsis 3 in the context of middle-income settings. *Revista Brasileira de Terapia Intensiva* **28**(4):361–365 [DOI 10.5935/0103-507X.20160068](https://doi.org/10.5935/0103-507X.20160068).
- Mansour DA, Eisha HAA, Asaad AE. 2019.** Validation of revised trauma score in the emergency department of Kasr Al Ainy. *The Egyptian Journal of Surgery* **38**(4):679–684 [DOI 10.4103/ejs.ejs\\_82\\_19](https://doi.org/10.4103/ejs.ejs_82_19).
- Mitsunaga T, Hasegawa I, Uzura M, Okuno K, Otani K, Ohtaki Y, Sekine A, Takeda S. 2019.** Comparison of the National Early Warning Score (NEWS) and the Modified Early Warning Score (MEWS) for predicting admission and in-hospital mortality in elderly patients in the pre-hospital setting and in the emergency department. *PeerJ* **7**:e6947 [DOI 10.7717/peerj.6947](https://doi.org/10.7717/peerj.6947).
- Nik A, Andalibi MSS, Ehsaei MR, Zarifian A, Karimiani EG, Bahadoorkhan G. 2018.** The efficacy of glasgow coma scale (GCS) score and acute physiology and chronic health evaluation (APACHE) II for predicting hospital mortality of ICU patients with acute traumatic brain injury. *Bulletin of Emergency & Trauma* **6**(2):141–145  
[DOI 10.29252/beat-060208](https://doi.org/10.29252/beat-060208).
- Nuttall AG, Paton KM, Kemp AM. 2018.** To what extent are GCS and AVPU equivalent to each other when assessing the level of consciousness of children with head injury? A cross-sectional study of UK hospital admissions. *BMJ Open* **8**(11):e023216  
[DOI 10.1136/bmjopen-2018-023216](https://doi.org/10.1136/bmjopen-2018-023216).
- Park HO, Kim JW, Kim SH, Moon SH, Byun JH, Kim KN, Yang JH, Lee CE, Jang IS, Kang DH, Kim SC, Kang C, Choi JY. 2017.** Usability verification of the Emergency Trauma Score (EMTRAS) and Rapid Emergency Medicine Score (REMS) in patients with trauma: a retrospective cohort study. *Medicine* **96**(44):e8449  
[DOI 10.1097/MD.00000000000008449](https://doi.org/10.1097/MD.00000000000008449).
- Quah SR. 2016.** International encyclopedia of public health. Cambridge, MA, USA: Academic Press.
- Raith EP, Udy AA, Bailey M, McGloughlin S, MacIsaac C, Bellomo R, Pilcher DV. 2017.** Prognostic accuracy of the SOFA score, SIRS criteria, and qSOFA score for in-hospital mortality among adults with suspected infection admitted to the intensive care unit. *JAMA* **317**(3):290–300 [DOI 10.1001/jama.2016.20328](https://doi.org/10.1001/jama.2016.20328).

- Raum MR, Nijsten MW, Vogelzang M, Schuring F, Lefering R, Bouillon B, Rixen D, Neugebauer EA, Henk J, PSG of the German Trauma Society. 2009.** Emergency trauma score: an instrument for early estimation of trauma severity. *Critical Care Medicine* 37(6):1972–1977 DOI 10.1097/CCM.0b013e31819fe96a.
- Salottolo K, Carrick M, Johnson J, Gamber M, Bar-Or D. 2017.** A retrospective cohort study of the utility of the modified early warning score for inter-facility transfer of patients with traumatic injury. *BMJ Open* 7(5):e016143 DOI 10.1136/bmjopen-2017-016143.
- Smith GB, Prytherch DR, Meredith P, Schmidt PE, Featherstone PI. 2013.** The ability of the National Early Warning Score (NEWS) to discriminate patients at risk of early cardiac arrest, unanticipated intensive care unit admission, and death. *Resuscitation* 84(4):465–470 DOI 10.1016/j.resuscitation.2012.12.016.
- Wang G, Pan Z, Pan L. 2017.** Analysis of the factors affecting the death of patients with multiple injuries in emergency. *Tianjin Medical Journal* 45:885–888.
- Wei X, Ma H, Liu R, Zhao Y. 2019.** Comparing the effectiveness of three scoring systems in predicting adult patient outcomes in the emergency department. *Medicine* 98(5):e14289 DOI 10.1097/MD.00000000000014289.
- Wong TH, Krishnaswamy G, Nadkarni NV, Nguyen HV, Lim GH, Bautista DCT, Chiu MT, Chow KY, Ong M. EH. 2016.** Combining the new injury severity score with an anatomical polytrauma injury variable predicts mortality better than the new injury severity score and the injury severity score: a retrospective cohort study. *Scandinavian journal of trauma, Resuscitation and Emergency Medicine* 24(1):25 DOI 10.1186/s13049-016-0215-6.
- Xie X, Huang W, Liu Q, Tan W, Pan L, Wang L, Zhang J, Wang Y, Zeng Y. 2018.** Prognostic value of modified early warning score generated in a Chinese emergency department: a prospective cohort study. *BMJ Open* 8(12):e024120 DOI 10.1136/bmjopen-2018-024120.
- Yin Q, Liang Y, Liu Z. 2015.** Analysis of the emergency death risk factors and death causes for multiple trauma. *Journal of Clinical Emergency* 16:591–593.
- Yingcheng X, Chaofang S, Yong X, Er L. 2014.** Clinical analysis of emergency assessment and treatment of 216 severe multiple trauma patients. *Modern Journal of Integrated Traditional Chinese and Western Medicine* 23(19):2094–2097.
- Zhang Z, Hong Y, Gregory JS. 2017.** Trauma care in China: a systematic review. *Journal of Emergency and Critical Care Medicine* 1(11):33 DOI 10.21037/jeccm.2017.10.05.