

## Effect of Eight Weeks of Quercetin Supplementation on Exercise Performance, Muscle Damage and Body Muscle in Male Badminton Players

Pooya Daneshvar<sup>3</sup>, Mitra Hariri<sup>1,2</sup>, Reza Ghasvand<sup>1,2</sup>, Gholamreza Askari<sup>1,2</sup>, Leila Darvishi<sup>1,2</sup>, Nafiseh Shokri Mashhadi<sup>1,2</sup>, Hossein Khosravi-boroujeni<sup>4</sup>

<sup>1</sup>Food Security Research Center, Isfahan University of Medical Sciences, Isfahan, Iran, <sup>2</sup>Department of Community Nutrition, School of Nutrition and Food Science, Isfahan University of Medical Sciences, Isfahan, Iran, <sup>3</sup>Child Growth and Development Research Center, Isfahan University of Medical Sciences, Isfahan, Iran, <sup>4</sup>Falavarjan Branch, Islamic Azad University, Falavarjan, Iran

### Correspondence to:

Mrs. Leila Darvishi,  
Food Security Research  
Center, Isfahan University of  
Medical Sciences, Isfahan, Iran.  
E-mail: leilad\_78@yahoo.com.

**Date of Submission:** Feb 24, 2012

**Date of Acceptance:** Jun 07, 2012

**How to cite this article:** Daneshvar P, Hariri M, Ghasvand R, Askari G, Darvishi L, Mashhadi NS, Khosravi-boroujeni H. Effect of eight weeks of quercetin supplementation on exercise performance, muscle damage and body muscle in male badminton players. *Int J Prev Med* 2013; Supplement 1: S53-7.

### ABSTRACT

**Background:** Quercetin is a bioflavonoid which occurs in many food items. Some previous studies on quercetin showed inconsistent results on exercise performance, muscle damage and body muscle in athletes. The aim of this study was to determine the effects of eight weeks of quercetin supplementation on exercise performance, muscle damage indices and body muscle in badminton players.

**Methods:** This placebo-controlled, double-blind clinical trial was conducted on 26 badminton players for eight weeks. The subjects were randomly assigned to one of two groups to receive quercetin (1000 mg) or placebo (1000 mg dextrose).  $VO_2$  and time to exhaustion (TTE) for measuring performance and body fat percentage (BFP) were measured before and after intervention. Plasma sample were obtained for the determination of plasma lactate before and after intervention.

**Results:** Lactate concentration, body fat percentage and  $VO_2$  max did not show any significant difference ( $P > 0/05$ ) after eight weeks of supplementation with placebo and quercetin between two groups and within one group. There was a significant increase in TTE after intervention in the quercetin group ( $P < 0/05$ ) but a significant change was not observed in the placebo group ( $P > 0/05$ ).

**Conclusions:** The current study shows that intake of quercetin may improve endurance exercise performance but may not reduce the body fat percentage.

**Keywords:** Body mass, exercise performance, muscle damage, quercetin

### INTRODUCTION

Polyphenolic compounds, or polyphenols, are hydroxylated phytochemicals, of which the two main classes include flavonoids and phenolic acids.<sup>[1]</sup> Quercetin is a flavonol that constitutes the aglycone of the plant glycosides rutin and quercetin.<sup>[2-4]</sup> Human subjects can absorb significant amounts of quercetin, with a reported half-life ranging from 3.5 to 28 h.<sup>[4,5]</sup> High concentrations are found in apples, red wine, onions and tea.<sup>[6]</sup>

Cardiorespiratory endurance exercise increases active skeletal muscle mitochondrial density by 20-100%.<sup>[7,8]</sup> Exercise-induced

mitochondrial biogenesis is mediated by expression of mitochondrial and nuclear genes.<sup>[9,10]</sup> A few recent studies have shown that quercetin may increase expression of mitochondrial and nuclear genes and effects on endurance performance and mitochondrial biogenesis<sup>[11,12]</sup> similar to some other nutrients.<sup>[13-15]</sup> Recent studies have examined whether quercetin supplement enhances endurance exercise capacity.

One study of 11 elite male cyclists reported 1.7% performance enhancement above placebo after six weeks of quercetin supplementation.<sup>[16]</sup> Another study using sedentary mice showed that quercetin feeding (both 12.5 and 25 mg/kg) increased treadmill run time to fatigue by 37%. Another study of trained cyclists showed no effect of 1000 mg of quercetin a day compared with placebo on cycling time trial performance.<sup>[17]</sup>

Quercetin also has non-selective anti-inflammatory actions that rely on a wide range of mechanisms of action.<sup>[18]</sup> For instance, quercetin may inhibit nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B) activity, inducible nitric oxide synthesis expression and lipoxygenase activity.<sup>[19]</sup> Many studies report that quercetin has potent anti-oxidative properties.<sup>[20,21]</sup> Aerobic exercise elevates free radical generation in muscle and free radicals increase fatigue and muscle damage.<sup>[22]</sup>

Badminton is both an aerobic and anaerobic sport with emphasis on the aerobic aspect so this sport generates free radicals. However, supplementation with an antioxidant may reduce oxidative stress, muscle damage and fatigue.<sup>[22]</sup>

*In vitro* studies suggest that quercetin affects adipogenesis and basal metabolism,<sup>[23]</sup> however, whether this metabolic effect translates into reductions in body mass or improvement in body composition in humans is unknown.

As per our knowledge, this is first time that the effect of quercetin on exercise performance, muscle damage and body muscle was investigated in male badminton players.

Given the potential of quercetin to decrease body fat percent and mediate anti-oxidative and anti-inflammatory activity the purpose of this study was to measure the influence of eight weeks of quercetin supplementation on exercise performance, muscle damage and body muscle in male badminton players.

## METHODS

A randomized double-blind placebo-controlled clinical trial was conducted on 26 male badminton players. The study was approved by the Ethics Committee, Isfahan Sport Medicine Association, Iran. The written informed consent form was obtained from all of them. Anthropometric indices including weight, height and body mass index (BMI) were obtained through physical examination. None of the participants had consumed quercetin, or any other dietary supplements, for a minimum of three months before the study. Athletes kept a record of their physical and dietary intakes and they did not have exhaustive exercise 24 h before the study. Participants filled a food record for two days before intervention.

Subjects were randomly allocated to one of the two groups according to pre-arranged balanced block randomization to receive quercetin (1000 mg) or placebo (1000 mg dextrose). Supplements and placebo had the same appearance and were supplied by Solaray<sup>®</sup>, USA, Inc. According to the findings of a previous study the supplementation procedure for quercetin was set at two capsules per day for two months—this amount provided a total daily dose of 1000 mg per day. Venous blood samples were taken from all participants between 5 and 6 PM, after intensive endurance exercising, before intervention and after intervention.

At pre- and post-supplementation protocol all participants performed a graded exercise test (GTX) on an electronically braked cycle ergometer (Lode, The Netherlands) to determine  $\text{Vo}_2$  peak and time to exhaustion (TTE). Pedal cadence was maintained at 70 rpm, while the power output was initially set at 30 watts and increased by 30 watts every 2 min, until the participants could no longer maintain the required power output.

Plasma samples were obtained for the determination of plasma lactate immediately prior to each GTX and 2 min post exercise. We used a lactometer (Lactate Pro LT-1710, Canada) to analyze lactate and a Body Composition Analyzer (JAWON IOI 353, Korea) to determine body fat percent pre and post supplementation.

Dietary analyses were performed using the Nutritionist IV software. We used SPSS Version 13 for statistical analyses. Data are presented as mean  $\pm$  standard deviation. Significant differences among and between the groups were determined

by independent *t*-test and paired sample *t*-test. An alpha of  $P < 0.05$  was considered statistically significant.

## RESULTS

General mean  $\pm$  SD for all study samples for age (years), weight (kg) and Body Mass Index (BMI) is shown in Table 1. As shown in Table 1 there was no significant difference between the two groups in age, weight and BMI ( $P > 0/05$ ). Table 2 shows the mean  $\pm$  SD values of time to exhaustion (TTE), lactate concentration,  $VO_2$  max and body fat percent pre and post supplementation. Dietary intake before trial initiation was not similar ( $P < 0.05$ ) for energy and macronutrients between the two groups [Table 1].

Lactate concentration, body fat percent and  $VO_2$  did not show any significant difference ( $P > 0/05$ ) after eight weeks of supplementation with placebo and quercetin between the two groups and within one group. There was a significant increase in TTE after intervention in the quercetin group ( $P < 0/05$ ) but significant change was not observed in the placebo group ( $P > 0/05$ ).

**Table 1:** General characteristics and dietary intakes of study participants separately by case and control groups before intervention<sup>1</sup>

Variables	Case (Quercetin) (n=14)	Control (Placebo) (n=12)	P
Age (years)	17.5 $\pm$ 2	17 $\pm$ 1.5	0.34
Weight (Kg)	54.5 $\pm$ 3.5	52.5 $\pm$ 3.2	0.21
BMI (kg/m <sup>2</sup> )			
Energy (kcal)	2,355 $\pm$ 470	2,225 $\pm$ 515	0.01
Protein (g)	120 $\pm$ 35	112 $\pm$ 23	0.01
Carbohydrates (g)	344 $\pm$ 82	331 $\pm$ 77	0.01
Fat (g)	61 $\pm$ 34	54 $\pm$ 26	0.01

<sup>1</sup>Data are means $\pm$ standard deviation, BMI=Body mass index

**Table 2:** Comparison of exercise performance indices, pre and post supplementation<sup>1</sup>

Variables	Case (Quercetin) (n=14)			Control (Placebo) (n=12)		
	Pre	Post	P values	Pre	Post	P values
$VO_{2max}$ (L.min <sup>-1</sup> )	3.15 $\pm$ 0.32	2.98 $\pm$ 0.58	0.9	3.08 $\pm$ 0.55	3.16 $\pm$ 0.51	0.6
TTE (min)	13.45 $\pm$ 1.29	14.84 $\pm$ 1.83*	0.48	13.02 $\pm$ 1.11	13.65 $\pm$ 1.14	0.21
Lactate (mg/dl)	13.5 $\pm$ 3.2	12.9 $\pm$ 3.4	0.07	12.8 $\pm$ 4.1	11.6 $\pm$ 5.5	0.06
Body fat (%)	16.7 $\pm$ 5.4	16.8 $\pm$ 5.2	2.1	18.1 $\pm$ 4.5	18.2 $\pm$ 4.4	0.35

<sup>1</sup>Data are means $\pm$ standard deviation

## DISCUSSION

The finding of our study suggests that eight-week supplementation with quercetin may improve the endurance exercise performance as measured by TTE. This result has been shown in several studies.

Dumke *et al.*, used a randomized crossover trial and showed that two weeks of supplementation with 1000 mg/day quercetin on 26 untrained men resulted in a significant increase in exercise performance.<sup>[24]</sup>

In the Davis study, treadmill run time to exhaustion increased by 37% in sedentary mice that fed with quercetin (12.5 and 25 mgkg<sup>-1</sup>) for 7 days.<sup>[25]</sup>

The one mechanism for enhanced exercise performance that is followed by quercetin is quercetin's potential to induce mitochondrial biogenesis and quercetin may increase  $VO_2$  max and endurance exercise performance.<sup>[26]</sup> Our study showed no significant difference in  $VO_2$  max in the quercetin group after intervention.

TEE and  $VO_2$  max are considered to be predictors of endurance exercise performance but subjects with already elevated mitochondrial density would be less likely to obtain benefit from quercetin, especially for  $VO_2$  max.<sup>[27]</sup>

In our study supplementation with quercetin did not cause reduction in BFP. Fat may be oxidized as a preferential fuel source if mitochondrial biogenesis is stimulated by quercetin ingestion so BFP may decrease.<sup>[24]</sup> However, studies in animals and humans have exhibited little success.<sup>[28]</sup> Studies indicate that quercetin has beneficial effects on adipocytes *in vitro* and causes apoptosis of adipocytes through the adenosine monophosphate-activated protein kinase pathway (AMPK).<sup>[29]</sup>

For example Stewart *et al.*, found that quercetin supplementation for eight weeks in mice caused no difference in body mass or body composition between the quercetin group versus placebo.<sup>[30]</sup>

Similar to our findings Egert *et al.*, showed that two-week supplementation with low doses

of quercetin did not alter BFP in healthy human subjects.<sup>[31-33]</sup>

## CONCLUSIONS

The current study shows that intake of quercetin may improve endurance exercise performance but may not reduce the body fat percent. Thus, more studies with longer periods of supplementation and larger doses that may increase quercetin bioactive effects are necessary.

## ACKNOWLEDGMENTS

This study was financially supported by grants from the Isfahan Sport Medicine Association. There are no conflicts of interest. The author would like to acknowledge the athletics who participated in this study.

## REFERENCES

1. Bieger J, Cermak R, Blank R, de Boer VC, Hollman PC, Kamphues J, *et al.* Tissue distribution of quercetin in pigs after long-term dietary supplementation. *J Nutr* 2008;138:1417-20.
2. Chun OK, Chung SJ, Song WO. Estimated dietary flavonoid intake and major food sources of U.S. adults. *J Nutr* 2007;137:1244-52.
3. Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of *in vivo* toxicity, including lack of genotoxic/carcinogenic properties. *Food Chem Toxicol* 2007;45:2179-205.
4. Manach C, Williamson G, Morand C, Scalbert A, Remesy C. Bioavailability and bioefficacy of polyphenols in humans. I. Review of 97 bioavailability studies. *Am J Clin Nutr* 2005;81:230S-42.
5. Moon YJ, Wang L, DiCenzo R, Morris ME. Quercetin pharmacokinetics in humans. *Biopharm Drug Dispos* 2008;29:205-17.
6. Williamson G, Manach C. Bioavailability and bioefficacy of polyphenols in humans. II. Review of 93 intervention studies. *Am J Clin Nutr* 2005;81:243S-55.
7. Chow LS, Greenlund LJ, Asmann YW, Short KR, McCrady SK, Levine JA, *et al.* Impact of endurance training on murine spontaneous activity, muscle mitochondrial DNA abundance, gene transcripts, and function. *J Appl Physiol* 2007;102:1078-89.
8. Hoppeler H, Fluck M. Plasticity of skeletal muscle mitochondria: Structure and function. *Med Sci Sports Exerc* 2003;35:95-104.
9. Hood DA, Irrcher I, Ljubcic V, Joseph AM. Coordination of metabolic plasticity in skeletal muscle. *J Exp Biol* 2006;209:2265-75.
10. Freyssenet D. Energy sensing and regulation of gene expression in skeletal muscle. *J Appl Physiol* 2007;102:529-40.
11. Lopez-Lluch G, Hunt N, Jones B, Zhu M, Jamieson H, Hilmer S, *et al.* Calorie restriction induces mitochondrial biogenesis and bioenergetic efficiency. *Proc Natl Acad Sci USA* 2006;103:1768-73.
12. Narkar VA, Downes M, Yu RT, Emblar E, Wang YX, Banayo E, *et al.* AMPK and PPAR delta agonists are exercise mimetics. *Cell* 2008;134:405-15.
13. Ghiasvand R, Askari Gh, Malekzadeh J, Hajishafiee M, Daneshvar P, Akbari F, *et al.* Effects of Six weeks of  $\beta$ -alanine Administration on  $VO_2$  max, time to exhaustion and lactate concentrations in physical education students. *Int J Prev Med* 2012;8:559-63.
14. Askari G, Ghiasvand R, Karimian J, Feizi A, Paknahad Z, Sharifirad G, *et al.* Does quercetin and vitamin C improve exercise performance, muscle damage, and body composition in male athletes? *J Res Med Sci* 2012;4:328-31.
15. Ghiasvand R, Djalali M, Djazayeri S, Keshavarz S, Hosseini M, Askari G, *et al.* Effect of Eicosapentaenoic Acid (EPA) and vitamin e on the blood levels of inflammatory markers, antioxidant enzymes, and lipid peroxidation in Iranian Basketball Players. *Iran J Public Health* 2010;1:15-21.
16. MacRae HS, Mefferd KM. Dietary antioxidant supplementation combined with quercetin improves cycling time trial performance. *Int J Sport Nutr Exerc Metab* 2006;16:405-19.
17. Nieman DC, Henson DA, Maxwell KR, Williams AS, McAnulty SR, Jin FX, *et al.* Effects of Quercetin and EGCG on Mitochondrial Biogenesis and Immunity. *Med Sci Sport Exerc* 2009;41:1467-75.
18. Camuesco D, Comalada M, Concha A, Nieto A, Sierra S, Xaus J, *et al.* Intestinal anti-inflammatory activity of combined quercitrin and dietary olive oil supplemented with fish oil, rich in EPA and DHA (n-3) polyunsaturated fatty acids, in rats with DSS-induced colitis. *Clin Nutr* 2006;25:466-76.
19. Liu J, Li X, Yue Y, Li J, He T, He Y. The inhibitory effect of quercetin on IL-6 production by LPS-stimulated neutrophils. *Cell Mol Immunol* 2005;2:455-60.
20. Ben Abdallah F, Zribi N, Ammar-Keskes L. Antioxidative potential of Quercetin against hydrogen peroxide induced oxidative stress in spermatozoa *in vitro*. *Andrologia* 2011;43:261-5.
21. Ishisaka A, Ichikawa S, Sakakibara H, Piskula MK, Nakamura T, Kato Y, *et al.* Accumulation of orally



- administered quercetin in brain tissue and its antioxidative effects in rats. *Free Radic Biol Med* 2011;51:1329-36.
22. Reid MB. Free radicals and muscle fatigue: Of ROS, canaries, and the IOC. *Free Radic Biol Med* 2008;44:169-79.
  23. Andersen C, Rayalam S, Della-Fera MA, Baile CA. Phytochemicals and adipogenesis. *Biofactors* 2010;36:415-22.
  24. Dumke CL, Nieman DC, Utter AC, Rigby MD, Quindry JC, Triplett NT, *et al.* Quercetin's effect on cycling efficiency and substrate utilization. *Appl Physiol Nutr Metab* 2009;34:993-1000.
  25. Davis JM, Carlstedt CJ, Chen S, Carmichael MD, Murphy EA. The dietary flavonoid quercetin increases VO (2 max) and endurance capacity. *Int J Sport Nutr Exerc Metab* 2010;20:56-62.
  26. Kressler J, Millard-Stafford M, Warren GL. Quercetin and endurance exercise capacity: A systematic review and meta-analysis. *Med Sci Sports Exerc* 2011;43:2396-404.
  27. Cureton KJ, Tomporowski PD, Singhal A, Pasley JD, Bigelman KA, Lambourne K, *et al.* Dietary quercetin supplementation is not ergogenic in untrained men. *J Appl Physiol* 2009;107:1095-104.
  28. Srujana R, Mary A, Clifton A. Phytochemicals and regulation of the adipocyte life cycle. *J Nutr Biochem* 2008;19:717-26.
  29. Ahn J, Lee H, Kim S, Park J, Ha T. The anti-obesity effect of quercetin is mediated by the AMPK and MAPK signaling pathways. *Biochem Biophys Res Commun* 2008;373:545-9.
  30. Stewart LK, Soileau JL, Ribnicky D, Wang ZQ, Raskin I, Poulev A, *et al.* Quercetin transiently increases energy expenditure but persistently decreases circulating markers of inflammation in C57BL/6J mice fed a high-fat diet. *Metabolism* 2008;57 (7 Suppl 1):S39-46.
  31. Egert S, Wolfram S, Bosy-Westphal A, Boesch-Saadatmandi C, Wagner AE, Frank J, *et al.* Daily quercetin supplementation dose-dependently increases plasma quercetin concentrations in healthy humans. *J Nutr* 2008;138:1615-21.
  32. Mirkheshti A, Aryani MR, Shojaei P, Dabbagh A. The effect of adding magnesium sulfate to lidocaine compared with paracetamol in prevention of acute pain in hand surgery patients under Intravenous Regional Anesthesia (IVRA). *Int J Prev Med* 2012;3:616-21.
  33. Djalalinia S, Tehrani FR, Afzali HM, Hejazi F, Peykari N. Parents or school health trainers, which of them is appropriate for menstrual health education? *Int J Prev Med* 2012;3:622-7.

**Source of Support:** Nil, **Conflict of Interest:** None declared.

### Staying in touch with the journal

#### 1) Table of Contents (TOC) email alert

Receive an email alert containing the TOC when a new complete issue of the journal is made available online. To register for TOC alerts go to [www.ijpm.ir/signup.asp](http://www.ijpm.ir/signup.asp).

#### 2) RSS feeds

Really Simple Syndication (RSS) helps you to get alerts on new publication right on your desktop without going to the journal's website. You need a software (e.g. RSSReader, Feed Demon, FeedReader, My Yahoo!, NewsGator and NewzCrawler) to get advantage of this tool. RSS feeds can also be read through FireFox or Microsoft Outlook 2007. Once any of these small (and mostly free) software is installed, add [www.ijpm.ir/rssfeed.asp](http://www.ijpm.ir/rssfeed.asp) as one of the feeds.